

Multiple drug resistance due to resistance to stem cells and stem cell treatment progress in cancer (Review)

CHONG DI and YAODONG ZHAO

Department of Neurosurgery, Shanghai 10th People's Hospital, Tongji University School of Medicine, Shanghai 200072, P.R. China

Received March 28, 2014; Accepted December 5, 2014

DOI: 10.3892/etm.2014.2141

Abstract. In recent years, the cancer stem cell (CSC) theory has provided a new angle in the research of cancer, and has gradually gained significance. According to this theory, the multiple drug resistance (MDR) of cancer is most likely due to the resistance of CSCs, and a significant quantity of research has been carried out into the MDR mechanisms of CSC. Over time, some of these mechanisms have been gradually accepted, including ATP-binding cassette transporters, aldehyde dehydrogenase, the CSC microenvironment and epithelial to mesenchymal transition. In the present review, we summarize these mechanisms in detail and review possible appropriate therapy plans against CSCs based on CSC theory.

Contents

1. Introduction
2. Cancer stem cells
3. Mechanisms of CSC MDR
4. Conclusion

1. Introduction

Cancer is a notable threat to human health with a rising incidence rate; however, there is still no known cure. In recent years, the cancer stem cell (CSC) theory was proposed, which provided a new angle in the research of cancer, and gradually gained significance. According to this theory, the multiple drug resistance (MDR) of cancer is most likely due to the

resistance of CSCs, which have received increasing attention from oncologists around the world (1). Therefore, we intend to review the common mechanisms of MDR in CSC, so as to discuss possible relevant therapeutic strategies against cancer.

2. Cancer stem cells

The concept of CSCs was first raised by Park *et al* (2) in 1971. The concept was that cancers are diseases driven by a subpopulation of self-renewing CSCs, which possess the ability to generate the diverse differentiated cell populations that comprise the cancer mass (3). Since then, CSCs have been identified and isolated from tumors of the hematopoietic system, breast, lung, prostate, colon, brain, head and neck, and pancreas.

Existing therapies have been developed largely against the bulk population of tumor cells to shrink tumors. However, most cancer cells only have limited proliferative potential; an ability to shrink a tumor mainly reflects an ability to kill these cells. Even therapies that cause complete regression of tumors might spare sufficient CSCs to allow regrowth of the tumors (4). Only therapies that are more specifically directed against CSCs might result in more durable responses and even a cure for metastatic tumors.

3. Mechanisms of CSC MDR

ATP-binding cassette (ABC) transporters. The ABC transporter superfamily in humans includes at least 49 genes grouped into seven families (from A to G) with various functions, and at least 16 of these proteins are implicated in cancer drug resistance (5). They bind ATP and use the energy to drive the transport of various molecules across the plasma membrane; since they can eject anti-cancer drugs from cells, it may lead to the drug's resistance (6). Among the ABC proteins, the most significant are glycoprotein P (P-gp), encoded by the ABCB1 [multidrug resistance protein 1 (MDR1)] gene, and breast cancer resistance protein (BCRP or ABCG2), which was cloned from a mitoxantrone-resistant subline of the breast cancer cell line MCF-7. BCRP confers resistance to a number of chemotherapeutics, including mitoxantrone. Other well-known proteins from the ABC family responsible for MDR are MDR-related protein (MRP) 1 (ABCC1) and MRP2 (ABCC2). ABC transporters are also expressed in

Correspondence to: Dr Yaodong Zhao, Department of Neurosurgery, Shanghai 10th People's Hospital, Tongji University School of Medicine, 301 Yanchangzhong Road, Shanghai 200072, P.R. China

E-mail: zhaoyd@aliyun.com

Key words: cancer stem cell, multiple drug resistance, adenosine triphosphate-binding cassette transporter, aldehydes dehydrogenase, microenvironment, epithelial to mesenchymal transition

normal stem cells to maintain a relatively stable intracellular environment and to keep them in a quiescent state. In addition, these transporters have certain other notable roles in normal physiology in the transport of drugs across the placenta and the intestine, and are essential components of the blood-brain and blood-testis barriers. By using the energy from ATP hydrolysis, these transporters actively efflux drugs from cells, serving to protect them from cytotoxic agents.

Evidence for MDR of CSCs mediated by ABC transporters. Numerous studies have demonstrated the role of ABC transporters in CSC resistance. CD133-positive glioma stem cells exhibited a notable effect on tumor resistance to chemotherapy, which is possibly as a result of their high expression of ABCG2/BCRP1 (7). Similar results also were observed in the CSCs of lung cancer (8), osteoma sarcomatosum (9), ovarian cancer (10), prostatic carcinoma (11) and nasopharyngeal carcinoma (12). Furthermore, the high ABC transporter levels were associated with increased Akt signaling in drug-resistant CSCs, and the Akt signaling was able to alter the subcellular localization of BCRP transporters, thus determining drug efflux in CSCs (13). In the same study, Akt signal inhibition by P13K inhibitors not only suppressed cancer cell proliferation, but also increased the sensitivity of drug-resistant cells (13).

Therapeutic measures targeting MDR mediated by ABC transporters. Theoretically, ABC transporter inhibitors could reverse the chemotherapy resistance of CSCs and eliminate tumors. Currently, its inhibitors have been developed to the third generation. First-generation compounds included drugs identified as ABCB1 inhibitors, including verapamil and cyclosporine, which were in clinical use to treat other diseases. These inhibitors were combined with a range of chemotherapy regimens for numerous types of cancer, but the results were not satisfactory. The clinical results for the second-generation inhibitors, including valspodar and biricodar, were also disappointing. Fumitremorgin C and its chemically synthesized derivatives including Ko143 have been developed; certain studies have demonstrated a positive effect *in vitro*, particularly for Ko143, which has high specificity and low toxicity (14). Studies with Ko143 have also revealed that inhibition of ABCG2 allows for a greater absorption of certain drugs across the intestine (14). In addition, the compound GF120918 is an ABCB1 inhibitor that has been demonstrated to inhibit ABCG2 *in vitro* as well as *in vivo* (15). Compared with those of previous generations, other inhibitors, including zosuquidar (LY335979) and tariquidar (XR9576), have higher inhibitory activity and selectivity, without affecting the metabolism of chemotherapeutic drugs themselves, and it is possible to overcome the resistance to CSCs. Moreover, they have already been approved for clinical studies. Certain drugs are in clinical use; for example, difluorinated curcumin enhances the sensitivity of CD44⁺CD166⁺ colon carcinoma stem cells to the combination of 5-fluorouracil and oxaliplatin by a mechanism that involves ABCG2 downregulation (15). The crude extract of traditional Chinese medicine *Schisandra* and its active ingredient deoxyschizandrin have the effect of reversing MDR (16), and enhance the curative effect of vincristine and adriamycin. *Schisandrae Lignans* inhibits the outflow of intracellular drugs (17,18). Chai *et al* (19) considered that

schizandrin had strong potential in terms of reversing MDR, and that it could induce apoptosis and decrease the expression of P-gp and protein kinase C. Other medicines, including salinomycin (20,21), nigericin and abamectin have a similar effect. Nicardipine is also an inhibitor of ABCG2 (7).

Aldehyde dehydrogenase (ALDH). The ALDHs are a family of nicotinamide adenine dinucleotide phosphate [NAD (P)⁺]-dependent enzymes involved in detoxifying a wide variety of aldehydes to weak carboxylic acids, leading to increased protection of the cell against insult by environmental chemicals and drugs. ALDHs are also regarded as a type of biomarker for stem cells. They include the ALDH1 family (ALDH1A1, 1A2, 1A3, 1L1, 1L2), ALDH2 and ALDH3A1. In particular, ALDH1A1 and ALDH3A1 have been shown to play significant functional roles in stem cells (22), which was demonstrated in hematopoietic stem cells, neural stem cells and adipose tissue stem cells.

Evidence for MDR of CSCs mediated by ALDHs. ALDH1 is a CSC marker and its presence is strongly correlated with tumor malignancy as well as self-renewal properties of stem cells in various tumors, including breast cancer, hepatoma, colon cancer and lung cancer (22). CSCs with a high expression of ALDH have the characteristics of being chemotherapy-resistant, which has been proven in breast cancer stem cells (23) and head/neck squamous cell carcinoma stem cells (24). The mechanism of ALDH expression of the gene may be associated with high Snail expression, since the knockdown of Snail expression significantly decreases the expression of ALDH1, inhibits cancer stem-like properties and blocks the tumorigenic abilities of CD44⁺CD24⁻ALDH1⁺ cells (24).

Therapeutic measures targeting MDR mediated by ALDHs. A number of natural compounds are effective against CSCs and lead to decreasing numbers of ALDH-positive cells (25). For example, exposure of breast cancer cells to the dietary polyphenols, curcumin and piperine, severely reduces mammosphere formation (26), as do sulforaphane (27) and butein (28).

In addition, the role of chemical synthesis of the drug in inhibiting ALDH has also been studied. The specific ALDH inhibitor diethylaminobenzaldehyde may increase the sensitization of ALDH^{br}CD44⁺ cells to chemotherapy/radiotherapy (23). Disulfiram is capable of inhibiting ALDH and enhancing the sensitization of cancer cells to chemotherapy (29). All-trans-retinoic acid treatment results in a 50-60% reduction in ALDH activity and proteins of both isozymes, ALDH1A1 and ALDH3A1, leading to the increased sensitization of ALDH1⁺ cancer cells to chemotherapy.

Microenvironment

Mechanisms of the microenvironment influencing MDR of CSCs. Normal adult stem cells are regulated by molecular cues provided by neighboring connective tissue cells, mainly mesenchymal (fibroblast-like) cells and vascular cells; these stromal cells contribute to the stem cell niche. CSCs often appear to be present in and influenced by a similar microenvironment, which has been proven in brain cancer (30), squamous cell carcinoma (31), bladder cancer (32), leukemia (33,34) and colorectal cancer (35). The tumor microenviron-

ment has been recognized as a major factor influencing the growth of cancer and impacting the outcome of therapy. Thus, niche cells have become an attractive target for chemotherapeutic agents (36). Targeted therapies through the production of secreted factors which might drive tumor growth and MDR may be available (37). Hypoxia has been considered as a major feature of the tumor microenvironment and a potential contributor to the MDR and enhanced tumorigenicity of CSCs (38). The acidic microenvironment around hypoxia cells is combined with the activation of a subset of proteases that contribute to metastasis (39). Due to aberrant angiogenesis and the inaccessible location, hypoxic cells are less likely to accumulate therapeutic concentrations of chemotherapeutics, which lead to MDR. Therefore, targeting the CSC niche in combination with chemotherapy may provide a promising strategy for eradicating CSCs.

Therapeutic measures targeting MDR mediated by the microenvironment. Treatment of animals with DC101, an antibody of the vascular endothelial growth factor (VEGF) receptor, leads to a transient increase in oxygenation and deeper penetration of molecules into experimental tumors. An increased response of tumors to combined treatment with chemotherapy and anti-angiogenic agents has been demonstrated (40). Bevacizumab, another type of antibody to VEGF receptor, was reported to have a similar effect (41). Agents that damage existing blood vessels in tumors may also influence the response to chemotherapy. Vascular disrupting agents (including tumor necrosis factor, flavone acetic acid and its derivatives, and tubulin-binding agents including combretastatin A-4 disodium phosphate) directly damage the established tumor endothelium and have been shown to increase vessel permeability and drug delivery (42). Certain other medicines, for example histamine, a selective endothelin receptor A antagonist and botulinum neurotoxin type A, increase tumor blood flow and have been shown to promote *in vivo* tumor perfusion and to delay tumor growth when combined with cyclophosphamide (10). However, there also is some dispute. Injecting vascular-disrupting agents prior to chemotherapy may be problematic since it could result in reduced blood flow and increased interstitial fluid pressure, which together could impair the delivery of drugs to tumors and lessen the curative effect (42).

Epithelial to mesenchymal transitions (EMT). EMT is a process involving the dissolution of endothelial cell-cell junctions and loss of apico-basolateral polarity of endothelial cells, resulting in the formation of migratory mesenchymal cells with invasive properties. The loss of polarity and gain of motile characteristics of mesenchymal cells during embryonic development has prompted comparisons with metastatic cancer cells during malignant progression (43). Notably, these properties have also been ascribed to normal stem cells and CSCs. Cancer is a disease of abnormal wound healing and tissue repair, which is accompanied by pathophysiological EMT in adult tissues. Accordingly, EMT is able to trigger reversion to a CSC-like phenotype, providing an association between EMT, CSCs and MDR (44). In addition, a previous study has demonstrated that CSCs and EMT-type cells could play critical roles in drug resistance (45).

AMP-activated kinase (AMPK) inhibits EMT; however, the diabetes drug metformin may activate AMPK, which results in the elective killing of CSCs in combination with chemotherapy (46). However, it was unclear whether the positive results were a result of the combination with metformin. In addition, histone deacetylase (HDAC) is another significant molecule in EMT, and HDAC promotes EMT by inhibiting E-cadherin (47). Certain HDAC inhibitors, including trichostatin A and vorinostat, induce the differentiation of mesenchymal-like cancer cells and CSCs, which could trigger apoptotic responses or chemosensitize these cells to other therapies. Furthermore, HDACs affect the activity of non-histone substrates, including HIF-1 and NF- κ B, which have been implicated in driving EMT and drug resistance.

Molecular signaling pathways

Hedgehog (Hh) signaling pathway. Previous studies have suggested that the Hh pathway is essential for the maintenance of CSCs in various human cancer types including pancreatic cancer (48), gastric cancer (49) and colorectal cancer (50). It is also responsible for treatment resistance of cancer cells. Thus, inhibitors such as cyclopamine and GDC-0449 (51), that obstruct the Hh signaling pathway, may cause depletion of CSCs, overcome MDR, and enhance the curative effect. Tang *et al* (48) observed that epigallocatechin-3-gallate (EGCG) inhibited the components of the sonic hedgehog (SHh) pathway (SMO, Ptch, Gli1 and Gli2) and Gli transcriptional activity, and the combination of quercetin with EGCG had synergistic inhibitory effects on the self-renewal capacity of CSCs through attenuation of TCF/LEF and Gli activities. The authors suggested that therapeutics targeting the SHh pathway might improve the therapeutic outcome of patients with pancreatic cancer by targeting CSCs.

Notch signaling pathway. The Notch signaling pathway plays crucial roles in embryonic development and the proliferation, differentiation and apoptosis of stem cells. Specifically, Notch functions as an oncogenic protein in most human cancers including cervical, lung, colon, head and neck, prostate and pancreatic cancer, while it acts as a tumor suppressor in skin cancer, hepatocellular carcinoma and small cell lung cancer (52).

Fan *et al* (53) used G-secretase inhibitors (GSIs) to block the Notch pathway in glioblastoma, resulting in reduced neurosphere growth and clonogenicity *in vitro*, reduced expression of putative CSC markers and reduced tumor growth *in vivo*. Thus, they suggested that GSIs which block the Notch pathway might be useful chemotherapeutic reagents to target CSCs. MRK-03 has a similar effect on breast cancer stem cells (54).

Wnt and other signaling pathways. The Wnt signaling pathway is another developmental pathway involved in multiple biological processes including embryogenesis, development, cell proliferation, survival and differentiation. However, oncogenic mutations or deactivation of the tumor suppressor may result in the dysregulation of the Wnt/ β -catenin pathway in CSCs, which induces neoplastic proliferation. Small molecule inhibitors include existing drugs such as nonsteroidal anti-inflammatory drugs (NSAIDs) or natural

compounds, and molecular-targeted agents such as the cAMP response-element binding protein (CBP)/ β -catenin antagonist ICG-001. These inhibitors interfere with the Wnt pathway and harvest a therapeutic effect. Takahashi-Yanaga *et al* (55) demonstrated that CBP/ β -catenin antagonist ICG-001 was able to target and eliminate drug-resistant leukemic stem cells *in vivo* and *in vitro*.

Other mechanisms. Other mechanisms are also involved in the MDR of CSCs; for example, microRNAs, which are single-stranded 19-25 nucleotide short RNAs, play a significant role in the regulation of MDR (56). CSCs differentiation-inducing agents also improve the efficacy of treatment in combination with chemotherapy (57).

4. Conclusion

Due to poor chemotherapy efficacy in most cancer types, particularly in the case of metastases, we should change our view of the development of drug resistance in cancer and try to develop chemotherapy based on the CSC model. However, there also are certain unresolved issues concerning therapy aimed directly at CSCs. For example, ABC transporters and ALDHs are markers of normal stem cells, and the inhibition of these molecules in CSCs may result in them occurring in normal stem cells, consequently leading to severe side effects. In addition, it is still unclear which ALDH isozymes contribute to the ALDH activity and whether this is the same among all tissue-specific stem cells. It is even claimed that before CSCs can be clearly identified, therapeutic approaches designated to target them may actually cause more harm than good in glioblastoma multiforme patients (58). All these issues remain to be solved in the future.

Acknowledgements

The current study was supported by the National Natural Science Foundation of China (grant no. 81101909) and the Doctoral Program of Higher Education (Specialized Research Fund) of China (grant no. 20110072120055).

References

- Reya T, Morrison SJ, Clarke MF and Weissman IL: Stem cells, cancer, and cancer stem cells. *Nature* 414: 105-111, 2001.
- Park CH, Bergsagel DE and McCulloch EA: Mouse myeloma tumor stem cells: a primary cell culture assay. *J Natl Cancer Inst* 46: 411-422, 1971.
- Gupta PB, Chaffer CL and Weinberg RA: Cancer stem cells: mirage or reality? *Nat Med* 15: 1010-1012, 2009.
- Merlos-Suárez A, Barriga FM, Jung P, *et al*: The intestinal stem cell signature identifies colorectal cancer stem cells and predicts disease relapse. *Cell Stem Cell* 8: 511-524, 2011.
- Fletcher JI, Haber M, Henderson MJ and Norris MD: ABC transporters in cancer: more than just drug efflux pumps. *Nat Rev Cancer* 10: 147-156, 2010.
- Gottesman MM: Mechanisms of cancer drug resistance. *Annu Rev Med* 53: 615-627, 2002.
- Jin Y, Bin ZQ, Qiang H, *et al*: ABCG2 is related with the grade of glioma and resistance to mitoxantone, a chemotherapeutic drug for glioma. *J Cancer Res Clin Oncol* 135: 1369-1376, 2009.
- Wang B, Yang H, Huang YZ, Yan RH, Liu FJ and Zhang JN: Biologic characteristics of the side population of human small cell lung cancer cell line H446. *Chin J Cancer* 29: 254-260, 2010.
- Adhikari AS, Agarwal N, Wood BM, *et al*: CD117 and Stro-1 identify osteosarcoma tumor-initiating cells associated with metastasis and drug resistance. *Cancer Res* 70: 4602-4612, 2010.
- Hu L, McArthur C and Jaffe RB: Ovarian cancer stem-like side-population cells are tumorigenic and chemoresistant. *Br J Cancer* 102: 1276-1283, 2010.
- Tokar EJ, Qu W, Liu J, *et al*: Arsenic-specific stem cell selection during malignant transformation. *J Natl Cancer Inst* 102: 638-649, 2010.
- Wang J, Guo LP, Chen LZ, Zeng YX and Lu SH: Identification of cancer stem cell-like side population cells in human nasopharyngeal carcinoma cell line. *Cancer Res* 67: 3716-3724, 2007.
- Hu C, Li H, Li J, *et al*: Analysis of ABCG2 expression and side population identifies intrinsic drug efflux in the HCC cell line MHCC-97L and its modulation by Akt signaling. *Carcinogenesis* 29: 2289-2297, 2008.
- Allen JD, van Loevezijn A, Lakhai JM, *et al*: Potent and specific inhibition of the breast cancer resistance protein multidrug transporter *in vitro* and in mouse intestine by a novel analogue of fumitremorgin C. *Mol Cancer Ther* 1: 417-425, 2002.
- Cisternino S, Mercier C, Bourasset F, Roux F and Schermann JM: Expression, up-regulation, and transport activity of the multidrug-resistance protein Abcg2 at the mouse blood-brain barrier. *Cancer Res* 64: 3296-3301, 2004.
- Qin XQ, Liang YG, Gao HZ, *et al*: Reversing mechanism of schizandrin A on multi-drug resistance of K562/ADR, HL60/ADR, MCF-7/ADR cell lines. *Chin Pharmacol Bull* 27: 329-334, 2011.
- Huang M, Jin J, Sun H, and Liu GT: Reversal of P-glycoprotein-mediated multidrug resistance of cancer cells by five schizandrins isolated from the Chinese herb *Fructus Schizandrae*. *Cancer Chemother Pharmacol* 62: 1015-1026, 2008.
- Slaninová I, Brezinová L, Koubíková L and Slanina J: Dibenzocyclooctadiene lignans overcome drug resistance in lung cancer cells - study of structure-activity relationship. *Toxicol In Vitro* 23:1047-1054, 2009
- Chai S, To KK and Lin G: Circumvention of multi-drug resistance of cancer cells by Chinese herbal medicines. *Chin Med* 5: 26, 2010.
- Gupta PB, Onder TT, Jiang G, *et al*: Identification of selective inhibitors of cancer stem cells by high-throughput screening. *Cell* 138: 645-659, 2009.
- Riccioni R, Dupuis ML, Bernabei M, *et al*: The cancer stem cell selective inhibitor salinomycin is a p-glycoprotein inhibitor. *Blood Cells Mol Dis* 45: 86-92, 2010.
- Ma I and Allan AL: The role of human aldehyde dehydrogenase in normal and cancer stem cells. *Stem Cell Rev* 7: 292-306, 2011.
- Croker AK and Allan AL: Inhibition of aldehyde dehydrogenase (ALDH) activity reduces chemotherapy and radiation resistance of stem-like ALDHhiCD44⁺ human breast cancer cells. *Breast Cancer Res Treat* 133: 75-87, 2012.
- Chen YC, Chen YW, Hsu HS, *et al*: Aldehyde dehydrogenase 1 is a putative marker for cancer stem cells in head and neck squamous cancer. *Biochem Biophys Res Commun* 385: 307-313, 2009.
- Januchowski R, Wojtowicz K and Zabel M: The role of aldehyde dehydrogenase (ALDH) in cancer drug resistance. *Biomed Pharmacother* 67: 669-680, 2013.
- Kakarala M, Brenner DE, Korkaya H, *et al*: Targeting breast stem cells with the cancer preventive compounds curcumin and piperine. *Breast Cancer Res Treat* 122: 777-785, 2010.
- Li Y, Zhang T, Korkaya H, *et al*: Sulforaphane, a dietary component of broccoli/broccoli sprouts, inhibits breast cancer stem cells. *Clin Cancer Res* 16: 2580-2590, 2010.
- Cioce M, Gherardi S, Viglietto G, *et al*: Mammosphere-forming cells from breast cancer cell lines as a tool for the identification of CSC-like- and early progenitor-targeting drugs. *Cell Cycle* 9: 2878-2887, 2010.
- Wickström M, Danielsson K, Rickardson L, *et al*: Pharmacological profiling of disulfiram using human tumor cell lines and human tumor cells from patients. *Biochem Pharmacol* 73: 25-33, 2007.
- Calabrese C, Poppleton H, Kocak M, *et al*: A perivascular niche for brain tumor stem cells. *Cancer Cell* 11: 69-82, 2007.
- Atsumi N, Ishii G, Kojima M, Sanada M, Fujii S and Ochiai A: Podoplanin, a novel marker of tumor-initiating cells in human squamous cell carcinoma A431. *Biochem Biophys Res Commun* 373: 36-41, 2008.
- He X, Marchionni L, Hansel DE, *et al*: Differentiation of a highly tumorigenic basal cell compartment in urothelial carcinoma. *Stem Cells* 27: 1487-1495, 2009.
- Colmone A, Amorim M, Pontier AL, Wang S, Jablonski E and Sipkins DA: Leukemic cells create bone marrow niches that disrupt the behavior of normal hematopoietic progenitor cells. *Science* 322: 1861-1865, 2008.

34. Lane SW, Scadden DT and Gilliland DG: The leukemic stem cell niche: current concepts and therapeutic opportunities. *Blood* 114: 1150-1157, 2009.
35. Vermeulen L, De Sousa E Melo F, van der Heijden M, *et al*: Wnt activity defines colon cancer stem cells and is regulated by the microenvironment. *Nature Cell Biol* 12: 468-476, 2010.
36. Basak GW, Srivastava AS, Malhotra R and Carrier E: Multiple myeloma bone marrow niche. *Curr Pharm Biotechnol* 10: 345-346, 2009.
37. Ostman A: The tumor microenvironment controls drug sensitivity. *Nat Med* 18: 1332-1334, 2012.
38. Vinogradov S and Wei X: Cancer stem cells and drug resistance: the potential of nanomedicine. *Nanomedicine (Lond)* 7: 597-615, 2012.
39. Harris AL: Hypoxia - a key regulatory factor in tumour growth. *Nat Rev Cancer* 2: 38-47, 2002.
40. Segers J, Di Fazio V, Ansiaux R, *et al*: Potentiation of cyclophosphamide chemotherapy using the anti-angiogenic drug thalidomide: importance of optimal scheduling to exploit the 'normalization' window of the tumor vasculature. *Cancer Lett* 244: 129-135, 2006.
41. Sandler A, Gray R, Perry MC, *et al*: Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med* 355: 2542-2550, 2006.
42. Horsman MR and Siemann DW: Pathophysiologic effects of vasculartargeting agents and the implications for combination with conventional therapies. *Cancer Res* 66: 11520-11539, 2006.
43. Yang J and Weinberg RA: Epithelial-mesenchymal transition: at the crossroads of development and tumor metastasis. *Dev Cell* 14: 818-829, 2008.
44. Polyak K and Weinberg RA: Transitions between epithelial and mesenchymal states: acquisition of malignant and stem cell traits. *Nat Rev Cancer* 9: 265-273, 2009.
45. Voulgari A and Pintzas A: Epithelial-mesenchymal transition in cancer metastasis: mechanisms, markers and strategies to overcome drug resistance in the clinic. *Biochim Biophys Acta* 1796: 75-90, 2009.
46. Rattan R, Ali Fehmi R and Munkarah A: Metformin: an emerging new therapeutic option for targeting cancer stem cells and metastasis. *J Oncol* 2012: 928127, 2012.
47. Singh A and Settleman J: EMT, cancer stem cells and drug resistance: an emerging axis of evil in the war on cancer. *Oncogene* 29: 4741-4751, 2010.
48. Tang SN, Fu J, Nall D, Rodova M, Shankar S and Srivastava RK: Inhibition of sonic hedgehog pathway and pluripotency maintaining factors regulate human pancreatic cancer stem cell characteristics. *Int J Cancer* 131: 30-40, 2012.
49. Song Z, Yue W, Wei B, *et al*: Sonic hedgehog pathway is essential for maintenance of cancer stem-like cells in human gastric cancer. *PLoS One* 6: e17687, 2011.
50. Merchant AA and Matsui W: Targeting Hedgehog - a cancer stem cell pathway. *Clin Cancer Res* 16: 3130-3140, 2010.
51. Singh BN, Fu J, Srivastava RK and Shankar S: Hedgehog signaling antagonist GDC-0449 (Vismodegib) inhibits pancreatic cancer stem cell characteristics: molecular mechanisms. *PLoS One* 6: e27306, 2011.
52. Xia J, Chen C, Chen Z, Miele L, Sarkar FH and Wang Z: Targeting pancreatic cancer stem cells for cancer therapy. *Biochim Biophys Acta* 1826: 385-399, 2012.
53. Fan X, Khaki L, Zhu TS, *et al*: NOTCH pathway blockade depletes CD133-positive glioblastoma cells and inhibits growth of tumor neurospheres and xenografts. *Stem Cells* 28: 5-16, 2010.
54. Kondratyev M, Kreso A, Hallett RM, *et al*: Gamma-secretase inhibitors target tumor-initiating cells in a mouse model of ERBB2 breast cancer. *Oncogene* 31: 93-103, 2011.
55. Takahashi-Yanaga F and Kahn M: Targeting Wnt signaling: can we safely eradicate cancer stem cells? *Clin Cancer Res* 16: 3153-3162, 2010.
56. Sarkar FH, Li Y, Wang Z, Kong D and Ali S: Implication of microRNAs in drug resistance for designing novel cancer therapy. *Drug Resist Updat* 13: 57-66, 2010.
57. Arrieta O, González-De la Rosa CH, Aréchaga-Ocampo E, *et al*: Randomized phase II trial of all-trans-retinoic acid with chemotherapy based on paclitaxel and cisplatin as first-line treatment in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 28: 3463-3471, 2010.
58. Sze CI, Su WP, Chiang MF, Lu CY, Chen YA and Chang NS: Assessing current therapeutic approaches to decode potential resistance mechanisms in glioblastomas. *Front Oncol* 3: 59, 2013.