

framework (27), *Sarcosuchus* and its closest relatives are understood as basal neosuchians, allied with but outside Crocodylia and the radiation that gave rise to all living crocodylians.

References and Notes

1. F. Broin, P. Taquet, *C. R. Acad. Sci. Paris* **262**, 2326 (1966).
2. P. Taquet, *Cah. Paléontol.* **1**, 1 (1975).
3. *Sarcosuchus imperator*, revised diagnosis (conditions unknown in *S. hartti*): crocodyliform with premaxillary narial bulla, maxillae that expand about 25% in width anteriorly, anterior maxillary teeth reduced in size. *Sarcosuchus*, revised diagnosis (*S. imperator*, *S. hartti*): crocodyliforms with dentary teeth 3 and 4 enlarged (28); diastema between dentary tooth 4 and 5; dentaries with fan-shaped distal expansion (28) reaching a width 50% greater than the narrowest portion of the rami; dentary symphysis extending posteriorly to about dentary tooth 20.
4. A. C. Pooley, in *Crocodyles and Alligators*, C. A. Ross, Ed. (Merehurst, London, 1989), pp. 76–91.
5. X.-C. Wu, D. B. Brinkman, A. P. Russell, *Can. J. Earth Sci.* **33**, 606 (1996).
6. R. W. Hooley, *Q. J. Geol. Soc. London* **63**, 50 (1907).
7. M. K. Hecht, in *Santana Fossils*, J. G. Maisey, Ed. (T.F.H. Publications, Neptune City, NJ, 1991), pp. 342–347.
8. F. D. Ross, G. C. Mayer, in *Advances in Herpetology and Evolutionary Biology*, A. G. Rhodin, K. Miyata, Eds. (Harvard Univ. Press, Cambridge, MA, 1983), pp. 305–331.
9. E. Buffetaut, *Mém. Soc. Geol. France Nov. Ser.* **60**, 1 (1981).
10. Skull length is measured from the anterior end of the premaxilla to the posterior end of the occiput in the midline.
11. H. Wermuth, *Senck. Biol.* **45**, 369 (1964).
12. A. E. Greer, *J. Herpetol.* **8**, 381 (1974).
13. G. J. W. Webb, H. Messel, *Aust. J. Zool.* **26**, 1 (1978).
14. L. A. K. Singh, H. R. Bustard, *Br. J. Herpatol.* **6**, 253 (1982).
15. G. J. W. Webb, R. Buckworth, S. C. Manolis, *Aust. Wildl. Res.* **10**, 383 (1983).
16. A. R. Woodward, J. H. White, S. B. Linda, *J. Herpetol.* **29**, 507 (1995).
17. The holotypic skull of *S. imperator* in the collections of the Musée National du Niger (MNN) has a length of 160 cm (2) and is the largest verified skull length on record (the same reference reported a 170-cm skull, but there are no figures of this specimen).
18. Regressions of body length (y) against head length (x) for *G. gangeticus* ($y = -69.369 + 7.4x$, $r^2 = 0.972$) and *C. porosus* ($y = -20.224 + 7.717x$, $r^2 = 0.98$) yield total body length estimates of 11.15 m and 12.15 m, respectively, for a skull length of 160 cm. Mean total body length equals 11.65 m (38 feet, 3 inches) (Fig. 4B). Data for *G. gangeticus* ($n = 17$) include measurements taken by P. Sereno from captive bred gharials in the Kukrail Picnic Center and Katerniaghat National Reserve in northern India and available measurements for individuals greater than 1.5 m long (11). Data for *C. porosus* ($n = 28$) come from wild individuals in northern Australia (A. Britton) and from available measurements for individuals greater than 1.5 m long (11) (excluding one suspicious record with body length of 4.91 m) (for measurement data, see supplementary material) (29).
19. J. J. Head, *J. Vert. Paleontol.* **21**, 59A (2001).
20. G. M. Erickson, C. A. Brochu, *Nature* **398**, 205 (1999).
21. D. R. Schwimmer, *J. Vert. Paleontol.* **19**, 74A (1999).
22. Using linear equations for *C. porosus* (13), we estimated snout-vent length in *S. imperator* from skull length (160 cm, yielding 572 cm) and from total body length (11.65 m, yielding 570 cm). We used mean snout-vent length (571 cm), in turn, to estimate a body weight (bw) of 7.96 metric tons [$\log bw = -2.0894 + 3.2613 (\log 571 \text{ cm})$].
23. S. Grenard, *Handbook of Alligators and Crocodiles* (Kreiger, Malabar, FL, 1991).
24. X.-C. Wu, A. P. Russell, S. L. Cumbaa, *J. Vert. Paleontol.* **21**, 492 (2001).
25. *Sarcosuchus* + *Terminonaris*: premaxillary palate with

26. C. W. Andrews, *Annu. Mag. Nat. Hist.* **11**, 485 (1913).
27. Phylogenetic definitions are provided below for key higher level taxa for stability of taxonomic content (30): Crocodyliformes, *Protosuchus richardsoni*, *Crocodylus niloticus*, and all descendants of their common ancestor; Protosuchia, all crocodyliforms more closely related to *Protosuchus richardsoni* than to *Crocodylus niloticus*; Mesoeucrocodylia, all crocodyliforms more closely related to *Crocodylus niloticus* than to *Protosuchus richardsoni*; Metasuchia, *Notosuchus terrestris*, *Crocodylus niloticus*, and all descendants of their common ancestor; Notosuchia, all crocodyliforms more closely related to *Notosuchus terrestris* than to *Crocodylus niloticus*; Neosuchia, all crocodyliforms more closely related to *Crocodylus niloticus* than to *Notosuchus terrestris*. Crocodylia is defined as *Gavialis gangeticus*, *Crocodylus niloticus*, and all descendants of their common ancestor [modified from (31)].
28. E. Buffetaut, P. Taquet, *Palaeontology* **20**, 203 (1977).
29. Supplementary data are available on Science Online at www.sciencemag.org/cgi/content/full/1066521/DC1.

30. P. C. Sereno, *Syst. Biol.* **48**, 329 (1999).
31. H. C. E. Larsson, dissertation, University of Chicago (2000).
32. A. G. Smith, D. G. Smith, B. M. Funnell, *Atlas of Mesozoic and Cenozoic Coastlines* (Cambridge Univ. Press, Cambridge, 1994).
33. D. L. Swofford, PAUP 3.1 (Illinois Natural History Survey, Champaign, IL, 1993).
34. Supported by the David and Lucile Packard Foundation, the National Geographic Society, and the Pritzker Foundation. We thank C. Abraczinskas for the finished illustrations; A. Beck, A. Britton, C. Brochu, J. Conrad, J. Head, W. Langston Jr., and J. Wilson for comments on the manuscript; B. Barr, B. C. Choudhury, and G. Martin for help in measuring *G. gangeticus*; A. Britton for measurements of *C. porosus*; E. Dong, A. Gray, L. Mahler, T. Keillor, R. Masek, and C. Noto for fossil preparation; A. Beck, D. Blackburn, J. Bradshaw, J.-P. Cavigelli, J. Conrad, E. Duneman, D. Duteil, M. Hettwer, G. Lyon, T. Lyman, J. Marcot, R. Sadleir, G. Wilson, J. Wilson, and D. Varrichio for their contributions during expeditions in 1997 and 2000; and I. Kouada and the government of the Niger Republic for their support of this research.

24 September 2001; accepted 16 October 2001
 Published online 25 October 2001;
 10.1126/science.1066521
 Include this information when citing this paper.

ARR1, a Transcription Factor for Genes Immediately Responsive to Cytokinins

Hiroe Sakai,^{1*} Takashi Honma,^{1†} Takashi Aoyama,¹ Shusei Sato,² Tomohiko Kato,² Satoshi Tabata,² Atsushiro Oka^{1‡}

Cytokinins are a class of phytohormones involved in various physiological events of plants. The *Arabidopsis* sensor histidine kinase CRE1 was recently reported to be a cytokinin receptor. We used a steroid-inducible system to show that the transcription factor-type response regulator ARR1 directs transcriptional activation of the *ARR6* gene, which responds to cytokinins without de novo protein synthesis. This fact, together with characteristics of *ARR1*-overexpressing plants and *arr1* mutant plants, indicates that the phosphorelay to ARR1, probably from CRE1, constitutes an intracellular signal transduction occurring immediately after cytokinin perception.

Cytokinins induce a variety of physiological events, including cell division, chloroplast development, and shoot formation (1, 2). These cytokinin responses in *Arabidopsis* are at least partly triggered through the recognition of cytokinins by the sensor protein CRE1, which is a member of the protein histidine kinase family (3, 4). In bacteria, histidine kinases participate in His-Asp phosphorelays, which respond to environmental stimuli, usually in association with cytoplasmic response regulators, the majority of which are transcription factors (5). If

the His-Asp phosphorelay is also the case in plants, an intracellular signal transduction pathway starting from the CRE1 sensor may involve response regulators. The *Arabidopsis* genome codes for 22 response regulators (ARRs), 12 of which contain a Myb-like DNA binding domain called ARRM (type B) (6–9). The remainder (type A) possess no apparent functional unit other than a signal receiver domain containing two aspartate and one lysine residues (DDK) at invariant positions, and their genes are transcriptionally induced by cytokinins without de novo protein synthesis (7, 9, 10). The type B members, ARR1 and ARR2, bind DNA in a sequence-specific manner and work as transcriptional activators (11, 12).

We analyzed the morphological characteristics of transgenic plants carrying *35S::ARR1* and *35S::ARR1ΔDDK* genes (13), in which the full-length ARR1 and its truncated version missing

¹Laboratory of Molecular Biology, Institute for Chemical Research, Kyoto University, Uji, Kyoto 611-0011, Japan. ²Kazusa DNA Research Institute, Kisarazu, Chiba 292-0812, Japan


*These authors contributed equally to this work.

†Deceased 14 July 2001.

‡To whom correspondence should be addressed. E-mail: oka@scl.kyoto-u.ac.jp

REPORTS

Fig. 1. Phenotypes of transgenic plants. (A and E) Wild-type plants grown without and with 1 μ M BA, respectively. (B and F) 35S::ARR1 plants grown without and with 1 μ M BA, respectively. (G) An enlargement of (F). (C and D) 35S::ARR1 Δ DDK plants without BA. (H and I) 35S::ARR1 Δ DDK::GR plants without and with 0.1 μ M DEX, respectively. Arrowheads indicate cotyledons. Scale bars, 1 mm.



the NH₂-terminal signal receiver domain, respectively, were directed by the cauliflower mosaic virus 35S promoter. On the one hand, transgenic 35S::ARR1 plants showed no phenotypic change under normal growth conditions, except for hypertrophic cotyledons, longer cotyledonary petioles, and shorter roots than wild-type plants (Fig. 1, A and B, and Fig. 2A). On the other hand, 35S::ARR1 Δ DDK plants did show a phenotypic change under normal growth conditions, forming ectopic shoots on the adaxial surface of their cotyledons (Fig. 1D). In lines expressing the transgene at relatively high levels, plant growth was occasionally inhibited, with concomitant disordered cell proliferation around the shoot apex (Fig. 1C). Because the transactivating function of ARR1 was masked by its signal receiver domain in transient expression experiments (12), the phenotypes generated by 35S::ARR1 Δ DDK likely resulted from the constitutive transactivating function of ARR1 Δ DDK, independent of a signal from an upstream component. To test whether the cytokinin signal regulates ARR1 through its signal receiver domain, we treated 35S::ARR1 plants with exogenous 6-benzylaminopurine (BA) at various concentrations and compared their morphologies with those of nontreated 35S::ARR1 Δ DDK plants. Transgenic 35S::ARR1 plants grown in the presence of 1 μ M BA showed growth inhibition concomitant with disordered cell proliferation around the shoot apex (Fig. 1, F and G). This phenotype was similar to a severe phenotype seen in nontreated 35S::ARR1 Δ DDK plants (Fig. 1C). Other kinds of cytokinins (kinetin, trans-zeatin, and isopentenyladenine) had similar effects on 35S::ARR1 plants, whereas a different phytohormone, auxin (3-indoleacetic acid), did not (14). These results suggest that the signal receiver domain of ARR1 suppresses the function of ARR1 in the absence of cytokinins and that a cytokinin signal releases the suppression. Treatment of wild-type plants with BA caused growth inhibition but not the disordered cell proliferation that was observed in 35S::ARR1 plants. (Fig. 1E). This difference

might imply that ectopic overexpression of ARR1 disturbs homeostasis in cytokinin responses.

Another finding from morphological analyses is that plants overexpressing ARR1 appear to be more sensitive to cytokinins than are wild-type plants (Fig. 1, E and F). To investigate the relationship between sensitivity to cytokinins and level of ARR1 expression, we examined wild-type, 35S::ARR1, and *arr1-1* mutant plants by two different procedures: root-elongation and callus-formation assays. The *arr1-1* mutant obtained by screening transferred DNA (T-DNA)-tagged lines (15) contains a T-DNA insertion within the first intron of the ARR1 gene and produces no mature ARR1 mRNA (see below). Apparent phenotypic alterations were not detected in *arr1-1* plants, except for longer roots than in wild-type plants under normal growth conditions (Fig. 2A). In the root-elongation assay, we grew seedlings on agar media containing various concentrations of BA. In the absence of BA, 35S::ARR1 and *arr1-1* plants had shorter and longer roots than did wild-type plants, respectively. BA inhibited root growth in all the plants at concentrations of 10 nM and higher (Fig. 2). However, roots of 35S::ARR1 and *arr1-1* plants were more and less sensitive, respectively, to BA than were those of wild-type plants (see IC₅₀, the concentration of BA needed to cause 50% inhibition of root growth, shown in Fig. 2B). In the callus-formation assay, we tested the response (green callus formation) of tissue-cultured explants to varying concentrations of the cytokinin kinetin and to the auxin 2,4-dichlorophenoxyacetic acid (2,4-D), using the procedure described by Kubo and Kakimoto (16). Like wild-type explants, 35S::ARR1 and *arr1-1* explants responded to increasing levels of kinetin with rapid proliferation and greening, and in the case of 35S::ARR1 explants, occasionally with shoot formation (Fig. 3). However, the sensitivity of 35S::ARR1 and *arr1-1* explants to kinetin was higher and lower, respectively, than that of wild-type explants. These results show that the level of ARR1 expression




Fig. 2. Root growth of wild-type (circles), 35S::ARR1 (triangles), and *arr1-1* (squares) plants in the presence of exogenous cytokinin. Seeds were germinated and grown on vertical agar medium containing various concentration of BA. (A) The primary root length of 8-day-old seedlings was measured, and the average and standard deviations from at least 15 individuals for each condition were calculated and plotted against the BA concentration. (B) Root growth is expressed relative to the mean root length of respective plants grown without BA. Dashed lines indicate the BA concentrations causing IC₅₀.

correlates with sensitivity to cytokinin, which supports the view that ARR1 is involved in the cytokinin signal transduction pathway.

To gain insight into how ARR1 contributes to cytokinin signal transduction at the molecular level, we examined transcription levels of a cytokinin-responsive marker gene in 35S::ARR1 and *arr1-1* plants, with or without BA treatment, by Northern blot analysis (17). The marker gene we chose was ARR6 (18), a type A response regulator gene, because its expression is rapidly induced by cytokinins without de novo protein synthesis (7, 10). As in wild-type plants, ARR6 responded rapidly to BA in both 35S::ARR1 and *arr1-1* plants, even in the presence of cycloheximide (CHX), a protein synthesis inhibitor (Fig. 4A). However, the transcription levels induced by the BA treatment of 35S::ARR1 and *arr1-1* plants were higher and lower, respectively, than in wild-type plants. This induction pattern, together with the appearance of an ARR1 recognition sequence (5' -AGATT-3') (12) three times in the 200-base pair region upstream from the ARR6 translation start site, suggests that the transactivation of ARR6 is targeted by ARR1. Consistent with this, 35S::ARR1 Δ DDK plants showing severe phenotypes expressed ARR6 at much higher levels than did wild-type plants

REPORTS




Fig. 3. Callus growth of *35S::ARR1*, wild-type, and *arr1-1* explants exposed to different cytokinin and auxin concentrations. Hypocotyl segments were excised and cultured on MS media containing different levels of kinetin and 2,4-D. After 21 days in culture, the induced calli were arranged and photographed. (A) *35S::ARR1* explants, (B) wild-type explants, and (C) *arr1-1* explants.

under normal growth conditions (14). Direct transactivation of *ARR6* by *ARR1* was shown with transgenic plants carrying a chimeric *35S::ARR1ΔDDK::GR* gene, whose translation product was expected to become activated by glucocorticoid without de novo protein synthesis (19). Such steroid-inducible systems using the hormone-binding domain of the glucocorticoid receptor (GR) have been used as powerful tools to identify direct target genes for various transcription factors (20–22). Transgenic *35S::ARR1ΔDDK::GR* plants were indistinguishable from wild-type plants under normal growth conditions (Fig. 1H), but the addition of a synthetic glucocorticoid, dexamethasone (DEX), caused graduated morphological changes, which were dependent on the DEX concentration. A plant treated with 0.1 μM DEX showed growth inhibition concomitant with disordered cell proliferation around the shoot apex (Fig. 1I), which was similar to a severe phenotype in *35S::ARR1ΔDDK* plants (Fig. 1C). These results indicate that *ARR1* function can be modulated by DEX instead of by cytokinins in this system. Treatment of *35S::ARR1ΔDDK::GR* plants with DEX for 1 hour enhanced *ARR6* transcription, even in the presence of CHX (Fig. 4B) (17). This provides in vivo evidence for direct interaction of *ARR1ΔDDK::GR* with the *ARR6* gene. This in vivo evidence, taken together with the higher and lower *ARR6* transcription levels in BA-treated *35S::ARR1* and *arr1-1* plants, respec-




Fig. 4. Northern blot analysis for *ARR6* transcription. (A) Wild-type, *35S::ARR1*, and *arr1-1* plants were treated with 5 μM BA and/or 30 μM CHX for 30 min and then subjected to Northern blot analysis for *ARR6* and *ARR1*, with *UBQ5* as a control. (B) Transgenic *35S::ARR1ΔDDK::GR* plants were treated with 30 μM DEX and/or 30 μM CHX for 1 hour. Numerals below the *UBQ5* signals indicate the level of *ARR6* mRNA that was quantified by measuring the radioactivity of the signal band with a Fujix BA100 Bio-Image analyzer, divided by the radioactivity of the *UBQ5* signal band with the same RNA sample for normalization and presented as the relative value to that for wild-type plants (A) or *35S::ARR1ΔDDK::GR* plants (B) without chemical treatment.

tively, versus wild-type plants, lead us to conclude that *ARR1* directly activates *ARR6* transcription in response to cytokinins.

In *Arabidopsis*, cytokinin receptors, ethylene receptors, and, possibly, osmosensors are histidine kinases (3, 4, 23–25). However, their cognate response regulators, working as transcription factors, have never been identified. Here we present evidence that *ARR1* mediates a cytokinin signal, probably through its NH₂-terminal signal receiver domain, and transactivates *ARR6*, which is immediately responsive to cytokinins. A paralogous response regulator, *ARR2*, shows almost identical characteristics to *ARR1* (12), suggesting a functional overlap. Residual cytokinin responses observed with the *arr1-1* mutant may have been provided by *ARR2*. In addition to *ARR6*, other type A member genes, including *ARR4*, *ARR5*, *ARR7*, *ARR8*, and *ARR9*, were also activated by DEX at various levels in *35S::ARR1ΔDDK::GR* plants (14), suggesting that all the immediate cytokinin-responsive genes belonging to this group are directly activated by *ARR1*. Also, other cytokinin-responsive genes whose promoter regions contain the *ARR1* recognition sequences are possibly transactivated by *ARR1*. A screening for *ARR1* target genes using transgenic *35S::ARR1ΔDDK::GR* plants will shed light on the whole view of the early cytokinin signal transduction pathway. We conclude that *ARR1* is a principal transcription factor-type response regulator that is involved in an early step of cytokinin signal transduction, possibly as a partner of the sensor histidine kinase CRE1.

References and Notes

1. F. Skoog, C. O. Miller, *Symp. Soc. Exp. Biol.* **11**, 118 (1957).
2. M. C. Mok, in *Cytokinins*, D. W. S. Mok, M. C. Mok, Eds. (CRC Press, Boca Raton, FL, 1994), pp. 155–166.
3. T. Inoue et al., *Nature*, **409**, 1060 (2001).
4. T. Suzuki et al., *Plant Cell Physiol.* **42**, 107 (2001).
5. J. B. Stock, A. M. Stock, J. M. Mottonen, *Nature* **344**, 395 (1990).
6. J. L. Riechmann et al., *Science* **290**, 2105 (2000).
7. I. B. D'Agostino, J. Derrière, J. J. Kieber, *Plant Physiol.* **124**, 1706 (2000).
8. H. Sakai, T. Aoyama, H. Bono, A. Oka, *Plant Cell Physiol.* **39**, 1232 (1998).
9. A. Imamura et al., *Plant Cell Physiol.* **40**, 733 (1999).
10. M. Taniguchi et al., *FEBS Lett.* **429**, 259 (1998).
11. J. Lohrmann et al., *Plant Biol.* **1**, 495 (1999).
12. H. Sakai, T. Aoyama, A. Oka, *Plant J.* **24**, 703 (2000).
13. The chimeric genes *35S::ARR1* and *35S::ARR1ΔDDK* were described previously (12). For the chimeric *35S::ARR1ΔDDK::GR* gene, *35S::ARR1ΔDDK* was ligated inframe to a cassette directing the hormone-binding domain of the rat GR (19). *Arabidopsis thaliana* (Columbia ecotype) plants were transformed by the vacuum infiltration method (26). More than 10 independent lines were obtained for each construct, and their T3 plants were used in experiments. All these plants were confirmed to transcribe the respective transgene by Northern blot analysis.
14. H. Sakai, T. Honma, T. Aoyama, A. Oka, unpublished results.
15. We obtained the T-DNA insertion mutant *arr1-1* (Columbia ecotype) using the screening system developed at the Kazusa DNA Research Institute. The principles of the screening method were as described (27). Synthetic primers 5'-ACCCAGAATTGAGCTCAAAGC-3', 5'-GTGCAATCCCTCTCTGTATATC-3', and 5'-CAGCAGCAACAACTGCTGATG-3' corresponded to the *ARR1* gene; and the primers 5'-AAGAAAATGCCGATACTTCAATGGC-3' and 5'-TTCCTTAATTCTCCGCTCATGATC-3' corresponded to the T-DNA left and right borders, respectively, of the pGTAC-LUS vector. Six pairs of primers, each consisting of an *ARR1*-specific primer and a T-DNA-specific primer, were used for polymerase chain reactions. The position of the T-DNA insert was determined by sequencing the junction between the T-DNA and the genome.
16. M. Kubo, T. Kakimoto, *Plant J.* **23**, 385 (2000).
17. Wild-type *35S::ARR1*, *35S::ARR1ΔDDK::GR*, and *arr1-1* plants were grown at 22°C under constant light on sterile filter paper on Murashige and Skoog (MS) medium (28) supplemented with 1% sucrose and 0.8% agar. Two-week-old plants, which were adapted to open-air conditions, were treated for the indicated lengths of time by immersing the filter paper and roots in a hydroponic growth medium containing 5 μM BA and 30 μM DEX and/or 30 μM CHX. Total RNA was isolated and subjected to Northern blot analysis as previously described (8).
18. A. Imamura et al., *Proc. Natl. Acad. Sci. U.S.A.* **95**, 2691 (1998).
19. T. Aoyama, N.-H. Chua, *Plant J.* **11**, 605 (1997).
20. R. W. M. Sablowski, E. M. Meyerowitz, *Cell* **92**, 93 (1998).
21. D. Wagner, R. W. M. Sablowski, E. M. Meyerowitz, *Science* **285**, 582 (1999).
22. A. Samach et al., *Science* **288**, 1613 (2000).
23. C. Chang, S. F. Kwok, A. B. Bleecker, E. M. Meyerowitz, *Science* **262**, 539 (1993).
24. J. Hua, C. Chang, Q. Sun, E. M. Meyerowitz, *Science* **269**, 1712 (1995).
25. T. Urao et al., *Plant Cell* **11**, 1743 (1999).
26. N. Bechtold, J. Ellis, G. Pelletier, *C. R. Acad. Sci. Paris. Life Sci.* **316**, 1194 (1993).
27. E. C. McKinney et al., *Plant J.* **8**, 613 (1995).
28. T. Murashige, F. Skoog, *Physiol. Plant* **15**, 493 (1962).
29. Supported by grants from the Ministry of Education, Culture, Sports, Science, and Technology of Japan to A.O. (Special Coordination Fund) and to T.A. (Grants-in-Aid for Scientific Research on Priority Areas, no. 12142205).

8 August 2001; accepted 3 October 2001