

**Size-relative Effectiveness of Clove Oil as an Anaesthetic for Rainbow Trout
(*Oncorhynchus mykiss* Walbaum, 1792) and Goldfish
(*Carassius auratus* Linnaeus, 1758)**

Costas Perdikaris¹, Cosmas Nathanaelides¹, Evangelia Gouva¹, Ugwemorubong Ujagwung Gabriel², Konstantina Bitchava³, Fotini Athanasopoulou³, Ageliki Paschou⁴, Ioannis Paschos¹

¹Department of Aquaculture and Fisheries, Technological Educational Institute of Epirus, Igoumenitsa, Greece

²Department of Fisheries and Aquatic Environment, Faculty of Agriculture, Rivers State University of Science and Technology, Nigeria

³Laboratory of Ichthyology and Aquatic Animal Medicine, Faculty of Veterinary Medicine, University of Thessaly, Karditsa, Greece

⁴Hatzikosta Hospital, Haematology and Microbiology Laboratory, Ioannina, Greece

Received June 1, 2009

Accepted January 19, 2010

Abstract

The purpose of this work was to investigate the size-relative effectiveness of clove oil as an anaesthetic for rainbow trout and goldfish. In total, 128 rainbow trout (*Oncorhynchus mykiss*) (two groups of 20–23 and 30–33 cm mean fork length) and 160 goldfish (*Carassius auratus*) (four size groups of 1.5–2.5, 5–7, 11–15 and 20–25 cm) were anaesthetized at different clove oil concentrations of 50, 100, 150 mg·l⁻¹ for trouts and 75, 100, 150 mg·l⁻¹ for goldfish. Rainbow trout exhibited total loss of balance and no response to external stimuli with shorter induction time as dosage increased (120.5 s, 64.4 s and 44.3 s, respectively). Goldfish exhibited total loss of balance and no response to external stimuli after induction time that varied with dosage used and body size of fish. The small fish (1.5–7 cm) exhibited shorter induction time which ranged from 84.28 s at 75 mg·l⁻¹ clove oil to 41.14 s at 150 mg·l⁻¹ clove oil. The larger fish had a longer induction time inversely related to the dosage. Recovery time was longer than induction time in both species. Both species recovered within 6 min after anaesthesia at 150 mg·l⁻¹ clove oil. Clove oil did not produce marked changes ($P < 0.05$) in the physiological indicators of goldfish compared to the control. However, marked changes ($P < 0.05$) were exhibited in the haematocrit of treated rainbow trout that also exhibited hyperkalaemia and hyperglycaemia ($P > 0.05$). For both fish species, clove oil was effective, producing minimum stress and zero mortalities, and can be recommended as an effective anaesthetic.

Syzygium aromaticum, fish anaesthesia, animal welfare

Rapid expansion of the aquaculture industry that occurred in previous decades prompted scientific debates on the potential suffering of fish being handled during common aquaculture procedures or during slaughtering. Research aimed at lessening the suffering of cultured fish is vital to meet the concern for farmed fish welfare (Ashley 2007). Handling stress and various manipulations in aquaculture can have a negative impact on fish health and their growth (Hoskonen and Pirhonen 2006). Anaesthetics are therefore applied to reduce these negative effects to the minimum. The dosage required to induce general anaesthesia varies according to the anaesthetic used and other factors such as water temperature, hardness, salinity, oxygen concentration, length of exposure, body weight, the ratio of gill area/body surface area and the species of fish. In general, small fish are more sensitive to anaesthesia than larger fish (Ross and Ross 1999).

An ideal anaesthetic for fish should induce anaesthesia in less than 3 to 5 min, with total loss of balance and muscle tone, allowing an uneventful and rapid (i.e. less than 10 min) recovery with low tissue residues after recovery, thus being safe to users and consumers. The anaesthetic should be inexpensive and easy to use (Gilderhus and Marking 1987;

Address for correspondence:

Cosmas Nathanaelides
Department of Aquaculture & Fisheries
Technological Educational Institute of Epirus
Irinis & Filias 1, 46100 Igoumenitsa, Greece

Phone: +30 26650 49852
Fax: +30 26650 49886
E-mail: cosmasfax@yahoo.com
<http://www.vfu.cz/acta-vet/actavet.htm>

Summerfelt and Smith 1990; Hseu et al. 1998; Ross and Ross 1999; Tsantilas et al. 2006; Klimankova et al. 2008).

Some traditionally used anaesthetics in aquaculture include tricaine methanesulphonate (MS-222), benzocaine, etomidate, metomidate, 2-phenoxyethanol and quinaldine. In farmed fish, MS-222 appeared to be the most widely used anaesthetic. However, recently there has been considerable interest in another fish anaesthetic, clove oil extracted from the clove tree, *Eugenia caryophyllus* (syn. *Syzygium aromaticum*) which has traditionally been used as human anaesthetic (Soto and Burhanuddin 1995). Chaieb et al. (2007) reported that the main active ingredients of clove oil are: eugenol (88.58%), eugenyl acetate (5.62%), β -caryophyllene (1.39%), 2-heptanone (0.93%), ethyl hexanoate (0.66%), humulenol (0.27%), α -humulene (0.19%), calacorene (0.11%) and calamenene (0.10%). Clove oil is generally considered a reasonable alternative of fish anaesthetic with low cost and of no risk to human health (Velisek et al. 2005a,b). In fact, clove oil is used for local application to reduce pain and promote healing and exhibits antimicrobial, antioxidant, antifungal and antiviral properties (Chaieb et al. 2007). According to the U.S. FDA, the constituting ingredients of clove oil are considered safe (substance that can be used in food industry), but none of them has been approved for fish anaesthesia yet (FDA/CVM, 2009).

Clove oil has recently been studied as a potential anaesthetic for some ornamental fish (Kaiser et al. 2006; Macova et al. 2008) and several farmed cold and warm water fish species (Wagner et al. 2003; Holloway 2004; Velisek et al. 2005a,b; Hajek et al. 2006, Gomulka et al 2008). However, results of several studies suggest that the physiology of fish can be affected by changes in the haematology and biochemistry of exposed fish. Therefore it is important to assess the extent of physiological changes of fish exposed to an anaesthetic (Inoue et al. 2005; Park et al. 2008). Results of the numerous studies on clove oil revealed that effective concentrations for anaesthesia vary with fish body size and water temperature (Oikawa et al. 1994; Hoskonen and Pirhonen 2006; Mylonas et al. 2005; Zahl et al. 2009) with the smaller fish being more responsive than the larger one (Holloway et al. 2004; Velisek et al. 2005a,b). Velisek et al. (2005b) studied the effect of clove oil on juvenile and brood stock of rainbow trout (*Oncorhynchus mykiss*) but there are no data for larger sizes of this species, in particularly for the final growing stages (200–300 g) where size grading prior to harvesting may be required. Goldfish (*Carassius auratus*) is a warm water ornamental fish species, which is frequently subjected to anaesthesia for spawning and vaccination and could benefit from the use of clove oil.

The aim of this study was to observe the response of two fish species of different sizes to clove oil used as an anaesthetic and to observe its effect on selected haematological and biochemical variables.

Materials and Methods

Experimental animals

Male rainbow trout ($n = 128$) with an average fork length 31.06 cm (± 2.47 , SD) and goldfish ($n = 160$) with mean body weight of 6.8 g (± 1.11 , SD) were used for independent experiments. The trout were maintained in a circular 2 m³ tank connected to a re-circulating system at mean temperature of 12 ± 0.6 °C. Goldfish were kept in a circular 200 l tank connected to a second re-circulating system at a mean temperature of 18 ± 0.7 °C. The trout were fed daily with trout pellets at 2% of body weight; the goldfish were provided carp pellets at a 3% daily ratio. Oxygen concentrations were maintained above 7 and 8.5 ppm in the tanks for goldfish and trout, respectively. In both groups, concentrations of pH, nitrate and ammonia were within the physiological/optimal range for these species (pH: 7.5–7.7; nitrate ($P < 0.01$ mg l⁻¹), ammonia ($P < 0.03$ mg l⁻¹); photoperiod 12 L:12 D. All fish were starved for 24 h before the experiment.

Before the experiment, the fish were observed daily to evaluate their health condition as indicated by their activity and external appearance. Four days before the experiment, ten fish were randomly selected from each tank and sampled for bacteriological and parasitological examinations. Kidney and spleen samples were inoculated onto Tryptone Soy Agar (TSA) and Thiosulphate Citrate Bile Salt Agar (TCBS) for bacteriology assessment (Roberts and Shepherd 1997). Squash imprints of gill, skin, gall bladder, liver, spleen, kidney, muscle, brain and gut tissues from freshly killed fish were examined for the presence of parasites.

Prior to anaesthesia, the goldfish were divided into four fork length classes (1.5-2.5, 5-7, 11-15 and 20-25 cm) and placed in separate glass aquaria (volume 70 l). The trout were divided into two size groups (20-23 and 30-33 cm) and maintained in different holding tanks. Seven to ten fish of each species and size group were exposed to three different concentrations of clove oil ($n = 7-10$ fish per size group/dosage tested).

AAnaesthesia preparation and procedures

Clove oil (Sigma Aldrich Co., St. Louis, USA) was initially dissolved in 95% ethanol (clove oil:ethanol at a ratio 1:9). In pilot experiments, it was confirmed that the volume of ethanol used in each experiment did not have a visible anaesthetic effect on fish. The various concentrations (50, 100 and 150 mg·l⁻¹ for trout and 75, 100 and 150 mg·l⁻¹ for goldfish) were prepared and mixed in 20 l observation tanks filled with water from the respective recirculating systems. Fish were randomly caught by hand-net in the holding tanks and transferred immediately into the anaesthetic in the observation tank. The fish were observed for opercular movements, balance and thereafter loss of response to stimulus. The fish were removed when they exhibited loss of equilibrium, no spinal reflexes and imperceptible opercular movements.

Induction and recovery times were obtained using a digital chronometer for fish. Induction of anaesthesia was assumed as complete when fish lost equilibrium and reflex reactivity (stage 5). Once a fish reached stage 5 of anaesthesia, it was removed from the experimental tanks, dried and measured. Then the fish was transferred to fresh water in identical tank without rinsing to remove traces of the anaesthetic. Recovery was considered complete in all groups when fish were able to regain upright position and swim normally.

Recovery and induction times were recorded for each fish. Following recovery, the fish were transferred to maintenance tanks and observed for 48 h for potential mortality. Control groups were fish that remained in the rearing tanks and left undisturbed until they were collected for blood sampling.

Blood sample collection

Blood was sampled from trout (30-33 cm group) and goldfish (20-25 cm group) with heparinized syringes from the caudal vein at stage 5 of anaesthesia. Haematocrit was calculated after centrifugation of microhaematocrit tubes (4,000 g for 5 min, Sigma 1-15 microcentrifuge). For plasma biochemical analysis, blood was collected (0.5-1.5 ml) in Eppendorff vials and centrifuged (9,000 g for 13 min, Sigma 1-15 microcentrifuge). The plasma was collected and stored at -80 °C for further analysis. Glucose and calcium plasma concentrations in trout and goldfish plasma were measured by Olympus 600 Medicon Analyzer, using Hexokinase enzymatic UV and Arsanazo III photometric colour tests, respectively.

Statistical analysis

Box plots were plotted for induction and recovery times for the various sizes of fish using SPSS 15.0 software. The above data and haematocrit, glucose and calcium were subjected to a one-way ANOVA, and Duncan multiple range test used to separate the means (Zar 1984). In all cases, confidence levels were set at 95%.

Results

All fish used in the present study were healthy as was indicated by their activity and exterior appearance and the bacteriological and parasitological examinations carried out in samples of ten fish from each holding tank. No mortality was observed during the acclimatization period. Furthermore, no deaths or other adverse effects occurred within 48 h following recovery from anaesthesia.

Anaesthetic effect of clove oil on rainbow trout

Figs 1a and 1b present the results of induction (1a) and recovery (1b) time of rainbow trout exposed to different doses of the anaesthetic, and the same results are presented in Figs 2a and 2b with fish grouped according to their size. Anaesthesia progressed rapidly but steadily in all treated groups increasing with the concentration of clove oil. Induction time was faster (120 s) and similar for all groups with concentration 50 mg·l⁻¹. Induction time decreased with the higher concentration used in large (30-33 cm) size group, whereas the reverse effect was observed for the smaller (20-23 cm) size group (Fig. 3a, Table 2). Compared to the large size class, the small size class exhibited longer induction time at all dosages tested ($P < 0.001$).

Recovery time was more variable at 100 mg·l⁻¹ ($P < 0.001$) compared to the other concentrations (Fig. 3b). This time increased with higher concentration used, ranging between 2 - 6.6 s (Table 2), but did not differ with size, $P > 0.05$ (Fig. 4b, Table 2). The higher dose (100 mg·l⁻¹) induced deep anaesthesia within 45 s, i.e. much faster compared to the other doses ($P < 0.001$). All fish recovered within 6 min, although in the lower doses recovery progressed much more rapidly ($P < 0.001$).




Fig. 1. Box plots of (a) induction and (b) recovery times for rainbow trout anaesthetized with various concentrations of clove oil. The upper and lower hinges of each plot represent the maximum and times, middle line represents the median, the shaded inner box represents the 25th and 75th percentiles and the whiskers represent lower and upper limits.

Anaesthetic effect of clove on goldfish

Figs 3a and 3b present the results of induction (3a) and recovery (3b) of goldfish exposed to different doses of the anaesthetic, and the same results are presented in Figs 4a and 4b for fish grouped according to size. Induction time of anaesthesia varied ($P < 0.01$) with clove concentration, decreasing with the increase of clove concentration (Fig. 1a, Table 1) and was more variable at 75 and 150 mg·l⁻¹ compared to 100 mg·l⁻¹ (Fig. 1b, Table 1). An increase in recovery time with clove concentration was recorded for the largest size class, a decrease for the smallest, and a variable response for the medium (Table 1). Induction time was rapid in small-size goldfish, irrespective of the clove oil dose applied. Anaesthesia was more size-dependent at the lower dose, compared to the higher dose ($P < 0.001$). Smaller fish required less time to lose the upright position than larger ones at the dose of




Fig. 2. Box plots of (a) induction and (b) recovery times for different sizes of rainbow trout anaesthetized with clove oil. The upper and lower hinges of each plot represent the maximum and times, middle line represents the median, the shaded inner box represents the 25th and 75th percentiles and the whiskers represent lower and upper limits.

75 mg·l⁻¹ ($P < 0.001$). For the higher dose, induction time was rather uniform in all size groups ($P > 0.05$).

Recovery time was more size-dependent at the lower dose (75 mg·l⁻¹), with larger fish requiring less time to recover ($P < 0.001$) (Table 1). However, smaller fish recovered earlier compared to the other size groups ($P < 0.001$).

Physiological responses in trout and goldfish

Clove oil did not produce marked changes ($P < 0.05$) in the physiological indicators of experimental goldfish compared to control (Table 3). However, it produced significant changes between haematocrit of the control for rainbow trout and the treated group; and




Fig. 3. Box plots of (a) induction and (b) recovery times for goldfish anaesthetized at various concentrations of clove oil. The upper and lower hinges of each plot represent the maximum and times, middle line represents the median, the shaded inner box represents the 25th and 75th percentiles and the whiskers represent lower and upper limits.

between the treated groups ($P < 0.001$). It also caused hyperkalaemia and hyperglycaemia ($P > 0.05$) in the trout compared to control, but not in the goldfish (Table 3).

Discussion

In this study, clove oil was found to be effective anaesthetic for both rainbow trout and goldfish. Both fish species exhibited normal behavior and remained calm during the induction time with no struggling or rapid swimming, which was a positive sign of their welfare. Additionally, no mortalities were observed for a 24 h post-recovery period.

Some species-specific differences were evident in the induction time. Rainbow trout exhibited a more rapid induction time compared to goldfish at all the doses tested. This may be the result of a higher uptake of eugenol via the gills (Keene et al. 1998) in rainbow trout which is a fish species with higher respiratory activity compared to the more sluggish ornamental goldfish. In goldfish, induction time was related to the size of fish, with more time needed for larger fish to be anaesthetized. This may be because of higher rate of uptake of the anaesthetic through the gills in the smaller fish compared to the larger ones. This is in accordance with the results reported by Woody et al. (2002) for sockeye




Fig. 4. Box plots of (a) induction and (b) recovery times for different sizes of goldfish anaesthetized with clove oil. The upper and lower hinges of each plot represent the maximum and times, middle line represents the median, the shaded inner box represents the 25th and 75th percentiles and the whiskers represent lower and upper limits.

salmon (*Oncorhynchus nerka*). The higher dose produced a rapid and uniform response in all size classes, suggesting that a dose of 150 mg·l⁻¹ might be well over the effective concentration.

For rainbow trout, recovery time increased with increased doses of clove. In goldfish, recovery time varied with the concentration of clove oil for various size groups. An increase in recovery time relative to clove concentration was recorded in roach (*Rutilus rutilus*) (Sudagara et al. 2009), cobia (*Rachycentron canadum*) (Gullan and Villanuera 2009) and rainbow trout (Velisek et al. 2005b), and to 2-phenoxyethanol in rainbow trout (Velisek et al. 2004). Woody et al. (2002) and Inoue et al. (2003) found no direct relationship between clove concentration and recovery times in adult sockeye salmon (*Oncorhynchus nerka*) and matrinxá (*Brycon cephalus*). Results of this study seem to confirm species-specific differences in recovery times of anaesthetized fish. In rainbow trout, recovery time was slightly longer 1.3 - 8.1 × compared to induction time. This could be explained by the fact that eugenol exerts an inhibitory effect on respiration, reducing the ability of fish to remove the excess of eugenol from the system via the

Table 1. Induction and recovery time of goldfish anaesthetized with clove oil

Fish length (cm)	Dosage of clove oil (mg·l ⁻¹)			Mean (± SD) for size	Dosage of clove oil (mg·l ⁻¹)		
	75	100	150		75	100	150
	Induction time (s)				Recovery time (s)		
1.5-2.0	84.28 (9.39)	111.0 (13.67)	41.15 (5.63)	78.81 (31.01) ^c	306.86 (12.90)	294.14 (16.43)	115.00 (4.90)
11.0-15	256.43 (34.22)	270.43 (16.66)	119.43 (7.16)	78.81 (31.01) ^b	189.00 (25.76)	364.00 (24.11)	257.71 (14.24)
20.0-25	475.00 (34.22)	366.14 (56.93)	117.14 (35.14)	215.43 (71.00) ^a	110.86 (23.17)	256.00 (28.33)	381.71 (38.94)
Mean (± SD) for each dosage	271.90 (165.19) ^c	249.19 (112.87) ^b	92.5714 (42.25) ^a		202.23 (85.00) ^a	304.71 (50.95) ^b	251.47 (114.00) ^c

(n = 10 for each data point, figures in parenthesis, SD; Mean in the respective row or column under induction or recovery with similar superscripts are not significantly different, $p > 0.05$)

Table 2. Induction and recovery time of rainbow trout anaesthetized with clove oil

Fish length (cm)	Dosage of clove oil (mg·l ⁻¹)			Mean (± SD) for size	Dosage of clove oil (mg·l ⁻¹)		
	75	100	150		75	100	150
	Induction time (s)				Recovery time (s)		
20.00-23.00	120.50 ± 10.55	340.17 ± 15.77	399.50 ± 53.24	283.53 ± 123.23 ^a	176.00 ± 10.34	340.17 ± 15.77	399.50 ± 53.24
30-33.00	120.50 ± 0.56 ^c	64.40 ± 3.20 ^b	44.30 ± 6.24 ^a	76.4000 ± 33.55 ^b	156.80 ± 5.88 ^c	240.40 ± 13.25 ^b	363.50 ± 9.32 ^a
Mean (± SD) for each dosage	121.27 ± 0.83 ^c	167.81 ± 138.21 ^b	145.79 ± 168.55 ^a		163.20 ± 11.86 ^c	277.81 ± 51.76 ^b	373.79 ± 31.61 ^a

(n = 10 for each data point, figures in parenthesis, SD; Mean in the respective row or column under induction or recovery with similar superscripts are not significantly different, $p > 0.05$)

Table 3. Haematocrit and biochemical variables of rainbow trout and goldfish anaesthetized with clove oil

Concentration of clove (mg/l)	Haematocrit (%)	Calcium (mmol/l)	Glucose (mmol/l)
Rainbow trout			
0.00	45.58 ± 3.23 ^c	8.23 ± 1.28 ^b	4.94 ± 0.72 ^b
50.00	50.90 ± 4.48 ^b	10.38 ± 0.79 ^a	6.05 ± 0.20 ^a
100.00	58.97 ± 2.67 ^a	9.99 ± 0.62 ^a	5.27 ± 1.35 ^{ab}
150.00	54.58 ± 2.97 ^b	10.51 ± 0.48	5.00 ± 0.42 ^{ab}
Goldfish			
0.00	36.08 ± 1.48 ^a	8.36 ± 1.23 ^a	5.08 ± 0.76 ^a
75.00	37.06 ± 3.22 ^a	8.48 ± 0.59 ^b	5.97 ± 0.29 ^a
100.00	36.03 ± 2.44 ^a	9.3457 ± 1.43 ^{ab}	5.27 ± 0.67 ^a
150.00	35.79 ± 2.32 ^a	8.9367 ± 503 ^{ab}	5.47 ± 0.828 ^a

Means in the respective column for each species with similar superscripts are not significantly different, $p > 0.05$

gills (Keene et al. 1998; Guénette et al. 2007). Our results indicate that clove concentration of 50 mg·l⁻¹ induces the minimum induction and recovery times in rainbow trout; thus, this dose may be suitable for sedation and for standard on-farm procedures that require anaesthesia. However,

the recommended concentrations for both rainbow trout and goldfish may be influenced by environmental factors quite different from those under which this study was conducted.

In some fish species exposed to clove oil, alteration of some blood biochemical and haematocrit values were detected (Cho and Heath 2000; Velisek et al. 2005ab). However, increased haematocrit concentration in the experimental group of rainbow trout compared to control does not suggest toxicity in haematopoietic centres (spleen and head kidney) of fish (Van der Weiden et al. 1994). The effects of toxicants on various tissues may vary with dosage, exposure, body size and species. Tort et al. (2002) reported clove oil altering haematocrit concentrations in rainbow trout. Velisek et al. (2005b) observed the same in rainbow trout and carp, both of them exhibited also significantly increased plasma glucose concentrations. Nevertheless, these findings were observed after longer exposure periods (10 min) compared to 0.69-7.92 min in the present study. These results reveal the significance of exposure time and dosage on some physiological indicators of anaesthetized fish. The short induction time after which blood was collected could explain the lack of biochemical alteration and post-exposure mortality in this study. These results indicate the need to study possible physiological changes occurring in different fish species exposed to different doses of clove oil.

Glucose and calcium plasma concentrations along with cortisol concentrations have been used as stress indicators in fish (Schreck 2000; Sink et al. 2006). In previous studies, clove oil did not induce changes ($P > 0.05$) in plasma glucose and calcium concentrations in trout and goldfish used for experiments (Andersen et al. 1991; Cho and Heath 2000; Wagner et al. 2003). However, mild hyperglycaemia and hypercalcaemia were recorded in some treated groups of the two species. A rise in glucose concentration is a second order reaction under stress (Barton and Iwama 1991) and is mediated by the rise in cortisol concentration induced by stress. Clove was found to block the activity of cortisol, although not completely, in matrxnā (*Brycon cephalus*) (Inoue 2005). Although the mechanism is not well known, Iverson (2003) suggested that it blocks transmission of impulses to the hypothalamus-pituitary interrenal axis (HPI). Cortisol is considered one of the mediators of the increase in plasma glucose under stress (Barton et al. 2002) and the hyperglycaemia recorded in rainbow trout may be explained by the increase in cortisol concentration.

In conclusion, clove oil was found to be safe and can be effectively and easily applied at 150 mg·l⁻¹ to anaesthetize various size groups of rainbow trout and goldfish with minimal disruption in the physiological indicators studied and with zero mortality. Clove oil was easy to handle, without producing unpleasant or irritating odor. Hence, serious considerations should be given to the use of clove oil as a replacement for synthetic forms of anaesthetics.

References

- Ashley PJ 2007. Fish welfare: Current issues in aquaculture. *Appl Anim Behav Sci* **104**: 199-235
- Barton BA, Iwama GK 1991. Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annu Rev Fish Dis* **1**: 3-26
- Chaiet K, Hajlaoui H, Zmantar T, Kahla-Nakbi AB, Rouabia M, Mahdouani K, Bakhrouf A 2007. The chemical composition and biological activity of clove essential oil, *Eugenia caryophyllata* (*Syzygium aromaticum* L. Myrtaceae): A short review. *Phytotherapy Res* **21**: 501-506
- Cho GK, Heath DD 2000. Comparison of tricaine methanesulphonate (MS222) and clove oil anaesthesia effects on the physiology of juvenile chinook salmon *Oncorhynchus tshawytscha* (Walbaum). *Aquac Res* **31**: 537-546
- FDA/CVM 2009. Concerns Related to the use of Clove Oil as an Anesthetic for Fish. FDA Guidance for Industry No 150. Center for Veterinary Medicine, Food and Drug Administration, USA.
- Gilderhus PA, Marking LL 1987. Comparative efficacy of 16 anaesthetic chemicals in rainbow trout. *N Am J Fish Manag* **7**: 288-292
- Gomulka P, Własow T, Velíšek J, Svobodová Z, Chmielinska E 2008. Effects of eugenol and MS-222 anaesthesia on siberian sturgeon acipenser baerii brandt. *Acta Vet Brno* **77**: 447-453
- Guénette SA, Uhlund FC, Heulie P, Beaudry F, Vachon P 2007. Pharmacokinetics of eugenol in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* **266**: 262-265
- Gullian M, Villanueva J 2009. Efficacy of tricaine methanesulphonate and clove oil as anaesthetics for juvenile cobia *Rachycentron canadum*. *Aquac Res* **40**: 852-860

- Hajek GJ, Klyszejko B, Dziaman R 2006. The anaesthetic effect of clove oil on common carp, *Cyprinus carpio* L. *Acta Ichthyol Piscat* **36**: 93-97
- Holloway AC, Keene JL, Noakes DG, Moccia RD 2004. Effects of clove oil and MS-222 on blood hormone profiles in rainbow trout *Oncorhynchus mykiss*, Walbaum. *Aquac Res* **35**: 1025-1030
- Hoskonen P, Pirhonen J 2006. Effects of repeated handling, with or without anaesthesia, on feed intake and growth in juvenile rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Aquac Res* **37**: 409-415
- Hseu JR, Yeh SL, Chu YT, Ting YY 1998. Comparison of efficacy of five anaesthetics in goldlined sea bream, *Sparus sarba*. *Acta Zool Taiwan* **9**: 35-41
- Inoue LAKA, Afonso LOB, Iwama GK, Moares G 2005. Effects of clove oil on the stress response of matrinxã (*Brycon cephalus*) subjected to transport. *Acta Amaz* **35**: 289-295
- Iverson M, Finstad B, McKinley RS, Eliassen RA 2003. The efficacy of metomidate, clove oil, Aqui-S and Benzoak as anaesthetics in Atlantic salmon (*Salmo salar* L.) smolts, and their potential stress-reducing capacity. *Aquaculture* **221**: 549-566
- Keene JL, Noakes DLG, Moccia RD, Soto CG 1998. The efficacy of clove oil as an anaesthetic for rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Aquac Res* **29**: 89-101
- Klimáneková E, Riddellová K, Hajšlová J, Pouštník J, Kolářová J, Kocourek V 2008. Development of an SPME-GC-MS/MS procedure for the monitoring of 2-phenoxyethanol in anaesthetised fish. *Talanta* **75**: 1082-1088
- Macova S, Dolezelova P, Pistekova V, Svobodova Z, Bedanova I, Voslarova E 2008. Comparison of acute toxicity of 2-phenoxyethanol and clove oil to juvenile and embryonic stages of *Danio rerio*. *Neuroendocrinol Lett* **29**: 680-684
- Mylonas CC, Cardinaletti G, Sigelaki I, Polzonetti-Magni A 2005. Comparative efficacy of clove oil and 2-phenoxyethanol as anaesthetics in the aquaculture of European sea bass (*Dicentrarchus labrax*) and gilthead sea bream (*Sparus aurata*) at different temperatures. *Aquaculture* **246**: 467-481
- Oikawa S, Takeda T, Itazawa Y 1994. Scale effects of MS-222 on a marine teleost, porgy *Pagrus major*. *Aquaculture* **121**: 369-379
- Park MO, Hur WJ, Im SY, Seol DW, Lee J, Park IS 2008. Anaesthetic efficacy and physiological responses to clove oil anaesthetized kelp grouper *Epinephelus bruneus*. *Aquac Res* **39**: 877-884
- Roberts RJ, Shepherd CJ 1997. Handbook of Trout and Salmon Diseases, 3rd ed. Fishing News Books, Oxford
- Ross L, Ross B 1999. Anaesthetic and sedative techniques for aquatic animals. Blackwell Science, London, 176 p.
- Schreck CB 2000. Accumulation and long term effects of stress in fish. Moberg GP and Mench JA (eds.): The biology of animal stress: basic principles and implications for animal welfare, CABI Publishing, Wallingford: 147-158
- Sink TD, Richard JS, Henry GK, Alan GM 2006. Linking stress to increased mortality of channel catfish at varying concentrations of *Edwardsiella ictaluri*. *J Anim Vet Adv* **5**: 208-214
- Soto CG, Burhanuddin S 1995. Clove oil as a fish anaesthetic for measuring length and weight of rabbitfish (*Siganus lineatus*). *Aquaculture* **136**: 149-152
- Sudagara M, Mohammadizarejabada A, Mazandarania R, Pooralimotlagha S 2009. The efficacy of clove powder as an anaesthetic and its effects on hematological parameters on roach (*Rutilus rutilus*). *J Aquacult Feed Sci Nutri* **1**: 1-5
- Summerfelt RC, Smith LS 1990. Anaesthesia, surgery, and related techniques. Schreck CB and Moyle PB (eds.), Methods for Fish Biology, American Fisheries Society, Bethesda, MD, pp. 213-272
- Tort L, Puigcerver M, Crespo S, Padrós F 2002. Cortisol and haematological response in sea bream and trout subjected to the anaesthetics clove oil and 2-phenoxyethanol. *Aquac Res* **33**: 907-910
- Tsantilas H, Galatos AD, Athanassopoulou F, Prassinos NN, Kousoulaki K 2006. Efficacy of 2-phenoxyethanol as an anaesthetic for two size classes of white sea bream, *Diplodus sargus* L., and sharp snout sea bream, *Diplodus puntazzo* C. *Aquaculture* **253**: 64-70
- Van der Weiden MEJ, Bleumink R, Seinen W, Van den Berg M 1994. Concurrence of P450 1A induction and toxic effects in the mirror carp (*Cyprinus carpio*), after administration of a low dose of 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Aquat Toxicol* **29**: 147-162
- Velišek J, Svobodová Z 2004. Anaesthesia of common carp (*Cyprinus carpio* L.) with 2-phenoxyethanol: Acute toxicity and effects on biochemical blood profile. *Acta Vet Brno* **73**: 247-252
- Velišek J, Svobodová Z, Piackova V, Groch L, Nepejchalova L 2005a. Effects of clove oil anaesthesia on common carp (*Cyprinus carpio* L.). *Vet Med Czech* **50**: 269-275
- Velišek J, Svobodová Z, Piackova V 2005b. Effects of clove oil anaesthesia on rainbow trout (*Oncorhynchus mykiss*). *Acta Vet Brno* **74**: 139-146
- Wagner GN, Singer TD, McKinley RS 2003. The ability of clove oil and MS-222 to minimize handling stress in rainbow trout (*Oncorhynchus mykiss* Walbaum). *Aquac Res* **34**: 1139-1146
- Woody CA, Nelson J, Ramstad K 2002. Clove oil as an anaesthetic for adult sockeye salmon: field trials. *J Fish Biol* **60**: 340-347
- Zahl IH, Kiessling A, Samuels OB, Hansen MK 2009. Anaesthesia of Atlantic cod (*Gadus morhua*) - Effect of pre-anaesthetic sedation, and importance of body weight, temperature and stress. *Aquaculture* **295**: 52-59