

$b \rightarrow s\gamma$ Branching Fraction and CP Asymmetry

CLEO Collaboration

(February 3, 2008)

Abstract

We present preliminary results on $b \rightarrow s\gamma$ from the CLEO experiment. An updated result on the branching fraction is reported at $\mathcal{B}(b \rightarrow s\gamma) = (3.15 \pm 0.35 \pm 0.32 \pm 0.26) \times 10^{-4}$, where the first uncertainty is statistical, the second is systematic, and the third for model dependence. We also describe a new analysis performed to search for CP asymmetry in $b \rightarrow s\gamma$ decays. We observe no such asymmetry and set conservative limits at $-0.09 < \mathcal{A} < 0.42$.

S. Ahmed,¹ M. S. Alam,¹ S. B. Athar,¹ L. Jian,¹ L. Ling,¹ A. H. Mahmood,^{1,*} M. Saleem,¹ S. Timm,¹ F. Wappler,¹ A. Anastassov,² J. E. Duboscq,² K. K. Gan,² C. Gwon,² T. Hart,² K. Honscheid,² H. Kagan,² R. Kass,² J. Lorenc,² H. Schwarthoff,² E. von Toerne,² M. M. Zoeller,² S. J. Richichi,³ H. Severini,³ P. Skubic,³ A. Undrus,³ M. Bishai,⁴ S. Chen,⁴ J. Fast,⁴ J. W. Hinson,⁴ J. Lee,⁴ N. Menon,⁴ D. H. Miller,⁴ E. I. Shibata,⁴ I. P. J. Shipsey,⁴ Y. Kwon,^{5,†} A.L. Lyon,⁵ E. H. Thorndike,⁵ C. P. Jessop,⁶ K. Lingel,⁶ H. Marsiske,⁶ M. L. Perl,⁶ V. Savinov,⁶ D. Ugolini,⁶ X. Zhou,⁶ T. E. Coan,⁷ V. Fadeyev,⁷ I. Korolkov,⁷ Y. Maravin,⁷ I. Narsky,⁷ R. Stroynowski,⁷ J. Ye,⁷ T. Wlodek,⁷ M. Artuso,⁸ R. Ayad,⁸ E. Dambasuren,⁸ S. Kopp,⁸ G. Majumder,⁸ G. C. Moneti,⁸ R. Mountain,⁸ S. Schuh,⁸ T. Skwarnicki,⁸ S. Stone,⁸ A. Titov,⁸ G. Viehhauser,⁸ J.C. Wang,⁸ A. Wolf,⁸ J. Wu,⁸ S. E. Csorna,⁹ K. W. McLean,⁹ S. Marka,⁹ Z. Xu,⁹ R. Godang,¹⁰ K. Kinoshita,^{10,‡} I. C. Lai,¹⁰ P. Pomianowski,¹⁰ S. Schrenk,¹⁰ G. Bonvicini,¹¹ D. Cinabro,¹¹ R. Greene,¹¹ L. P. Perera,¹¹ G. J. Zhou,¹¹ S. Chan,¹² G. Eigen,¹² E. Lipeles,¹² M. Schmidtler,¹² A. Shapiro,¹² W. M. Sun,¹² J. Urheim,¹² A. J. Weinstein,¹² F. Würthwein,¹² D. E. Jaffe,¹³ G. Masek,¹³ H. P. Paar,¹³ E. M. Potter,¹³ S. Prell,¹³ V. Sharma,¹³ D. M. Asner,¹⁴ A. Eppich,¹⁴ J. Gronberg,¹⁴ T. S. Hill,¹⁴ D. J. Lange,¹⁴ R. J. Morrison,¹⁴ T. K. Nelson,¹⁴ J. D. Richman,¹⁴ R. A. Briere,¹⁵ B. H. Behrens,¹⁶ W. T. Ford,¹⁶ A. Gritsan,¹⁶ H. Krieg,¹⁶ J. Roy,¹⁶ J. G. Smith,¹⁶ J. P. Alexander,¹⁷ R. Baker,¹⁷ C. Bebek,¹⁷ B. E. Berger,¹⁷ K. Berkelman,¹⁷ F. Blanc,¹⁷ V. Boisvert,¹⁷ D. G. Cassel,¹⁷ M. Dickson,¹⁷ P. S. Drell,¹⁷ K. M. Ecklund,¹⁷ R. Ehrlich,¹⁷ A. D. Foland,¹⁷ P. Gaidarev,¹⁷ L. Gibbons,¹⁷ B. Gittelman,¹⁷ S. W. Gray,¹⁷ D. L. Hartill,¹⁷ B. K. Heltsley,¹⁷ P. I. Hopman,¹⁷ C. D. Jones,¹⁷ D. L. Kreinick,¹⁷ T. Lee,¹⁷ Y. Liu,¹⁷ T. O. Meyer,¹⁷ N. B. Mistry,¹⁷ C. R. Ng,¹⁷ E. Nordberg,¹⁷ J. R. Patterson,¹⁷ D. Peterson,¹⁷ D. Riley,¹⁷ J. G. Thayer,¹⁷ P. G. Thies,¹⁷ B. Valant-Spaight,¹⁷ A. Warburton,¹⁷ P. Avery,¹⁸ M. Lohner,¹⁸ C. Prescott,¹⁸ A. I. Rubiera,¹⁸ J. Yelton,¹⁸ J. Zheng,¹⁸ G. Brandenburg,¹⁹ A. Ershov,¹⁹ Y. S. Gao,¹⁹ D. Y.-J. Kim,¹⁹ R. Wilson,¹⁹ T. E. Browder,²⁰ Y. Li,²⁰ J. L. Rodriguez,²⁰ H. Yamamoto,²⁰ T. Bergfeld,²¹ B. I. Eisenstein,²¹ J. Ernst,²¹ G. E. Gladding,²¹ G. D. Gollin,²¹ R. M. Hans,²¹ E. Johnson,²¹ I. Karliner,²¹ M. A. Marsh,²¹ M. Palmer,²¹ C. Plager,²¹ C. Sedlack,²¹ M. Selen,²¹ J. J. Thaler,²¹ J. Williams,²¹ K. W. Edwards,²² R. Janicek,²³ P. M. Patel,²³ A. J. Sadoff,²⁴ R. Ammar,²⁵ P. Baringer,²⁵ A. Bean,²⁵ D. Besson,²⁵ R. Davis,²⁵ S. Kotov,²⁵ I. Kravchenko,²⁵ N. Kwak,²⁵ X. Zhao,²⁵ S. Anderson,²⁶ V. V. Frolov,²⁶ Y. Kubota,²⁶ S. J. Lee,²⁶ R. Mahapatra,²⁶ J. J. O'Neill,²⁶ R. Poling,²⁶ T. Riehle,²⁶ and A. Smith²⁶

¹State University of New York at Albany, Albany, New York 12222

²Ohio State University, Columbus, Ohio 43210

³University of Oklahoma, Norman, Oklahoma 73019

⁴Purdue University, West Lafayette, Indiana 47907

⁵University of Rochester, Rochester, New York 14627

*Permanent address: University of Texas - Pan American, Edinburg TX 78539.

†Permanent address: Yonsei University, Seoul 120-749, Korea.

‡Permanent address: University of Cincinnati, Cincinnati OH 45221

- ⁶Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309
- ⁷Southern Methodist University, Dallas, Texas 75275
- ⁸Syracuse University, Syracuse, New York 13244
- ⁹Vanderbilt University, Nashville, Tennessee 37235
- ¹⁰Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061
- ¹¹Wayne State University, Detroit, Michigan 48202
- ¹²California Institute of Technology, Pasadena, California 91125
- ¹³University of California, San Diego, La Jolla, California 92093
- ¹⁴University of California, Santa Barbara, California 93106
- ¹⁵Carnegie Mellon University, Pittsburgh, Pennsylvania 15213
- ¹⁶University of Colorado, Boulder, Colorado 80309-0390
- ¹⁷Cornell University, Ithaca, New York 14853
- ¹⁸University of Florida, Gainesville, Florida 32611
- ¹⁹Harvard University, Cambridge, Massachusetts 02138
- ²⁰University of Hawaii at Manoa, Honolulu, Hawaii 96822
- ²¹University of Illinois, Urbana-Champaign, Illinois 61801
- ²²Carleton University, Ottawa, Ontario, Canada K1S 5B6
and the Institute of Particle Physics, Canada
- ²³McGill University, Montréal, Québec, Canada H3A 2T8
and the Institute of Particle Physics, Canada
- ²⁴Ithaca College, Ithaca, New York 14850
- ²⁵University of Kansas, Lawrence, Kansas 66045
- ²⁶University of Minnesota, Minneapolis, Minnesota 55455

I. INTRODUCTION

Because flavor changing neutral currents are forbidden in the Standard Model, electroweak penguins such as $b \rightarrow s\gamma$ give a direct look at loop and box processes. The inclusive branching fraction for $b \rightarrow s\gamma$ is important for restricting physics beyond the Standard Model, and CLEO's 1995 published result [1] of $(2.32 \pm 0.57 \pm 0.35) \times 10^{-4}$ generated much theoretical interest [2]. The branching fraction result presented here is an improved measurement with 60% additional data and enhanced analysis techniques. The theoretical branching fraction prediction from the Standard Model has also improved, with a full next-to-leading-log calculation [3] of $(3.28 \pm 0.33) \times 10^{-4}$. We also introduce a new analysis: a search for CP asymmetry in $b \rightarrow s\gamma$.

II. THE CLEO DETECTOR

Data for these analyses were taken with the CLEO detector at the Cornell Electron Storage Ring (CESR) with center of mass energy at the $\Upsilon(4S)$ resonance (10.58 GeV). Additional data were taken 60 MeV below the $\Upsilon(4S)$ (off-resonance) for continuum background subtraction. The CLEO detector [4] measures charged particles over 95% of 4π steradians with a system of cylindrical drift chambers. The barrel and endcap CsI calorimeters cover 98% of 4π , and the energy resolution for photons near 2.5 GeV in the central angular region ($|\cos\theta_\gamma| < 0.7$) is 2%. In 1995, a silicon vertex detector replaced the inner most tracking chamber, though it is not used in the analyses described here.

III. BRANCHING FRACTION ANALYSIS

This analysis is discussed in detail elsewhere [5] and is briefly covered here. The signal for $b \rightarrow s\gamma$ is a photon from B meson decay with $2.1 < E_\gamma < 2.7$ GeV (the published analysis [1] used $2.2 < E_\gamma < 2.7$ GeV). The Fermi momentum of the b quark in the B meson and the momentum of the B meson in the lab frame (B mesons are produced from $\Upsilon(4S)$ decays with approximately 300 MeV/ c of momentum at CESR) Doppler broadens the photon line. Calculations using a spectator model [6] indicate that 85-94% of the signal lies in the photon energy range.

To obtain a photon energy spectrum, we select hadronic events with a high energy calorimeter cluster in the central region ($|\cos\theta_\gamma| < 0.7$). We reject a cluster if, when paired with another calorimeter cluster in the event, it forms a combined $\gamma\gamma$ mass consistent with a π^0 or η . We also require the cluster shape be consistent with that of a single high energy photon.

A naïve approach to this analysis would be to measure the photon spectrum in on-resonance data and use the off-resonance data to subtract the continuum background. But the backgrounds from continuum with initial state radiation ($e^+e^- \rightarrow q\bar{q}\gamma$) (ISR) and from continuum processes ($e^+e^- \rightarrow q\bar{q}$) involving a high energy π^0 , η , or ω where one of the daughter photons is not detected dominate to such an extent that extraction of a signal is impossible. We therefore suppress the continuum background with two separate methods and subtract what remains with off-resonance data.

The first continuum suppression scheme [7] involves exploiting differences in the event shapes between signal ($B\bar{B}$) and continuum. While continuum events appear jetty and continuum with ISR appear jetty when transformed to the rest frame of the system after the initial state radiation,

the signal events are spherical due to the other B meson decaying in the event. We characterize the event shape with eight variables[6]: the Fox-Wolfram second moment (R_2), ratio of the sum of transverse momenta of particles more than 45° away from the high energy photon axis to the total scalar sum of particle momenta in the event excluding the momentum of the high energy photon (S_\perp), R_2 transformed to the rest frame of the system after ISR (assuming the high energy photon in the event is initial state radiation) (R'_2), $\cos\theta$ between the photon and the thrust axis of the rest of the event transformed into the rest frame of the system after ISR ($\cos\theta'$), and the energies of particles found in 20° and 30° cones parallel and anti-parallel to the high energy photon direction. Since no one variable provides enough discriminating power, we combine them with a neural network into a single variable r . The distribution of r tends towards -1 for continuum and ISR events and $+1$ for signal $b \rightarrow s\gamma$ events. This shape method is approximately 30% efficient for signal.

The other method of suppressing continuum and ISR events is a pseudo-reconstruction of the parent B meson involved in the $b \rightarrow s\gamma$ decay. We look for a charged or neutral kaon with up to four pions, one of which may be a π^0 , which when combined with the high energy photon are consistent with being the decay products of a B meson. We try all combinations of kaons and pions given the above restrictions and choose the one that minimizes a χ^2 that includes,

$$\chi_B^2 = \left(\frac{M - M_B}{\sigma_M} \right)^2 + \left(\frac{E - E_{\text{beam}}}{\sigma_E} \right)^2 \quad (1)$$

as well as contributions from particle identification such as dE/dx and π^0 and K_s^0 mass deviations. In Eq. (1), $M = \sqrt{E_{\text{beam}}^2 - P^2}$ is the beam-constrained mass, where P is the vector sum of particle momenta comprising the B meson candidate, M_B is the B mass ($5.28 \text{ GeV}/c^2$), E is the energy of the candidate B , and σ_M and σ_E are the resolutions on the beam constrained mass and the measured B energy respectively. Events that have a minimum $\chi_B^2 < 20$ are considered pseudo-reconstructed (“pseudo” is used because this method is only for continuum suppression, and we are not concerned that the parent B reconstruction be exactly correct). If the minimum $\chi_B^2 > 20$ or not one combination of kaons and pions was found in the event, then the shape analysis described above is used.

For events that are pseudo-reconstructed, we calculate the angle between the thrust axis of the inferred signal B meson and the thrust axis of the rest of the event ($\cos\theta_{tt}$). Events that are in actuality continuum typically have $|\cos\theta_{tt}|$ peaked at 1, while signal events have a flat distribution. To get the most discriminating power, we combine χ_B^2 and $\cos\theta_{tt}$ from the pseudo-reconstruction and r from the shape neural network into a new variable r_c using another neural network. As with r , the distribution of r_c tends towards $+1$ for signal events and -1 for continuum. While this combined method is 10% efficient for signal, it reduces the continuum background by an additional factor of four over the shape method alone.

We then weight each event according to its value of r_c if it is pseudo-reconstructed or r if not. The weighting scheme [5] is designed so that the weighted yield is equivalent to the event yield in the absence of continuum backgrounds. This procedure gives the smallest statistical uncertainty on this background subtracted yield. Note that there are non-continuum backgrounds discussed below, but these are small compared to continuum. The continuum background remaining after the suppression efforts is determined from off-resonance data.

$B\bar{B}$ backgrounds are dominated by $b \rightarrow c$ (e.g. $B \rightarrow X\pi^0$ and $B \rightarrow X\eta$) though $b \rightarrow u$ and $b \rightarrow sg$ may have a small contribution. We estimate the $B\bar{B}$ backgrounds with Monte Carlo (MC),




FIG. 1. Photon energy spectra. (a) On-resonance data (points), luminosity scaled off-resonance data (double hatched), and $B\bar{B}$ background (single hatched). (b) Background-subtracted data (points) and Monte Carlo prediction of the shape of the $b \rightarrow s\gamma$ signal from a spectator model calculation ($\langle m_b \rangle = 4.88 \text{ GeV}/c^2$) and Fermi momentum set at $250 \text{ MeV}/c$. Only statistical uncertainties are shown.

but correct the π^0 and η momentum spectra for any differences between data and MC. We obtain the momenta spectra from data by treating π^0 's and η 's as if they were photons and follow the same analysis procedure using weights described above. Thus, the MC is used only for the π^0 and η veto efficiencies and for any other small $B\bar{B}$ backgrounds not originating from missing a γ daughter from a π^0 or η .

The weighted yields from 3.1 fb^{-1} of on-resonance data plotted with respect to photon energy are shown in Fig. 1, with all of the backgrounds shown in (a) and the background subtracted weighted yield, which is the signal, shown in (b). 1.6 fb^{-1} of off-resonance data were used for the continuum subtraction. The expected spectrum for $b \rightarrow s\gamma$ from a spectator model is also shown in (b). In the region of interest ($2.1 < E_\gamma < 2.7 \text{ GeV}$) we measure 500.5 ± 7.4 on-resonance weighted events, 382.3 ± 7.0 scaled off-resonance weighted events, and estimate the $B\bar{B}$ background to be 20.6 weighted events (all uncertainties statistical). Applying the π^0 and η corrections to the estimate (1.9 ± 0.5 and 3.6 ± 1.1 weighted events respectively), we derive the weighted, background-subtracted yield of $92.2 \pm 10.3 \pm 6.5$, where the first uncertainty is statistical and the second is systematic.

For a typical sample of MC $b \rightarrow s\gamma$ events, approximately 50% pass all analysis requirements and of these, 40% are pseudo-reconstructed ($\chi_B^2 < 20$). But since we use weighted events for the signal yield, we need a weighted efficiency; the sum of weights passing all requirements. Because of our definition of weights, the weighted efficiency tends to be smaller than the efficiencies stated above. To model the signal, we use the spectator model of Ali and Greub [6], which includes gluon bremsstrahlung and other higher order radiative effects. Details of MC production and parameters used are given in ref [1]. The weighed efficiency is $(4.43 \pm 0.29 \pm 0.22 \pm 0.03 \pm 0.31) \times 10^{-2}$, where the first uncertainty is due to spectator model inputs, the second due to recoil system hadronization, the third due to uncertainty in the production ratio of neutral and charged B meson pairs, and the last to detector modeling.

The weighted efficiency is combined with the background subtracted weighted yield to give the inclusive branching fraction of $b \rightarrow s\gamma$ as $(3.15 \pm 0.35 \pm 0.32 \pm 0.26) \times 10^{-4}$, where the uncertainties are statistical, systematic and for model dependence respectively. Our results are in agreement with the Standard Model prediction. Conservatively allowing for the systematic uncertainty, we find that the branching fraction must be between 2.0×10^{-4} and 4.5×10^{-4} (each limit at 95% CL).

IV. CP ASYMMETRY

The SM predicts no CP asymmetry in $b \rightarrow s\gamma$ decays, but some recent theoretical work [8] suggests that non-SM physics may significantly contribute to a CP asymmetry. Furthermore, if new physics has a weak phase difference near 90° with respect to the SM and the strong phase is non-zero, the new physics would only slightly alter the $b \rightarrow s\gamma$ inclusive branching ratio, but could produce a large effect in the asymmetry. Since we can reuse much of the machinery from the branching fraction measurement, a search for CP asymmetry is an obvious extension to the $b \rightarrow s\gamma$ analysis.

The pseudo-reconstruction procedure described above is used to tag the b quark flavor (b or \bar{b}), because it determines the particles decayed from the signal B meson. But since the pseudo-reconstruction was designed for continuum suppression, we must determine how often it reconstructs the b flavor correctly. If a signal B meson decays to a neutral kaon and overall neutral pions (*e.g.* $B^0 \rightarrow K_s^0 \pi^+ \pi^-$), then we cannot determine the b flavor from pseudo-reconstruction. But for the majority of cases where the b flavor is determinable, it can be deduced by the charge of the kaon, or the sum pion charge if the kaon is neutral. From MC, we estimate the probability of reconstructing the incorrect b flavor when the b flavor is determinable to be $8.3 \pm 1.6\%$. The quoted uncertainty includes a combination of MC statistics and uncertainties derived from varying spectator model inputs to the MC and varying the ratio of charged to neutral B meson decays. We also take into account the probability of tagging the b flavor when it should be indeterminable and vice-versa, though these effects are small and will not be discussed further here.

The asymmetry we measure is $(N_1 - N_2)/(N_1 + N_2)$, where N_1 is the weighted yield tagged as b quarks from pseudo-reconstruction and determinable as such, and N_2 is similarly the weighted yield tagged as \bar{b} . From the misreconstruction rates described above, we apply a multiplicative correction factor of 1.22 ± 0.04 to the asymmetry. In our misreconstruction determination, we assumed that the rate of mistagging a b as a \bar{b} was the same as mistagging a \bar{b} as a b . Initial studies lead us to apply a conservative additive systematic error on the asymmetry of 5% to account for unequal mistagging rates, though we observe no measurable asymmetry in the off-resonance data and other control samples.

Using the weighted yield from pseudo-reconstructed events with photons within ($2.2 < E_\gamma < 2.7$ GeV), we observe 36.8 ± 5.2 weights reconstructed as b quarks and 28.4 ± 5.2 weights reconstructed as \bar{b} quarks (uncertainties are statistical only), leading to a raw asymmetry measurement of 0.13 ± 0.11 . Applying the correction factors, we obtain our preliminary CP asymmetry measurement in $b \rightarrow s\gamma$ decays: $(0.16 \pm 0.14 \pm 0.05) \times (1.0 \pm 0.04)$, where the first number is the central value, the second is the statistical uncertainty, the third is the additive systematic described above, and the multiplicative factor is from the uncertainty on the mistagging rate correction. Within the uncertainties, we measure no CP asymmetry. From this result, we derive 90% CL limits on the CP asymmetry (\mathcal{A}) of $-0.09 < \mathcal{A} < 0.42$.

V. SUMMARY AND OUTLOOK

We presented CLEO's preliminary results on the $b \rightarrow s\gamma$ branching ratio and the CP asymmetry in $b \rightarrow s\gamma$ decays with approximately 3.1 fb^{-1} of on-resonance data. CLEO has since accumulated a total of 10 fb^{-1} of on-resonance data. Work is underway to determine the $b \rightarrow s\gamma$ inclusive branching

fraction and CP asymmetry with all of the data. We are also exploring extending both $b \rightarrow s\gamma$ analyses by tagging the b flavor with a lepton from the other B meson in the event it undergoes semileptonic decay. This procedure offers further continuum suppression and may improve the asymmetry measurement by allowing the use of events where the b flavor is undeterminable by pseudo-reconstruction.

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity and running conditions. We also thank M. Neubert for bringing the theoretical interest in $b \rightarrow s\gamma$ CP asymmetry to our attention. J.R. Patterson and I.P.J. Shipsey thank the NYI program of the NSF, M. Selen thanks the PFF program of the NSF, M. Selen and H. Yamamoto thank the OJI program of the DOE, J.R. Patterson, K. Honscheid, M. Selen and V. Sharma thank the A.P. Sloan Foundation, M. Selen and V. Sharma thank the Research Corporation, F. Blanc thanks the Swiss National Science Foundation, and H. Schwarthoff and E. von Toerne thank the Alexander von Humboldt Stiftung for support. This work was supported by the National Science Foundation, the U.S. Department of Energy, and the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

- [1] M.S. Alam *et al.* (CLEO), Phys. Rev. Lett. **74**, 2885 (1995).
- [2] For example, F. Borzumati and C. Greub, Phys. Rev. D **58**, 074004 (1998); W. de Boer *et al.*, Phys. Lett. B **438**, 281 (1998); H. Baer *et al.*, Phys. Rev. D **58**, 15007 (1998). For implications on non-standard model physics see J. Hewett, SLAC-PUB-6521 (1994).
- [3] K. Chetyrkin *et al.*, Phys. Lett. B **400**, 206 (1997); Phys. Lett. B **425**, 414 (1998); A. Kagan and M. Neubert, Eur. Phys. J. **C7**, 5 (1999).
- [4] Y. Kubota *et al.* (CLEO), Nucl. Instrum. Meth. A **320**, 66 (1992).
- [5] CLEO Conference report 98-17 (ICHEP98 1011).
<http://www.lns.cornell.edu/public/CONF/1998/CONF98-17/>
- [6] A. Ali and C. Greub, Phys. Lett. B **259**, 182 (1991).
- [7] J.A. Ernst, Ph.D. thesis, Univ. of Rochester (1995).
- [8] A.L. Kagan and M. Neubert, Phys. Rev. D **58**, 094012 (1998) and M. Aoki, G. Cho and N. Oshimo, Phys. Rev. D **60**, 035004 (1999).