



$\gamma\delta$ T Cells Are Required for Pulmonary IL-17A Expression after Ozone Exposure in Mice: Role of TNF α

Joel A. Mathews*, Alison S. Williams, Jeffrey D. Brand, Allison P. Wurmbrand, Lucas Chen, Fernanda M.C. Ninin, Huiqing Si, David I. Kasahara, Stephanie A. Shore

Molecular and Integrative Physiological Sciences Program, Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts, United States of America

Abstract

Ozone is an air pollutant that causes pulmonary symptoms. In mice, ozone exposure causes pulmonary injury and increases bronchoalveolar lavage macrophages and neutrophils. We have shown that IL-17A is important in the recruitment of neutrophils after subacute ozone exposure (0.3 ppm for 24–72 h). We hypothesized that $\gamma\delta$ T cells are the main producers of IL-17A after subacute ozone. To explore this hypothesis we exposed wildtype mice and mice deficient in $\gamma\delta$ T cells (TCR $\delta^{-/-}$) to ozone or room air. Ozone-induced increases in BAL macrophages and neutrophils were attenuated in TCR $\delta^{-/-}$ mice. Ozone increased the number of $\gamma\delta$ T cells in the lungs and increased pulmonary *Il17a* mRNA expression and the number of IL-17A⁺ CD45⁺ cells in the lungs and these effects were abolished in TCR $\delta^{-/-}$ mice. Ozone-induced increases in factors downstream of IL-17A signaling, including G-CSF, IL-6, IP-10 and KC were also decreased in TCR $\delta^{-/-}$ versus wildtype mice. Neutralization of IL-17A during ozone exposure in wildtype mice mimicked the effects of $\gamma\delta$ T cell deficiency. TNFR2 deficiency and etanercept, a TNF α antagonist, also reduced ozone-induced increases in *Il17a* mRNA, IL-17A⁺ CD45⁺ cells and BAL G-CSF as well as BAL neutrophils. TNFR2 deficient mice also had decreased ozone-induced increases in Ccl20, a chemoattractant for IL-17A⁺ $\gamma\delta$ T cells. *Il17a* mRNA and IL-17A⁺ $\gamma\delta$ T cells were also lower in obese *Cpe^{fat}* versus lean WT mice exposed to subacute ozone, consistent with the reduced neutrophil recruitment observed in the obese mice. Taken together, our data indicate that pulmonary inflammation induced by subacute ozone requires $\gamma\delta$ T cells and TNF α -dependent recruitment of IL-17A⁺ $\gamma\delta$ T cells to the lung.

Citation: Mathews JA, Williams AS, Brand JD, Wurmbrand AP, Chen L, et al. (2014) $\gamma\delta$ T Cells Are Required for Pulmonary IL-17A Expression after Ozone Exposure in Mice: Role of TNF α . PLoS ONE 9(5): e97707. doi:10.1371/journal.pone.0097707

Editor: Shama Ahmad, University of Colorado, Denver, United States of America

Received: January 7, 2014; **Accepted:** April 22, 2014; **Published:** May 13, 2014

Copyright: © 2014 Mathews et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by: F32ES02256, NIH-HL007118, NIEHS: ES-013307 and ES-000002. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: jmathews@hsph.harvard.edu

Introduction

$\gamma\delta$ T cells are a key component of the innate immune response, especially at mucosal surfaces. These cells are found throughout the lung, particularly in the subepithelial region, where they may regulate other immune cells including macrophages and dendritic cells [1]. $\gamma\delta$ T cells are an important source of IL-17A, a key cytokine involved in neutrophilic inflammation [2]. In mice, the number of pulmonary $\gamma\delta$ T cells increases following infection with certain bacteria [3]. Mice deficient in $\gamma\delta$ T cells (TCR $\delta^{-/-}$ mice) have attenuated pulmonary clearance of these bacteria, likely as a result of loss of IL-17A production by $\gamma\delta$ T cells and consequent reduced neutrophil recruitment [4]. The number of $\gamma\delta$ T cells in the lung also increases under conditions associated with oxidative stress, including smoking, bleomycin instillation, and allergen challenge [5–8]. Moreover, the pulmonary inflammation induced by such agents requires $\gamma\delta$ T cells.

Inhalation of ozone (O₃), a common air pollutant, has a significant impact on human health. O₃ causes respiratory symptoms and reductions in lung function [9–13]. O₃ also increases the risk of respiratory infections and is a trigger for asthma [14–16]. Exposure to O₃ induces oxidative stress in the lung, damages lung epithelial cells, and causes the release of

numerous cytokines and chemokines that recruit neutrophils and macrophages to the lung [9,17]. We have reported increased *Il17a* mRNA expression and increased numbers of IL-17A⁺ $\gamma\delta$ T cells in the lungs after subacute O₃ exposure (0.3 ppm O₃ for 24–72 h) [18]. Hence, we tested the hypothesis that $\gamma\delta$ T cells, via their ability to produce IL-17A, are involved in orchestrating the inflammatory response to subacute O₃ exposure. We examined IL-17A expression in WT and TCR $\delta^{-/-}$ mice after exposure to air or to O₃ (0.3 ppm for 24–72 h). We also examined the effect of IL-17A neutralizing antibodies on O₃-induced inflammation. Our results indicate an important role for IL-17A⁺ $\gamma\delta$ T cells in the inflammatory cell recruitment induced by subacute O₃ exposure.

TNF α a pleiotropic pro-inflammatory cytokine, enhances the recruitment of neutrophils to the lungs in response to a variety of noxious stimuli, including LPS [19], cigarette smoke [20], and enterobacteria [21]. TNF α is also required for neutrophil recruitment after subacute O₃ exposure [22,23]. However, TNF α does not have direct chemoattractant activity for neutrophils [24]. Instead, TNF α recruits neutrophils in part by inducing expression of other cytokines and chemokines [24,25]. In several pathological states, TNF α induces the expression of IL-17A [26,27]. Hence, we hypothesized that TNF α contributes to neutrophil recruitment following subacute O₃ exposure by promoting recruitment to or

activation of IL-17A⁺ $\gamma\delta$ T cells in the lungs. We used two methods to test this hypothesis. First, we assessed the effect of O₃ exposure on pulmonary *Il17a* expression and recruitment of IL-17A⁺ $\gamma\delta$ T cells in WT mice and in mice deficient in TNFR2 (TNFR2^{-/-} mice). Others have established that either TNFR1 or TNFR2 deficiency reduces the inflammatory response to subacute O₃, and there is no further impact of combined TNFR1/TNFR2 deficiency [22]. Second, we examined the impact of the TNF α antagonist, etanercept, on *Il17a* expression. Our data suggest that TNF α is required for the recruitment of IL-17A⁺ $\gamma\delta$ T cells to the lung after subacute O₃ exposure.

Approximately one third of the US population is obese and another third is overweight, but our understanding of how obesity impacts pulmonary responses to O₃ is still rudimentary. Such an understanding may have broad reaching implications since oxidative stress also contributes to responses to a variety of other noxious stimuli [5–8], many of which are affected by obesity [28,29]. In mice, the impact of obesity on responses to O₃ depends on the nature of the exposure: the pulmonary inflammation induced by acute O₃ exposure (2 ppm for 3 h) is augmented in all types of obese mice examined to date [30–33], whereas the pulmonary inflammation induced by subacute O₃ exposure (0.3 ppm for 24–72 h) is reduced [34]. Given our findings of the requirement for TNF α -recruitment of IL-17A producing $\gamma\delta$ T cells in the induction of pulmonary inflammation after subacute O₃, we sought to determine if changes in the activation of $\gamma\delta$ T cells might explain the reduced responses to subacute O₃ we observed in obese *Cpe*(carboxypeptidase E)^{fat} mice. Data described below indicate that the reduced O₃-induced neutrophil recruitment observed in obese mice is likely the result of reduced *Il23* expression leading to reduced IL-17A⁺ $\gamma\delta$ T cells. Given the importance of IL-17⁺ $\gamma\delta$ T cells for responses to viral and bacterial pathogens (see above), these observations might explain the altered response of the obese to bacteria and virus (see review by Peter Mancuso [35]).

Methods

Animals

This study was approved by the Harvard Medical Area Standing Committee on Animals. Male age-matched WT and TCR δ ^{-/-} mice were either purchased from The Jackson Laboratory (Bar Harbor, ME) and acclimated for 4 weeks, or bred in house. *Cpe*^{fat} mice are deficient in carboxypeptidase E, an enzyme involved in processing neuropeptides involved in eating behaviors [36]. The breeding strategy used to generate *Cpe*^{fat}/TNFR2^{-/-} mice from *Cpe*^{fat} and TNFR2^{-/-} mice (also originally purchased from The Jackson Laboratory) was previously described [37]. All mice were on a C57BL/6J background, fed a standard mouse chow diet, and were 10–13 weeks old at the time of study.

Protocol

For comparisons of WT and TCR δ ^{-/-} mice, mice were exposed to O₃ (0.3 ppm) or to air, for 24–72 hours, as previously described [18]. Mice were exposed in normal cages without the microisolator top, but with free access to water and food throughout exposure. Mice were checked daily. At least two mice were placed in each cage to limit stress. After exposure, mice were euthanized with an overdose of sodium pentobarbital. The trachea was cannulated and bronchoalveolar lavage (BAL) was performed. After BAL, the lungs were flushed of blood by injecting 10 ml of cold PBS through the right ventricle, after creating a large excision in the left ventricle. One lung was excised and used for flow cytometry. The other was excised and placed in RNAlater

(Qiagen, Germantown, MD) for preparation of RNA for real time PCR. In another cohort, WT mice were injected i.p. with 100 μ g of anti-IL-17A neutralizing monoclonal antibody (Ab) (Rat IgG2A, clone 50104, MAB421; R&D Systems, Minneapolis, MN) or isotype control Ab (clone 54447, MAB006; R&D Systems) in 100 μ l of sterile saline 24 hours before O₃ exposure. Mice were exposed to O₃ for 72 hours, euthanized, and tissues were harvested as described above. In a separate series of experiments, WT, TNFR2^{-/-}, *Cpe*^{fat}, and *Cpe*^{fat}/TNFR2^{-/-} mice were exposed to room air or O₃ (0.3 ppm) for 48 h followed by BAL and tissue harvest. In other experiments, WT and *Cpe*^{fat} mice were treated twice (48 h and 1 h prior to O₃ exposure) with the TNF α blocking drug, etanercept (30 mg/kg s.c.) (Immunex, Thousand Oaks, CA), or vehicle. A similar etanercept dosing regimen has been shown to be effective in inhibiting TNF α in mice over the time course of O₃ exposures we used (48 h) [38,39].

Bronchoalveolar Lavage

BAL was performed and cells counted as previously described [18]. BAL supernatant was stored at -80°C until assayed. BAL KC, IL-6, MCP-1, IP-10 and G-CSF were measured by ELISA (R&D Systems). In mice treated with anti-IL-17A, BAL cytokines and chemokines were measured by multiplex assay (Eve Technologies, Calgary, Alberta). Total BAL protein was measured by Bradford assay (Bio-Rad, Hercules, CA).

Flow Cytometry

Left lungs were harvested and placed on ice in RPMI 1640 media containing 2% FBS and HEPES. Lungs were digested and prepared for flow cytometry as previously described [18]. Cells were stained using the following antibodies: Alexa Fluor 647 anti-IL-17A (clone: TC11-18H10.1), PE anti-TCR δ (clone: GL3), PE-cy7 anti-CD45 (clone: 30-F11), and APC-cy7 anti-CD3 (clone: 17A2) (all antibodies from Biolegend). Isotype control antibodies were used to set all gates. Cells were visualized using a Canto II (BD Biosciences) and the data was analyzed using Flowjo (Tree Star; Ashland, OR).

To determine if TNF α impacted IL-12R β 1 expression on lung $\gamma\delta$ T cells, lungs from WT mice were digested as above and then cultured in complete RPMI media (RPMI 1640 (Corning, Tewksbury, MA), 10% FBS (Life Technologies), 2 Mm L-glutamine (Life Technologies), 100 units/ml Pen/Strep (Lonza, Hopkinton, MA) and 20 Mm Hepes (Thermo Scientific, Tewksbury, MA)). Cells were plated at a concentration of 10⁶ cells/ml in 24 well plates with or without 100 ng/ml of recombinant murine TNF α (R&D Systems) [40]. Cells were harvested after 24 h, washed with PBS, and stained using the following antibodies: anti-CD16/32 (True Stain biolegend), Strep-APC (Biolegend), PE anti-CD212 (IL-12R β 1) (BD Biosciences), Biotin anti-TCR δ (clone: GL3, biolegend), PE-cy7 anti-CD45 (clone: 30-F11) and analyzed by flow cytometry as described above.

Real-time PCR

RNA was extracted from lung tissue and prepared for qPCR using the SYBR method as previously described [18]. All expression values were normalized to 36B4 expression using the $\Delta\Delta\text{Ct}$ method. The primers for *Il17a* and *36B4* were previously described [37]. Primers for *Ccl20*, *Il23* (p19) and *Il12R β 1* are described in Table 1. For each set of primers, melt curve analysis yielded a single peak. *Il12R β 1* expression was measured at baseline in order to tease apart the effects of genotype (deficiency of TNF α signaling versus sufficient signaling) versus O₃ exposure.

Table 1. Primers used for real time PCR.

<i>Il23p19</i>	F: CCC ATG GAG CAA CTT CAC AC R: GCT GCC ACT GCT GAC TAG AAC
<i>Ccl20</i>	F: AAG ACA GAT GGC CGA TGA AG R: AGG TTC ACA GCC CTT TTC AC
<i>Il12Rb1</i>	F: GTG CTC GCC AAA ACT CGT TT R: GGA TGT CAT GTT GCC TCC CA

doi:10.1371/journal.pone.0097707.t001

Statistical Analysis

Data were analyzed by factorial ANOVA using STATISTICA software (Statistica, StatSoft; Tulsa, OK) with mouse genotype and exposure as main effects. Fisher's least significant difference test was used as a post-hoc test. BAL cells and flow cytometry data were normalized by log transformed prior to analysis. A p value < 0.05 was considered significant.

Results

O₃-induced Inflammation is Reduced in TCR $\delta^{-/-}$ Mice

In WT mice, O₃ exposure caused a time-dependent increase in BAL neutrophils, macrophages, and protein (a measure of O₃-induced lung injury [41]) (Fig. 1A–C), consistent with previous reports by ourselves and others [18,22,23,41,42]. Increases in BAL inflammatory cells were significantly reduced in TCR $\delta^{-/-}$ versus WT mice after 48 (neutrophils) and 72 (neutrophils and macrophages) hours of exposure (Fig. 1A,B). BAL protein was also reduced in TCR $\delta^{-/-}$ versus WT mice after 72 hours exposure, but not at earlier times (Fig. 1C).

Several cytokines, including KC, IL-6, IP-10 (CXCL10), G-CSF, MCP-1 and IL-17A [17,18,22,23,41–44], can contribute to inflammatory cell recruitment to the lungs after O₃ exposure. BAL IL-17A expression was below the limits of detection of ELISA. Consequently, we used q-RT-PCR to measure IL-17A. *Il17a* mRNA abundance increased after 24, 48 and 72 hours of O₃ in WT but not TCR $\delta^{-/-}$ mice (Fig. 1D). O₃-induced increases in BAL concentrations of BAL G-CSF, IL-6, KC and IP-10 were each reduced in TCR $\delta^{-/-}$ versus WT mice at 72 hours of exposure (Fig. 1E–H). For G-CSF and IP-10, there was a similar trend at 24 and 48 hours (Fig. 1E,G). $\gamma\delta$ T cell deficiency had no effect on O₃-induced changes in BAL MCP-1, although MCP-1 trended lower in TCR $\delta^{-/-}$ versus WT mice at 72 hours.

IL-17A⁺ $\gamma\delta$ T Cells are Increased by O₃ Exposure

Flow cytometry indicated that the number of IL-17A⁺ CD45⁺ cells was significantly increased by O₃ in WT mice. This effect was ablated in TCR $\delta^{-/-}$ mice (Fig. 2A). Further analysis indicated that in WT mice, the numbers of IL-17A⁺ $\gamma\delta$ T cells as well as the total number of $\gamma\delta$ T cells were increased by O₃ (Fig. 2B, C), as reported previously using a similar gating strategy [18].

Effect of Anti-IL-17A Treatment

Compared to isotype control, anti-IL-17A treatment of WT mice caused a significant reduction in BAL neutrophils and macrophages (Fig. 3A). Anti-IL-17A treatment also significantly decreased BAL protein (Fig. 3B) and BAL G-CSF (Fig. 3C). Given this key role for IL-17A, these data indicate that the decreased inflammatory response observed in the TCR $\delta^{-/-}$ mice was likely due to the lack of *Il17a* expression (Fig. 1D) and demonstrate that G-CSF likely contributes to the effect of IL-17A on neutrophil recruitment.

Role of TNF α

BAL neutrophils were significantly lower in TNFR2 $^{-/-}$ versus WT mice exposed to O₃ for 48 h (Fig. 4A), consistent with the results of Cho et al [22]. Similar results were obtained in WT mice treated with etanercept versus vehicle (Fig. 4D). O₃ exposure caused a significant increase in pulmonary *Il17a* expression in WT mice (Fig. 4B), consistent with results described above (Fig. 1D). However in TNFR2 $^{-/-}$ mice, no such increase in *Il17a* mRNA abundance was observed (Fig. 4B). Similar results were obtained in mice treated with etanercept (Fig. 4E). Flow cytometry also indicated a decrease in IL-17A⁺CD45⁺ cells in O₃-exposed TNFR2 $^{-/-}$ versus WT mice (Fig. 5A). This change was due to decreased numbers of IL-17A⁺ $\gamma\delta$ T cells (Fig. 5B). BAL G-CSF was also significantly lower in O₃-exposed TNFR2 $^{-/-}$ versus WT mice (Fig. 4C) and in etanercept treated versus vehicle treated WT mice (Fig. 4F).

The requirement of IL-23 and IL-6 for IL-17A expression in $\gamma\delta$ T cells [45,46], suggested that reductions in IL-17A⁺ $\gamma\delta$ T cells in TNFR2 $^{-/-}$ mice might be the result of loss of TNF α -induced expression of IL-23 or IL-6. O₃ increased BAL IL-6 in WT mice (Fig. 1F) and O₃ also increased pulmonary *Il23* (p19) mRNA abundance (Fig. 6B), but neither IL-6 nor IL-23 were affected by TNFR2 deficiency or etanercept treatment (Fig. 6A, C). In contrast, TNFR2 $^{-/-}$ mice had reduced expression at baseline of *Il12Rb1* (Fig. 6H), a component of the IL-23 receptor. A similar trend was observed in etanercept treated mice (data not shown). O₃ exposure had no effect on *Il12Rb1* (data not shown). Expression of the other component of the IL-23 receptor, *Il23R*, was not affected by TNFR2 deficiency (data not shown). To determine if TNF α was having direct effects on *Il12Rb1* expression on $\gamma\delta$ T cells, we isolated total lung cells from WT mice, stimulated them overnight with TNF α and examined IL-12R β 1 expression on $\gamma\delta$ T cells by flow cytometry (Fig. 6I,J). TNF α had no effect on the levels of IL-12R β 1 on $\gamma\delta$ T cells as measured by MFI and did not affect the percentage of $\gamma\delta$ T cells expressing IL-12R β 1, suggesting that other cells in the lung accounted for differences in *Il12Rb1* mRNA expression.

We also considered the possibility that TNF α might impact the recruitment of $\gamma\delta$ T cells to the lung. In WT mice, O₃ exposure caused an increase in pulmonary mRNA expression of *Ccl20* (Fig. 6E), a chemoattractant for IL-17A⁺ cells [47,48], whereas no such increase was observed in mice treated with etanercept (Fig. 6F), suggesting that the role of TNF α is in the CCL20 dependent recruitment of IL-17⁺ $\gamma\delta$ T cells to the lungs. Similarly, there was a trend towards reduced *Ccl20* mRNA abundance in O₃-exposed TNFR2 $^{-/-}$ versus WT mice (Fig. 6G), although the effect did not reach statistical significance.

Response to O₃ in Obese Mice

Cpe^{fat} mice, regardless of their TNFR2 genotype or exposure, weighed almost twice as much as controls (data not shown). BAL neutrophils were significantly lower in *Cpe^{fat}* versus WT mice exposed to O₃ (Fig. 4A,D), consistent with our previous observations using this exposure regimen [34]. In contrast to the




Figure 1. Effect of $\gamma\delta$ T cell deficiency on pulmonary inflammation and injury. (A–C) BAL neutrophils, macrophages, and protein; (D) pulmonary *Il17a* mRNA expression; (E–I) BAL G-CSF, IL-6, IP-10, KC, and MCP-1. Results are mean \pm SEM of 4–11 mice per group. * $p < 0.05$ versus genotype-matched air-exposed mice. # $p < 0.05$ versus WT mice with the same exposure. doi:10.1371/journal.pone.0097707.g001

substantial reduction in BAL neutrophils observed in TNFR2 $^{-/-}$ versus WT mice, TNFR2 deficiency had no significant effect on BAL neutrophils in O₃-exposed *Cpe^{fat}* mice (Fig. 4A). Similar results were obtained in etanercept treated WT mice (Fig. 4D). *Cpe* genotype had no impact on the number of BAL or lung macrophages (data not shown).

Il17a expression was significantly lower in O₃ exposed *Cpe^{fat}* versus WT mice (Fig. 4B,E). The number of IL-17A⁺ CD45⁺ cells was also significantly lower in O₃-exposed *Cpe^{fat}* than WT mice (Fig. 5A). The total number of $\gamma\delta$ T cells and the number of IL-17A⁺ $\gamma\delta$ T cells was also reduced in the lungs of *Cpe^{fat}* versus WT mice (Fig. 5B,C). O₃-induced increases in BAL G-CSF were also




Figure 2. Effect of O₃ exposure on IL-17A positive lung cells assessed by flow cytometry. (A) lung IL-17A⁺CD45⁺; (B) lung IL-17A⁺ $\gamma\delta$ T cells; (C) total lung $\gamma\delta$ T cells. Results are mean \pm SEM for 3–6 air-exposed and 4–11 O₃-exposed mice. * $p < 0.05$ versus genotype-matched air-exposed mice. # $p < 0.05$ versus WT mice with same exposure. doi:10.1371/journal.pone.0097707.g002




Figure 3. Effect of anti-IL-17A on O₃-induced pulmonary inflammation and injury. WT mice were injected with anti-IL-17A or isotype 24 h prior to O₃ (0.3 ppm O₃ for 72 h). (A) BAL macrophages and neutrophils; (B) BAL protein; (C) BAL cytokines determined by multiplex assay. Results are mean \pm SEM of 5–7 mice per group. #p < 0.05 versus isotype control. doi:10.1371/journal.pone.0097707.g003

lower in *Cpe^{fat}* versus WT mice (Fig. 4C, E) consistent with the reductions in IL-17A expression. Both BAL IL-6 and pulmonary *Il23* mRNA expression were lower in *Cpe^{fat}* versus WT mice (Fig. 6A, C, D). Reductions in these cytokines would be expected to reduce IL-17A expression, as observed (Fig. 4B, E). Whereas TNFR2 deficiency and etanercept reduced *Il17a* mRNA, IL-17A⁺ $\gamma\delta$ T cells, and BAL G-CSF in lean WT mice, neither TNFR2 deficiency or etanercept affected these outcomes in obese *Cpe^{fat}* mice (Fig. 4B, C and 5A–C).

Discussion

Our data indicate a key role for IL-17A⁺ $\gamma\delta$ T cells in the pulmonary inflammation induced by subacute O₃. Our data also indicate that TNF α promotes pulmonary inflammation after subacute O₃ by inducing recruitment of IL-17A⁺ $\gamma\delta$ T cells, likely via *Ccl20* expression. Finally, our data suggest that the attenuated

pulmonary inflammation observed in obese mice after subacute O₃ is the result of reduced pulmonary IL-17A⁺ $\gamma\delta$ T cells, consequent to reduced IL-23 and IL-6 expression.

Inflammatory cell recruitment to the lungs after subacute O₃ exposure required $\gamma\delta$ T cells (Fig. 1A, B). $\gamma\delta$ T cells have also been shown to be required for the pulmonary inflammation observed 24 but not 8 hours after acute exposure to much higher O₃ concentrations (2 ppm) [49, 50], consistent with the time needed for recruitment and activation of $\gamma\delta$ T cells. However, in those studies, the precise role of these $\gamma\delta$ T cells was not assessed. Our data indicate that after exposure to lower concentrations of O₃ for much longer periods of time, the role of $\gamma\delta$ T cells involved IL-17A expression. Both lung *Il17a* mRNA and lung IL-17A⁺ $\gamma\delta$ T cells increased after subacute O₃ exposure with a time course similar to that of neutrophil recruitment (Figs. 1A, 1D, 2B). Furthermore, O₃-induced increases in *Il17a* mRNA abundance were abolished in TCR $\delta^{-/-}$ mice (Fig. 1D). In addition, both BAL neutrophils




Figure 4. Impact of TNFR2 deficiency (A–C) or etanercept (D–F) on O₃-induced inflammation in obese (*Cpe^{fat}*) and lean (WT) mice. (A, D) BAL neutrophils; (B, E) *Il17a* mRNA expression; (C, F) BAL G-CSF. Results are mean \pm SE of data from 3–11 mice in each group. *p < 0.05 versus air-exposed mice of same genotype and treatment; #p < 0.05 versus exposure matched lean mice with same TNFR2 genotype or treatment; & p < 0.05 versus TNFR2 sufficient (A–C) or vehicle treated mice (D–F) with same exposure and *Cpe* genotype. doi:10.1371/journal.pone.0097707.g004




Figure 5. Role of TNF α for IL-17A expression in $\gamma\delta$ T cells. Total number of (A) lung IL-17A⁺CD45⁺ cells; (B) lung IL-17A⁺ $\gamma\delta$ T cells; and (C) total lung $\gamma\delta$ T cells. Results are mean \pm SE of data from 5–6 mice in each group. #p<0.05 compared to lean mice with same TNFR2 genotype; & p<0.05 compared to TNFR2^{+/+} Cpe genotype matched mice. doi:10.1371/journal.pone.0097707.g005

and macrophages were reduced in mice treated with anti-IL-17A versus isotype control antibody (Fig. 3A). This ability of IL-17A⁺ $\gamma\delta$ T cells to control the influx of macrophages and neutrophils is consistent with the findings in other models of lung infection and injury [4,51–54]. While our data indicate that IL-17⁺ $\gamma\delta$ T cells are required for O₃-induced inflammatory cell recruitment, they are not sufficient. For example, O₃ is highly reactive and macrophages and epithelial cells are the initial targets of its action. These cells are the likely source of TNF α which is required for neutrophil recruitment (Fig. 4) perhaps via induction of CCL20 and consequent recruitment IL-17A⁺ $\gamma\delta$ T cells (Figs. 5,6). Epithelial cells are also the likely source of CCL20. Furthermore, macro-

phages also produce IL-17A after O₃ exposure [18], and the role of $\gamma\delta$ T cells may be to promote these effects. Macrophages and epithelial cells are also the likely source of other chemokines that interact with IL-17A (see below) to promote neutrophil recruitment.

IL-17A has direct chemoattractant effects on macrophages [55], which likely explains the ability of anti-IL-17A to attenuate O₃-induced increases in BAL macrophages (Fig. 3A). In contrast, IL-17A induces neutrophil recruitment to the lungs by inducing expression of other neutrophil chemotactic and survival factors. With subacute O₃ exposure, G-CSF appears to be one of these factors. In WT mice, the time courses of induction of BAL G-CSF




Figure 6. TNF α signaling is required for expression of *Il12rβ1* and *Ccl20*. (A) BAL IL-6; (B–D) *Il23* (*p19*) mRNA; (E–G) *Ccl20* mRNA; (H) *Il12rβ1* mRNA; (I) MFI and (J) % of $\gamma\delta$ T cells positive for IL-12R β 1 after stimulation with TNF α . Results are mean \pm SE of data from 3–11 mice in each group. *p<0.05 versus air exposed mice of the same genotype; #p<0.05 versus exposure matched lean mice with the same TNFR2 genotype or treatment; & p<0.05 versus WT; %<0.05 obese versus lean regardless of TNFR2 genotype. doi:10.1371/journal.pone.0097707.g006

and *Il17a* expression were similar (Fig. 1D,E). Importantly, anti-IL-17A and $\gamma\delta$ T cell deficiency each caused a marked and significant reduction in BAL G-CSF in O₃ exposed mice (Fig. 1E, 3C). The data are also consistent with our previous observations showing reductions in BAL G-CSF in O₃-exposed adiponectin-deficient mice treated with anti-IL-17A [18]. The observed role of IL-17A in G-CSF expression is in agreement with previous reports indicating that IL-17A signaling increases the transcription and stability of the *Gcsf* mRNA [56,57], via effects on ERK1/2 activation [58]. G-CSF causes neutrophil release from bone marrow and promotes neutrophil survival [59]. Since serum G-CSF did not increase after subacute O₃ exposure (data not shown), G-CSF is unlikely to act via effects on bone marrow in this model. Instead, G-CSF likely contributes by increasing the survival of neutrophils recruited to the lungs in response to other factors such as IP-10 (Fig. 1G).

TNF α is not directly chemotactic for neutrophils [24]. However, in lean WT mice, TNFR2 deficiency or the TNF α antagonist, etanercept, reduced the O₃-induced increase in BAL neutrophils (Fig. 4A,D) consistent with previous reports [22,23,60] indicating a role for TNF α in neutrophil recruitment induced by subacute O₃. TNF α also contributes to neutrophil recruitment in other conditions (reviewed in [61]), though the mechanism is not well understood. Our data suggest that at least in the setting of O₃ exposure, the ability of TNF α to recruit neutrophils involves IL-17A and that the source of this IL-17A is $\gamma\delta$ T cells (Fig. 5). O₃-induced increases in pulmonary *Il17a* expression were attenuated in TNFR2^{-/-} versus WT mice (Fig. 4B) and in etanercept versus vehicle treated WT mice (Fig. 4E). The number of IL-17A⁺ $\gamma\delta$ T cells in the lung was also lower in TNFR2^{-/-} versus WT mice exposed to O₃ (Fig. 5A,B). The ability of TNF α to promote pulmonary IL-17A expression after O₃ exposure is consistent with the role of TNF α in other pathogenic states. For example, etanercept reduces the elevated blood and skin Th17 cells observed in patients with psoriasis [26]. Similarly, another anti-TNF α therapy, infliximab, reduces IL-17A in ocular fluid from uveitis patients with Behcet's disease [27].

To better understand the role of TNF α , we examined IL-6 and IL-23 expression. Both these cytokines can contribute to induction of IL-17A in $\gamma\delta$ T cells [45,62]. Both IL-6 and IL-23 were induced in the lungs after O₃ exposure, but were not affected by TNFR2 deficiency or by etanercept (Fig. 6A,C,D), indicating that TNF α is not required for their expression. We did observe that mRNA expression of one of the two subunits of the IL-23 receptor, *Il12rb1*, was decreased (Fig. 6H) in unexposed lungs from TNFR2^{-/-} mice. Similar trends were observed after etanercept treatment (data not shown). Since others have reported that TNF α can act directly on $\gamma\delta$ T cells [40,63], we considered the possibility that TNF α was acting to increase *Il12rb1* expression on $\gamma\delta$ T cells, thus increasing their ability to respond to IL-23. However, culture of lung cells with TNF α resulted in no change in surface bound IL-12R β 1 on $\gamma\delta$ T cells (Fig. 6I,J). Instead, our data suggest that effects of TNF α on *Ccl20* expression (Fig. 6F,G) account for the observed effects of TNF α /TNFR blockade on IL-17A⁺ $\gamma\delta$ T cells. *Ccl20* acts via CCR6, a receptor expressed by IL-17A⁺ $\gamma\delta$ T cells that promotes chemotaxis of these cells [64]. TNF α is also required for pulmonary *Ccl20* expression after acute O₃ exposure (2 ppm for 3 h) [37]. A role for TNF α in *Ccl20* expression has also been demonstrated in dermal lesions of psoriasis patients based on treatment with the TNF α antagonist infliximab [65].

We observed fewer neutrophils in BAL fluid of obese *Cpe^{fat}* versus lean WT mice after subacute O₃ exposure (Fig. 4A,D), consistent with previous observations [34]. Reduced responses are observed in *Cpe^{fat}* mice not only after 48 h exposure (Fig. 4A,D), but also after 24 or 72 h exposures [34]. Pulmonary *Il17a* expression and IL-17A⁺ $\gamma\delta$ T cells were also reduced in the obese mice, as was the total number of $\gamma\delta$ T cells (Fig. 4). BAL G-CSF was also lower in *Cpe^{fat}* versus lean WT mice (Fig. 4C,F). Moreover, O₃-induced increases in BAL IL-6 and pulmonary *Il23* expression were also reduced in *Cpe^{fat}* versus WT mice (Fig. 6C,D). TNFR2 deficiency or etanercept treatment in *Cpe^{fat}* mice did not further reduce BAL neutrophils or pulmonary *Il17a* expression, in contrast to what was observed in WT mice (Fig. 4B,E). Given the already reduced numbers of total $\gamma\delta$ T cells in *Cpe^{fat}* mice exposed to O₃ (Fig. 5C), and our observations indicating the key role for IL-17A⁺ $\gamma\delta$ T cells in the effects of TNF α on neutrophil recruitment, it is not surprising that TNF α had no further effect on the response to O₃ in obese mice. Taken together, the data suggest that obesity-related reductions in neutrophil recruitment induced by subacute O₃ exposure are the result of reduced IL-17A-dependent G-CSF release, consequent to reduced IL-6 and IL-23 expression. However, we cannot rule out the possibility that other factors also contributed. For example, neutrophils from obese mice exhibit reduced chemotactic activity towards CXCR2 ligands [66]. Such defects in neutrophil chemotaxis would also be expected to reduce O₃-induced neutrophil recruitment in *Cpe^{fat}* mice.

In addition to affecting responses to O₃, obesity also impacts responses to bacterial and viral infections [67–71]. As described above, IL-17⁺ $\gamma\delta$ T cells contribute to neutrophil recruitment and pathogen clearance after certain bacterial infections [3,4]. IL-17⁺ $\gamma\delta$ T cells are also required for clearance of secondary infections after influenza [72]. Hence, obesity-related changes in IL-17⁺ $\gamma\delta$ T cells (Figs. 4b, 5a,b) may contribute not only to obesity-related alterations in responses to O₃, but may have broader implications for effects of obesity on host defense. In support of this, obese mice compared to lean mice have fewer skin $\gamma\delta$ T cells number and the few $\gamma\delta$ T cells they have are dysfunctional [73], which leads to impairment in wound healing. These decreases in $\gamma\delta$ T cells numbers and impairment in function of the skin in obese mice are due to altered STAT5 signaling and chronic TNF α signaling [74].

In summary, our data indicate that $\gamma\delta$ T cells are required for the pulmonary inflammation that occurs after subacute O₃ exposure in mice via their ability to produce IL-17A. IL-17A then leads to G-CSF expression. Our data also indicate that TNF α is required for recruitment IL-17A⁺ $\gamma\delta$ T cells to the lungs likely through its ability to induce *Ccl20*. These results emphasize the importance of $\gamma\delta$ T cells not only for pathogen clearance, but also for responses to other insults that induce oxidative stress, and describe a new role for TNF α in these events. Finally, our data indicate that obesity-related reductions in the ability of subacute O₃ to promote neutrophil recruitment to the lungs are the result of reduced IL-17A⁺ $\gamma\delta$ T cells. These results suggest that other conditions that impact $\gamma\delta$ T cell recruitment or activation will also impact responses to this common pollutant.

Author Contributions

Conceived and designed the experiments: JAM ASW JDB HS DIK SAS. Performed the experiments: JAM ASW JDB APW LC FMCN. Analyzed the data: JAM ASW JDB SAS. Contributed reagents/materials/analysis tools: JAM ASW. Wrote the paper: JAM ASW SAS.

References

- Wands JM, Roark CL, Aydin MG, Jin N, Hahn Y-S, et al. (2005) Distribution and leukocyte contacts of $\gamma\delta$ T cells in the lung. *Journal of Leukocyte Biology* 78: 1086–1096.
- Laan M, Cui Z-H, Hoshino H, Lötvalld J, Sjöstrand M, et al. (1999) Neutrophil Recruitment by Human IL-17 Via C-X-C Chemokine Release in the Airways. *The Journal of Immunology* 162: 2347–2352.
- Skeen MJ, Ziegler HK (1993) Induction of murine peritoneal gamma/delta T cells and their role in resistance to bacterial infection. *The Journal of Experimental Medicine* 178: 971–984.
- Cheng P, Liu T, Zhou W-Y, Zhuang Y, Peng L-s, et al. (2012) Role of gamma-delta T cells in host response against *Staphylococcus aureus*-induced pneumonia. *BMC Immunology* 13: 38.
- Koohsari H, Tamaoka M, Campbell H, Martin J (2007) The role of gammadelta T cells in airway epithelial injury and bronchial responsiveness after chlorine gas exposure in mice. *Respiratory Research* 8: 21.
- McMenamin C, Pimm C, McKersey M, Holt PG (1994) Regulation of IgE responses to inhaled antigen in mice by antigen-specific gamma delta T cells. *Science* 265: 1869–1871.
- Pociask DA, Chen K, Mi Choi S, Oury TD, Steele C, et al. (2011) $\gamma\delta$ T Cells Attenuate Bleomycin-Induced Fibrosis through the Production of CXCL10. *The American Journal of Pathology* 178: 1167–1176.
- Pons J, Sauleda J, Ferrer JM, Barceló B, Fuster A, et al. (2005) Blunted $\gamma\delta$ T-lymphocyte response in chronic obstructive pulmonary disease. *European Respiratory Journal* 25: 441–446.
- Devlin RB, McDonnell WF, Mann R, Becker S, House DE, et al. (1991) Exposure of Humans to Ambient Levels of Ozone for 6.6 Hours Causes Cellular and Biochemical Changes in the Lung. *American Journal of Respiratory Cell and Molecular Biology* 4: 72–81.
- Bell ML, Dominici F, Samet JM (2005) A Meta-Analysis of Time-Series Studies of Ozone and Mortality With Comparison to the National Morbidity, Mortality, and Air Pollution Study. *Epidemiology* 16: 436–445. doi:10.1097/000165817.0000140152.0000165885.
- Levy JI, Chemerynski SM, Sarnat JA (2005) Ozone Exposure and Mortality: An Empirical Bayes Meta-regression Analysis. *Epidemiology* 16: 458–468. doi:10.1097/000165817.0000108301.0000165823.
- Triche EW, Gent JF, Holford TR, Belanger K, Bracken MB, et al. (2006) Low-level ozone exposure and respiratory symptoms in infants. *Environ Health Perspect* 114: 911–916.
- Chiu H-F, Cheng M-H, Yang C-Y (2009) Air Pollution and Hospital Admissions for Pneumonia in a Subtropical City: Taipei, Taiwan. *Inhalation Toxicology* 21: 32–37.
- Peden DB (1996) Effect of Air Pollution in Asthma and Respiratory Allergy. *Otolaryngology – Head and Neck Surgery* 114: 242–247.
- Charpin D, Pascal L, Birnbaum J, Armengaud A, Sambuc R, et al. (1999) Gaseous air pollution and atopy. *Clin Exp Allergy* 29: 1474–1480.
- Boutin-Forzano S, Hammou Y, Gouitaa M, Charpin D (2005) Air pollution and atopy. *Eur Ann Allergy Clin Immunol* 37: 11–16.
- Zhao Q, Simpson LG, Driscoll KE, Leikauf GD (1998) Chemokine regulation of ozone-induced neutrophil and monocyte inflammation. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 274: L39–L46.
- Kasahara DI, Kim HY, Williams AS, Verbout NG, Tran J, et al. (2012) Pulmonary inflammation induced by subacute ozone is augmented in adiponectin-deficient mice: role of IL-17A. *J Immunol* 188: 4558–4567.
- Shimizu M, Hasegawa N, Nishimura T, Endo Y, Shiraiishi Y, et al. (2009) Effects of TNF-alpha-converting enzyme inhibition on acute lung injury induced by endotoxin in the rat. *Shock* 32: 535–540.
- Chung A, Dai J, Tai H, Xie C, Wright JL (2002) Tumor Necrosis Factor- α Is Central to Acute Cigarette Smoke-Induced Inflammation and Connective Tissue Breakdown. *American Journal of Respiratory and Critical Care Medicine* 166: 849–854.
- Malaviya R, Ikeda T, Ross E, Abraham SN (1996) Mast cell modulation of neutrophil influx and bacterial clearance at sites of infection through TNF- α . *Nature* 381: 77–80.
- Cho H-Y, Zhang L-Y, Kleberger SR (2001) Ozone-induced lung inflammation and hyperreactivity are mediated via tumor necrosis factor- α receptors. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 280: L537–L546.
- Kleberger SR, Levitt RC, Zhang LY, Longphre M, Harkema J, et al. (1997) Linkage analysis of susceptibility to ozone-induced lung inflammation in inbred mice. *Nat Genet* 17: 475–478.
- Yonemaru M, Stephens KE, Ishizaka A, Zheng H, Hogue RS, et al. (1989) Effects of tumor necrosis factor on PMN chemotaxis, chemiluminescence, and elastase activity. *J Lab Clin Med* 114: 674–681.
- Pober JS (1987) Effects of tumour necrosis factor and related cytokines on vascular endothelial cells. *Ciba Found Symp* 131: 170–184.
- Antiga E, Volpi W, Cardilicchia E, Maggi L, Fili L, et al. (2012) Etanercept Downregulates the Th17 Pathway and Decreases the IL-17+/IL-10+ Cell Ratio in Patients with Psoriasis Vulgaris. *Journal of Clinical Immunology* 32: 1221–1232.
- Sugita S, Kawazoe Y, Imai A, Yamada Y, Horie S, et al. (2012) Inhibition of Th17 differentiation by anti-TNF-alpha therapy in uveitis patients with Behcet's disease. *Arthritis Research & Therapy* 14: R99.
- Cazzola M, Calzetta L, Lauro D, Bettoncelli G, Cricelli C, et al. (2013) Asthma and COPD in an Italian adult population: role of BMI considering the smoking habit. *Respir Med* 107: 1417–1422.
- Ehrlich SF, Quesenberry CP, Van Den Eeden SK, Shan J, Ferrara A (2010) Patients Diagnosed With Diabetes Are at Increased Risk for Asthma, Chronic Obstructive Pulmonary Disease, Pulmonary Fibrosis, and Pneumonia but Not Lung Cancer. *Diabetes Care* 33: 55–60.
- Johnston RA, Theman TA, Lu FL, Terry RD, Williams ES, et al. (2008) Diet-induced obesity causes innate airway hyperresponsiveness to methacholine and enhances ozone-induced pulmonary inflammation. *Journal of Applied Physiology* 104: 1727–1735.
- Johnston RA, Theman TA, Shore SA (2006) Augmented responses to ozone in obese carboxypeptidase E-deficient mice. *Am J Physiol Regul Integr Comp Physiol* 290: R126–133.
- Lu FL, Johnston RA, Flynt L, Theman TA, Terry RD, et al. (2006) Increased pulmonary responses to acute ozone exposure in obese db/db mice. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 290: L856–L865.
- Shore SA, Rivera-Sanchez YM, Schwartzman IN, Johnston RA (2003) Responses to ozone are increased in obese mice. *J Appl Physiol* 95: 938–945.
- Shore SA, Lang JE, Kasahara DI, Lu FL, Verbout NG, et al. (2009) Pulmonary responses to subacute ozone exposure in obese vs. lean mice. *Journal of Applied Physiology* 107: 1445–1452.
- Mancuso P (2010) Obesity and lung inflammation. *Journal of Applied Physiology* 108: 722–728.
- Coleman DL, Eicher EM (1990) Fat (fat) and Tubby (tubby): Two Autosomal Recessive Mutations Causing Obesity Syndromes in the Mouse. *Journal of Heredity* 81: 424–427.
- Williams AS, Mathews JA, Kasahara DI, Chen L, Wurmbrand AP, et al. (2013) Augmented Pulmonary Responses to Acute Ozone Exposure in Obese Mice: Roles of TNFR2 and IL-13. *Environ Health Perspect* 121: 551–557.
- Skerry C, Harper J, Klunk M, Bishai WR, Jain SK (2012) Adjunctive TNF inhibition with standard treatment enhances bacterial clearance in a murine model of necrotic TB granulomas. *PLoS ONE* 7: e39680.
- Grounds M, Davies M, Torrisi J, Shavlakadze T, White J, et al. (2005) Silencing TNF α activity by using Remicade or Enbrel blocks inflammation in whole muscle grafts: an in vivo bioassay to assess the efficacy of anti-cytokine drugs in mice. *Cell and Tissue Research* 320: 509–515.
- Lahn M, Kalatardi H, Mittelstadt P, Plüm E, Vollmer M, et al. (1998) Early Preferential Stimulation of $\gamma\delta$ T Cells by TNF- α . *The Journal of Immunology* 160: 5221–5230.
- Bhalla DK (1999) Ozone-induced lung inflammation and mucosal barrier disruption: toxicology, mechanisms, and implications. *J Toxicol Environ Health B Crit Rev* 2: 31–86.
- Backus GS, Howden R, Fostel J, Bauer AK, Cho HY, et al. (2010) Protective role of interleukin-10 in ozone-induced pulmonary inflammation. *Environ Health Perspect* 118: 1721–1727.
- Johnston RA, Schwartzman IN, Flynt L, Shore SA (2005) Role of interleukin-6 in murine airway responses to ozone. *American Journal of Physiology - Lung Cellular and Molecular Physiology* 288: L390–L397.
- Michalec L, Choudhury BK, Postlethwait E, Wild JS, Alam R, et al. (2002) CCL7 and CXCL10 orchestrate oxidative stress-induced neutrophilic lung inflammation. *J Immunol* 168: 846–852.
- Sutton CE, Lalor SJ, Sweeney CM, Brereton CF, Lavelle EC, et al. (2009) Interleukin-1 and IL-23 Induce Innate IL-17 Production from $\gamma\delta$ T Cells, Amplifying Th17 Responses and Autoimmunity. *Immunity* 31: 331–341.
- Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B (2006) TGF β in the Context of an Inflammatory Cytokine Milieu Supports De Novo Differentiation of IL-17-Producing T Cells. *Immunity* 24: 179–189.
- Li Z, Burns AR, Byeseda Miller S, Smith CW (2011) CCL20, $\gamma\delta$ T cells, and IL-22 in corneal epithelial healing. *The FASEB Journal* 25: 2659–2668.
- Mabuchi T, Singh TP, Takekoshi T, Jia G-f, Wu X, et al. (2013) CCR6 Is Required for Epidermal Trafficking of [gamma][delta]-T Cells in an IL-23-Induced Model of Psoriasisiform Dermatitis. *J Invest Dermatol* 133: 164–171.
- Matsubara S, Takeda K, Jin N, Okamoto M, Matsuda H, et al. (2009) Vgamma1+ T cells and tumor necrosis factor-alpha in ozone-induced airway hyperresponsiveness. *Am J Respir Cell Mol Biol* 40: 454–463.
- King DP, Hyde DM, Jackson KA, Novosad DM, Ellis TN, et al. (1999) Cutting Edge: Protective Response to Pulmonary Injury Requires $\gamma\delta$ T Lymphocytes. *The Journal of Immunology* 162: 5033–5036.
- Umecoma M, Yahagi A, Hamada S, Begum MD, Watanabe H, et al. (2007) IL-17-Mediated Regulation of Innate and Acquired Immune Response against Pulmonary *Mycobacterium bovis* Bacille Calmette-Guérin Infection. *The Journal of Immunology* 178: 3786–3796.
- Braun RK, Ferrick C, Neubauer P, Sjöding M, Sterner-Kock A, et al. (2008) IL-17 producing gammadelta T cells are required for a controlled inflammatory response after bleomycin-induced lung injury. *Inflammation* 31: 167–179.

53. Wozniak K, Kolls J, Wormley F (2012) Depletion of neutrophils in a protective model of pulmonary cryptococcosis results in increased IL-17A production by gamma/delta T cells. *BMC Immunology* 13: 65.
54. Lo Re S, Dumoutier L, Couillin I, Van Vyve C, Yakoub Y, et al. (2010) IL-17A-Producing $\gamma\delta$ T and Th17 Lymphocytes Mediate Lung Inflammation but Not Fibrosis in Experimental Silicosis. *The Journal of Immunology* 184: 6367–6377.
55. Sergejeva S, Ivanov S, Lotvall J, Linden A (2005) Interleukin-17 as a recruitment and survival factor for airway macrophages in allergic airway inflammation. *Am J Respir Cell Mol Biol* 33: 248–253.
56. Cai X-Y, Gommoll Jr CP, Justice L, Narula SK, Fine JS (1998) Regulation of granulocyte colony-stimulating factor gene expression by interleukin-17. *Immunology Letters* 62: 51–58.
57. Jones CE, Chan K (2002) Interleukin-17 stimulates the expression of interleukin-8, growth-related oncogene-alpha, and granulocyte-colony-stimulating factor by human airway epithelial cells. *Am J Respir Cell Mol Biol* 26: 748–753.
58. Hirai Y, Iyoda M, Shibata T, Kuno Y, Kawaguchi M, et al. (2012) IL-17A stimulates granulocyte colony-stimulating factor production via ERK1/2 but not p38 or JNK in human renal proximal tubular epithelial cells. *American Journal of Physiology - Renal Physiology* 302: F244–F250.
59. Cox G, Gauldie J, Jordana M (1992) Bronchial epithelial cell-derived cytokines (G-CSF and GM-CSF) promote the survival of peripheral blood neutrophils in vitro. *Am J Respir Cell Mol Biol* 7: 507–513.
60. Bauer AK, Travis EL, Malhotra SS, Rondini EA, Walker C, et al. (2010) Identification of novel susceptibility genes in ozone-induced inflammation in mice. *Eur Respir J* 36: 428–437.
61. Vassalli P (1992) The Pathophysiology of Tumor Necrosis Factors. *Annual Review of Immunology* 10: 411–452.
62. Korn T, Petermann F (2012) Development and function of interleukin 17-producing $\gamma\delta$ T cells. *Annals of the New York Academy of Sciences* 1247: 34–45.
63. Ueta C, Kawasumi H, Fujiwara H, Miyagawa T, Kida H, et al. (1996) Interleukin-12 activates human gamma delta T cells: synergistic effect of tumor necrosis factor-alpha. *Eur J Immunol* 26: 3066–3073.
64. Kim CH (2009) Migration and function of Th17 cells. *Inflamm Allergy Drug Targets* 8: 221–228.
65. Brunner PM, Koszik F, Reininger B, Kalb ML, Bauer W, et al. (2013) Infliximab induces downregulation of the IL-12/IL-23 axis in 6-sulfo-LacNac (slan)+ dendritic cells and macrophages. *Journal of Allergy and Clinical Immunology* 132: 1184–1193.e1188.
66. Kordonowy LL, Burg E, Lenox CC, Gauthier LM, Petty JM, et al. (2012) Obesity Is Associated with Neutrophil Dysfunction and Attenuation of Murine Acute Lung Injury. *American Journal of Respiratory Cell and Molecular Biology* 47: 120–127.
67. Smith AG, Sheridan PA, Harp JB, Beck MA (2007) Diet-Induced Obese Mice Have Increased Mortality and Altered Immune Responses When Infected with Influenza Virus. *The Journal of Nutrition* 137: 1236–1243.
68. Mancuso P, Gottschalk A, Phare SM, Peters-Golden M, Lukacs NW, et al. (2002) Leptin-Deficient Mice Exhibit Impaired Host Defense in Gram-Negative Pneumonia. *The Journal of Immunology* 168: 4018–4024.
69. Wieland CW, Florquin S, Chan ED, Leemans JC, Weijer S, et al. (2005) Pulmonary Mycobacterium tuberculosis infection in leptin-deficient ob/ob mice. *International Immunology* 17: 1399–1408.
70. Milner JJ, Sheridan PA, Karlsson EA, Schultz-Cherry S, Shi Q, et al. (2013) Diet-Induced Obese Mice Exhibit Altered Heterologous Immunity during a Secondary 2009 Pandemic H1N1 Infection. *The Journal of Immunology* 191: 2474–2485.
71. Morgan OW, Bramley A, Fowlkes A, Freedman DS, Taylor TH, et al. (2010) Morbid Obesity as a Risk Factor for Hospitalization and Death Due to 2009 Pandemic Influenza A(H1N1) Disease. *PLoS ONE* 5: e9694.
72. Li W, Moltedo B, Moran TM (2012) Type I interferon induction during influenza virus infection increases susceptibility to secondary Streptococcus pneumoniae infection by negative regulation of gammadelta T cells. *J Virol* 86: 12304–12312.
73. Taylor KR, Costanzo AE, Jameson JM (2011) Dysfunctional gammadelta T cells contribute to impaired keratinocyte homeostasis in mouse models of obesity. *J Invest Dermatol* 131: 2409–2418.
74. Taylor KR, Mills RE, Costanzo AE, Jameson JM (2010) Gammadelta T cells are reduced and rendered unresponsive by hyperglycemia and chronic TNF α in mouse models of obesity and metabolic disease. *PLoS ONE* 5: e11422.