

A new species of genus *Pseudaspidapion* Wanat, 1990 (Coleoptera, Apionidae) from China

M. A. Alonso-Zarazaga^{1,2,†}, Zhiliang Wang^{1,‡}, Runzhi Zhang^{1,§}

1 Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Beijing, People's Republic of China **2** Depto. de Biodiversidad y Biología Evolutiva, Museo Nacional de Ciencias Naturales (CSIC), José Gutiérrez Abascal, 2, E-28006 Madrid, Spain

† [urn:lsid:zoobank.org/author:EA3695DA-2C2E-4021-95CB-11D24ED253F0](https://doi.org/urn:lsid:zoobank.org/author:EA3695DA-2C2E-4021-95CB-11D24ED253F0)

‡ [urn:lsid:zoobank.org/author:982BF33F-BB3B-4D0C-945F-8F0AACA6369B](https://doi.org/urn:lsid:zoobank.org/author:982BF33F-BB3B-4D0C-945F-8F0AACA6369B)

§ [urn:lsid:zoobank.org/author:9C171E03-04E6-40FB-8336-FF89ED387136](https://doi.org/urn:lsid:zoobank.org/author:9C171E03-04E6-40FB-8336-FF89ED387136)

Corresponding author: Runzhi Zhang (zhangrz@ioz.ac.cn)

Academic editor: Terry Erwin | Received 26 April 2011 | Accepted 4 July 2011 | Published 25 July 2011

[urn:lsid:zoobank.org/pub:0273339E-155D-4628-9375-3E30945A2C2F](https://doi.org/urn:lsid:zoobank.org/pub:0273339E-155D-4628-9375-3E30945A2C2F)

Citation: Alonso-Zarazaga MA, Wang Z, Zhang R (2011) A new species of genus *Pseudaspidapion* Wanat, 1990 (Coleoptera, Apionidae) from China. ZooKeys 120: 41–54. doi: 10.3897/zookeys.120.1434

Abstract

Pseudaspidapion botanicum **sp. n.** from China is described and figured. Its host plant is *Grewia biloba* G. Don var. *parviflora* (Bunge) Hand.-Mazz (Malvaceae: Grewioideae). The genus *Harpapion* Voss, 1966 is recorded as new for China and Vietnam and two **comb. n.** are proposed: *Harpapion vietnamense* (Korotyaev, 1985) (from *Aspidapion*) and *H. coelebs* (Korotyaev, 1987) (from *Pseudaspidapion*). A key to the known species of the genus *Pseudaspidapion* from China is presented.

Keywords

Weevils, *Pseudaspidapion botanicum*, *Harpapion*, Beijing, Shaanxi, new species, new combination, morphology, systematics, biology, key

Introduction

M. Wanat (1990) erected the genus *Pseudaspidapion* relating it to the genus *Aspidapion* Schilsky, 1901. Alonso-Zarazaga and Lyal (1999) placed it in the tribe Aspidapiini

Alonso-Zarazaga, 1990 of the subfamily Apioninae Schoenherr, 1823 (Coleoptera, Curculionioidea). This genus is quite similar to *Aspidapion*, but it can be distinguished from the latter by relatively equal width of elytral striae and interstriae, the presence of fenestrae in the tegminal plate, and the absence of protibial mucrones in males, among other features (Wanat 1990). Apart from the type species, *Apion spadiceum* Wagner, 1908, Wanat also proposed 15 new combinations for species coming from East Africa, India and South China. At present, there are 17 paleotropical species included in the genus. In the Chinese fauna, there were 4 species recorded only from Yunnan province, and these specimens were all collected during the 1950's (Korotyaev 1985, 1987).

In 2008, we collected a series of specimens of a *Pseudaspidapion*. Further specimens were later found. We consider them to represent a species new to science that we describe below.

In addition to the description of the adult characters, we also provide some biological data for the new species after one year of survey in the Beijing area.

Material and methods

Materials examined of new species for this study are to be deposited in the Institute of Zoology, Chinese Academy of Sciences, Beijing (IZCAS), the Museo Nacional de Ciencias Naturales, Madrid (MNCN), the Zoological Institute of Russian Academy of Sciences, Moscow (ZIN), the Museum of Natural History, University of Wrocław (MNHW) and the Beijing Botanical Garden, Beijing (BBG).

Type specimens were obtained from ZIN on loan or belong to IZCAS and their data are summarised in Table 1. Information about their condition and taxonomy is given in the Discussion section.

Descriptions were made and photographs were taken with a CCD Qimage MicroPublisher 5.0 RTV mounted on a Zeiss SteREO Discovery V.12. Extended focus images were generated with Auto-Montage Pro 5.03.0061 and edited with Adobe Photoshop CS 5.0 if required. Microscopic slides were studied under a Leica DM 2500 microscope and photos were taken with a Nikon CoolPix 5400. The map was made with the software ArcGIS 9.3. Drawings were made from the original photographs by using the software Adobe Illustrator CS5.0, or directly by using a drawing tube linked to the microscope.

Nomenclature of the rostral parts follows Alonso-Zarazaga (1989) and that of genitalia follows Alonso-Zarazaga (1990).

The dissecting method used follows Alonso-Zarazaga (1990). Abdomens were put into 10% NaOH for several hours until the inner tissues were digested, and the resultant structures were placed on a temporary microscope slide for examination.

After description, the genitalia and other parts of each specimen were placed in DMHF on a plastic card for long term conservation (Steedman 1958; Bameul 1990).

Labels are described as they are (in Chinese), with pinyin romanization or comments in square brackets; labels are separated by semicolons and lines by slashes.

Table 1. List of studied type specimens and depositories.

Species	Depository	Examined type material
<i>Aspidapion inarmatum</i> Korotyaev, 1985	IZCAS	2 paratypes
<i>A. panfilovi</i> Korotyaev, 1985	IZCAS	Holotype and 1 paratype
<i>Pseudaspidapion coelebs</i> (Korotyaev, 1987)	ZIN	1 paratype
<i>P. kryzhanovskii</i> (Korotyaev, 1987)	ZIN	1 paratype
<i>P. medvedevi</i> (Korotyaev, 1985)	ZIN	Holotype
<i>P. topali</i> (Korotyaev, 1985)	ZIN	1 paratype
<i>P. vietnamense</i> (Korotyaev, 1987)	ZIN	Holotype
<i>P. yunnanicum</i> (Korotyaev, 1985)	IZCAS	Holotype and 3 paratypes
<i>P. zagulajevi</i> (Korotyaev, 1987)	ZIN	2 paratypes

Taxonomic treatment

Pseudaspidapion botanicum Alonso-Zarazaga & Wang, sp. n.

urn:lsid:zoobank.org:act:88656E61-918F-4F1B-A3BD-6A3FA714A237

http://species-id.net/wiki/Pseudaspidapion_botanicum

Figs 1–28 (from paratypes)

Diagnosis. This new species resembles *P. yunnanicum*, but it can be distinguished from the latter by the characters in the table 2.

Description (male holotype, except where indicated). *Measurements* (in mm): Standard length: 1.82. Rostrum: length: 0.71, maximum width: 0.17. Pronotum: median length: 0.49, maximum width: 0.59. Elytra: median length: 1.38, maximum width: 1.08.

With the general characters of genus *Pseudaspidapion* as described in Wanat (1990). *Integument.* Generally piceous black (Figs. 1–2).

Vestiture composed of thick, longer white piliform scales with truncate apex and thin, shorter brownish to greyish acute hairs; scales present on coxae, femora and tibiae, sides of pronotum, meso- and metaventricle, mesoventral process, base of 3rd elytral interstria, metarostrum, and weak subocular patch, sometimes intermediate types present. Antennal scape apex and pedicel with sparse short greyish hairs, club and desmomerer 2–7 covered with dark hairs only. Pronotal vestiture centrifugal, scales both on apex and base perpendicular to the margins, hairs on disc distinctly longer than those on elytra, reaching middle of preceding hair. Elytral vestiture white at base and sides, brownish on disc, interstitial scales large, longer than 2/3 of the interstitial width, in two rather regular rows per interstria, one specialized seta on apical region of 9th interstria.

Rostrum cylindrical and moderately robust, in dorsal view 4.18× as long as maximum width, 1.45× as long as pronotum in midline, widest at mesorostrum, pro- and metarostrum with sides almost parallel, metarostrum with two very fine, punctulate dorsal submedial sulci shortened at mesorostral level, and two weak dorsal sublateral sulci, dorsal submedial sulci prolonged on frons, separated, dorsal sublateral sulci prolonged close to ocular margins, metarostrum microreticulate, matt, pro- and

Table 2. Comparative diagnostic characters for *P. botanicum* sp. n. and *P. yunnanicum*.

	<i>P. yunnanicum</i>	<i>P. botanicum</i> sp. n.
Scales on pronotal disc	Finer, sparser, shorter, tips of posterior scales not or hardly reaching the base of anterior ones	Thicker, denser and longer, tips of posterior scales surpassing the anterior by almost the basal half
Scape	Slender, ca. 0.26 mm long, 7.20–7.60× as long as wide, 1.66–1.85× as long as mesorostral width	Robust, ca. 0.20 mm long, 5.00–5.80× as long as wide, 1.24–1.67× as long as mesorostral width
Rostrum (♀)	Longer, 1.91–2.25× as long as pronotum, thinner, 7.35–10.30× as long as maximum width	Shorter, 1.35–1.53× as long as pronotum, more robust, 4.00–5.07× as long as maximum width
Metarostrum (dorsal view)	Microreticulate	Nearly smooth
Pronotum	Strong subapical constriction and sides moderately rounded behind it	Weaker subapical constriction, sides weakly rounded behind it
Prescutellar fovea	Deep, distinct, lanceolate, not prolonged apicad	Shallow, indistinct, sublinear, prolonged apicad to middle
Scutellum	Two basal tubercles separated medially by a distinctly deep notch, in anterior view tubercles not fused basally	Two basal tubercles separated medially by a weak notch, in anterior view tubercles basally fused

mesorostrum smooth, shining, sparsely punctulate; in lateral view weakly and evenly curved, ventral margin forming a weak angle at mesorostrum, each side with a very thin low dorsal sublateral keel running from front margin of eye to upper margin of scrobe and beyond, limiting ventrally the dorsal sublateral sulcus, prostrum with a marked ventral sublateral sulcus under this keel.

Head transverse, frons weakly convex with four rows of relatively deep punctures running from metarostrum and hardly surpassing hind margin of eyes, occiput almost reaching hind eye level, subocular keel reaching middle of eyes, the area between subocular keels microreticulate and impunctate, with a low, fine median keel. Eyes round, moderately convex.

Antennae inserted at basal 0.26 of rostral length, scape 5.00× as long as wide, as long as club, 1.18× as long as mesorostral width. Pedicel 2.14× as long as wide, as long as desmomer 2+3, desmome 2 1.67× as long as wide, desmomer 3–5 1.33× as long as wide, desmome 6 1.14× as long as wide, desmome 7 1.25× as long as wide, shortly obconical. Club oblong, compact, 2.5× as long as wide, as long as the last 5.5 desmomer, sutures well marked (Fig 7).

Pronotum campaniform, transverse, 0.83× as long as wide, widest just behind middle, constrictions weak, sides weakly dilated at middle, base 1.31× as wide as apex, bisinuate with moderate medial rounded projection towards scutellum, basal flange moderately developed. Prescutellar fovea shallow, as broad as one puncture's diameter, as long as 2–3 diameters, prolonged in a very fine sulcus reaching middle of pronotum. Discal punctures relatively deep, ca. 0.5–1× diameter apart, interspaces slightly convex, microreticulate.




Figures 1–4. *Pseudaspidapion botanicum* Alonso-Zarazaga & Wang, sp. n. **1** male paratype, dorsal view **2** male paratype, lateral view **3** female paratype, head and rostrum, dorsal view **4** female paratype, head and rostrum, lateral view. Scales: 1–2: 1000 μm ; 3–4: 500 μm .

Scutellum large, elongated, triangular, $1.63\times$ as long as wide, with two basal tubercles separated by a weak median depression, in lateral view subacutely prominent, in front view tubercles fused basally; apex constricted and moderately raised, distinctly visible in lateral view.

Elytra $1.23\times$ as long as wide, $2.82\times$ as long as pronotum, widest almost at the middle, humeri distinct; striae deep, about as wide as interstriae at base and apex, about half as wide at elytral disc, distinctly catenulate-punctate, punctures round to oblong, space between punctures about $2.00\times$ as long as puncture length, striae apically connected 1+2+9, 3+4, 5+6, 7+8. Interstriae evidently convex with small punctures, surface distinctly wrinkled, not microreticulate, shining. Macropterous (wing of paratype, Fig 8).

Ventral areas. Mesocoxae and metacoxae narrowly separated by a distance of $0.26\times$ and $0.38\times$ of their transverse diameter, respectively. Metaventricle ca. as long as mesocoxae. Mesoventral process more prominent than metaventral process. Anterior metaventral rim present, weak. Abdominal ventrites microreticulate, with length ratios along midline: 31-17-6-7-16. Ventrites 1–2 coarsely and densely punctate, 3–4 and base of 5 very sparsely and minutely punctate, apical half of 5 densely punctate with a median convexity. Suture I distinct, distance from hind margin of metacoxae, as long as ventrite 2. Ventrite 5 subsemicircular, transverse, $0.41\times$ as long as wide.

Pygidium (from paratype) semicircular, $0.69\times$ as long as wide, apical edge of arc with a flat and glabrous side, punctate medially with interspace microreticulate (Figs 9–10).



Figures 5–15. *Pseudaspidapion botanicum* Alonso-Zarazaga & Wang, sp. n., male paratype **5** front leg **6** hind tibia **7** antenna **8** hind wing **9** pygidium, lateral view **10** pygidium, dorsal view **11** tegmen, dorsal view **12** tegmen, lateral view **13** penis, dorsal view **14** penis, lateral view **15** spiculum gastrale. Scales: 5: 500 μ m; 6–7, 9–15: 200 μ m; 8: 1000 μ m.

Legs. Metafemora a little more robust than pro- and mesofemora, profemur $2.15\times$ as long as wide, widest at middle, minutely punctate. Protibia almost straight, $6.50\times$ as long as wide (Fig 5). Protarsomere 1 $2.29\times$ as long as wide, tarsomere 2 $1.22\times$ as long as wide, tarsomere 3 bilobed, $0.8\times$ as long as wide, lobes narrow, not much dilated outwards, onychium $3.25\times$ as long as wide, projecting from lobes of tarsomere 3 for $0.54\times$ its length. Meso- and metatibiae similarly mucronate, mucro subconical (Fig 6), about $0.50\times$ as long as tibial apical width. Tarsal claws with conspicuous obtuse basal tooth.

Genitalia and terminalia (from topotypic paratypes). Ninth sternite (spiculum gastrale) Y-shaped and slightly winged near arm base, manubrium ca. $2.00\times$ as long as arms (Fig 15). Penis depressed, moderately curved, pedon with apical plate straight in side view, truncate and slightly constricted apicad in side view; temones about $0.50\times$ as long as pedon; endophallus without large structures (Figs 13–14). Tegminal plate articulated with free ring, laterally developed, enveloping, with parameroid lobes not notched apically, membranous area rhombic in general outline, microsetose apically, reaching middle of parameroid lobes, basal sclerotized area large, each side with 5 short macrochaetae, 2 latero-apical and 3 latero-medial, several sensilla distributed on middle part and posterior margin; fenestrae short, transverse, narrowly separated; linea arquata visible; prostegium bidentate, apex of projection rounded; median unsclerotized strip moderately elongate, reaching hind margin of fenestrae; manubrium with apex not broadened (Figs 11–12).

Variation. Male paratypes. Measurements (in mm) ($n=10$): Standard length: 1.38–2.05 (mean= 1.735). Rostrum: length: 0.52–0.82 (mean= 0.722), maximum width: 0.13–0.19 (mean= 0.164). Pronotum: median length: 0.36–0.53 (mean= 0.479), maximum width: 0.44–0.68 (mean= 0.594). Elytra: median length: 1.06–1.54 (mean= 1.406), maximum width: 0.78–1.18 (mean= 1.038).

Rostrum 1.44 – $1.58\times$ as long as pronotum, 4.00 – $4.63\times$ as long as wide. *Ventrite 5* sometimes slightly subtriangular. Otherwise as in holotype.

Female paratypes (Figs 3–4). Measurements (in mm) ($n=10$): Standard length: 1.56–2.05 (mean= 1.831). Rostrum: length: 0.52–0.76 (mean= 0.669), maximum width: 0.130–0.155 (mean= 0.143). Pronotum: median length: 0.37–0.53 (mean= 0.460), maximum width: 0.48–0.65 (mean= 0.571). Elytra: median length: 1.24–1.62 (mean= 1.444), maximum width: 0.88–1.14 (mean= 1.032).


Sexual dimorphism rather weak in this species, females differ from males by a weakly thinner and more evenly curved rostrum with a brighter and less sculptured metarostrium, and all the tibiae without mucros. *Rostrum* 1.35 – $1.53\times$ as long as pronotum, 4.00 – $5.07\times$ as long as wide. *Antennae* inserted at ca. basal 0.27 of rostrum. *Ventrite 5* almost flat, hardly punctured. Otherwise practically as in male.

Material examined. Holotype: ♂: (white, printed): 北京植物园樱桃沟 [Běijīng zhīwùyuán Yīngtáogōu] / 2008.VI.1 / Leg. 王志良 [Wáng Zhiliáng]; (white, printed): IOZ(E)1638667, deposited in IZCAS. This is the Beijing Botanical Garden at Yingtaogou ($40^{\circ} 0' 51.41''$ N $116^{\circ} 12' 14.34''$ E), Haidian district, Beijing (P. R. of China). Paratypes (132♂137♀): 4♂7♀: (white, printed and handwritten): 北京三堡 [Běijīng Sānpù]; (white, printed and handwritten): 1964.VII.20 / leg. 马文珍 [Mǎ Wénzhēn]; (white, printed): IOZ(E)1638668–1638678; 3♀: (white, printed and handwritten):



Figures 16–17. *P. botanicum* Alonso-Zarazaga & Wang sp. n. **16** female resting on underside of leaf **17** female boring a bud for oviposition, with guarding male.

北京三堡 [Běijīng Sānpù] / 600m; (white, printed and handwritten): 1964.VIII.21 / leg. 李铁生 [Lǐ Tiěshēn]; (white, printed): IOZ(E)1638679-1638681; 1♂1♀: (white, printed and handwritten): 北京三堡 [Běijīng, Sānpù] / 1979.VIII.8; (white, handwritten): leg. 廖素柏 [Liào Sùbǎi]; (white, printed): IOZ(E)1638682-1638683; 5♂5♀: (white, printed and handwritten): 北京上方山 [Běijīng Shàngfāngshān]; (white, printed and handwritten): 1980.VIII.2 / leg. 廖素柏 [Liào Sùbǎi]; (white, printed) IOZ(E)1638684-1638693; 2♂: (white, printed and handwritten): 北京上方山 [Běijīng Shàngfāngshān]; (white, printed and handwritten): 1979.VII.25 / leg. 陈元清 [Chén Yuánqīng]; (white, printed): IOZ(E)1638694-1638695; 2♂2♀: (white, printed and handwritten): 北京居庸关 [Běijīng Jūyōngguān] / 500m; (white, printed and handwritten): 1964.VIII.20 / leg. 李铁生 [Lǐ Tiěshēn]; (white, printed): IOZ(E)1638696-1638699; 1♂: (white, printed and handwritten): 北京八达岭 [Běijīng Bādǎlíng] / 700m; (white, printed and handwritten): 1963.VII.25 / leg. 李铁生 [Lǐ Tiěshēn]; (white, printed): IOZ(E)1638700; 8♂8♀: (white, printed): 北京香山 [Běijīng Xiāngshān]; (white, printed and handwritten): 1963.V.30 / leg. 李铁生 [Lǐ Tiěshēn]; (white, printed): IOZ(E)1638701-1638716; 1♀: (white, printed and handwritten): 北京香山 [Běijīng Xiāngshān]; (white, printed and handwritten): 1964.V.5 / leg. 马文珍 [Mǎ Wénzhēn]; (white, printed): IOZ(E)1638717; 1♀: (white, handwritten): 北京香山 [Běijīng Xiāngshān] / 1957.VIII.16; (white, printed): IOZ(E)1638718; 1♂: (white, printed and handwritten): 北京卧佛寺 [Běijīng Wòfósi] / 50m; (white, printed and handwritten): 1962.VI.25 / leg. 王春光 [Wáng Chūnguāng]; (white, printed): IOZ(E)1638719; 1♀: (white, printed and handwritten): 北京卧佛寺 [Běijīng Wòfósi]; (white, printed and handwritten): 1963.IX.3 / leg. 姜胜巧 [Jiāng Shèngqiǎo]; (white, printed): IOZ(E)1638720; 1♀: (white, printed and handwritten): 北京卧佛寺 [Běijīng Wòfósi] / 50m; (white, printed and handwritten): 1962.VIII.31 / leg. 谢汝忠 [Xiè Rǔzhōng]; (white, printed): IOZ(E)1638721; 1♂: (white, printed and handwritten): 北京潭柘寺 [Běijīng Tánzhèsi] / 1975.VII.24; (white, printed and handwritten): leg. 王书永 [Wáng Shūyǒng]; (white, printed): IOZ(E)1638722; 1♀: (white, printed and handwritten): 1964.VII.20 / leg. 马文珍



Figures 18–33. Types of *Pseudaspidapion*, *Aspidapion* and *Harpapion*, in lateral and dorsal views **18–19** *A. panfilovi* Korotyaev, 1985, female holotype **20–21** *P. zagulajevi* (Korotyaev, 1987), female paratype **22–23** *P. kryzhanovskii* (Korotyaev, 1987), female paratype **24–25** *P. yunnanicum* (Korotyaev, 1985), female holotype **26–27** *P. topali* (Korotyaev, 1985), male paratype **28–29** *P. medvedevi* (Korotyaev, 1985), male holotype **30–31** *H. coelebs* (Korotyaev, 1987), male paratype **32–33** *H. vietnamense* (Korotyaev, 1987), male holotype. Scales: 18–33: 1000 μ m.

[Mǎ Wénzhēn]; (white, printed): IOZ(E)1638723; 2♂3♀: (white, printed and handwritten): 北京圆明园 [Běijīng Yuánmíngyuán] / 1980. VII.7 / leg. 姜胜巧 [Jiāng Shèngqiáo]; (white, printed): IOZ(E)1638724-1638728; 2♂10♀: (white, printed): 北京门头沟军庄西杨坨村 [Běijīng Méntóugōu Jūnzhuāng Xiyángtún] / 2008. VII.27 / leg. 王志良 [Wáng Zhìliáng]; (white, printed): 寄主: 小花扁担木 [Jìzhǔ: Xiǎohuābiǎndàn mù] / *Grewia biloba* G. Don var. *parviflora* (Bunge) Hand.-Mazz; (white, printed): IOZ(E)1638729-1638740; 5♂8♀: (white, printed): 北京怀柔九渡河镇怀九河 [Běijīng Huáiróu Jiùdùhézhèn Huáijiùhé] / 2008.VI.15 / leg. 王志良 [Wáng Zhìliáng]; (white, printed): IOZ(E)1638741-1638753; 1♀: (white, printed): 北京怀柔三渡河 [Běijīng Huáiróu Sāndùhé] / 2008.V.24 / leg. 王志良 [Wáng Zhìliáng]; (white, printed) IOZ(E)1638754; 46♂24♀: (white, printed): 北京香山植物园樱桃沟 [Běijīng Xiāngshān Zhíwùyuán Yīngtáogōu] / 2008.VIII.19 / leg. 王志良 [Wáng Zhìliáng]; (white, printed): IOZ(E)1638755-1638770, IOZ(E)1638784-1638790, IOZ(E)1638804-1638850; 28♂41♀: (white, printed): 北京植物园樱桃沟 [Běijīng zhíwùyuán Yīngtáogōu] / 寄主: 小花扁担木 [Jìzhǔ: Xiǎohuābiǎndàn mù] / 2008. VI.1 / Leg. 王志良 [Wáng Zhìliáng]; (white, printed): IOZ(E)1638851-1638900, IOZ(E)1638911, IOZ(E)1638928-1638945; 7♂7♀: (white, printed): 北京海淀百望山 [Běijīng Hǎidiàn Bǎiwàngshān] / 2009.VI.21 / leg. 杨干燕 [Yáng Gānyàn]; (white, printed): IOZ(E)1639912-1638918, IOZ(E)16398920-1638926; 2♂1♀: (white, printed and handwritten): 陕西华阴县孟塬 [Shǎnxī Huáyīn Mèngyuán] / 450m / 1972.VIII.9; (white, printed): IOZ(E)1639562-1639563, IOZ(E)1639540, deposited in IZCAS.

10♂10♀: (white, printed): 北京香山植物园樱桃沟 [Běijīng Xiāngshān Zhíwùyuán Yīngtáogōu] / 2008.VIII.19 / leg. 王志良 [Wáng Zhìliáng]; (white, printed): IOZ(E)1638781, IOZ(E)1638801, IOZ(E)1638771-1638779, IOZ(E)1638791-1638799, to be deposited in MNCN.

1♂1♀: (white, printed): 北京香山植物园樱桃沟 [Běijīng Xiāngshān Zhíwùyuán Yīngtáogōu] / 2008.VIII.19 / leg. 王志良 [Wáng Zhìliáng]; (white, printed): IOZ(E)1638780, IOZ(E)1638800, to be deposited in BBG.

1♂1♀: (white, printed): 北京香山植物园樱桃沟 [Běijīng Xiāngshān Zhíwùyuán Yīngtáogōu] / 2008.VIII.19 / leg. 王志良 [Wáng Zhìliáng]; (white, printed): IOZ(E)1638783, IOZ(E)1638803, to be deposited in ZINM.

1♂1♀: (white, printed): 北京香山植物园樱桃沟 [Běijīng Xiāngshān Zhíwùyuán Yīngtáogōu] / 2008.VIII.19 / leg. 王志良 [Wáng Zhìliáng]; (white, printed): IOZ(E)1638782, IOZ(E)1638802, to be deposited in MNHW.

The male holotype has not been dissected to avoid any damage, since there are many specimens collected with it.

Etymology. The new species is named after the first locality where it was found during a collecting visit: the Beijing Botanical Garden. It is a Latin adjective.

Distribution. The species is known for the moment only from the municipality of Beijing and the province of Shaanxi.

Biology. *P. botanicum* sp. n. was collected from *Grewia biloba* G. Don var. *parviflora* (Bunge) Hand.-Mazz (Malvaceae: Grewioideae) (Stevens 2011), which is a com-

mon shrub in Beijing and adjacent provinces, blooming from June to July (Tang et al. 1989), this being the season in which the parasite was found on the plant. The adults feed on leaves and flower buds of their host, while they mate and oviposit in the bud. The egg is located in the androecium, and then the larva feeds on the pistil and pupates there.

Discussion

During this research, we were able to study types of several of the species placed in the genera *Pseudaspidapion* and *Aspidapion* in China in the recent Palaearctic catalogue (Alonso-Zarazaga 2011) (table 1). Until now, females of *P. coelebs* and *A. vietnamense* are unknown, and the same can be said of the males of *P. kryzhanovskii* and *A. panfilovi*. We found that the holotype (and only known specimen) of *Aspidapion vietnamense*, which has been previously dissected and whose pygidium and tegmen were not conserved, shows several differences from *Aspidapion*, namely, the metatibial mucros are evidently elongate and knicked at their apices (subdentate at the outer margin), the rostrum is clearly dilated at the antennal insertion and distinctly constricted apicad, the setae on the front margin of pronotum are parallel to it, and the apex of the penis is distinctly curved in lateral view, dorsally dentate near the apex. All these characters suggest it is a *Harpapion* Voss, 1966. After a study of the type species of this genus, *Harpapion considerandum* (Fåhraeus, 1871), we consider that, even in the absence of these diagnostic parts, it can be unmistakably considered a member of this genus and consequently is here transferred: *Harpapion vietnamense* (Korotyaev), comb. n. The same can be said about *Pseudaspidapion coelebs* (Korotyaev, 1987), which show also the characters of the latter genus, and is here formally transferred as well: *Harpapion coelebs* (Korotyaev, 1987), comb. n. This genus will be the subject of a forthcoming paper.

The systematic placement of *A. inarmatum* is unclear at the moment, but it belongs neither to *Aspidapion* nor to *Pseudaspidapion* or *Harpapion*.

Aspidapion panfilovi, known from females until now, possesses several coincident characters with some *Pseudaspidapion*: its more elongate and apparently glabrous elytra, the shallow striae with interstriae about twice as broad as striae on the disc of the elytra, etc. Although it resembles the female of *P. zagulajevi* very much, we think it is better to postpone any decision on its placement until the male is discovered. This highlights the difficulty of placing in their correct genera the species of Apionidae described only from females, a practice that should be avoided, as that of designating females as name bearing types.

Previously, the genus *Pseudaspidapion* was not known to exceed the Tropic of Cancer. However, the new species *P. botanicum* reaches 40°N latitude, an area clearly belonging to the Eastern Palaearctic region (Fig. 34). Thus this genus shows a broader distributional range extending from the Afrotropical across the Oriental to the Palaearctic region. On the other hand, we identified in the collection of IZCAS several specimens of *Aspidapion validum* (Germar, 1817), from Altay prefecture in northern Xinjiang




Figure 34. Distribution of the species of *Pseudaspidapion*, *Harpation* and *Aspidapion* mentioned in the text.

(China). Apart from the species *Aspidapion panfilovi*, whose placement may need a revision as mentioned above, these specimens represent the easternmost distribution record of the otherwise Western Palearctic genus *Aspidapion*. Therefore, the distribution pattern of *Pseudaspidapion* is mainly Palearctic but has expanded to Philippines (Wanat 1990) and to North China while *Aspidapion* is distributed in the Palearctic region but has also expanded to northeast tropical Africa (Eritrea, Ethiopia). The two genera can still be distinctly separated from each other. Also, we suppose that the blank distributional area of *Pseudaspidapion* between Shaanxi and South China represents a lack of collection records rather than an actual distributional range limit, because the vegetation between these two points includes most of the range of *G. biloba* (which includes also Korea) and many other congeneric species of the host genus.

Some of these species are described only from one sex, which makes it difficult to key all of the Chinese species. However, we have tried to provide a key to the known Chinese species (see below). Consequently, more field work is needed to gather specimens of both sexes, and complete our knowledge. We have not studied specimens of *P. rufopiceum* (Wagner, 1909), a related species, which is included below from the data in its original description.

Key to the Chinese species of *Pseudaspidapion*

This key includes also *Aspidapion panfilovi* Korotyaev, 1985 (only known from female), which seems to be related to these species. Body length excludes rostrum.

- 1 Meso- and metatibiae mucronate. (males).....**2**
- Meso- and metatibiae unarmed. (females).....**4**
- 2 Body length 3.1–3.6 mm. Elytral interstitial scales minute, shorter than 1/3 width of interstria, subtransparent, elytra appearing glabrous... ***P. zagulajevi***
- Body length 1.3–2.2 mm. Elytral interstitial scales larger, longer than 2/3 of the interstitial width, white or dark brown, elytra appearing pubescent.....**3**
- 3 Body length in general smaller: 1.3–1.6 mm. Body dark reddish-brown, appendages light reddish-brown. Pedicel little longer than wide...***P. rufopiceum***
- Body length in general larger: 1.5–2.2 mm. Body and appendages dark brown to piceous black. Pedicel at least 2 × as long as wide.....
.....***P. yunnanicum* / *P. botanicum* sp. n.** (see Table 2 for separation)
- 4 Body length more than 3.7 mm. Rostrum ca. 2.5 × as long as pronotum. Elytra short, rounded, in side view strongly convex. Scutellum large, base about as wide as interocular distance, triangular, sides straight; in side view completely prominent above elytral outline ***P. kryzhanovskii***
- Body length less than 3.7 mm. Elytra oblong-elliptical, in side view moderately convex. Scutellum small, base at most 2/3 as wide as interocular distance, sides concave, forming a basal subpentagonal area prolonged apically in a shaft; in side view, at most some of the tubercles prominent above elytral outline **5**
- 5 Body length more than 2.5 mm. Elytral interstitial scales minute, shorter than 1/3 width of interstria, subtransparent, elytra appearing glabrous.....**6**
- Body length less than 2.5 mm. Elytral interstitial scales larger, longer than 2/3 width of interstria, white or dark brown, elytra appearing pubescent **3**
- 6 Rostrum ca. 1.8 × as long as pronotum, robust, mesorostrum wider than interocular distance. Basal tubercles of scutellum large, acutely prominent in side view. Body length: 3.1–3.6 mm ***P. zagulajevi***
- Rostrum ca. 2.0 × as long as pronotum, fine, mesorostrum at most as wide as interocular distance. Basal tubercles of scutellum small, hardly visible in side view. Body length: 2.7 mm ***A. panfilovi***

Acknowledgements

We thank Marek Wanat for kindly providing us references and many good advices, Boris A. Korotyaev for lending paratypes, Steven R. Davis for doing the linguistic review before submission, Yang Ganyan for collecting specimens, Zhou Dakang for providing information about the host plants, Yin Ziwei and Nico M. Franz for pro-

viding the brush-pencil Plug-ins of Adobe Illustrator. We also thank two anonymous reviewers whose suggestions considerably improved our work. This research has been supported by an Invited Professor Award (2009) of the Chinese Academy of Sciences to the senior author and by project grant CGL2010–15786 (Ministerio de Ciencia e Innovación, Spain).

References

- Alonso-Zarazaga MA (1989) Revision of the supraspecific taxa in the Palearctic Apionidae Schoenherr, 1823. 1. Introduction and subfamily Nanophyinae Seidlitz, 1891 (Coleoptera, Curculionoidea). *Fragmenta Entomologica*, Roma 21(2): 205–262.
- Alonso-Zarazaga MA (1990) Revision of the supraspecific taxa in the Palearctic Apionidae Schoenherr, 1823 (Coleoptera, Curculionoidea). 2. Subfamily Apioninae Schoenherr, 1823: introduction, keys and descriptions. *Graellsia* 46: 19–156.
- Alonso-Zarazaga MA (2011) Apionidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palearctic Coleoptera*. Vol. 7: Curculionoidea I. Apollo Books, Stenstrup, 373 pp.
- Alonso-Zarazaga MA, Lyal CHC (1999) A world catalogue of families and genera of Curculionoidea (Insecta: Coleoptera) (excepting Scolytidae and Platypodidae). Barcelona: Entomopraxis, 315 pp.
- Bameul F (1990) Le DMHF: un excellent milieu de montage en entomologie. *L'Entomologiste* (Paris) 46(5): 233–239.
- Korotyaev BA (1985) Novye vidy dolgonosikov podsemejstva Apioninae (Coleoptera, Apionidae) iz V'etnama, Kitaâ i Indii. In: Medvedev LN (Ed) *Nasekomye V'etnama*. Nauka, Moskva, 133–156.
- Korotyaev BA (1987) Novye vidy dolgonosikov podsemejstva Apioninae (Coleoptera, Apionidae) iz tropičeskikh i subtropičeskikh rajonov Azii. In: Medvedev LN (Ed) *Entomofauna V'etnama*. Nauka, Moskva, 94–120.
- Steedman HF (1958) Dimethyl Hydantoin Formaldehyde: a new water-soluble resin for use as a mounting medium. *Quarterly Journal of Microscopical Science* 99(4): 451–452.
- Stevens PF (2011). *Angiosperm Phylogeny Website*. Version 9, June 2008. <http://www.mobot.org/MOBOT/research/APweb/> [accessed 20.IV.2011]
- Tang Y, Gilbert MG, Dorr LJ (2007) Tiliaceae. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China*. Vol. 12 (Hippocastanaceae through Theaceae). Science Press, Beijing; Missouri Botanical Garden Press, St. Louis, 240–263.
- Wagner H (1909) Fünf neue Apioniden der alten Welt. *Societas Entomologica* 24(4): 25–28.
- Wanat M (1990) Apionidae (Coleoptera, Curculionoidea) of the Arabian Peninsula. *Fauna of Saudi Arabia* 11: 55–81.