

Nutrient enrichment effect on macroinvertebrates in a lowland stream of Argentina

Agustina Cortelezzi^{1,3}, Carolina Ocón^{2,3}, María V. López van Oosterom^{2,3},
Rosana Cepeda¹ & Alberto Rodrigues Capítulo^{2,3}

1. Instituto Multidisciplinario sobre Ecosistemas y Desarrollo Sustentable, UNCPBA, Paraje Arroyo Seco s/n, Tandil, Argentina. (aguscorle@gmail.com)
2. Instituto de Limnología Dr. Raúl A. Ringuelet (ILPLA), CCT La Plata – CONICET, UNLP, Bv. 120 s/n, La Plata, Argentina.
3. Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

ABSTRACT. One of the most important effects derived from the intensive land use is the increase of nutrient concentration in the aquatic systems due to superficial drainage. Besides, the increment of precipitations in South America connected to the global climate change could intensify these anthropic impacts due to the changes in the runoff pattern and a greater discharge of water in the streams and rivers. The pampean streams are singular environments with high natural nutrient concentrations which could be increased even more if the predictions of global climate change for the area are met. In this context, the effect of experimental nutrient addition on macroinvertebrates in a lowland stream is studied. Samplings were carried out from March 2007 to February 2009 in two reaches (fertilized and unfertilized), upstream and downstream from the input of nutrients. The addition of nutrients caused an increase in the phosphorus concentration in the fertilized reach which was not observed for nitrogen concentration. From all macroinvertebrates studied only two taxa had significant differences in their abundance after fertilization: *Corbicula fluminea* and Ostracoda. Our results reveal that the disturbance caused by the increase of nutrients on the benthic community depends on basal nutrients concentration. The weak response of macroinvertebrates to fertilization in the pampean streams could be due to their tolerance to high concentrations of nutrients in relation to their evolutionary history in streams naturally enriched with nutrients. Further research concerning the thresholds of nutrients affecting macroinvertebrates and about the adaptive advantages of taxa in naturally eutrophic environments is still needed. This information will allow for a better understanding of the processes of nutrient cycling and for the construction of restoration measures in natural eutrophic ecosystems.

KEYWORDS. Phosphorus, nitrogen, fertilization, invertebrate abundance, climate change.

RESUMEN. Efecto del enriquecimiento de nutrientes en macroinvertebrados en un arroyo pampeano de Argentina. Uno de los efectos más importantes derivados del uso intensivo de la tierra es el aumento de la concentración de nutrientes en los sistemas acuáticos debido al escurrimiento superficial. Además, el incremento de las precipitaciones en América del Sur vinculado al cambio climático global podría intensificar estos impactos antrópicos debido a los cambios en los patrones de escurrimiento y una mayor descarga de agua en los arroyos y ríos. Los arroyos pampeanos son ambientes particulares, con altas concentraciones basales de nutrientes, que podrían incrementarse aún más si las predicciones del cambio climático global se cumplen. En este contexto, se estudió el efecto de la adición experimental de nutrientes en un arroyo pampeano sobre los macroinvertebrados. Los muestreos se llevaron a cabo entre marzo de 2007 y febrero de 2009 en dos tramos seleccionados, aguas arriba y aguas abajo de la entrada de nutrientes. La adición de nutrientes causó un aumento en la concentración de fósforo en el tramo fertilizado, lo que no se observó para la concentración de nitrógeno. De todos los macroinvertebrados estudiados sólo dos taxa tuvieron diferencias significativas en su abundancia después de la fertilización: *Corbicula fluminea* y ostrácodos. Nuestros resultados revelaron que la perturbación causada por el aumento de nutrientes en una comunidad bentónica depende de la concentración basal de nutrientes. La respuesta débil de macroinvertebrados a la fertilización en las corrientes pampeanas podría ser debido a su tolerancia a altas concentraciones de nutrientes de acuerdo a su historia evolutiva en arroyos enriquecidos naturalmente con nutrientes. Se necesitan nuevas investigaciones sobre los umbrales a partir de los cuales los macroinvertebrados podrían verse afectados y sobre las ventajas adaptativas de los taxones en ambientes eutróficos naturales. Esta información nos permitirá comprender mejor los procesos de reciclaje de nutrientes y así poder pensar medidas de restauración de ecosistemas eutróficos naturales.

PALABRAS CLAVE. Fósforo, nitrógeno, fertilización, abundancia de invertebrados, cambio climático.

Given that benthic macroinvertebrates represent an important fraction of stream biodiversity, they are frequently used as indicators in aquatic environments and their assemblage composition and richness exhibits a strong correlation with environmental change at reach and basin levels. However, while tolerance levels of a wide range of taxa have been established along gradients of organic pollution, little is known about how macroinvertebrates respond to elevated nutrient levels (FRIBERG *et al.*, 2010). While nutrient studies in lowland streams showed impacts on primary producers (ACUÑA *et al.*, 2011) and on benthic microbial community (COCHERO *et al.*, 2013), researches showing direct and indirect effects on higher trophic levels have been less conclusive due to the complexity of food webs that can buffer the potential transference of bottom-up effects after nutrient addition (ARTIGAS *et al.*, 2013).

Besides, most studies report changes in macroinvertebrates because the effect of increased nutrients has been conducted primarily in oligotrophic aquatic systems (BIGGS *et al.*, 2000; ROBINSON & GESSNER, 2000).

The pampean area of Buenos Aires Province (Argentina) holds the highest demographic and industrial concentration of the country, the greatest agriculture and livestock production as well as the most intense use of agrochemicals, as a result of the great expansion of agriculture in the last century (SALAZAR *et al.*, 1996; GOMEZ & RODRIGUES CAPÍTULO, 2001). Because of such intensive human activity, many rivers and streams are impacted by diffuse pollution associated with crop cultivation and cattle raising (RODRIGUES CAPÍTULO *et al.*, 2010). However, high nutrient levels have been reported for these streams even in sites with low or moderate cattle-raising and agriculture

(FEIJOÓ *et al.*, 1999; BAUER *et al.*, 2002). The occurrence of high nutrient concentrations is characteristic of pampean streams (FEIJOÓ & LOMBARDO, 2007). Indeed, nutrient levels in stream water show a relatively high concentration when compared with other lotic systems of the world (FEIJOÓ *et al.*, 2014). Moreover, pampean streams can be classified as eutrophic when phosphorus is considered and as meso-eutrophic when nitrogen is taken into account (BARBOUR *et al.*, 1999). This high natural nutrient loading of the Pampean streams, could be further increased if predictions of global climate change for the area are met. Thus, changes in the runoff patterns and a greater discharge of water in the streams and rivers (HULME & SHEARD, 1999) would lead to a greater input of nutrients, especially nitrogen and phosphorous, and contaminants into streams and rivers (RODRIGUES CAPÍTULO *et al.*, 2010; FERREIRO *et al.*, 2011). This situation may have harmful effects on the whole ecosystem with accompanying economic repercussions (DODDS *et al.*, 2009; ARTIGAS *et al.*, 2013).

As part of the GlobRio project (“Global changes in river systems: effects on biodiversity, food web and system functioning”) our main objective was to explore experimentally the effect of increased nutrient concentrations on the parameters of macroinvertebrate assemblages in a pampean stream. We also examined the temporal variability of the community descriptors over a study period. We hypothesized that density and dominance of macroinvertebrates would increase in the reach under nutrient enrichment while taxa richness would decrease.

MATERIAL AND METHODS

Study area. Pampean streams run through a vast grassy plain that shelters central Argentina, denominated the

Pampas. The pampean steppe is composed of gramineous grasses, which is the typical biome of the area (CABRERA, 1976). The climate is humid and temperate with a mean annual temperature of 20 °C and a mean precipitation of 900 mm/year. La Choza stream is a 2nd order stream located in the Pampas, Buenos Aires Province, Argentina (Fig. 1). Like most lotic systems of the region, it is characterized by a low slope (<1%), a slow flow velocity, a high turbidity, abundant organic matter, and an elevated content of suspended solids (GIORGI *et al.*, 2005). The streambed consists of fine sediments (primarily silt and clay) without stones or pebbles (FEIJOÓ *et al.*, 2014). Besides, even in the upper-stream reaches, the absence of riparian vegetation, results in high levels of irradiance along the course, favoring the development of dense macrophyte stands and phitoperiphyton growth (FEIJOÓ & LOMBARDO, 2007). Moreover, the heterogeneity of the pampean streams, resides not in the different types and sizes of the substrates present but rather in the diverse architecture of the aquatic plants which shelter a rich and dense invertebrate community (GIORGI *et al.*, 2005).

La Choza watershed has a surface of 15,200 ha and features intense agricultural activity, with 64% of the area for cattle raising, 34% for extensive agriculture, and the remaining 2% divided between poultry farming and different other urban uses (RODRIGUES CAPÍTULO *et al.*, 2010).

Experimental design. Two reaches of 100 m in length were selected, the first site was located upstream (control site, named “C” in Fig. 1: 34°39’14”S, 59°10’00”W), and the second downstream from the experimental input of nutrients (treatment site, named “T” in Fig. 1: 34°47’51”S, 58°03’17”W). The distance from the source is 2.5 km for control site and 7.5 km for the treatment site. Bimonthly




Fig. 1. Map of the study area showing La Choza stream, Buenos Aires Province, Argentina and the location of the sampling stations: control (C) and treatment (T).

samplings, from March 2007 through February 2009, took place at each site. On each sampling date the following physicochemical parameters were measured: temperature, pH, conductivity, dissolved-oxygen level (DO) with a Hach HQ40d18 portable meter (Hach, Loveland, Colorado) and water samples (250 ml) in triplicate were collected and filtered through glass-fiber filters of 0.7 µm pore size (Whatman GF/F, Whatman International Ltd., Maidstone, England). The whole filtrate samples were analyzed for dissolved inorganic nitrogen (DIN) and soluble reactive phosphorus (SRP) by standard methods (APHA, 1998). The process of experimental fertilization initiated at the end of October 2007 and involved a continuous dissolution of nutrients at the treatment site of the stream. Due to the low water velocity, to obtain a homogeneous fertilization along the reach, nutrient additions consisted of mesh bags with 750 g of commercial fertilizer Compo Nitrofoska® (12% nitrogen, 12% phosphorus, 17% potassium), a mixture of frequent use in farming, and 250 g of urea. Twelve in-water bags were distributed along the treatment reach, and replaced 2 or 3 times a week during the fertilization period to maintain a constant fertilization rate (FEIJÓ *et al.*, 2014). Water samples in triplicate also were collected and analyzed during enrichment period every 5 to 10 d to better capture changes in nutrient concentrations (ARTIGAS *et al.*, 2013).

Macroinvertebrate sampling. Macroinvertebrates were collected following RODRIGUES CAPÍTULO *et al.* (2009) and PLAFKIN *et al.* (1989). At each site, three replicates of benthic macroinvertebrates were collected with an Ekman grab (100 cm²). The samples were fixed in situ in 5% formaldehyde and the organisms sorted, identified and counted under a stereomicroscope following FERNÁNDEZ & DOMÍNGUEZ (2001) taxonomic keys. The invertebrates were identified at the lowest possible level.

Data analysis. The physicochemical variables that may be affected by fertilization in the control and treatment sites were analyzed through descriptive statistics. In addition, macroinvertebrates density was described during the study period. Dominance index (Simpson), diversity (Shannon) and richness (such as number of taxa) were evaluated to describe the structure of the community at each site. The variation of these metrics on sites (spatial variation) and along the study (temporal variation) was tested by repeated measures ANOVA. We consider two situations: comparing the control and treatment site (C1 vs T1) in 4 dates before fertilization and comparison between control and treatment in 6 dates after fertilization (C2 vs T2).

Repeated measures ANOVA were also used to evaluate time and fertilization effect for the density of each taxon. Although this method is commonly used in experimental studies of nutrient fertilization, in our case it was not useful due to the taxa abundance showing a high variation between dates even before the start of fertilization. It is why we decided to use this information to perform a Student t test. For each taxon the dates as independent

replicates in two moments before (b) and after fertilization (a) were used. If the abundance taxa was statistically the same in treatment site and control site before fertilization between dates ($p > 0.05$), and after fertilization this situation changed ($p < 0.05$), at that time the differences detected could be due to treatment. Similarly, if before treatment, the abundance was variable between sites ($p < 0.05$) and after the fertilization the abundances are similar ($p > 0.05$) that is, they stabilize; might it be an effect of fertilization. After this analysis, we described variations of the abundance of taxa that presented differences before and after fertilization using average graphs by date and sites. These values were presented on a logarithmic scale. Analyses were performed with InfoStat statistical software (DI RIENZO *et al.*, 2008). We considered significative analyzes at $p < 0.05$.

RESULTS

The main physical and chemical characteristics of the sampling stations before and after fertilization of La Choza stream are shown in Table I. In general, conductivity, temperature and pH were similar before and after fertilization. The values of DO in both control and treatment sites remained stable, however after fertilization a decrease of this parameter was recorded in the treated site. The experimental fertilization increased the average SRP concentration by about 5-fold in water column of treatment site above the natural value. However, the nutrient addition in treatment reach slightly altered the average DIN levels. At the beginning of fertilization a slight increase in the values of DIN was recorded which then decreased and remained low in the following samplings.

In the present study 25 taxa were identified, 23 of which were common to both sites, while the remaining 2 (*Perithemis* sp. and *Belostoma* sp.) were found only at the treatment site (Tab. II). In the control site, the total abundance was 70,345 ind/m². In the treatment site, the total invertebrate abundance decreased from 181,466 to 100,722 ind/m² after fertilization. In relation to community composition, the control site was dominated by *Heleobia* (25.3%), *Oligochaeta* (21.8%) and *Chironomidae* (19.3%). In the treatment site, before fertilization, the community was dominated by oligochaetes (75.5%), and after fertilization oligochaetes decreased their abundance (37.2%), whereas *Heleobia* sp. (24.8%) and chironomids (9.6 %) increased their representation resembling in composition to the control site.

Most of the metrics analyzed (Shannon, Simpson, and total density of macroinvertebrates) show a significant interaction between time and treatment but not between the site and time throughout the study. Metrics along sampling dates do not show the same performance in control and treatment sites, that is, there are no significant interaction between sites and time. There are significant differences between dates from the same site and between sites on the same date partitioning the study before and after fertilization. In the case of richness, the interaction also was

Tab. I. Physicochemical parameters and nutrient concentration in the control and treatment reaches of La Choza stream, Buenos Aires Province, Argentina. Values are mean and standard deviations measured before and after fertilization.

Parameters	Control		Treatment	
	Before	After	Before	After
Conductivity (uS/cm)	866.66 (511.8)	1272.66 (101.53)	1748.44 (500.62)	1159.58 (96.48)
T° (°C)	13.97 (7.88)	19.99 (7.57)	13.84 (6.01)	19.22 (8.31)
DO (mg/l)	9.8 (5.04)	9.47 (3.86)	9.85 (4.12)	8.58 (2.25)
pH	7.37 (0.77)	8.27 (0.44)	7.60 (0.41)	8.09 (0.29)
DIN (ug/l)	650.75 (435.21)	925.72 (1182.72)	884.78 (572.41)	883.26 (985.37)
SRP (ug/l)	205.62 (178.96)	93.47 (25.41)	247.59 (136.59)	431.60 (246.08)

Tab. II. Mean density and standard deviation of the taxa present in La Choza stream, Buenos Aires Province, Argentina in control and treatment reaches.

	Control	Treatment	
	(n=30)	Before (n=12)	After (n=18)
Hirudinea	570 (242)	142 (50)	494 (284)
Nematoda	5232 (1502)	2150 (1479)	2106 (532)
Oligochaeta	15304 (4158)	137094 (84970)	37517 (12729)
<i>Pomacea canaliculata</i>	68 (20)	33 (26)	172 (92)
<i>Heleobia parchappei</i>	17814 (5131)	12983 (4882)	24944 (4653)
<i>Hebetancylus moricardi</i>	276 (108)	233 (143)	161 (84)
<i>Drepanotrema kermatoides</i>	118 (111)	167 (104)	6 (6)
<i>Pisidium vile</i>	13 (13)	8 (8)	156 (134)
<i>Pisidium sterkianum</i>	8350 (4925)	283 (122)	517 (220)
<i>Corbicula fluminea</i>	360 (333)	219 (98)	6328 (2123)
Elmidae	42 (29)	1636 (815)	2233 (822)
Hydrophilidae	8 (5)	0	61 (56)
Coenagrionidae	8 (7)	75 (66)	72 (46)
<i>Perithemis</i> sp.	0	1633 (868)	0
<i>Caenis</i> sp.	3541 (1104)	1794 (795)	5133 (1571)
<i>Americabaetis</i> sp.	12 (8)	242 (166)	161 (155)
<i>Belostoma</i> sp.	0	7900 (5631)	267 (267)
<i>Hyallela curvispina</i>	141 (69)	144 (102)	339 (138)
<i>Aegla</i> sp.	3 (3)	50 (42)	56 (27)
Chironomidae	13608 (7765)	9805 (5252)	9650 (3594)
Ceratopogonidae	1215 (996)	69 (45)	228 (64)
Ostracoda	289 (96)	267 (205)	5861 (1474)
Cladocera	1025 (308)	4233 (2370)	1067 (594)
Copepoda	2111 (642)	136 (75)	3083 (1385)
Acari	237 (222)	167 (104)	111 (52)

significant between time and treatment before treatment. However, after fertilization, richness was always higher in the sites treated than in the control site.

Although the general fertilization effects were weak, some specific invertebrate taxa were distinctly affected in abundance. Certain taxa such as *Oligochaeta*, *Pomacea canaliculata*, *Hebetancylus moricardi*, *Pisidium sterkianum*, *Corbicula fluminea* and *Ostracoda* showed significant differences in abundance at the sampling sites before and after fertilization (Tab. III). However, when comparing the control and treatment sites by dates, before and after fertilization, only *Corbicula fluminea* and *Ostracoda* showed an increase in abundance after fertilization (Fig. 2). For the rest of the taxa the differences could not be attributed to fertilization.

DISCUSSION

In this study, the experimental addition of phosphorus and nitrogen to a pampean stream caused the SRP concentration to be slightly higher in the fertilized reach, as expected, although this was not observed for DIN.

Tab. III. Results of the Student's t test for taxa abundance in the treatment reach site before (b) and after (a) fertilization, La Choza stream, Buenos Aires Province, Argentina.

Taxa	Treatment	p
<i>Oligochaeta</i>	b	0.4963
	a	0.0063
<i>Pomacea canaliculata</i>	b	0.041
	a	0.444
<i>Hebetancylus moricardi</i>	b	0.037
	a	0.531
<i>Pisidium sterkianum</i>	b	0.031
	a	<0.0001
<i>Corbicula fluminea</i>	b	0.492
	a	<0.0001
<i>Ostracoda</i>	b	0.498
	a	<0.0001

The dissolved nitrogen added to the stream could have been either rapidly assimilated by algae, macrophytes, or bacteria, or could have left the system by denitrification processes caused by anaerobic bacteria in anoxic conditions (COCHERO *et al.*, 2013). According to LIN & WEBSTER (2012) all these organisms can quickly take up nutrients from the water column and use these nutrients for production.




Fig. 2. Taxa abundance by date in control and treatment sites in La Choza stream, Buenos Aires Province, Argentina during 2007-2008. The dotted line indicates the time when fertilization begins.

In the two reaches studied, the invertebrate assemblage composition was similar although the abundance of taxa presents differences being lower in the control site. In the treatment site, the invertebrate density after fertilization decreased. Moreover, most of the metrics analyzed (Shannon, Simpson, and total density of macroinvertebrates) did not show a significant interaction between site and time throughout the study. These results indicate that neither diversity nor dominance and total macroinvertebrate density were affected by fertilization. This result contradicts our initial hypothesis that density and dominance of macroinvertebrates would increase in the

reach under nutrient enrichment. In coincidence, SABATER *et al.* (2005) addressed that macroinvertebrate densities in the control reach of two mediterranean streams were similar to those in the impacted reach before an experimental fertilization took place and the observed effects were noticeable only for particular taxa. These authors found more significant variations for certain Oligochaeta and Mollusca species. In La Choza stream the selected treatment site also showed high dominance of oligochaete. However, after fertilization, the oligochaete density decreased significantly allowing greater representation of other taxa. According to ARMENDARIZ *et al.* (2012) a shift

in Oligochaeta composition in favor of herbivores and detritivores and the replacement of sensitive species of the community could occur in nutrient-enriched systems, resulting in a greater tolerance of the overall population to environmental stress.

Our hypothesis about the taxa richness decreasing in the fertilized reach is rejected because after fertilization this metric has always been higher in the sites treated than in the control site. In this sense, McCORMICK *et al.* (2004) did not register variations for species richness between control and fertilized sites in an experimental study carried out in Florida (USA). ARTIGAS *et al.* (2013) did not detect reduced species richness but a change in the dominance of taxa in favor of the more tolerant.

In La Chozá stream, only 2 of 24 taxa of macroinvertebrates showed a response that could be attributable to fertilization. The enrichment favored taxa more tolerant to the new conditions and therefore increased their relative abundances. This was the case for *C. fluminea* and Ostracoda that increased their abundances, while Oligochaeta showed an increase but not sustained after addition. ARMENDÁRIZ *et al.* (2012) studied the response of Oligochaeta taxa but at family and species levels in the same lotic system. These authors found that abundance, biomass, diversity, evenness and richness increased significantly in the treatment site after fertilization in the sediments and in the aquatic vegetation, especially between Naidinae and Pristininae.

Corbicula fluminea increased significantly its abundance after fertilization. This clam is one of the most invasive species in freshwater aquatic ecosystems. The rapid growth, earlier sexual maturity, short life span, high fecundity and its association with human activities makes *C. fluminea* a non-indigenous invasive species likely to colonize new environments (SOUSA *et al.*, 2008). This adaptive advantage over the other species could generate a rapid response to changes in the environment caused by the nutrient addition. Moreover, *C. fluminea* is an important coupler between the water column and the stream sediments because this bivalve dominates filter-feeding in this stream and also pedal-feeding on buried organic material. OCÓN *et al.* (2013) suggests that the increase in P concentration may have caused an increase in phytoplankton density on treatment site, making more resources available to filter feeders, which increased in proportion after fertilization. According to COHEN *et al.* (1984), this species contributed to clarify the water bodies, thus decreasing the density of phytoplankton. Future studies are needed to explore the response of introduced species on the dynamics of the nutrients of the Pampean ecosystems.

MURKIN *et al.* (1994) found no significant differences in numbers, biomass of total invertebrates, and invertebrate functional groups in a similar experience in Canadian wetlands. These authors concluded that the nutrient additions did increase community productivity; however, the levels used in their study were insufficient to yield a sustained increase in primary or secondary

productivity. According to CROSS *et al.* (2007) the lack of a response of the macroinvertebrates assemblage could be due to the fact that a longer period of fertilization would be needed. These authors concluded that the effects on community composition require at least three years of continuous fertilization in order to be observed. However, effects could be seen in the short term when it comes to sensitive taxa, especially in environments with low nutrient concentration (SLAVIK *et al.*, 2004; SABATER *et al.*, 2011) or assemblages of primary producers (GÓMEZ *et al.*, 2009) and bacterial biomass (COCHERO *et al.*, 2013) in eutrophic environments. Finally, ARTIGAS *et al.* (2013) showed that the pampean stream communities were less affected by enrichment than those of the Andean and Mediterranean streams despite wide windows of opportunity (periods when environmental constraints other than nutrients do not limit biomass growth).

In this study, the weak response of macroinvertebrates to fertilization in the pampean streams could be due to their tolerance to high concentrations of nutrients in accordance with their evolutionary history in streams naturally enriched with nutrients. We consider that in the Pampean streams the effect of nutrient enrichment may be less predictable in the short term due to their high basal nutrient concentration, in agreement with ARTIGAS *et al.* (2013). We suggest that further research should be focused on the study of the limits from which macroinvertebrates may be affected and on the adaptive advantage of taxa in naturally eutrophic environments. This information will allow for a better understanding of the processes of nutrient cycling and for the construction of restoration measures in natural eutrophic ecosystems.

Acknowledgments. This study was supported by BBVA Foundation's GlobRio Project and CONICET. This is Scientific Contribution n° 946 from the Instituto de Limnología "Dr. Raúl A. Ringuet" (CONICET La Plata-UNLP). We wish to thank Mónica Caviglia for editing the final version of the manuscript.

REFERENCES

- ACUÑA, V.; VILCHES, C. & GIORGI, A. 2011. As productive and slow as a stream can be: the metabolism of a Pampean stream. **Journal of the North American Benthological Society** 30:71-83.
- APHA. 1998. **Standard methods for the examination of water**. 20ed. Washington D.C., American Public Health Association. 1134p.
- ARMENDÁRIZ, L.; OCON, C. & RODRIGUES CAPÍTULO, A. 2012. Potential responses of oligochaetes (Annelida, Clitellata) to global changes: Experimental fertilization in a lowland stream of Argentina (South America). **Limnologia** 42:118-126.
- ARTIGAS, J.; GARCÍA BERTHOU, E.; BAUER, D. E.; CASTRO, M. I.; COCHERO, J.; COLAUTTI, D.; CORTELEZZI, A.; DONATO, J.; ELOSEGI, A.; FEJOÓ, C.; GIORGI, A.; GÓMEZ, N.; LEGGIERI, L.; MUÑOZ, I.; RODRIGUES CAPÍTULO, A.; ROMANÍ, A. M. & SABATER, S. 2013. Global pressures, specific responses: effects of nutrient enrichment in streams from different biomes. **Environmental Research Letters** 8:1-13.
- BARBOUR, M. T.; GERRITSEN, J.; SNYDER, B. D. & STRIBLING, J. B. 1999. **Rapid bioassessment protocols for use in streams and wadeable rivers: periphyton, benthic macroinvertebrates and fish**. 2ed. Washington, D.C., U.S. Environmental Protection Agency, Office of Water. 197p.
- BAUER, D. E.; DONADELLI, J.; GÓMEZ, N.; LICURSI, M.; OCON, C.; PAGGI, A. C.; RODRIGUES CAPÍTULO, A. & TANGORRA, M. 2002.

- Ecological status of the Pampean plain streams and rivers (Argentina). **Verhandlungen des Internationalen Verein Limnologie** 28:259-262.
- BIGGS, B. J. F.; FRANCOEUR, S. N.; HURY, A. D.; YOUNG, R.; ARBUCKLE, C. J. & TOWNSEND, C. R. 2000. Trophic cascades in streams: effects of nutrient enrichment on autotrophic and consumer benthic communities under two different fish predation regimes. **Canadian Journal of Fisheries and Aquatic Sciences** 57:1380-1394.
- CABRERA, A. L. 1976. Regiones Fitogeográficas Argentinas. 2ed. **Enciclopedia Argentina de Agricultura y Jardinería**. Tomo II, Fasc. 1. Buenos Aires, ACME. 85p.
- COCHERO, J.; ROMANÍ, A. & GÓMEZ, N. 2013. Delayed response of microbial epipellic biofilm to nutrient addition in a Pampean stream. **Aquatic Microbial Ecology** 69:145-155.
- COHEN, R. R. H.; DRESLER, P. V.; PHILLIPS, E. J. P. & CORY, R. L. 1984. The effect of the Asiatic clam, *Corbicula fluminea*, on phytoplankton of the Potomac River, Maryland. **Limnology & Oceanography** 29:170-180.
- CROSS, W. F.; WALLACE, J. B. & ROSEMOND, A. D. 2007. Nutrient enrichment reduces constraints on material flows in a detritus-based food web. **Ecology** 88:2563-2575.
- DI RIENZO, J. A.; CASANOVES, F.; BALZARINI, M. G.; GONZALEZ, L.; TABLADA, M. & ROBLEDO, C. W. 2008. **InfoStat, versión 2013**. Córdoba, Grupo InfoStat, FCA, Universidad Nacional de Córdoba.
- DODDS, W.; BOUSKA, W.; EITZMANN, J.; PILGER, T.; PITTS, K.; RILEY, A.; SCHOESSER, J. & THORNBRUGH, D. 2009. Eutrophication of U.S. freshwaters: analysis of potential economic damages. **Environmental Science and Technology** 43:12-19.
- FEIJOÓ, C. S. & LOMBARDO, R. J. 2007. Baseline water quality and macrophyte assemblages in Pampean streams: a regional approach. **Water Research** 41:1399-1410.
- FEIJOÓ, C. S.; GIORGI, A.; GARCÍA, M. E. & MOMO, F. 1999. Temporal and spatial variability in streams of a pampean basin. **Hydrobiologia** 394:41-52.
- FEIJOÓ, C.; LEGGIERI, L.; OCON, C.; RODRIGUES CAPÍTULO, A.; GIORGI, A.; COLAUTTI, D.; MUÑOZ, I.; FERREIRO, N.; LICURSI, M.; GÓMEZ, N. & SABATER, S. 2014. Stoichiometric homeostasis in the food web of a chronically nutrient-rich stream. **Freshwater Science** 33(3):820-831.
- FERNÁNDEZ, H. & DOMÍNGUEZ, E. eds. 2001. **Guía para la determinación de los artrópodos bentónicos sudamericanos**. San Miguel de Tucumán, EudeT (Serie Investigaciones de la UNT). 282p.
- FERREIRO, N.; GIORGI, A.; LEGGIERI, L.; FEIJOÓ, C. & VILCHES, C. 2011. Phosphorus enrichment affects immobilization but no litter decomposition or exoenzymatic activities in a Pampean stream. **International Review of Hydrobiology** 96:209-220.
- FRIBERG, N.; SKRIVER, J.; LARSEN, S. E.; PEDERSEN, M. L. & BUFFAGNI, A. 2010. Stream macroinvertebrate occurrence along gradients in organic pollution and eutrophication. **Freshwater Biology** 55:1405-1419.
- GIORGI, A.; FEIJOÓ, C. & TELL, G. 2005. Primary producers in a Pampean stream: temporal variation and structuring role. **Biodiversity and Conservation** 14:1699-1718.
- GÓMEZ, N. & RODRIGUES CAPÍTULO, A. 2001. Los bioindicadores y la salud de los ríos. In: **Anales del V Seminario Internacional de Ingeniería y Ambiente**. La Plata, Facultad de Ingeniería, Universidad Nacional de La Plata, p. 109-118.
- GÓMEZ, N.; SIERRA, M. V.; COCHERO, J.; LICURSI, M. & BAUER, D. 2009. Epipellic biofilms as indicators of environmental changes in lowland fluvial systems. In: BAILEY, W. ed. **Biofilms: Formation, Development and Properties**. Hauppauge, Nova Science Publishers. 691p.
- HULME, M. & SHEARD, N. 1999. **Climate Change Scenarios for Australia**. Available at <http://www.cru.uea.ac.uk>. Accessed on May 2013.
- LIN, L. & WEBSTER, J. R. 2012. Sensitivity analysis of a pulse nutrient addition technique for estimating nutrient uptake in large streams. **Limnology and Oceanography: Methods** 10:718-727.
- MCCORMICK, P.; SHUFORD III, R. & RAWLIK, P. 2004. Changes in macroinvertebrate community structure and function along a phosphorus gradient in the Florida Everglades. **Hydrobiologia** 529:113-132.
- MURKIN, H.; POLLARD, J.; STANTON, M.; BOUGHEN, J. & TITMAN, R. 1994. Nutrient additions to wetlands in the Interlake region of Manitoba, Canada: effects of periodic additions throughout the growing season. **Hydrobiologia** 279/280:483-495.
- OCÓN, C. S.; LÓPEZ VAN OOSTEROM, M. V.; MUÑOZ, I. & RODRIGUES CAPÍTULO, A. 2013. Macroinvertebrate trophic responses to nutrient addition in a temperate stream in South America. **Fundamental and Applied Limnology** 182:17-30.
- PLAFKIN, J. L.; BARBOUR, M. T.; PORTER, K. D.; GROSS, S. K. & HUGHES, R. M. 1989. Rapid bioassessment protocols for use in streams and rivers. Benthic macroinvertebrates and fish. Available at <http://water.epa.gov/scitech/monitoring/rsl/bioassessment/>. Accessed on May 2014.
- ROBINSON, C. T. & GESSNER, M. O. 2000. Nutrient addition accelerates leaf breakdown in an alpine springbrook. **Oecologia** 122:258-263.
- RODRIGUES CAPÍTULO, A.; GÓMEZ, N.; GIORGI, A. & FEIJOÓ, C. 2010. Global changes in Pampean lowland streams (Argentina): implications for biodiversity and functioning. **Hydrobiologia** 657:53-70.
- RODRIGUES CAPÍTULO, A.; MUÑOZ, I.; BONADA, N.; GAUDES, A. & TOMANOVA, S. 2009. La biota de los ríos: los invertebrados. In: ELOSEGI, A. & SABATER, S. eds. **Conceptos y técnicas para el estudio de la ecología de ríos**. Bilbao, Fundación BBVA, p. 248-264.
- SABATER, S.; ACUÑA, V.; GIORGI, A.; GUASCH, H.; GUERRA, E.; MUÑOZ, I. & ROMANÍ, A. 2005. Assessing the ecological integrity after nutrient inputs in streams: the relevance of the descriptor and its associated scale. **Aquatic Ecosystem Health & Management** 8:397-403.
- SABATER, S.; ARTIGAS, J.; GAUDES, A.; MUÑOZ, I.; URREA, G. & ROMANÍ, A. 2011. Long-term moderate nutrient inputs enhance autotrophy in a forested Mediterranean stream. **Freshwater Biology** 56:1266-1280.
- SALAZAR, R. H.; LUZZI, D. & LACOSTE, C. 1996. **Cuencas Hídricas Contaminación, evaluación de riesgo y saneamiento**. Buenos Aires, Instituto Provincial de Medio Ambiente. 184p.
- SLAVIK, K.; PETERSON, B.; DEEGAN, L.; BOWDEN, B.; HERSHEY, A. & HOBBI, J. 2004. Long-term responses of the Kuparuk River ecosystem to phosphorus fertilization. **Ecology** 85:939-954.
- SOUSA, R.; ANTUNES, C. & GULHERMINO, L. 2008. Ecology of the invasive Asian clam *Corbicula fluminea* (Muller, 1774) in aquatic ecosystems: an overview. **Annales de Limnologie - International Journal of Limnology** 44:85-94.