

LOCAL SPECTRAL THEORY FOR 2×2 OPERATOR MATRICES

H. ELBJAOUI and E. H. ZEROUALI

Received 13 February 2001 and in revised form 11 July 2001

We discuss the spectral properties of the operator $M_C \in \mathcal{L}(X \oplus Y)$ defined by $M_C := \begin{pmatrix} A & C \\ 0 & B \end{pmatrix}$, where $A \in \mathcal{L}(X)$, $B \in \mathcal{L}(Y)$, $C \in \mathcal{L}(Y, X)$, and X, Y are complex Banach spaces. We prove that $(S_{A^*} \cap S_B) \cup \sigma(M_C) = \sigma(A) \cup \sigma(B)$ for all $C \in \mathcal{L}(Y, X)$. This allows us to give a partial positive answer to Question 3 of Du and Jin (1994) and generalizations of some results of Houimdi and Zguitti (2000). Some applications to the similarity problem are also given.

2000 Mathematics Subject Classification: 47A10.

1. Introduction. Let X and Y be complex Banach spaces and let $\mathcal{L}(X)$, $\mathcal{L}(Y)$, and $\mathcal{L}(Y, X)$ be the algebras of all continuous linear operators on X , Y , and from Y to X , respectively. For $T \in \mathcal{L}(X)$, we denote by $\sigma(T)$ its spectrum, $\sigma_p(T)$ its point spectrum, T^* its adjoint operator, and R_T its resolvent map. Let $x \in X$ and $\lambda_0 \in \mathbb{C}$, we say that λ_0 is in the *local resolvent* of T at x , denoted by $\rho_T(x)$, if the equation

$$(T - \lambda)f(\lambda) = x \tag{1.1}$$

admits an analytic solution in a neighborhood of λ_0 . The set $\sigma_T(x) = \mathbb{C} \setminus \rho_T(x)$ is called the *local spectrum* of T at x .

If for every $x \in X$ any two solutions of (1.1) agree on their common domain, T is said to have the *single-valued extension property* (SVEP). It is obvious that T has the SVEP if and only if the zero function is the only analytic function which satisfies $(T - \lambda)f(\lambda) = 0$.

For $T \in \mathcal{L}(X)$ and F a closed set of \mathbb{C} , denote by the set $X_T(F) := \{x \in X, \sigma_T(x) \subset F\}$ the *analytic spectral space*. The analytic residuum S_T is the set of $\lambda_0 \in \mathbb{C}$ for which there exist a neighborhood G_{λ_0} and $f: G_{\lambda_0} \rightarrow X$ a nonzero analytic function such that $(T - \lambda)f(\lambda) = 0$ for all $\lambda \in G_{\lambda_0}$. We say that the operator T has the *Dunford condition C* (DCC) if $X_T(F)$ is closed whenever F is closed. It is clear that T has the SVEP if and only if $S_T = \emptyset$. The *surjective spectrum* of T is given by $\sigma_{\text{su}}(T) := \{\lambda \in \mathbb{C} / \mathbb{T} - \lambda \text{ is not surjective}\}$. It is known that $\sigma_{\text{su}}(T) = \cup_{x \in X} \sigma_T(x)$ and that $\sigma(T) = S_T \cup \sigma_{\text{su}}(T)$ (see [6, 7, 8]). In particular, if T has the SVEP, we obtain that $\sigma(T) = \sigma_{\text{su}}(T)$. A complete study of basic notions of local spectral theory can be found in [1, 4].

The operator $M_C \in \mathcal{L}(X \oplus Y)$ is defined by $M_C := \begin{pmatrix} A & C \\ 0 & B \end{pmatrix}$, where $A \in \mathcal{L}(X)$, $B \in \mathcal{L}(Y)$, $C \in \mathcal{L}(Y, X)$, and X, Y are complex Banach spaces. In [2], a number of natural questions concerning the relation between the spectrums of these operators have been considered. Our interest in this paper is to develop some new conditions under which we have the equality $\sigma(M_C) = \sigma(A) \cup \sigma(B)$. Houimdi and Zguitti [3] give a positive answer to this question when the operator B has the SVEP.

Our main result gives a partial answer to [2, Question 3], and generalizes results from [3]. At the end, we give a generalization of [3, Proposition 3.2] related to the DCC for the operator M_C and some applications to the similarity of orbits.

We collect in the following proposition some useful spectral properties of the operator M_C from [2, 3], that can be also obtained easily.

PROPOSITION 1.1. *If $A \in \mathcal{L}(X)$, $B \in \mathcal{L}(Y)$, and $C \in \mathcal{L}(Y, X)$, then*

- (1) $\sigma_p(A) \subset \sigma_p(M_C) \subset \sigma_p(A) \cup \sigma_p(B)$,
- (2) $\sigma_{ap}(A) \subset \sigma_{ap}(M_C) \subset \sigma_{ap}(A) \cup \sigma_{ap}(B)$,
- (3) $S_A \subset S_{M_C} \subset S_A \cup S_B$,
- (4) $\sigma_{su}(B) \subset \sigma_{su}(M_C) \subset \sigma_{su}(A) \cup \sigma_{su}(B)$,
- (5) $\sigma_{ap}(A) \cup \sigma_{su}(B) \subset \sigma(M_C)$,
- (6) $\sigma(A) \cup \sigma(B) = S_{A^*} \cup S_B \cup \sigma(M_C)$.

2. Spectral theory of the operator M_C . We study in the sequel spectral theory of M_C , we refine the inclusions given in Proposition 1.1, and provide some properties of local spectrum.

Using Leiterer's theorem (see [6, Theorem 3.2.1, page 212] and [5]), we derive the following proposition.

PROPOSITION 2.1. *If A, B , and C are given, then*

$$S_B \subset \sigma_{su}(A) \cup S_{M_C}. \tag{2.1}$$

PROOF. Let $\lambda_0 \in S_B \setminus \sigma_{su}(A)$. There exist a neighborhood V_{λ_0} and a nonzero analytic function $g : V_{\lambda_0} \rightarrow Y$ satisfying

$$(B - \mu)g(\mu) = 0, \quad V_{\lambda_0} \cap \sigma_{su}(A) = \emptyset. \tag{2.2}$$

By Leiterer's theorem, there exists an analytic function $f : V_{\lambda_0} \rightarrow X$ satisfying

$$(A - \mu)f(\mu) = -Cg(\mu) \quad \forall \mu \in V_{\lambda_0}. \tag{2.3}$$

The nonzero analytic function $f \oplus g : V_{\lambda_0} \rightarrow X \oplus Y$ defined by $(f \oplus g)(\mu) = f(\mu) \oplus g(\mu)$ satisfies

$$(M_C - \mu)(f(\mu) \oplus g(\mu)) = 0 \quad \forall \mu \in V_{\lambda_0}, \tag{2.4}$$

hence $\lambda_0 \in S_{M_C}$. □

The following corollary is immediate from [Proposition 2.1](#) and [Proposition 1.1\(4\)](#).

COROLLARY 2.2. *For given operators $A, B,$ and $C,$*

$$\sigma(B) \subset \sigma_{\text{su}}(A) \cup \sigma(M_C). \tag{2.5}$$

To establish our main theorem, we first claim the following proposition related to the local spectrum of the operator $M_C,$ which generalizes [[3](#), Proposition 2.1].

PROPOSITION 2.3. *If $A, B,$ and C are given,*

$$S_B \cup \sigma_A(x) = S_B \cup \sigma_{M_C}(x \oplus 0) \quad \forall x \in X. \tag{2.6}$$

PROOF. If $\lambda_0 \notin (S_B \cup \sigma_{M_C}(x \oplus 0)),$ then there exist a neighborhood V_{λ_0} and a nonzero analytic function $h : V_{\lambda_0} \rightarrow X \oplus Y$ satisfying

$$(M_C - \mu)h(\mu) = x \oplus 0 \quad \forall \mu \in V_{\lambda_0}. \tag{2.7}$$

Let $h = h_1 \oplus h_2$ with $h_1 : V_{\lambda_0} \rightarrow X$ and $h_2 : V_{\lambda_0} \rightarrow Y$ analytic functions. We obtain

$$(A - \mu)h_1(\mu) + Ch_2(\mu) = x, \quad (B - \mu)h_2(\mu) = 0. \tag{2.8}$$

As $\lambda_0 \notin S_B,$ we conclude that $h_2 = 0$ on $V_{\lambda_0}.$ Hence

$$(A - \mu)h_1(\mu) = x \quad \forall \mu \in V_{\lambda_0}, \tag{2.9}$$

thus $\lambda_0 \notin \sigma_A(x).$ The reverse inclusion is clear. □

The following result generalizes [[3](#), Corollary 2.2].

COROLLARY 2.4. *For given operators $A, B,$ and $C,$*

$$S_B \cup \sigma_{\text{su}}(M_C) = \sigma_{\text{su}}(A) \cup \sigma(B). \tag{2.10}$$

PROOF. By [Proposition 2.1](#), we have $\sigma_{\text{su}}(M_C) \subset \sigma_{\text{su}}(A) \cup \sigma_{\text{su}}(B).$ Hence

$$S_B \cup \sigma_{\text{su}}(M_C) \subset \sigma_{\text{su}}(A) \cup \sigma(B). \tag{2.11}$$

For the converse, we obtain by [Proposition 2.3](#) that

$$S_B \cup \sigma_A(x) = S_B \cup \sigma_{M_C}(x \oplus 0) \quad \forall x \in X, \tag{2.12}$$

hence

$$S_B \cup \sigma_{\text{su}}(A) \subset S_B \cup \sigma_{\text{su}}(M_C). \tag{2.13}$$

Since $\sigma_{\text{su}}(B) \subset \sigma_{\text{su}}(M_C)$, we check that

$$\sigma_{\text{su}}(A) \cup \sigma(B) \subset S_B \cup \sigma_{\text{su}}(M_C). \tag{2.14}$$

□

We are now in a position to derive a generalization of [3, Theorem 2.1].

THEOREM 2.5. *For given operators A, B , and C ,*

$$(S_{A^*} \cap S_B) \cup \sigma(M_C) = \sigma(A) \cup \sigma(B). \tag{2.15}$$

PROOF. In the first step, we prove that

$$S_B \cup \sigma(M_C) = \sigma(A) \cup \sigma(B). \tag{2.16}$$

By [Corollary 2.4](#) and [Proposition 1.1\(3\)](#), we obtain that

$$\sigma(A) \cup \sigma(B) \subset S_B \cup \sigma(M_C). \tag{2.17}$$

The converse is clear since the inclusion $\sigma(M_C) \subset \sigma(A) \cup \sigma(B)$ is always true. In the second step, remark that

$$M_C^* := \begin{pmatrix} A^* & 0 \\ C^* & B^* \end{pmatrix}. \tag{2.18}$$

Thus, in the same way, we obtain that

$$S_{A^*} \cup \sigma(M_C) = \sigma(A) \cup \sigma(B). \tag{2.19}$$

Hence the theorem is proved. □

We give in what follows a new condition under which we have the desired equality.

COROLLARY 2.6. *For given operators A, B , and C , if $S_{A^*} \cap S_B = \emptyset$, then $\sigma(M_C) = \sigma(A) \cup \sigma(B)$.*

3. Applications. In this section, we give some necessary conditions such that M_C has the SVEP and provide some results on similarity orbits and the range of generalized derivation using [Theorem 2.5](#). The following result gives a partial characterization of the property of the single extension property of the operator M_C and discusses the converse of [3, Proposition 3.1].

PROPOSITION 3.1. *For given operators A, B , and C such that the surjective spectrum of the operator A has an empty interior,*

$$S_{M_C} = \emptyset \iff S_A = S_B = \emptyset. \tag{3.1}$$

In particular M_C has SVEP if and only if A and B have the SVEP.

PROOF. If A and B have the SVEP, so is the case of M_C . (See [Proposition 2.1\(3\)](#).) On the other hand, the relations

$$S_B \subset \sigma_{\text{su}}(A) \cup S_{M_C}, \quad S_A \subset S_{M_C} \subset S_A \cup S_B \tag{3.2}$$

imply that $S_A = S_B = \emptyset$. The proof is complete. □

The next proposition provides a generalization of [[3](#), Proposition 3.2].

PROPOSITION 3.2. *If A and B are given and F is a closed set such that $S_B \subset F$, then the following assertion holds true: if there exists $C \in \mathcal{L}(Y, X)$ such that the analytic subspace $(X \oplus Y)_{M_C}(F)$ is closed, then $X_A(F)$ is also closed.*

PROOF. Let (x_n) be a sequence of elements of $X_A(F)$ which converges to $x \in X$. By [Proposition 2.3](#), we have

$$S_B \cup \sigma_A(x_n) = S_B \cup \sigma_{M_C}(x_n \oplus 0) \quad \forall x \in X. \tag{3.3}$$

We deduce that $\sigma_{M_C}(x_n \oplus 0) \subset F \cup S_B$. Thus, $\sigma_{M_C}(x_n \oplus 0) \subset F$. Since $(X \oplus Y)_{M_C}(F)$ is closed, we have $\sigma_{M_C}(x \oplus 0) \subset F$. Then

$$\sigma_A(x) \subset S_B \cup \sigma_A(x) = S_B \cup \sigma_{M_C}(x \oplus 0). \tag{3.4}$$

We derive that $\sigma_A(x) \subset F \cup S_B \subset F$, then $x \in X_A(F)$, hence $X_A(F)$ is closed. □

Our second application is related to the similarity problem. Let A, B , and M_C be as above and consider $\delta_{A,B}$ the *derivation operator* defined by $\delta_{A,B}(X) = AX - XB$ for $X \in \mathcal{L}(X)$. Let $\text{Im}(\delta_{A,B})$ be its range and denote by H_1, H_2 , and H_3 the following classes of operators:

$$\begin{aligned} H_1 &:= \{C \in \mathcal{L}(Y, X) \text{ such that } C \in \text{Im}(\delta_{A,B})\}, \\ H_2 &:= \{C \in \mathcal{L}(Y, X) \text{ such that } M_C \text{ is similar to } M_O\}, \\ H_3 &:= \{C \in \mathcal{L}(Y, X) \text{ such that } \sigma(M_C) = \sigma(M_O)\}. \end{aligned} \tag{3.5}$$

It is obvious that

$$H_1 \subset H_2 \subset H_3. \tag{3.6}$$

The above inclusions received a lot of interest (see [[9](#), [10](#)]). In general, any of these inclusions can be strict. We construct some classes of operators such that $H_1 \neq H_2$ and $H_2 \neq H_3$.

Remark first by [Corollary 2.6](#) that if $S_{A^*} \cap S_B = \emptyset$, then $H_3 = \mathcal{L}(Y, X)$. Suppose in the sequel that $X = Y$ and let S be the unilateral shift. We consider, respectively, the operators $A := \begin{pmatrix} S & 0 \\ 0 & 0 \end{pmatrix}$ and $M_C := \begin{pmatrix} A & C \\ 0 & A \end{pmatrix}$.

By [[9](#), Theorem 6], if M_C and M_O are similar, then C is a commutator, that is, $C \neq \lambda I + K$ for any $\lambda \in \mathbb{C}$ and K a compact operator. Hence, choosing A so

that $S_{A^*} \cap S_A = \emptyset$ and considering $C = \lambda I + K$ for some $\lambda \in \mathbb{C}$ and K a compact operator, we get $M_C \in H_3 \setminus H_2$. Moreover, for $C = \begin{pmatrix} 0 & I \\ 0 & 0 \end{pmatrix}$, we conclude by [9] that $M_C \in H_2 \setminus H_1$. Consequently, the inclusions in (3.6) are all strict for $M_C = \begin{pmatrix} A & C \\ 0 & A \end{pmatrix}$ with $A = \begin{pmatrix} S & 0 \\ 0 & 0 \end{pmatrix}$, $C = \begin{pmatrix} 0 & I \\ 0 & 0 \end{pmatrix}$, and S is the unilateral shift.

ACKNOWLEDGMENTS. This work was partially supported by Abdus Salam ICTP Centre. The authors are specially grateful to Professor C. E. Chidume, Mathematics Section, and Dr. H. Mahzouli for fruitful discussions.

REFERENCES

- [1] I. Colojoară and C. Foiaş, *Theory of Generalized Spectral Operators*, Gordon and Breach Science Publishers, New York, 1968.
- [2] H. K. Du and P. Jin, *Perturbation of spectrums of 2×2 operator matrices*, Proc. Amer. Math. Soc. **121** (1994), no. 3, 761–766.
- [3] M. Houimdi and H. Zguitti, *Propriétés spectrales locales d'une matrice carrée des opérateurs*, Acta Math. Vietnam. **25** (2000), no. 2, 137–144 (French).
- [4] R. Lange and S. W. Wang, *New Approaches in Spectral Decomposition*, Contemporary Mathematics, vol. 128, American Mathematical Society, Rhode Island, 1992.
- [5] K. B. Laursen and M. M. Neumann, *On analytic solutions of the equation $(T - \lambda)f(\lambda) = x$* , LEU Seminar Notes in Funct. Anal. PDEs, Louisiana State University, Louisiana, 1994, pp. 256–265.
- [6] ———, *An Introduction to Local Spectral Theory*, London Mathematical Society Monographs. New Series, vol. 20, The Clarendon Press, Oxford University Press, New York, 2000.
- [7] K. B. Laursen and P. Vrbová, *Some remarks on the surjectivity spectrum of linear operators*, Czechoslovak Math. J. **39(114)** (1989), no. 4, 730–739.
- [8] M. M. Neumann, *On local spectral properties of operators on Banach spaces*, Rend. Circ. Mat. Palermo (2) Suppl. (1998), no. 56, 15–25.
- [9] A. Schweinsberg, *The operator equation $AX - XB = C$ with normal A and B* , Pacific J. Math. **102** (1982), no. 2, 447–453.
- [10] ———, *Similarity orbits and the range of the generalized derivation $X \rightarrow MX - XN$* , Trans. Amer. Math. Soc. **324** (1991), no. 1, 201–211.

H. Elbjaoui: Département de Mathématiques et Informatique, Faculté des Sciences de Rabat, Université Mohamed V, BP 1014, Rabat, Morocco

E-mail address: hicham.elbjaoui@caramail.com

E. H. Zerouali: Département de Mathématiques et Informatique, Faculté des Sciences de Rabat, Université Mohamed V, BP 1014, Rabat, Morocco

E-mail address: zerouali@fsr.ac.ma

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk