

# Gastric Adenocarcinoma with Thymic Metastasis after Curative Resection: A Case Report

Tomoyuki Matsunaga, Hiroaki Saito, Koza Miyatani, Seigo Takaya,  
Yoji Fukumoto, Tomohiro Osaki, and Masahide Ikeguchi

*Division of Surgical Oncology, Department of Surgery, Tottori University School of Medicine, Yonago, Japan*

The peritoneum is the most frequent site of recurrence for gastric cancer after gastrectomy, followed by the liver and lymph nodes. In contrast, metastasis to the thymus is rare. Annual surveillance with computed tomography was performed on a 67-year-old man who previously underwent a distal gastrectomy and D2 lymph node dissection for gastric cancer at Tottori University. Five years after the initial operation, an anterior mediastinal tumor was detected by computed tomography. The patient underwent video-assisted thoracic surgery to remove the tumor. Histopathology revealed adenocarcinoma cells similar to those of the gastric cancer resected 5 years previously. Thymic metastasis was considered likely based on the location of the tumor. The recognition that gastric cancer can metastasize to unusual anatomic locations, such as the thymus, can facilitate an accurate, prompt diagnosis and appropriate treatment.

**Key Words:** Stomach neoplasms; Neoplasm metastasis

## Introduction

It is well known that the most common sites of gastric carcinoma recurrence are the liver, lungs, lymph nodes, and peritoneum.<sup>1</sup> Although metastasis to the posterior mediastinal lymph nodes is occasionally encountered after gastrectomy, recurrence in the anterior mediastinum is extremely rare. We report a very rare case of a single thymic metastasis from gastric cancer occurring 5 years after a distal gastrectomy with lymph node dissection.

## Case Report

An asymptomatic 67-year-old man had a history of gastric ad-

enocarcinoma (Fig. 1), for which he had undergone a distal partial gastrectomy 5 years previously. Histological examination of the resected specimen had revealed moderately differentiated adenocarcinoma invading the muscularis propria with 5 regional lymph node metastases at stations 1 (3/7), 3 (1/1), and 7 (1/1). The disease was stage IIB as defined by the Japanese Gastric Cancer Association.<sup>2</sup> The patient received postoperative adjuvant chemotherapy with the oral fluoropyrimidine S-1 (100 mg/body). However, he could only take 1 course of chemotherapy because of severe diarrhea.

Four years and 8 months after the initial operation, a computed tomography scan from the chest to the pelvis demonstrated a tumor in the anterior mediastinum (Fig. 2). With 18F-fluorodeoxyglucose positron emission tomography computed tomography, a focal hypermetabolic lesion with a maximum standardized uptake value of 3.6 was identified in this lesion (Fig. 3). The patient underwent an anterior mediastinal tumor resection by video-assisted thoracic surgery (Fig. 4); the preoperative diagnosis was thymoma, thymic carcinoma, or malignant lymphoma in the thymus. Histopathology of the tumor revealed adenocarcinoma cells that were similar to those of the gastric cancer resected 5 years previously. Further-

Correspondence to: Hiroaki Saito  
Division of Surgical Oncology, Department of Surgery, Tottori University  
School of Medicine, 36-1 Nishi-cho, Yonago 683-8504, Japan  
Tel: +81-859-38-6567, Fax: +81-859-38-6569  
E-mail: sai10@med.tottori-u.ac.jp  
Received June 3, 2014  
Revised June 25, 2014  
Accepted July 7, 2014

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

more, immunochemistry indicated that tumor cells were positive for cytokeratin 20 and CDX-2 (C and D) and negative for cytokeratin 7.

Thymic metastasis was considered likely based on the location of the tumor (Fig. 5). The patient is currently taking the oral


fluoropyrimidine S-1 (100 mg/body) as adjuvant chemotherapy to prevent recurrence.

## Discussion


In this report, we describe a solitary thymic metastasis from a previously resected gastric cancer, which is an extremely rare occurrence. The thymus is an essential organ that controls cellular




**Fig. 1.** Gastrofibroscopy showed a type 0-IIa+IIc lesion with a maximum size of 20 mm in the posterior side of the lower third of the stomach.




**Fig. 2.** Enhanced computed tomography scan revealed an anterior mediastinal tumor.



**Fig. 3.** Positron emission tomography computed tomography showed a focal hypermetabolic lesion with a maximum standardized uptake value of 3.6 in the anterior mediastinum. There was no accumulation of  $^{18}\text{F}$ -fluorodeoxyglucose in the liver, lungs, lymph nodes, or peritoneum.



**Fig. 4.** The tumor in the anterior mediastinum was removed, including the surrounding fat tissue.



**Fig. 5.** Histopathological examination of the anterior mediastinal tumor showed a poorly differentiated adenocarcinoma that was similar to the gastric cancer removed 5 years previously. (A) Histopathological examination of gastric carcinoma (H&E,  $\times 10$ ). (B) Histopathological examination of the anterior mediastinal tumor (H&E,  $\times 10$ ). Furthermore, immunocytochemistry indicated that tumor cells were positive for cytokeratin 20 and CDX-2 (C, D) but negative for cytokeratin 7 (data not shown) (C: cytokeratin 20,  $\times 10$ ; D: CDX-2,  $\times 10$ ).

immune function; therefore, it has been considered almost impossible for a tumor to metastasize to it. However, detailed study of the thymic structure has revealed that it is not absolutely safe from tumor metastasis.<sup>3</sup> As described by Clark,<sup>4</sup> the parenchyma of the thymus is associated with a blood–thymus barrier that prevents it from making direct contact with antigens or cancerous cells, thereby seemingly excluding the occurrence of cancer metastasis. However, the septum of the thymus comprises interlobular connective tissue with blood vessels, lymph ducts, and nerves that theoretically enable contact with other parts of the body through the blood and lymphatic systems.

There are relatively few reports<sup>3,5–7</sup> of thymic metastases, but gastric, breast, laryngeal, and lung carcinoma metastasis to the thymus has been reported at autopsy.<sup>8</sup> In an autopsy series by Middle-

ton,<sup>8</sup> 7 cases of thymic metastasis occurred in 102 carcinoma cases, which is an incidence of approximately 7%. There was only 1 case of gastric cancer among these 7 cases. To our knowledge, no antemortem cases of thymic metastasis from gastric cancer have been previously reported.

In our case, thymic structures could not be recognized around the tumor. This is reasonable because the thymus degenerates after adolescence and is replaced by fatty tissue. Therefore, our conclusion that the tumor originated in the thymus is based on the metastatic tumor's location.

It is important to detect metastatic tumors as soon as possible because some patients can be successfully palliated with appropriate treatment and may even achieve prolonged survival. The recognition that gastric cancer can metastasize to unusual sites, such as the

thymus, can facilitate an accurate, prompt diagnosis and appropriate treatment.

## References

1. Saito H, Tsujitani S, Kondo A, Ikeguchi M, Maeta M, Kaibara N. Expression of vascular endothelial growth factor correlates with hematogenous recurrence in gastric carcinoma. *Surgery* 1999;125:195-201.
2. Japanese Gastric Cancer Association. Japanese classification of gastric carcinoma: 3rd English edition. *Gastric Cancer* 2011;14:101-112.
3. Hayashi S, Hamanaka Y, Sueda T, Yonehara S, Matsuura Y. Thymic metastasis from prostatic carcinoma: report of a case. *Surg Today* 1993;23:632-634.
4. Clark SL Jr. The reticulum of lymph nodes in mice studied with the electron microscope. *Am J Anat* 1962;110:217-257.
5. Phillips CJ. Case report: metastatic malignant testicular teratoma of the thymus. *Br J Radiol* 1994;67:203-204.
6. Nam MS, Chu YC, Choe WS, Kim SJ, Hong SB, Kim YJ, et al. Metastatic follicular thyroid carcinoma to the thymus in a 35-year-old woman. *Yonsei Med J* 2002;43:665-669.
7. Park SB, Kim HH, Shin HJ, Paik MH, Kim DB, Gong G. Thymic metastasis in breast cancer: a case report. *Korean J Radiol* 2007;8:360-363.
8. Middleton G. Involvement of the thymus by metastatic neoplasms. *Br J Cancer* 1966;20:41-46.