

EFI-95-65
 FERMILAB-PUB-95/345-T
 ANL-HEP-PR-95-80
 WIS-95/50-Oct.-Ph
 hep-ph/9511363
 November 1995

**ANGULAR DISTRIBUTIONS AND LIFETIME DIFFERENCES
 IN $B_s \rightarrow J/\psi\phi$ DECAYS**¹

*Amol S. Dighe*²

*Isard Dunietz*³

Harry J. Lipkin^{4, 5}

and

Jonathan L. Rosner^{2, 3}

ABSTRACT

The strange B meson $B_s \equiv \bar{b}s$ and its charge-conjugate $\bar{B}_s \equiv b\bar{s}$ are expected to mix with one another in such a way that the mass eigenstates B_s^H (“heavy”) and B_s^L (“light”) may have a perceptible lifetime difference of up to 40%, with the CP-even eigenstate being shorter-lived. A simple transversity analysis permits one to separate the CP-even and CP-odd components of $B_s \rightarrow J/\psi\phi$, and thus to determine the lifetime difference. The utility of a similar analysis for $B^0 \rightarrow J/\psi K^{*0}$ is noted.

The Cabibbo-Kobayashi-Maskawa picture of weak charge-changing transitions [1] predicts the strange B meson $B_s \equiv \bar{b}s$ and its charge-conjugate $\bar{B}_s \equiv b\bar{s}$ to mix with one another with a large amplitude. The mass eigenstates B_s^H (“heavy”) and B_s^L (“light”) with masses $m(B_s^H) \equiv m_H$ and $m(B_s^L) \equiv m_L$ are expected to be split by $\Delta m \equiv m_H - m_L \approx 25\bar{\Gamma}$, give or take a factor of two [2], where $\bar{\Gamma} \equiv (\Gamma_H + \Gamma_L)/2 \approx \Gamma(B^0)$ ($B^0 \equiv \bar{b}d$) and $\Gamma_{H,L} \equiv \Gamma(B_s^H, B_s^L)$. The measurement of such a large mass difference poses an experimental challenge.

To a good approximation, CP violation can be neglected in calculating the mass eigenstates, in which case they correspond to those $B_s^{(\pm)}$ of even and odd CP, with $B_s^L = B_s^{(+)}$ and $B_s^H = B_s^{(-)}$ as we shall see. The decay of a \bar{B}_s meson via the quark

¹To be submitted to Physics Letters B

²Enrico Fermi Institute and Department of Physics, University of Chicago, Chicago, IL 60637

³Theoretical Physics Division, Fermi National Accelerator Laboratory, Batavia, IL 60510

⁴High Energy Physics Division, Argonne National Laboratory, Argonne, IL 60439

⁵Department of Particle Physics, Weizmann Institute of Science, Rehovoth, Israel

subprocess $b(\bar{s}) \rightarrow c\bar{c}s(\bar{s})$ gives rise to predominantly CP-even final states [3]. Thus the CP-even eigenstate should have the greater decay rate. An explicit calculation [4] gives

$$\frac{\Gamma(B_s^{(+)}) - \Gamma(B_s^{(-)})}{\bar{\Gamma}} \simeq 0.18 \frac{f_{B_s}^2}{(200 \text{ MeV})^2} \quad , \quad (1)$$

where f_{B_s} is the B_s decay constant (in a normalization in which $f_\pi = 132 \text{ MeV}$). In one estimate [2], $f_{B_s} = 225 \pm 40 \text{ MeV}$, while a compilation of lattice results [5] obtains $f_{B_s} = 201 \pm 40 \text{ MeV}$ (90% c.l. limits). The upper limit of 40% for (1) is based on an estimate of the maximum possible contribution from the $b(\bar{s}) \rightarrow c\bar{c}s(\bar{s})$ subprocess [6, 7].

The ratio of the mass splitting to the width difference of strange B 's is predicted to be large and independent of CKM matrix elements [6, 8] (to lowest order, neglecting QCD corrections which may be appreciable):

$$\frac{\Delta m}{\Delta \Gamma} \simeq -\frac{2}{3\pi} \frac{m_t^2 h(m_t^2/M_W^2)}{m_b^2} \left(1 - \frac{8m_c^2}{3m_b^2}\right)^{-1} \simeq -200 \quad , \quad (2)$$

where $\Delta \Gamma \equiv \Gamma_H - \Gamma_L$. Here $h(x)$ decreases monotonically from 1 at $x = 0$ to $1/4$ as $x \rightarrow \infty$; it is about 0.54 for $m_t = 180 \text{ GeV}/c^2$. In view of the sign in Eq. (1) and since $\Delta m > 0$ by definition, we then identify $B_s^L = B_s^{(+)}$ and $B_s^H = B_s^{(-)}$ [6]. If the mass difference Δm turns out to be too large to measure at present because of the rapid frequency of $B_s - \bar{B}_s$ oscillations it entails, the width difference $\Delta \Gamma$ may be large enough to detect. The possibility of a value of $\Delta \Gamma/\Gamma$ for strange B mesons large enough to measure experimentally has been stressed previously [3, 4, 9].

One can measure $\bar{\Gamma}$ using semileptonic decays, while the decays to CP eigenstates can be measured by studying the correlations between the polarization states of the vector mesons in $B_s^{(\pm)} \rightarrow J/\psi\phi$. (For similar methods applied to decays of other spinless mesons see, e.g., Ref. [10].) In the present note we describe a means by which the $J/\psi\phi$ final states of definite CP in B_s decays may be separated from one another using a simple angular distribution based on a *transversity* variable [11, 12, 13]. This transversity variable allows one to directly separate the summed contribution of the even partial waves (S, D) from the odd one (P) by means of their opposite parities. The CDF Collaboration [14] has recently reported the first angular distribution analysis of the decay $B_s \rightarrow J/\psi\phi$, obtaining a separation into longitudinal and transverse helicity amplitudes without making a statement yet about the CP-even and CP-odd contributions.

We summarize our main result. Consider the final state $J/\psi\phi \rightarrow \ell^+\ell^-K^+K^-$, where $\ell = e$ or μ . In the rest frame of the J/ψ let the direction of the ϕ define the x axis. Let the plane of the K^+K^- system define the y axis, with $p_y(K^+) > 0$, so the normal to that plane defines the z axis. (We assume a right-handed coordinate system.) We define the angle θ as the angle between the ℓ^+ and the z axis. Then the time-dependent rate for the $J/\psi\phi$ mode is given by

$$\frac{d^2\Gamma}{d\cos\theta dt} = \frac{3}{8}p(t) (1 + \cos^2\theta) + \frac{3}{4}m(t) \sin^2\theta$$

$$= \frac{3}{8} [p(t) + 2m(t)] + \frac{3}{8} [p(t) - 2m(t)] \cos^2 \theta \quad , \quad (3)$$

where

$$p(t) = p(0)e^{-\Gamma_L t} \quad (\text{CP even}) \quad , \quad m(t) = m(0)e^{-\Gamma_H t} \quad (\text{CP odd}) \quad , \quad (4)$$

so that the probability of having a CP-even [CP-odd] state at proper time t is given by $p(t)/(p(t) + m(t))$ [$m(t)/(p(t) + m(t))$]. The angular distribution is normalized in such a way that

$$\frac{d\Gamma}{dt} = \int_{-1}^1 d(\cos \theta) \frac{d^2\Gamma}{d \cos \theta dt} = p(t) + m(t) \quad . \quad (5)$$

As t increases, one should see a growth of the $\sin^2 \theta$ component. The angle θ is an example of a transversity variable, whose utility for the determination of CP properties of multi-particle systems was pointed out some time ago [15].

The zero-angular-momentum states of two massive neutral vector mesons such as J/ψ and ϕ , both with the same CP (in this case, even) consist of two with even CP and one with odd CP. One can form states with orbital angular momenta $L = 0$ (CP even), $L = 1$ (CP odd), and $L = 2$ (CP even).

Alternatively, one can decompose the decay amplitude A into three independent components [16], corresponding to linear polarization states of the vector mesons which are either longitudinal (0), or transverse to their directions of motion and parallel (\parallel) or perpendicular (\perp) to one another. The states 0 and \parallel are P-even, while the state \perp is P-odd. Since J/ψ and ϕ are both C-odd eigenstates, the properties under P are the same as those under CP.

Consider the polarization three-vectors $\epsilon_{J/\psi}$ and ϵ_ϕ in the J/ψ rest frame. The independent decay amplitudes are the rotationally invariant quantities linear in $\epsilon_{J/\psi}^*$ and ϵ_ϕ^* and involving possible powers of $\hat{\mathbf{p}}$, a unit vector in the direction of the momentum of ϕ in the J/ψ rest frame.

The two CP-even decay amplitudes are the combinations $\epsilon_{J/\psi}^* \cdot \epsilon_\phi^*$ (contributing to A_0 and A_\parallel) and $\epsilon_{J/\psi}^* \cdot \hat{\mathbf{p}} \epsilon_\phi^* \cdot \hat{\mathbf{p}} = \epsilon_{J/\psi}^{*L} \epsilon_\phi^{*L}$ (contributing only to A_0), where $\epsilon^L \equiv \hat{\mathbf{p}} \cdot \epsilon$. Equivalently, one can subtract off the longitudinal component of the polarization vectors to replace $\epsilon_{J/\psi}^* \cdot \epsilon_\phi^*$ by $\epsilon_{J/\psi}^{*T} \cdot \epsilon_\phi^{*T}$, contributing only to A_\parallel , where the superscripts T refer to projections perpendicular to $\hat{\mathbf{p}}$. The CP-odd amplitude $\epsilon_{J/\psi}^* \times \epsilon_\phi^* \cdot \hat{\mathbf{p}}$ contributes only to A_\perp . The case of transverse (\parallel or \perp) polarization states is reminiscent of photon polarization correlations [17] in neutral pion decay. Thus we may write the decay amplitude as

$$A(B_s \rightarrow J/\psi\phi) = A_0(m_\phi/E_\phi)\epsilon_{J/\psi}^{*L}\epsilon_\phi^{*L} - A_\parallel\epsilon_{J/\psi}^{*T}\cdot\epsilon_\phi^{*T}/\sqrt{2} - iA_\perp\epsilon_{J/\psi}^*\times\epsilon_\phi^*\cdot\hat{\mathbf{p}}/\sqrt{2} \quad , \quad (6)$$

where E_ϕ is the energy of the ϕ in the J/ψ rest frame, and the individual amplitudes are real in the absence of final-state interactions. The amplitudes for the corresponding decays of $\bar{B}_s \equiv CP(B_s)$ are $\bar{A}_0 = A_0$, $\bar{A}_\parallel = A_\parallel$, and $\bar{A}_\perp = -A_\perp$. (We can see directly by counting powers of $\hat{\mathbf{p}}$ that A_0 and A_\parallel are P-even while A_\perp is P-odd.) We have normalized the partial widths for the three independent polarization states in such a way that

$$d\Gamma(B_s \rightarrow J/\psi\phi)/dt = |A_0|^2 + |A_\parallel|^2 + |A_\perp|^2 \quad , \quad (7)$$

and we may identify

$$p(t) = |A_0|^2 + |A_{\parallel}|^2 \quad , \quad m(t) = |A_{\perp}|^2 \quad . \quad (8)$$

For purposes of comparing with other notations, we can express the helicity amplitudes A_{λ} (where $\lambda = 1, 0, -1$ is the projection of the ϕ angular momentum on the x axis) in terms of the linear polarization basis by $A_{\pm 1} = (A_{\parallel} \pm A_{\perp})/\sqrt{2}$, with A_0 the same in either basis, and in terms of S-, P-, and D-wave amplitudes by

$$A_{\pm 1} = \sqrt{\frac{1}{3}}S \pm \sqrt{\frac{1}{2}}P + \sqrt{\frac{1}{6}}D \quad , \quad A_0 = -\sqrt{\frac{1}{3}}S + \sqrt{\frac{2}{3}}D \quad . \quad (9)$$

With these normalizations,

$$d\Gamma(B_s \rightarrow J/\psi\phi)/dt = |A_0|^2 + |A_1|^2 + |A_{-1}|^2 = |S|^2 + |P|^2 + |D|^2 \quad , \quad (10)$$

and

$$A_{\parallel} = \sqrt{\frac{2}{3}}S + \sqrt{\frac{1}{3}}D \quad , \quad A_{\perp} = P \quad . \quad (11)$$

The longitudinal and transverse partial widths are given, respectively, by

$$d\Gamma_0/dt = |A_0|^2 \quad , \quad d\Gamma_T/dt = |A_1|^2 + |A_{-1}|^2 \quad . \quad (12)$$

In terms of partial-wave amplitudes, one has

$$\begin{aligned} \frac{d\Gamma_0}{dt} &= |-\sqrt{1/3}S + \sqrt{2/3}D|^2 \quad , \quad \frac{d\Gamma_T}{dt} = |\sqrt{2/3}S + \sqrt{1/3}D|^2 + |P|^2 \quad , \\ \frac{d\Gamma_{\parallel}}{dt} &= |\sqrt{2/3}S + \sqrt{1/3}D|^2 \quad , \quad \frac{d\Gamma_{\perp}}{dt} = |P|^2 \quad , \end{aligned} \quad (13)$$

while

$$\frac{p(t)}{p(t) + m(t)} = \frac{|S|^2 + |D|^2}{|S|^2 + |P|^2 + |D|^2} \quad ; \quad \frac{m(t)}{p(t) + m(t)} = \frac{|P|^2}{|S|^2 + |P|^2 + |D|^2} \quad . \quad (14)$$

Finally, we note that in the covariant expression [18]

$$A_{\lambda} = \epsilon_{1\mu}^* \epsilon_{2\nu}^* \left[ag^{\mu\nu} + \frac{b}{m_1 m_2} p_2^{\mu} p_1^{\nu} + \frac{ic}{m_1 m_2} \epsilon^{\mu\nu\alpha\beta} p_{1\alpha} p_{2\beta} \right] \quad (15)$$

for the decay $B \rightarrow V_1 V_2$, where $\epsilon^{0123} \equiv +1$ and V_1 and V_2 are vector mesons with masses m_1 and m_2 and four-momenta p_1 and p_2 , the helicity amplitudes are

$$A_{\pm 1} = a \pm c\sqrt{x^2 - 1} \quad , \quad A_0 = -ax - b(x^2 - 1) \quad , \quad (16)$$

where $x \equiv p_1 \cdot p_2 / (m_1 m_2)$. We thus identify

$$S = \frac{1}{\sqrt{3}} [a(2+x) + b(x^2 - 1)] \quad , \quad P = c\sqrt{2(x^2 - 1)} \quad , \quad D = \sqrt{\frac{2}{3}} [a(1-x) - b(x^2 - 1)] \quad . \quad (17)$$

Note that S and D both involve a and b .

The derivation of Eq. (3) is elementary. The ϕ is coupled to K^+K^- through an amplitude $\epsilon_\phi \cdot (p_{K^+} - p_{K^-})$, where the quantities denote 4-vectors. Thus the plane of (linear) ϕ polarization is related to that of the K^+K^- system in the J/ψ rest frame. By definition, we have taken the ϕ linear polarization vector to lie in the $x - y$ plane. We may define an angle ψ as that of the K^+ in the ϕ rest frame relative to the helicity axis (the negative of the direction of the J/ψ in that frame). The spatial components of the ϕ and J/ψ polarizations must be correlated since the decaying strange B is spinless. The J/ψ then has a single linear polarization state ϵ for each amplitude: In the J/ψ rest frame,

$$A_{\parallel} : \quad \epsilon = \hat{y} \quad ; \quad A_0 : \quad \epsilon = \hat{x} \quad ; \quad A_{\perp} : \quad \epsilon = \hat{z} \quad . \quad (18)$$

A unit vector n in the direction of the ℓ^+ in J/ψ decay may be defined to have components

$$(n_x, n_y, n_z) = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta) \quad (19)$$

where φ is the angle between the projection of the ℓ^+ on the K^+K^- plane in the J/ψ rest frame and the x axis. The sum over lepton polarizations then leads to a tensor in the J/ψ rest frame with spatial components (in the limit of zero lepton mass, assumed here)

$$\sum_{\ell^{\pm}\text{pol}} [\bar{u}\gamma_i v]^* [\bar{u}\gamma_j v] \sim L_{ij} \equiv \delta_{ij} - n_i n_j \quad . \quad (20)$$

Physically this tensor simply expresses the fact that massless lepton pairs couple only to transverse polarization states of the J/ψ , as expected from the structure of the electromagnetic interactions.

Taking account of the definition (6), we then find that the probability for the decay $B_s \rightarrow (\ell^+ \ell^-)_{J/\psi} (K^+ K^-)_{\phi}$ is proportional to

$$\sum_{\ell^{\pm}\text{pol}} |A|^2 = A_i A_j^* L_{ij} \quad , \quad (21)$$

where

$$A_i = A_0 \delta_{ix} \cos \psi - A_{\parallel} \delta_{iy} \sin \psi / \sqrt{2} + i A_{\perp} \delta_{iz} \sin \psi / \sqrt{2} \quad . \quad (22)$$

Consequently, when we use the definitions (19), we find

$$\begin{aligned} \frac{d^4 \Gamma [B_s \rightarrow (\ell^+ \ell^-)_{J/\psi} (K^+ K^-)_{\phi}]}{d \cos \theta \, d \varphi \, d \cos \psi \, dt} &= \frac{9}{32 \pi} [2 |A_0|^2 \cos^2 \psi (1 - \sin^2 \theta \cos^2 \varphi) \\ &+ \sin^2 \psi \{ |A_{\parallel}|^2 (1 - \sin^2 \theta \sin^2 \varphi) + |A_{\perp}|^2 \sin^2 \theta - \text{Im}(A_{\parallel}^* A_{\perp}) \sin 2 \theta \sin \varphi \} \\ &+ \frac{1}{\sqrt{2}} \sin 2 \psi \{ \text{Re}(A_0^* A_{\parallel}) \sin^2 \theta \sin 2 \varphi + \text{Im}(A_0^* A_{\perp}) \sin 2 \theta \cos \varphi \}] \quad . \quad (23) \end{aligned}$$

The overall normalization has been chosen to agree with our previous conventions when one integrates over angles. For \bar{B}_s decays the interference terms involving A_{\perp} amplitudes are of opposite sign and all other terms are unchanged.

Integration over $\cos \psi$ leads to the distribution

$$\frac{d^3\Gamma[B_s \rightarrow (\ell^+\ell^-)_{J/\psi}(K^+K^-)_\phi]}{d\cos\theta d\varphi dt} = \frac{3}{8\pi}[|A_0|^2(1 - \sin^2\theta \cos^2\varphi) + |A_{\parallel}|^2(1 - \sin^2\theta \sin^2\varphi) + |A_{\perp}|^2 \sin^2\theta - \text{Im}(A_{\parallel}^*A_{\perp}) \sin 2\theta \sin\varphi] \quad . \quad (24)$$

Performing the integrals over φ and taking account of the differing time-dependences of the decays of $B_s^{(\pm)}$, we obtain the result (3). This is a suitable single-angle distribution to employ if one wishes to disentangle the CP-even and CP-odd components of the B_s . The two-angle distribution (24) allows one to separate out the individual quantities $|A_0|^2$, $|A_{\parallel}|^2$, and $|A_{\perp}|^2$.

An interesting oscillation appears in the interference terms between CP-even and CP-odd decays. For example, in the two-angle distribution (24), since the respective time-dependences of the $B_s^{(+)}$ and $B_s^{(-)}$ decay amplitudes are $e^{-im_L t - \Gamma_L t/2}$ and $e^{-im_H t - \Gamma_H t/2}$, the term $-\text{Im}(A_{\parallel}^*A_{\perp})$ behaves as $|A_{\parallel}(0)A_{\perp}(0)| \sin(\Delta m t - \delta)e^{-\bar{\Gamma}t}$, where δ is a strong final-state phase shift difference: $A_{\parallel}(0)^*A_{\perp}(0) = |A_{\parallel}(0)A_{\perp}(0)|e^{i\delta}$. Thus, if one tags the flavor of the decaying B_s , one can observe the effects of Δm in the $J/\psi\phi$ final state through the interference of final states of opposite CP (if both are present). The oscillation term averages out to zero if the initial numbers of B_s and \bar{B}_s are equal.

The distributions (3), (23), and (24) also permit one to separate out the components $|A_0|^2$, $|A_{\parallel}|^2$, $|A_{\perp}|^2$, and the interference terms $\text{Im}(A_{\parallel}^*A_{\perp})$, $\text{Re}(A_0^*A_{\parallel})$, and $\text{Im}(A_0^*A_{\perp})$ for the decays $B^0 \rightarrow J/\psi K^{*0}$. (Here and subsequently we imply the sum over a process and its charge-conjugate.) Moreover, in the limit of flavor SU(3) symmetry, one expects the ratios of the relative components in $B^0 \rightarrow J/\psi K^{*0}$ to be the same as those *at proper time* $t = 0$ in the decays $B_s \rightarrow J/\psi\phi$ [19]. Thus, an analysis of $B^0 \rightarrow J/\psi K^{*0}$ can provide an independent estimate of the relative contributions of CP-even and CP-odd final states at $t = 0$ to the decays $B_s \rightarrow J/\psi\phi$, enhancing the ability to determine Γ_H and Γ_L .

If the K^* is observed to decay to the CP eigenstate $K_S\pi^0$, the amplitudes A_0 and A_{\parallel} refer (as in the case of $J/\psi\phi$) to the CP-even eigenstate, while A_{\perp} refers to the CP-odd eigenstate. The expected dominance of the CP-even eigenstate (see below) means that in $B^0 \rightarrow J/\psi K_S\pi^0$ events the CP asymmetry will tend to be opposite to that in $B^0 \rightarrow J/\psi K_S$ [12, 20]. Since the rates for observing the processes $B^0 \rightarrow J/\psi K_S$ and $B^0 \rightarrow J/\psi K^{*0} \rightarrow J/\psi K_S\pi^0$ are comparable (taking account of branching ratios and typical detection efficiencies), the incorporation of $J/\psi K_S\pi^0$ data may add statistical power to any experiment studying the $J/\psi K_S$ final state, even when the π^0 is not observed directly but its existence inferred.

The distribution (3) permits one to separate amplitudes of opposite parity from one another even if the K^+K^- system in $B_s \rightarrow J/\psi K^+K^-$ or the $K\pi$ system in $B \rightarrow J/\psi K\pi$ is not a vector meson [12]. This is easily seen by considering the density matrix ρ_{ij} of the J/ψ , expressed in terms of linear polarization states, so that the decay rate is proportional to $\rho_{ij}L_{ij}$, with L_{ij} defined in (20). If we integrate over φ , we find

$$\frac{d^2\Gamma}{d\cos\theta dt} \sim (\rho_{xx} + \rho_{yy}) \left(\frac{1 + \cos^2\theta}{2} \right) + \rho_{zz} \sin^2\theta \quad , \quad (25)$$

where ρ_{xx} and ρ_{yy} correspond to linear J/ψ polarization states in the plane of the two pseudoscalar mesons, while ρ_{zz} corresponds to J/ψ polarization perpendicular to this plane, and thus represents an amplitude with parity opposite to those contributing to ρ_{xx} and ρ_{yy} . For cases where each particle in the final state is a C eigenstate as in $B^0 \rightarrow J/\psi K^{*0} \rightarrow J/\psi K_S \pi^0$ and $B_s \rightarrow J/\psi \phi \rightarrow J/\psi K_S K_L$ the parity separation is also a CP separation and the transversity analysis can be used without the need to extract the vector resonance from nonresonant or other background.

The dominance of the $|A_0|^2$ contribution in $B^0 \rightarrow J/\psi K^{*0}$ decays [14, 21, 22] implies via flavor SU(3) that the $|A_0|^2$ contribution should also dominate $B_s \rightarrow J/\psi \phi$, and hence that $B_s^{(-)} \rightarrow J/\psi \phi$ is likely to be suppressed in comparison with $B_s^{(+)} \rightarrow J/\psi \phi$. Thus the initial angular distribution is very likely to be dominated by the $1 + \cos^2 \theta$ component. As time increases, the fraction of the angular distribution proportional to this component will decrease while that proportional to $\sin^2 \theta$ will increase. It should be possible to separate out the two components by a combined analysis in θ and proper decay time. If the $\sin^2 \theta$ component does not show up even at large times, a single-exponential fit to the decay should provide a good estimate of the lifetime of the CP-even eigenstate.

The angular distribution in (3) has the form of an ellipsoid which is prolate if $p(t) > 2m(t)$ and oblate if $p(t) < 2m(t)$. (Here we imagine an average over φ to have been performed.) If all three partial waves are equally populated $p(t) = 2m(t)$ since there are two partial waves with even CP and only one with odd CP. For this case the angular distribution is isotropic in $\cos \theta$ as expected.

If $p(0) > 2m(0)$ (i.e., if the CP-even decay is initially more than 2/3 dominant), if the CP-even eigenstate $B_s^{(+)}$ has the greater decay rate as expected, and if there is a non-zero odd-CP component $m(0) \neq 0$, then the angular distribution in transversity will be initially prolate but will eventually become oblate as the quantity $p(t) - 2m(t)$ changes sign. This effect can be noted by dividing the events into two bins with $|\cos \theta| < 1/2$ and $|\cos \theta| > 1/2$, denoted by E (equatorial) and P (polar) respectively:

$$E \equiv \int_{-1/2}^{1/2} d(\cos \theta) \frac{d^2 \Gamma}{d \cos \theta dt} = \frac{13}{32} p(t) + \frac{11}{16} m(t) \quad , \quad (26)$$

$$P \equiv \left[\int_{-1}^{-1/2} + \int_{1/2}^1 \right] d(\cos \theta) \frac{d^2 \Gamma}{d \cos \theta dt} = \frac{19}{32} p(t) + \frac{5}{16} m(t) \quad , \quad (27)$$

$$P - E = \frac{3}{16} [p(t) - 2m(t)] \quad (28)$$

The difference between the numbers in the two bins provides an experimental number whose sign will change in time under the assumptions noted above. This two-bin analysis of data can be adjusted for optimum statistics by changing the sizes of the bins to correspond to the initial values of $p(0)$ and $m(0)$.

The analysis performed by CDF [14] for $B_s \rightarrow J/\psi \phi$ was also based on a single-angle distribution, but it separated out the transverse component from the longitudinal component. In the absence of vertex cuts these would be, respectively, $\Gamma_T \equiv \int_0^\infty dt (|A_{\parallel}|^2 + |A_{\perp}|^2)$ and $\Gamma_0 \equiv \int_0^\infty dt (|A_0|^2)$. With a minimum vertex cut of 50 μm , the

result obtained was $\Gamma_0/(\Gamma_0 + \Gamma_T) = 0.56 \pm 0.21$ (stat) $^{+0.02}_{-0.04}$ (sys). The transverse component contains both CP-even and CP-odd contributions, while the longitudinal component is CP-even. The separation of transverse and longitudinal components makes sense only if the $B_s^{(-)} \rightarrow J/\psi\phi$ decay is negligible, or if the lifetime difference between $B_s^{(+)}$ and $B_s^{(-)}$ can be ignored.

Corresponding determinations of $\Gamma_0/(\Gamma_0 + \Gamma_T)$ for the decay $B^0 \rightarrow J/\psi K^{*0}$ are $0.65 \pm 0.10 \pm 0.04$ (CDF) [14], $0.97 \pm 0.16 \pm 0.15$ (ARGUS) [21], $0.80 \pm 0.08 \pm 0.05$ (CLEO) [22], and 0.74 ± 0.07 (world average) [14]. This last value is compatible with the corresponding one for $B_s \rightarrow J/\psi\phi$. A discrepancy would have indicated either a violation of SU(3) or the lifetime effect mentioned above.

The presence of two eigenstates with possibly differing lifetimes can affect any determination of $\tau(B_s)$. When observing the B_s in a final state of definite flavor, such as $D_s\pi$ or $D_s\ell\nu_\ell$, one will be effectively measuring the average lifetime $\bar{\Gamma}$ of the CP-even and CP-odd states. Most measurements reported up to now, including a recent CDF determination [23] leading to a world average [24] of $\tau(B_s) = 1.58 \pm 0.10$ ps, are of this type. This quantity is expected [4] to be very close to the B^0 lifetime, for which the world average [24] is $\tau(B^0) = 1.57 \pm 0.05$ ps. However, minimum-lifetime cuts can bias the sample against the CP-even (shorter-lived) component, leading to results which depend on the cut if a single-exponential fit is adopted.

To summarize, we have found that a combined analysis with respect to proper decay time and a single transversity angle in the decay $B_s \rightarrow J/\psi\phi$ can determine the lifetime of at least the CP-even and possibly the CP-odd mass eigenstates of the $B_s - \bar{B}_s$ system. Additional information about the properties of the $J/\psi\phi$ mode at proper time $t = 0$ can be obtained by a similar analysis of the decays $B^0 \rightarrow J/\psi K^{*0}$. These analyses can already be attempted with the data sample [14] now in hand.

We thank B. Kayser, R. Kutschke, K. Ohl, M. Paulini, E. A. Paschos, M. P. Schmidt, and W. Wester for useful discussions. This work was supported in part by the United States Department of Energy under Contracts No. DE AC02 76CH03000, DE FG02 90ER40560 and W-31-109-ENG-38.

References

- [1] N. Cabibbo, Phys. Rev. Lett. **10** (1963) 531; M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49** (1973) 652.
- [2] See, e.g., J. L. Rosner, Enrico Fermi Institute Report No. EFI 95-36, hep-ph/9506364, lectures presented at the VIII J. A. Swieca Summer School, Rio de Janeiro, Feb. 7–11, 1995, proceedings to be published by World Scientific.
- [3] R. Aleksan, A. Le Yaouanc, L. Oliver, O. Pène, and J. C. Raynal, Phys. Lett. B **316** (1993) 567.
- [4] M. B. Voloshin, N. G. Uraltsev, V. A. Khoze, and M. A. Shifman, Yad. Fiz. **46** (1987) 181 [Sov. J. Nucl. Phys. **46** (1987) 112]; I. I. Bigi *et al.*, in *B Decays*, edited by S. Stone (World Scientific, Singapore, 1994), p. 132; I. I. Bigi, Univ. of Notre Dame Report No. UND-HEP-95-BIG02, June, 1995, to be published in Physics Reports.
- [5] A. Soni, Brookhaven National Laboratory report BNL-62284, October, 1995, to be published in Proceedings of Lattice 95, Melbourne, Australia, July 10-13, 1995.
- [6] I. Dunietz, Phys. Rev. D **52** (1995) 3048.
- [7] G. Buchalla, I. Dunietz, and H. Yamamoto, FERMILAB-PUB-95/167-T, July, 1995, to be published in Phys. Lett. B.
- [8] T. E. Browder and S. Pakvasa, Phys. Rev. D **52** (1995) 3123.
- [9] L. L. Chau, W.-Y. Keung, and M. D. Tran, Phys. Rev. D **27** (1983) 2145; L. L. Chau, Phys. Rep. **95** (1983) 1; A. J. Buras, W. Slominsky, and H. Steger, Nucl. Phys. **B245** (1984) 369; A. Datta, E. A. Paschos, and U. Türke, Phys. Lett. B **196** (1987) 382; A. Datta, E. A. Paschos, and Y. L. Wu, Nucl. Phys. **B311** (1988) 35; I. Dunietz, Ann. Phys. (N.Y.) **184** (1988) 350.
- [10] N.-P. Chang and C. A. Nelson, Phys. Rev. Lett. **40** (1978) 1617; Phys. Rev. D **20** (1978) 2923; C. A. Nelson, Phys. Rev. D **30** (1984) 107, 1937; **32** (1985) 1848(E).
- [11] H. J. Lipkin, in *Proceedings of the SLAC Workshop on Physics and Detector Issues for a High-Luminosity Asymmetric B Factory*, edited by David Hitlin, published as SLAC, LBL and Caltech reports SLAC-373, LBL-30097 and CALT-68-1697, 1990, p. 49.
- [12] I. Dunietz, H. R. Quinn, A. Snyder, W. Toki and H. J. Lipkin, Phys. Rev. D **43** (1991) 2193.
- [13] I. Dunietz, Appendix in *B Decays*, edited by S. Stone (World Scientific, Singapore, 1994), p. 550.

- [14] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **75** (1995) 3068.
- [15] A. Bohr, Nucl. Phys. **10** (1959) 486.
- [16] J. L. Rosner, Phys. Rev. D **42** (1990) 3732.
- [17] L. D. Landau, Dokl. Akad. Nauk SSSR **60** (1948) 207; C. N. Yang, Phys. Rev. **77** (1950) 242, 722.
- [18] G. Kramer and W. F. Palmer, Phys. Rev. D **45** (1992) 193; **46** (1992) 2969, 3197; Zeit. Phys. C **55** (1992) 497.
- [19] I. Dunietz, in *Proceedings of the Workshop on B Physics at Hadron Accelerators*, Snowmass, Colorado, June 21 – July 2, 1993, edited by P. McBride and C. S. Mishra (Fermilab report Fermilab-CONF-93/267, 1993), p. 83.
- [20] B. Kayser, M. Kuroda, R. D. Peccei, and A. I. Sanda, Phys. Lett. B **237** (1990) 508.
- [21] ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. B **340** (1994) 217.
- [22] CLEO Collaboration, M. S. Alam *et al.*, Phys. Rev. D **50** (1994) 43.
- [23] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **74** (1995) 4988.
- [24] I. J. Kroll, presented at 17th International Symposium on Lepton and Photon Interactions, Beijing, 11 August 1995.