

Taxonomic evaluation of eleven species of *Microcyclops* Claus, 1893 (Copepoda, Cyclopoida) and description of *Microcyclops inarmatus* sp. n. from America

Martha Angélica Gutiérrez-Aguirre¹, Adrián Cervantes-Martínez¹

¹ Universidad de Quintana Roo (UQROO), Unidad Cozumel, Av. Andrés Quintana Roo s/n, 77600, Cozumel, Quintana Roo México

Corresponding author: Martha Angélica Gutiérrez-Aguirre (margutierrez@uqroo.edu.mx)

Academic editor: D. Defaye | Received 10 December 2015 | Accepted 2 June 2016 | Published 6 July 2016

<http://zoobank.org/E0086EEB-7B77-4F71-B6BE-E8637207AF21>

Citation: Gutiérrez-Aguirre MA, Cervantes-Martínez A (2016) Taxonomic evaluation of eleven species of *Microcyclops* Claus, 1893 (Copepoda, Cyclopoida) and description of *Microcyclops inarmatus* sp. n. from America. ZooKeys 603: 33–69. doi: 10.3897/zookeys.603.7480

Abstract

Description and meristic analysis of eleven species of *Microcyclops* recorded in America were performed based on the examination of type specimens and fresh material. Microscopic analysis of oral appendages, such as the shape and armature of the distal coxal endite of the maxilla, the ornamentation on the caudal surface of the antenna, and the intercoxal sclerites and armament of the inner basis of all swimming appendages, were characteristics that allowed the differentiation between species. Among these species, our study confirmed the synonymy of *M. diversus* Kiefer, 1935 with *M. ceibaensis* (Marsh, 1919). The results of our observations showed that *M. alius* (Kiefer, 1935) is a junior synonym of *M. dubitabilis* Kiefer, 1934; the latter being confirmed as a valid species. Also, it is proposed that the records of *M. rubellus* (Lilljeborg, 1901) and *M. varicans* (Sars, 1863) in America should be revised as there are serious doubts about their distribution in America. The analysis suggested that *M. anceps pauxensis* Herbst, 1962 is distinct from *M. anceps* var. *minor* Dussart, 1984 and that both are likely different from *M. anceps anceps* (Richard, 1897). Finally a full morphological description of adult females of *Microcyclops inarmatus* sp. n. is presented.

Keywords

Diversity, Mexico, morphology, species richness

Introduction

In America, 16 species and subspecies of *Microcyclops* Claus, 1893 have been described and recorded: *M. alius* (Kiefer, 1935), *M. anceps anceps* (Richard, 1897), *M. anceps pauxensis* Herbst, 1962, *M. anceps* var. *minor* Dussart, 1984, *M. ceibaensis* (Marsh, 1919), *M. dubitabilis* Kiefer, 1934, *M. echinatus* Fiers, Ghenné & Suárez-Morales, 2000, *M. elongatus* (Lowndes, 1934), *M. finitimus* Dussart, 1984, *M. furcatus* (Daday, 1905), *M. mediasetosus* Dussart & Frutos, 1985, *M. medioides* Dussart & Frutos, 1985, *M. pumilis* Pennak & Ward, 1985, *M. rubellus* (Lilljeborg, 1901), *M. diversus* Kiefer, 1935, and *M. varicans* (Sars, 1863).

In her publication of an identification key for South American cyclopoids, Reid (1985) proposed that *Microcyclops diversus* is a probable synonym of *M. ceibaensis* (in 1986, this opinion was based on similarities in the fourth leg observed by the same author) and that *M. anceps* var. *minor* is a synonym of *M. anceps pauxensis*. Rocha (1998) proposed a set of morphological features that would be useful for distinguishing five species previously recorded in Brazil and suggested that *M. alius* is a possible synonym of *M. dubitabilis*. However, Reid (1990) had previously suggested that *M. dubitabilis* is a possible synonym of *M. varicans*.

Therefore, some questions on the taxonomic status of some species of *Microcyclops* in America remain unresolved. These taxonomic problems may be related to the lack of thorough and rigorous species descriptions. Rocha (1998), Mirabdullayev (1998, 2007), and Mirabdullayev and Urazova (2006) have documented morphological features that are useful for differentiating some species of the genus. For instance, they proposed the following morphological features as diagnostic: ornamentation of dorsal margins of prosomites, presence or absence of pores on second endopodite of first leg, ornamentation of the inner margin of basipodite of first leg, ornamentation of caudal ramus and caudal setae, relative lengths of caudal setae, proportions of second endopodite of fourth leg, and general ornamentation of fourth leg.

In Mexico, some species with uncertain taxonomic status have been recorded, including *M. ceibaensis*, *M. anceps*, and *M. dubitabilis* (Elías-Gutiérrez et al. 2008). In this paper, we propose a set of morphological features that are useful for distinguishing between these species, which have been documented by biological inventories of the country. These features include the mouth appendages, the ornamentation of intercoxal sclerites, and the inner margin of the basis of the first to fourth swimming legs.

Methods

The morphological analysis was performed following current standards for the taxonomic study of cyclopoid copepods (see Williamson and Reid 2001).

Material examined. The evaluation included analyses of holotypes, paratypes, and museum specimens deposited in different collections: the Collection of Zoo-plankton of ECOSUR at Chetumal (**ECO-CH-Z**), the collection of Copepoda of

the Muséum National d'Histoire Naturelle, Paris (**MNHN**), the Staatliches Museum für Naturkunde, Karlsruhe (**SMNK**) and the National Museum of Natural History, Smithsonian Institution, Washington, DC (**USNM**) (Table 2, as Suppl. material 1).

Terminology used for the armament of each appendage(s) follows Huys and Boxshall (1991):

A1	Antennule
A2	Antenna
BspA2	Antennal basipodite
Bsp	Basipodite
Enp₁-Enp_n	First to "n" endopodal segment
Exp₁-Exp_n	First to "n" exopodal segment
P1, P2, P3, P4	First, second, third, and fourth swimming legs
P5	Free segment of fifth leg

Lateral, outermost terminal, outer median terminal, inner median terminal, innermost terminal, and dorsal caudal setae are coded as setae II, III, IV, V, VI, and VII, respectively.

The morphology of several species was examined using light microscopy: A1, A2, the mouthparts, the entire structure of all of the swimming legs, and other taxonomically relevant structures, such as the frontal or caudal ornamentation of BspA2, the ornamentation of the distal coxal endite of the maxilla, and the ornamentation of maxillular palp, were illustrated with the aid of a camera lucida.

Sources for the morphological data were the types, paratypes, and other museum specimens (Table 2, as Suppl. material 1), and original descriptions of eleven named species and two named subspecies recorded in America. Detailed descriptions based on the microscopic and morphometric analyses of the adult females of each species are presented.

Results

Descriptive section

Below those morphological structures which are shared by all the species examined herein are described.

Antennule 11- or 12-segmented (Fig. 1). In the basic 12-segmented structure (Fig. 1A), each segment was armed with setae (s), spines (sp) or aesthetascs (ae) in the following order: (1) 8s; (2) 4s; (3) 2s; (4) 6s; (5) 3s; (6) 1s + 1sp; (7) 2s; (8) 3s; (9) 2s + 1ae; (10) 2s; (11) 2s + 1ae; (12) 7s + 1ae. In the case of the 11-segmented antennule the third and fourth segments are entirely or partially fused (Fig. 1B); then, the third segment bears 8s.

Antenna with coxa (without seta), Bsp (with 2 medial setae + one lateral seta representing Exp), and 3-segmented Enp (Fig. 3B). Labrum with strong teeth on distal



Figure 1. Morphology of antennules. **A** Antennule with 12 segments (*Microcycllops ceibaensis* from km 51-2) **B** Morphological variation in antennules with 11 segments (*M. dubitabilis* from km 51-2). Scale bars: 50 µm.

Table 1. Setation formula of the swimming legs in the *Microcycllops* species here examined (spine in Roman numerals, seta in Arabic numerals).

	Coxa	Basis	Exp	Enp
P1	0-1	1-I, or 1-0	I-1; III-5	0-1; 1-I-4
P2	0-1	1-0	I-1; IV-5	0-1; 1-I-5
P3	0-1	1-0	I-1; IV-5	0-1; 1-I-5
P4	0-1	1-0	I-0; III-5	0-1; 1-II-3

rim and strong, distal hairs overhanging distal rim (Fig. 6D). Mandible with toothed gnathobase; the innermost teeth bi-toothed. Innermost margin of mandibular gnathobase with one spinulose seta, palp with two long and one short seta. No spinules next to mandibular palp (Fig. 6E).

Praecoxal arthrite and palp of maxillule naked; praecoxal arthrite with 3 chitinized distal claws, and one spinulose seta on caudal side. Inner margin with one biserially plumose seta plus six naked setae (Fig. 6F). Maxilla with praecoxa and coxa partially fused, praecoxal endite with two setae, coxa naked with proximal endite bearing one seta (Fig. 6L) and distal endite with two armed long setae (Fig. 6M). Claw-like basal endite armed, and Enp one- or two-segmented.

Maxilliped with syncoxa bearing 2 or 3 spiniform setae, Bsp with two setae; and Enp two-segmented, first segment with 1 seta, second segment with 3 setae (Fig. 6N).

Armature formula of P1–P4 as in Table 1, endopods and exopods two-segmented in all swimming legs. Urosome five-segmented, fifth pediger bearing one free segment with one apical seta (fifth leg), and one lateral seta inserted on pediger (Fig. 8C). Detailed description of the species is provided in the next section. The material examined for each species is provided in Table 2, as Suppl. material 1.

Order: Cyclopoida Burmeister, 1835

Family: Cyclopidae Rafinesque, 1815

Subfamily Cyclopinae Rafinesque, 1815

Genus *Microcycllops* Claus, 1893

Microcycllops inarmatus sp. n.

<http://zoobank.org/687BDBC3-853D-437E-9310-4146F210094A>

Figures 2–5

Microcycllops varicans Reid, 1992; Trans. Am. Microsc. Soc. 111(3), p: 249–250, figs 8d, 9c.

Holotype. One adult female dissected on two slides: A1, A2 (slide 1, ECOCH-Z-09337); mandible, maxillule, maxilla, maxilliped, P1–P4, and urosome (slide 2, ECOCH-Z-09337). Collected 13.I.1998.




Figure 2. *Microcyclops inarmatus* sp. n. Adult female, holotype (except **A**). **A** Habitus of one paratype specimen (ECOCH-Z-09338) **B** Urosome ventral, note that the last fraction of the inner median terminal caudal seta is separated (ECOCH-Z-09337) **C** P1, caudal (ECOCH-Z-09337) **D** P4, caudal (ECOCH-Z-09337). Scale bars: 50 µm.

Paratypes. 10 adult females preserved in 90% ethanol with a drop of glycerine. ECOCH-Z-09338. Collected 13.I.1998.

Type locality. A pond in km 51 lado 1, Villahermosa-Frontera highway 18°23'16"N; 92°47'00"W.

Etymology. the name of the species means un-armed in Latin; it refers to the absence of ornamentation on the intercoxal sclerites, the lack of spinules at base of caudal furcal setae, the reduced number of setae on second antennal endopod, and the reduced ornamentation on antennal basis.

Additional material. One adult female collected 1.02.1935 from Laguna Rincon, Haiti (slide SMNK-2391; labelled as *Microcyclops dubitabilis* with A1, maxilla, P1-P4). One adult female collected from Laguna Rincon, Haiti (slide SMNK-2392; labelled as *M. dubitabilis* with urosome).

One adult female collected 05.1986 from Shark river slough, Everglades National Park, Florida, USA (slide 2 of 7, USNM-251321; labelled as *M. varicans* with A1, A2, P1-P4, and urosome).

Diagnosis. Adult female: Dorsal margin of prosomal somites smooth; body length 565 to 615 µm in paratypes. Antennule 12-segmented, not reaching the distal margin of the first prosomal segment (Fig. 2A). Fifth pediger nude; cylindrical free segment of P5 more than 3 times as long as wide, with tiny inner spine; genital double somite expanded proximally. Anal somite with strong spines on ventral distal margin; length to width ratio of caudal ramus less than 3; no spinules at base of lateral and outermost terminal caudal setae (Fig. 2B). Outer median terminal and inner median terminal caudal setae with heteronomous setulation (Fig. 2B). Endopodites and exopodites of P1-P4 bisegmented with setation formula as in Table 1, inner basis of P1 with long spine (long arrow in Fig. 2C), Enp2P1 with one pore on lateral margin (short arrow in Fig. 2C). Intercoxal sclerites of P1-P4 unarmed, long setules on inner margin of basipodites of P1-P4, medial spine of Enp2P4 almost as long as the segment and twice the length of the lateral spine (Fig. 2D).

Adult male: unknown.

Description of female. *Antennule* 12-segmented; antenna with 3-segmented Enp armed with 1, 6, and 7 setae respectively (Fig. 3A, B –position of missing setae in specimens is arrowed). Antennal basis with one group of spinules on the basal-outer margin in caudal view (Fig. 3A, B); frontal surface of the antennal basis with two rows of tiny spinules (Fig. 3C). Nine teeth on mandibular gnathobase (Fig. 3D). Maxillule (Fig. 3E) with unarmed palp; apical region of maxillular palp with two armed setae plus one smooth seta, three setae (one armed) on lateral lobe, proximal seta smooth (Fig. 3F). Distal coxal endite of the maxilla with two long setae: the proximal seta with two tiny spines at its base and bifurcated, distal seta with one row of tiny spines along one margin (Fig. 3G, H). Basipodite with one claw-like projection bearing 5-7 strong spines on the concave margin and one long, armed seta on its base; two-segmented Enp bearing 2 and 3 setae respectively (Fig. 3G). Because of the condition of the microscope slide preparatum we could not verify one basal seta on maxillar Enp1 (arrowed in Fig. 3H). Maxilliped with syncoxa (3 setae), basis (2 setae), and two-segmented Enp bearing 1 and 3 setae. Basis of maxilliped with a row of spinules on frontal and caudal surfaces (Fig. 3I).

Basipodites of P1-P3 with long hair-like setules on the inner margins; one row of tiny spinules along the lateral margins of coxa; intercoxal sclerites naked (Fig. 4A-D).




Figure 3. *Microcycllops inarmatus* sp. n. Adult female. **A** Antenna, caudal (USNM-251321), note that the position of missing seta is arrowed **B** Antenna, caudal (ECOCH-Z-09337), note that the position of missing seta is arrowed **C** Antennal basipodite, frontal (ECOCH-Z-09337) **D** Mandible (ECOCH-Z-09337) **E** Maxillule (ECOCH-Z-09337) **F** Maxillular palp (ECOCH-Z-09337) **G** Maxilla (ECOCH-Z-09337) **H** Maxilla (SMNK-2391) **I** Maxilliped (ECOCH-Z-09337). Scale bars: 50 µm.




Figure 4. *Microcycllops inarmatus* sp. n. Adult female. **A** P1, frontal (USNM-251321) **B** P1, frontal (SMNK-2391) **C** P2, caudal (ECOCH-Z-09337) **D** P3 coxa, basis, and sclerite, caudal (ECOCH-Z-09337) **E** P4, caudal, Exp unfigured (USNM-251321) **F** P4, frontal, Exp unfigured (SMNK-2391) **G** Fifth pediger and P5 (USNM-251321) **H** Fifth pediger and P5 (SMNK-2392). Scale bars: 50 µm.




Figure 5. *Microcyclops inarmatus* sp. n. Adult female. **A** Anal somite and caudal rami, ventral (USNM-251321) **B** Terminal caudal setae (USNM-251321) **C** Urosome, ventral (SMNK-2392) **D** Anal somite and caudal rami, ventral (SMNK-2392) **E** Anal somite and caudal rami (ECOCH-Z-0679) **F** Hyaline fringes of urosome (Pajonal). Scale bars: 50 µm.

Basis of P1 with one long spine on inner margin; spine reaching distal middle of Enp2P1 and armed with heteronomous setulation: hair-like setules on its base, tiny spinules distally (Figs 2C; 4A, B). One pore on the lateral margin of Enp2P1 (Fig. 4A, B). Basipodite of P4 with long hair-like setules on inner margin; P4 intercoxal sclerite quadrangular (Figs 2D; 4E, F), Enp2P4 2.14 ± 0.2 times as long as wide; and medial spine 1.97 ± 0.25 as long as lateral spine and 0.91 ± 0.04 as long as the segment (Figs 2D; 4E, F).

Fifth pediger nude; P5 with one cylindrical free segment, 3.23 ± 0.4 times as long as wide, bearing one tiny medial spinule. Free segment 0.27 ± 0.01 times as long as distal seta (Figs 2B; 4G, H). Hyaline fringes of urosomites serrated (Fig. 2B), petaloid or rounded (Fig. 5 A, C–F). Length to width ratio of caudal ramus 2.54 ± 0.44 , inner margin naked; no spinules at base of lateral caudal (II) and outermost terminal caudal setae (III) (Fig. 2B). Only 5–8 strong spinules present ventrally on the distal margin of anal somite, no spinules dorsally (Fig. 5 A, C–E). Lateral caudal seta (II) inserted at $58.6 \pm 3.9\%$ of caudal ramus.

Dorsal caudal seta (VII) 0.9 ± 0.1 times as long as caudal ramus, and innermost terminal caudal seta (VI) 1.4 ± 0.04 times as long as caudal rami (Fig. 2B). Relative lengths of terminal caudal seta from outermost caudal seta to innermost caudal seta is $1.0 : 4.9 : 7.3 : 1.6$ (Figs 2B; 5B, C). Outer median terminal caudal seta (IV) and inner median terminal caudal seta (V) with heteronomous setulation: proximally with spinule-like setules and distally with long and fine setules (Figs 2B; 5B, C).

Microcyclops dubitabilis Kiefer, 1934

Figures 6–8

Description of female. *Antennule* 11, or 12-segmented (intra- and interpopulation variation); 3-segmented endopod of antenna bearing 1, 9, and 7 setae, respectively (Fig. 6A, B). Antennal basis with three long rows of spinules on caudal surface (Fig. 6B): two basal rows, and one median row; frontal surface of antennal basis with two rows of tiny spinules (Fig. 6C). Labrum with 6–7 teeth between two curved lateral teeth, and 3–4 strong spinules on each round projections of the plate (Fig. 6D). Eight teeth on mandibular gnathobase (Fig. 6E). *Maxillule* as in Fig. 6F, maxillular palp with one armed seta plus two smooth setae apically, three smooth setae on lateral lobe, and one proximal nude seta (Fig. 6G, H). Maxilla with armed setae on distal coxal endite: proximal seta with one long spine-like setule at its base and bifurcated apically, distal seta with one row of tiny spines along inner margin (Fig. 6I–K, M). *Basipodite* with claw-like projection bearing 6–8 thin spinules on concave margin and one long seta on its base; this seta armed with two rows of spinules (long spinules on inner margin, and short spinules on outer margin) (Fig. 6I–L). Maxilla with two-segmented Enp bearing 2 and 3 setae respectively (Fig. 6I–L). Maxilliped with syncoxa (3 setae), basis (2 setae), and two-segmented Enp bearing 1 and 3 setae, respectively. Basis of the maxilliped nude, two spinules present on frontal surface of Enp1 (Fig. 6N).




Figure 6. *Microcyclops dubitabilis* Kiefer, 1934. Adult female. **A** Antenna, frontal (SMNK-2204) **B** Antenna, caudal (km 51-1) **C** Antennal basipodite, frontal (km 51-1) **D** Labrum (km 51-1) **E** Mandible (km 51-1) **F** Maxillule (km 51-1) **G** Maxillular palp (km 51-1) **H** Maxillular palp (USNM-251322) **I** Maxilla (USNM-251322) **J** Maxilla (SMNK-2081) **K** Maxilla (SMNK-2204) **L** Maxilla (km 51-1) **M** Distal coxal endite **N** Maxilliped (km 51-1). Scale bar: 50 µm.




Figure 7. *Microcycllops dubitabilis* Kiefer, 1934. Adult female. **A** First and second prosomal somite, dorsal (SMNK-2189) **B** P1, caudal (USNM-251322) **C** P1, frontal (SMNK-2081) **D** P1, frontal (km 51-1) **E** P4, caudal (USNM-251322) **F** P4, frontal (SMNK-2189) **G** P4, caudal (MNHN-Cp6764) **H** P4, caudal (km 51-1) **I** Fifth pediger, P5, genital double somite, ventral (USNM-251322) **J** Fifth pediger, P5, genital double somite, ventral (MNHN-Cp5398) **K** Fifth pediger, P5, genital double somite, ventral (SMNK-2204). Scale bars: 50 µm.




Figure 8. *Microcycllops dubitabilis* Kiefer, 1934. Adult female. **A** Caudal rami, ventral (USNM-251322) **B** Anal somite and caudal rami, ventral (SMNK-2204) **C** Urosome, ventral (SMNK-2081) **D** P6 (km 51-1). Scale bars: 50 µm.

Dorsal margin of prosomal somites smooth (Fig. 7A). Basis of P1 medially hairy. One short spine present on inner margin, spine biseriately armed with spinule-like setules (homonomous ornamentation) and reaching slightly beyond distal margin of the Enp1P1. Intercoxal sclerite of P1 naked (Fig. 7B–D). Pore on lateral margin of Enp2P1 sometimes present (interpopulation variation). Inner margin of basis of P2 and P3 hairy, and intercoxal sclerites of these swimming legs naked (unfigured). Inner margin of P4 basis with short hairs; intercoxal sclerite naked, rectangular (Fig. 7E–H), Enp2P4 1.9 ± 0.1 times as long as wide; medial spine 1.8 ± 0.3 times as long as lateral spine, and 0.8 ± 0.1 times as long as segment. Apical spines of Enp2P4 are subequal only in female USNM-251322 (Fig. 7E).

Fifth pediger nude; P5 free segment cylindrical, 3.6 ± 0.8 times as long as wide, without inner spine. Free segment 0.4 ± 0.1 times the length of the distal seta (Figs 7I–K; 8C). Length to width ratio of caudal ramus 2.4 ± 0.2 , inner margin naked. With or without spines at base of seta II (intrapopulation variation), spinules always present at base of setae III. Distal margin of anal somite bearing spinules: medial spinules are longer than lateral ones on ventral surface; spinule row can extend laterally or dorsally (Fig. 8 A–C). Seta II inserted at $71 \pm 5.7\%$ of caudal ramus.

Seta VII 1.02 ± 0.3 times as long as caudal ramus, and seta VI 1.4 ± 0.2 times longer than caudal ramus. Relative lengths of terminal caudal seta from outermost caudal seta to innermost caudal seta are $1.0 : 4.9 : 7.1 : 1.6$ (Fig. 8A, C). Seta IV and seta V with heteronomous setulation: proximally with spine-like setules and distally with long and fine setules (Fig. 8A–C). Sixth leg with two medial spines and one lateral seta (Fig. 8D).

Microcyclops ceibaensis (Marsh, 1919)

Figures 9–11

Description of female. *Antennule* 12-segmented (Fig. 9A). Antenna with 3-segmented endopod with 1, 9, and 7 setae, respectively (Fig. 9F). Frontal surface of antennal basis with one basal row of spinules arranged in arc next to medial (inner) margin, and one median row of spinules next to lateral (outer) margin. (Fig. 9D, E). Caudal surface of antennal basis with two basal rows of spinules arranged in arc, plus two rows of long spinules on outer margin (Fig. 9F). Labrum with 7 marginal teeth between two lateral curved teeth, and two rows of long spinules (6) overhanging distal margin (Fig. 9B, C). Gnathobase of the mandible with eight teeth (Fig. 9G). Maxillular palp with three apical setae, three setae on lateral lobe, and one proximal seta. The proximal seta armed on both margins, one seta on lateral lobe and one apical seta with setules (Fig. 9H). Distal coxal endite of the maxilla with two long setae: the proximal seta with two long basal spinules and bifurcated apically, distal seta smooth (Fig. 9I). Basipodite with one claw-like projection bearing thin spinules on concave margin, and one long smooth seta on its base. One-segmented Enp bearing 5 setae (Fig. 9I). Maxilliped with syncoxa (2 setae), basis (2 setae), and two-segmented Enp bearing 1 and 3 setae, respectively. Ornamentation of setae on syncoxa and Enp1 variable (intrapopulation variation, arrowed in Fig. 9J). Syncoxa and basis of maxilliped with a row of spinules on caudal surface next to lateral margin (Fig. 9J).

Dorsal margin of *prosomal somites* slightly serrated (Fig. 10A). Basis of P1–P3 with pilose inner margin. Intercoxal sclerites of P1–P3 with one row of short spinules (Fig. 10B–G), in some populations the sclerite of P3 with two rows of spinules (Fig. 10H). Enp2P1 with two pores on lateral margin (Fig. 10C). Because of the condition of the specimen in slide USNM-222299, it was not possible to verify the presence of spinules on the sclerite as well as the pores on the second endopodal segment of P1 (Fig. 10B).

P1 basis with long medial spine reaching distal third of Enp2P1. Spine ornamented with long setules near base and with short spinule-like setules more distally (Fig.




Figure 9. *Microcyclops ceibaensis* (Marsh, 1919). Adult female. **A** Antennule, segments 2-12 (USNM-222299) **B** Labrum (USNM-222299) **C** Labrum (km 51-2) **D** Antennal basipodite, frontal (USNM-222299) **E** Antennal basipodite frontal (km 154) **F** Antenna, caudal (km 154) **G** Mandible (km 51-1) **H** Maxillule (km 154) **I** Maxilla (km 51-2) **J** Maxilliped (km 154). Scale bars: 50 µm.




Figure 10. *Microcycllops ceibaensis* (Marsh, 1919). Adult female. **A** Prosomal fringes, dorsal **B** P1, frontal (USNM-222299) **C** P1, caudal (km 51-2) **D** P2 intercoxal sclerite, inner coxa and basis, caudal (USNM-222299) **E** P2, frontal (km 51-2) **F** P3 sclerite, inner coxa and basis, frontal (USNM-222298) **G** P3 intercoxal sclerite, coxa and basis, caudal (km 51-1) **H** P3 intercoxal sclerite, caudal (km 51-2) **I** P4 intercoxal sclerite, coxa and basis, caudal (USNM-222299) **J** Enp3P4 (USNM-222299) **K** P4, caudal (km 51-2) **L** P6. Scale bars: 50 μ m.




Figure 11. *Microcyclops ceibaensis* (Marsh, 1919). Adult female. **A** Fifth pediger, dorsal (USNM-222299) **B** Anal somite and caudal rami, dorsal (USNM-222299) **C** Urosome, ventral (km 51-1) **D** Caudal ramus and caudal setae, ventral (km 51-1) **E** Detail of inner median caudal seta (km 154) **F** Detail of inner median caudal seta (km 51-1). Scale bars: 50 µm.

10B, C). Inner margin of P4 basis with strong spinules. Intercoxal sclerite rectangular, and ornamented with two rows of spinules (Fig. 10I, K). Enp2P4 2.2 ± 0.1 times as long as wide; medial spine 1.5 ± 0.1 times as long as the lateral spine, and 0.6 ± 0.06 times as long as segment (Fig. 10J, K). Sixth leg with one long seta plus two short spines (Fig. 10L).

Fifth pediger nude (Fig. 11A). Urosomal somites with serrated hyaline fringes (Fig. 11B, C). Fifth leg with one cylindrical free segment 3.8 ± 1.4 times as long as wide;

tiny spinule present on inner margin. P5 free segment 0.3 ± 0.1 times the length of the distal seta (Fig. 11C). Distal margin of anal somite with a continuous row of strong spinules on ventral and dorsal surfaces (Fig. 11B, C). Caudal ramus 3.6 ± 0.4 times longer than wide, inner margin naked. Spinules present at base of caudal setae II and III; seta II inserted at $69 \pm 3.2\%$ of the caudal ramus (Fig. 11C).

Seta VII 0.7 ± 0.1 times as long as caudal ramus, seta VI 0.8 ± 0.1 times as long as caudal ramus. Relative lengths of terminal caudal seta from outermost caudal seta to innermost caudal seta: $1.0 : 5.7 : 9.5 : 1.8$ (Fig. 11C, D). Seta IV and seta V with homonomous setulation, with long and fine setules at whole length (Fig. 11D); inner median terminal caudal seta (V) with interrupted row of setules along the proximal, lateral margin (Fig. 11E, F).

***Microcyclops echinatus* Fiers, Ghenné & Suárez-Morales, 2000**

Figures 12, 13

Description of female. This description is a complement to the original description of Fiers et al. (2000). Frontal surface of antennal basis with one basal, inner row of spinules arranged in an arc, and one longitudinal row of spinules near lateral margin (Fig. 12A). Antenna with three-segmented endopod bearing 1, 9, and 7 setae, respectively (Fig. 12B). Caudal surface of antennal basis with two rows of long spinules next to exopodal seta, one group of long spinules at basal position, one basal row on inner margin and another basal row on outer margin (Fig. 12B). Nine teeth on the distal margin of the labrum (Fig. 12C). Eight teeth present on gnathobase of mandible (Fig. 12D). Maxillular palp with three apical setae (one of these setae armed with long setules); lateral lobe with three setae, the longer seta armed; proximal seta nude (Fig. 12E). Maxillar basipodite with one claw-like projection bearing thin spines on concave margin and one long seta with one (Fig. 12F) or four tiny spinules (Fiers et al. 2000); maxilla with two-segmented Enp bearing 2 and 3 setae on first and second endopodal segments, respectively (Fig. 12F). Maxilliped with syncoxa (3 setae), basis (2 setae), and two-segmented Enp bearing 1 and 3 setae. Syncoxa and basis of maxilliped with rows of spinules on caudal surface (Fig. 12G).

Two pores on lateral margin of second endopodal segment of P1, very long spinules present at insertion of apical spine of Enp2P1. Long medial spine of P1 basis with heteronomous setulation (Fig. 13A). Inner margin of P1–P3 basis with long hair-like setae (Fig. 13A, B), inner margin of P4 basis with one row of tiny spinules and one row of long setules (Fig. 13C). Intercoxal sclerites of all swimming legs ornamented on caudal surface: P1 with one row of spinules and P2 to P4 with two rows of spinules (Fig. 13A–C). Enp2P4 2.5 ± 0.1 times as long as wide; medial spine is 2.0 ± 0.1 times as long as lateral spine, and 0.8 ± 0.1 times as long as the segment.

Fifth pediger with rows of spinules on ventro-lateral surfaces. Fifth leg with tiny spinule on inner margin (Fig. 13D); cylindrical free segment 3.7 ± 0.1 times longer than wide and 0.45 ± 0.01 times as long as distal seta of P5. Caudal ramus 5.9 ± 0.4




Figure 12. *Microcyclops echinatus* (Fiers et al., 2000). Adult female (km 51-2). **A** Antennal basipodite, frontal **B** Antenna, caudal **C** Labrum **D** Mandible **E** Maxillule **F** Maxilla **G** Maxilliped. Scale bar 50 μm .




Figure 13. *Microcycllops echinatus* (Fiers et al., 2000). Adult female (Guanal). **A** P1, frontal **B** P3, caudal **C** P4 coxa, basis, sclerite, caudal **D** Fifth pediger, P5 and genital double-somite, ventral **E** Anal somite, caudal rami, and caudal setae, ventral. Scale bars: 50 µm.

times longer than wide. Seta VII 0.5 ± 0.1 times as long as caudal ramus, seta VI 0.5 ± 0.05 times as long as caudal ramus. Relative lengths of terminal caudal seta from outermost to innermost caudal seta, 1.0 : 6.5 : 10.4 : 2.0. Seta IV and seta V with homonomous setulation, bearing long and fine setules (Fig. 13E).

Microcyclops finitimus Dussart, 1984

Figure 14

Description of female. The following description is complementary to the original description of Dussart (1984). Antennule 12-segmented. Dorsal margin of prosomal somites 1 to 3 smooth (unfigured); hyaline fringe of fifth pediger serrated dorsally (Fig. 14A). Intercoxal sclerite of P1 smooth, inner margin of P1 basis with long hair-like setules, without spine on inner margin (Fig. 14B). Enp2P1 with one pore on lateral margin. Row of long spinules present at base of apical spine and lateral seta of Enp2P1 (Fig. 14C). Inner margin of P4 basis with long setules, intercoxal sclerite quadrangular, with one row of long spinules on caudal surface (Fig. 14D). Enp2P4 2.2 times as long as wide; medial spine 1.4 times as long as lateral spine, and 0.8 times as long as segment.

Anal somite with continuous row of spinules along distal margin (ventrally and dorsally), but on ventral surface medial spinules are longer and stronger than lateral spinules (Fig. 14E). No spinules at base of caudal seta II, but spinules present at base of caudal seta III; length to width ratio of caudal ramus 2.7. Relative lengths of terminal caudal setae from outermost to innermost seta, 1.0 : 6.1 : 8.9 : 2.1. Caudal setae IV, and V with homonomous setulation, bearing long and fine setules (Fig. 14E). Dorsal caudal seta (VII) 0.7 times as long as caudal rami, and innermost caudal seta (VI) 1.2 times longer than caudal rami. Lateral caudal seta (II) inserted at 75.5% of caudal ramus length.

Microcyclops anceps anceps (Richard, 1897)

Figures 15–17

Description of female. Dorsal posterior margin of second prosomal somite with crenulated hyaline fringe (Fig. 15A, B), posterior margin of fourth prosomal somite wrinkled (Fig. 15B). Caudal surface of antennal basis with three oblique rows of tiny spinules near inner margin and two basal (proximal) rows of long spinules near outer margin (Fig. 15C, D, F). Antenna with three-segmented endopod bearing 1, 9, and 7 setae, respectively. Frontal surface of antennal basis with three rows of spinules: one proximal oblique, one near lateral (outer) margin in middle of segment, and one next to exopod seta (Fig. 15E). Mandible with nine teeth on gnathobase (Fig. 15G). Maxillular palp with two armed and one naked setae apically; one armed seta plus two nude setae on lateral lobe, proximal seta with tiny spinules (Fig. 15H). Distal coxal endite of maxilla with two long setae: proximal seta with two long, basal setules and bifurcated




Figure 14. *Microcycllops finitimus* Dussart, 1984. Adult female (MNHN-Cp7294). **A** Fifth pediger and hyaline fringe, dorsal **B** P1, coxa, basis and sclerite, frontal **C** P1, Enp, frontal **D** P4, coxa, basis and intercoxal sclerite, frontal **E** Anal somite, caudal rami, and caudal setae, ventral. Scale bars: 50 µm.




Figure 15. *Microcycllops anceps anceps* (Richard, 1897). Adult female. **A** Second prosomal somite, dorsal (SMNK-2832) **B** Second and fourth prosomal somites, dorsal (Matillas) **C** Antennal basipodite, caudal (MNHN-Cp6876) **D** Antennal basipodite, caudal (MNHN-Cp7296) **E** Antenna, frontal (Matillas) **F** Antennal basipodite, caudal (Matillas) **G** Mandible (Matillas) **H** Maxillule (Matillas) **I** Maxilla (Matillas) **J** Maxilliped (Matillas). Scale bars: 50 µm.




Figure 16. *Microcycllops anceps anceps* (Richard, 1897). Adult female. **A** P1, frontal (Pajonal) **B** P2, frontal (Pajonal) **C** P3 intercoxal sclerite, caudal (Pajonal) **D** P4, caudal (Pajonal) **E** Fifth pediger, genital double-somite (Matillas). Scale bars: 50 μ m.




Figure 17. *Microcyclops anceps* (Richard, 1897). Adult female. **A** Fifth pediger, genital double-somite (SMNK-2833) **B** Anal somite, caudal rami and caudal setae (Matillas), ventral. *Microcyclops anceps* **C** Adult male. **C** P4 coxa, basis, and intercoxal sclerite, caudal (MNHN-Cp6877) **D** Fifth pediger and genital somite (MNHN-Cp6877) **E** Fifth pediger and genital somite (MNHN-Cp7295) **F** Anal somite, caudal rami and caudal setae (MNHN-Cp6876). Scale bars: 50 μ m.

apically; distal seta armed with a continuous row of tiny spinules along one (inner) margin (Fig. 15I). Basipodite with claw-like projection bearing two stout teeth followed by a row of tiny spinules, and one long smooth seta on its base; two-segmented Enp bearing 2 and 3 setae, respectively (Fig. 15I). Maxilliped with a row of spinules in syncoxa, Bsp, and Enp1, on frontal view (Fig. 15J).

Inner margin of *basipodite* with long and fine hairs in P1–P3 (Fig. 16A, B). One pore present on lateral margin of Enp2P1. Spine absent on inner margin of BspP1 (Fig. 16A). Inner margin of BspP4 with long. Inner margin of BspP4 with long spinules (Fig. 16D). Intercoxal sclerites naked in P1 and P2 (Fig. 16A, B). Usually one distal row or sometimes two rows of spinules present on intercoxal sclerite of P3 (Fig. 16C). P4 sclerite with two rows of spinules (Fig. 16D): spinules in distal row larger and stronger than those in proximal row (Fig. 16D). Medial apical spine of Enp2P4 1.3 ± 0.1 times as long as lateral apical spine, and 0.7 ± 0.03 times as long as segment; length to width ratio of segment 2.5 ± 0.1 .

Strong spinules present (Figs 16E, 17D) or absent (Fig. 17A) on *fifth pediger* near base of lateral seta of P5. This character shows both inter- and intrapopulation variation; in one population, the females do not have spinules next to lateral seta (MNHN-Cp7296, unfigured here), while the males of the same population do (Fig. 17E).

Free segment of *fifth leg* 2.5 ± 0.2 times longer than wide, with relatively large spinule in distal position (Figs 16E, 17A); free segment 0.4 ± 0.08 times as long as apical seta. Distal margin of anal somite with continuous row of strong spinules on ventral and dorsal surfaces (Figs 17B, F). Caudal ramus 3.7 ± 0.3 times longer than wide, inner margin naked. Spinules present at base of caudal seta III. Caudal seta II inserted at $71.1 \pm 1.15\%$ of caudal ramus length (Fig. 17B).

Seta VII and VI 0.5 ± 0.1 and 0.8 ± 0.1 times as long as caudal ramus, respectively. Relative lengths of terminal caudal seta from outermost to innermost, 1.0 : 4.9 : 7.1 : 1.3 (Fig. 17B). Caudal setae IV and V with homonomous setulation, with hair-like setules only (Fig. 17B).

Discussion

Microcyclops anceps anceps showed the least variation in the qualitative and morphometric characters even though specimens were examined from a wide latitudinal range (Venezuela, Mexico, Guyana, Brazil, Guatemala, and Uruguay).

Microcyclops echinatus (from southeastern Mexico) and *M. ceibaensis* (from Honduras and southeastern Mexico) appeared morphologically similar. Similarities between these species are in: the length and width ratio of Enp2P4; the P4 sclerite with two rows of spines; the length ratio of the terminal caudal setae III and IV; the presence of spines at the insertion of setae II, and III; and the heteronomous ornamentation of the spine on the inner margin of Bsp P1. But the features that separate the specimens of these species were the insertion of the caudal seta II (69.5% in *M. ceibaensis* vs. 73.2% in *M. echinatus*); the length and width ratio of the caudal ramus is 3.6 in *M. ceibaensis*

while 5.9 in *M. echinatus*; the presence of spines on the fifth pediger in *M. echinatus* vs. absence of these spines in *M. ceibaensis*; and the ornamentation of the inner basis of P4 with spine-like setae in *M. ceibaensis* vs. short setae plus hair-like setae in *M. echinatus*.

In 1935, Kiefer described two new species *Cyclops (Microcyclops) diversus* and *Cyclops (Microcyclops) alius* from Uruguay. The microscopic observations performed here, support the opinion of Reid (1986) on the synonymy of *M. ceibaensis* and *M. diversus*. The specimens labelled as *M. diversus* sp. n. share all the morphometric features of the type series of *M. ceibaensis*. Additionally, the structure of P1, P3, P4, and P5; the armament of the caudal surface of the antenna, the number of setae on each endopodal segment of the antenna, and the entire morphology of the urosome, and the caudal ramus in *M. diversus* are indistinguishable from the states found in *M. ceibaensis*.

The type specimens labelled as *M. dubitabilis* (from Trou Caiman, Haiti) and *Microcyclops alius* (from Barra Sta. Luzia, Uruguay) were morphologically similar to: 1) specimens identified as *M. rubellus* [including the specimen analysed by Reid (1992)], 2) some specimens from Southeastern Mexico, 3) *M. alius* from Brazil described by Rocha (1998), and 4) another specimen labelled as *M. dubitabilis* from Guadeloupe. The micro-structural analysis showed that the following features are common in all of the above mentioned specimens: the number of setae on the endopodal segments of the antenna and the number of rows of spines on the caudal surface of antennal basis; the ornamentation of the setae of the maxillular palp; the shape of the maxilla, and in particular, the structure of the distal coxal endite, the basipodite, and the seta on the claw-like projection; the shape and length of the spine on the inner margin of P1 basis; the shape and ornamentation of the intercoxal sclerites and the inner basis of P1, and P4; the meristic characters of P4, all traits of the caudal rami setae; the structure of P5, the anal somite, and the caudal rami. Therefore, *M. alius* is considered here as a junior synonym of *M. dubitabilis*, as it was suggested by Rocha (1998). Also, several specimens recorded under the name *M. rubellus* in the Americas likely refer to *M. dubitabilis*.

Microcyclops dubitabilis clearly differs from *M. varicans* s. str. at least in the next features: the spines at the insertion of caudal seta III are present in *M. dubitabilis*, but absent in *M. varicans*; caudal ramus is 3.5–4 times as long as wide in *M. varicans* and shorter in *M. dubitabilis*. Medial spine of Enp2P4 is around 0.8 times as long as the segment in *M. dubitabilis*, whereas that in *M. varicans* is shorter (around 0.5); and the seta inserted at base of claw-like projection in the maxilla is armed only with strong teeth at its base in *M. varicans*, but this armament is more complex in *M. dubitabilis*. Therefore *M. dubitabilis* is not a synonym of *M. varicans*.

Microcyclops rubellus and *M. varicans* have been recorded in several regions of the world and were thought to be likely cosmopolitan (Reid 1992) and highly variable in morphology. This is especially the case for *M. varicans* (Franke 1989, Alekseev 2002). The type material of *M. rubellus* and *M. varicans* is probably lost and both species were originally described from North Europe (Sars 1863, Lilljeborg 1901). Our review of the descriptions and drawings of *M. rubellus* from some European localities however revealed differences between the European and American specimens here examined.

in the medial surface ornamentation of the basis of the fourth swimming leg, in the length proportion of the medial apical spine and the Enp2P4, and the ornamentation of the distal margin of the anal somite (see Einsle 1993). Hence, *M. rubellus* s. str. probably is not distributed in America.

Remarks about the new species

The specimens from southeastern Mexico assigned to the new species *M. inarmatus* were morphologically similar to that from Laguna Rincon, Haiti identified as *M. dubitabilis* (SMNK-2391, 2392) and to the specimen examined by Reid (1992) and classified as *M. varicans*. The shared morphology of the antenna, maxilla, P1 to P5, and urosome is obvious in all of these specimens (figured and described in the descriptive section before).

Microcycllops inarmatus sp. n. can be distinguished from *M. varicans* by the following characters (see also Rylov 1948, Einsle 1993): *M. varicans* has a more elongated caudal rami (3.5–4 times as long as wide), the basipodite of P4 bears short spinules on inner margin, there are more setae on the second endopodite of A2, on the distal margin of anal somite the spinules are present ventral, lateral and dorsally; and the medial spine of the second endopod of fourth leg has around the half length of the segment. All these features clearly differ to *M. inarmatus* sp. n.

The analysis that included specimens from a wide latitudinal range showed a highly conserved morphology primarily in the inner region of each swimming leg and oral appendages. Thus, we may speculate that some reports of *M. varicans* and *M. rubellus* recorded in the Americas are in fact *M. inarmatus* sp. n. and *M. dubitabilis*, respectively.

Microcycllops inarmatus sp. n. has some similarities also to *M. dubitabilis*, but the following features differentiate these two species: setal formula of the antennal endopod (1, 9, 7 in *M. dubitabilis* vs. 1, 6, 7 in *inarmatus*); the ornamentation on the caudal surface of the antennal basis is less complex in *M. inarmatus* than in *M. dubitabilis*; setae on maxillular palp are more armed in *M. inarmatus* than in *M. dubitabilis*; and the basal seta inserted at base of claw-like projection in the maxilla is more simple in *M. inarmatus*, whereas *M. dubitabilis* has two opposite rows of different spines.

The inner margin of the basis of the first swimming leg has a long spine with heteronomous ornamentation in *M. inarmatus*, in comparison to the short, homonomously setulated spine on this site in *M. dubitabilis*. In addition, the inner margin of the basis of the fourth swimming leg bears long hair-like setae and the fourth sclerite is almost as long as wide in *M. inarmatus*, whereas in *M. dubitabilis*, this inner margin bears short setae and the fourth sclerite is wider than long. The free segment of the fifth leg has a tiny spine on the medial margin in *M. inarmatus* (not described in Reid (1992), but clearly observed in the slide USNM-251321), and this spine is absent in *M. dubitabilis*. Finally, the lateral caudal seta is located near the middle of the caudal ramus in *M. inarmatus*, whereas in *M. dubitabilis*, this seta is located in the distal third. In all of the material analysed, no spines were observed at insertion of outermost terminal

caudal seta in *M. inarmatus*, whereas in *M. dubitabilis*, these spines were observed in every specimen.

Other species of *Microcyclops* which has 12-segmented antenna, caudal rami with innermost terminal caudal setae longer than outermost terminal caudal setae, spines present only ventrally on the distal margin of the anal somite, caudal rami short (no more than 3 times as long as wide), one spine on inner basis of P1, and the intercoxal sclerite of P4 quadrangular and naked, are *M. davidi* (Chappuis, 1922) and *M. richardi* (Lindberg, 1942). *Microcyclops inarmatus* sp. n. differs from these species in the surface ornamentation of P4 basipodite: long hair-like setules vs. short spine-like setules in *M. davidi* (*sensu* Mirabdullayev et al. 2002) and *M. richardi* (see Lindberg 1942). The genital double-somite in *M. inarmatus* is short around 0.8 times as long as wide – similar in *M. davidi* –, but it is elongated in *M. richardi*, around 1.4 times longer than wide; and the second endopodite of A2 bears 9 setae in *M. davidi*, but only 6 setae in *M. inarmatus*.

Additionally, in *M. richardi* the free segment of P5 has no spine on medial margin, and the medial spine of Enp2P4 is short (0.6 times the length of the segment) in comparison with the new species. Finally, the seta on the base of the claw-like projection of the maxillar basipodite, is armed with tiny spinules in *M. inarmatus*, whereas in *M. davidi* it bears strong teeth on its base.

Remarks on *M. anceps pauxensis* Herbst, 1962 and *M. anceps* var. *minor* Dussart, 1984

Microcyclops anceps pauxensis and *M. anceps* var. *minor*, described from the Amazonian region and Venezuela respectively, are similar in the number of segments of A1 (12-segmented); the length ratio of Enp2P4 (2.35 vs. 2.46); the intercoxal sclerite of P4 with two rows of spines; the length ratio of the lateral and medial apical spines of Enp2P4 (0.64 vs. 0.51); the length ratio of the medial apical spine of En2P4 and the segment (0.7 in both species), the insertion of the caudal seta II (lateral) is at 68% of the caudal rami length in *M. a. pauxensis*, and 70% in *M. a. minor*, and the continuous row of spines along the ventral and dorsal margins of the anal somite.

However, according to Herbst (1962) and Dussart (1984), the inner basis of P1 is naked in *M. a. pauxensis*, and hairy in *M. a. minor*; the inner basis of P4 bears small setules in *M. a. pauxensis* but this is naked in *M. a. minor*. The ratio between the lengths of the caudal setae VI and III is lower in *M. a. pauxensis* than in *M. a. minor* (1.81 vs. 2.52); the ratio between the lengths of caudal seta VI and caudal rami is higher in *M. a. pauxensis* (1.44) than in *M. a. minor* (1.07); the length ratio between the free segment of P5 and distal seta of P5 is 0.18 in *M. a. pauxensis*, and 0.34 in *M. a. minor*. Other differences between both species were observed in the length ratio between the dorsal caudal seta and caudal ramus (1.5 in *M. a. pauxensis* vs. 0.57 in *M. a. minor*), and the spines on the base of the outer caudal seta (spinules present in

M. a. pauxensis vs. absent in *M. a. minor*). All of these differences suggest that *M. a. minor* is distinct from *M. a. pauxensis*.

Therefore, these taxa may constitute different species. The evaluation of mouth-parts and the ornamentation of the inner region of each swimming leg in the type material may facilitate species delimitation. Unfortunately, the type material of both “subspecies” was not available to us.

Conservative characters among species

Based upon morphological and morphometric features, eleven species and two subspecies of *Microcyclops* recorded in America were recognized. The following set of characters distinguishes between species: the ornamentation of the caudal surface of the antennal basis; the ornamentation of the setae of the maxillular palp; the shape and armature of the distal coxal endite of maxilla; and the basal seta in front of the claw-like projection of the maxillar basis. Previously, similar structures have been useful for differentiating other Cyclopinae species, such as *Mesocyclops* (Van de Velde 1984a, 1984b, Hołyńska 2000).

Among the specimens examined, the organization of the spine pattern on the antennal basipodite is similar to that proposed by Van de Velde (1984b) for *Mesocyclops* which is more complex on the caudal side than on the frontal side. Additionally, the caudal surface ornamentation of the antennal basis in *Microcyclops* here examined is similar to that in most New World *Mesocyclops*: the simple ornamentation pattern found in Neotropical *Mesocyclops* was considered by Hołyńska (2000) and Wyngaard et al. (2010) as an ancestral state. The pattern observed in *Microcyclops* is much less complex in comparison to those reported for some eucyclopinae species from the genus *Macrocylops* (Karanovic and Krajicek 2012), *Paracyclops* (Karaytug 1999) and *Eucyclops* (Alekseev et al. 2006, Mercado-Salas et al. 2015).

The micro-structures of the swimming legs as diagnostic characters have been explored in *Mesocyclops*. In *Mesocyclops*, the coxal and basis armament of the first and fourth trunk limbs are important (Van de Velde 1984a, 1984b). In *Eucyclops*, the coxal seta of P4 or the intercoxal sclerites of all trunk limbs are informative (Alekseev et al. 2006). Our results show that features such as the medial surface ornamentation of basis of all four legs, the shape and ornamentation of the sclerites of P1 to P4, the presence/absence or length and armature of the spine on the inner basis of P1, and the shape or armature of the free segment of P5 were useful for differentiating between species.

Important diagnostic morphometric features for *Microcyclops* were the relative position of the lateral seta on the caudal ramus; the relative length of the outermost terminal caudal seta (III) and the outer median terminal caudal seta (IV); the relative length of caudal seta III and the inner median terminal caudal seta (V); and the length: width ratio of caudal ramus. Traditionally, the length ratio of the second endopod and its apical spines of the fourth trunk limb have been used as features to separate species of

Microcyclops; however, as in another genus such as *Eucyclops* or *Mesocyclops*, the surface micro-structures together with the integumental armature and the meristic characters of the caudal rami are more informative.

This study is the first attempt to clarify the taxonomy of the species of *Microcyclops* recorded in America using detailed morphological analysis.

Conclusion

The microscopic analysis of oral and thoracic appendages facilitated better delineation of *Microcyclops* species recorded in America. The characters that better distinguish between species are the ornamentation of antennal basipodite, the armature of the coxal endite and basipodite of the maxilla, the surface ornamentation of the inner basis of P1, the structure of intercoxal sclerites of the trunk limbs, the length: width ratio of caudal ramus, the length proportion of the caudal setae, and the relative position of the lateral seta on the caudal ramus.

The analysis performed here show that *M. alius* is a junior synonym of *M. dubitabilis*, and support the opinion about the synonymy of *M. ceibaensis* and *M. diversus*.

Microcyclops inarmatus sp. n. can be distinguished from other known species of the genus by the unique combination of several characters such as: morphometric characters of the second endopodite of fourth trunk limb and caudal ramus, presence of 6 setae on the second endopodal segment of antenna, antennal basipodite with just one group of spinules on caudal surface, lack of ornaments on the intercoxal sclerites of all swimming appendages, absence of spinules at base of lateral caudal and outermost terminal caudal setae, and basipodites of first to fourth swimming legs with long hair-like setules on inner margin.

Key to the American species of *Microcyclops* (females)

The key is mainly based on the analysis performed in the descriptive section of this manuscript. Original descriptions were consulted in those species in which no microscopic observations could be made [*Microcyclops anceps pauxensis* (Herbst 1962); *M. anceps* var. *minor* (Dussart 1984); *M. mediasetosus* (Dussart and Frutos 1985); *M. pumilis* (Pennak and Ward 1985); and *M. medioides* (Dussart and Frutos 1986)].

- | | | |
|---|---|--------------------------------------|
| 1 | Cylindrical free segment of P5 smooth, without inner spine (Fig. 7I–K) | 2 |
| — | Cylindrical free segment of P5 with inner spine (Figs 4G, H; 16E)..... | 5 |
| 2 | Base of the outermost caudal seta (III) with a row of spines (Fig. 8A–C) | 3 |
| — | Base of the outermost caudal seta (III) without a row of spines..... | 4 |
| 3 | Length (L): width (W) ratio of caudal ramus is 4.35; lateral caudal seta inserted at 69% of the total caudal ramus length; inner basis of P4 naked..... | |
| | | <i>Microcyclops medioides</i> |

- L: W ratio of caudal ramus is 2.48 ± 0.2 ; lateral caudal seta inserted at $71 \pm 5.7\%$ of the total caudal ramus length (Fig. 8A–C); inner basis of P4 with short hair-like setae (Fig. 7E–H) *M. dubitabilis*
- 4 L: W ratio of caudal ramus is 5 ± 1 ; lateral caudal seta inserted at 80 % of the total caudal ramus length *M. furcatus*
- L: W ratio of caudal ramus is 2.3 ± 0.6 ; lateral caudal seta inserted at 55 % of the total caudal ramus length *M. pumilis*
- 5 Inner spine of the cylindrical free segment of P5 tiny, articulated, inserted medially, and does not reach the distal margin of the segment (Figs 4H, 11C, 13D) 6
- Inner spine of the cylindrical free segment of P5 strong, unarticulated; inserted terminally, projected beyond the distal margin of the segment (Figs 16E, 17A) 11
- 6 Length ratio of the innermost (VI): outermost (III) caudal setae is 3.0; L: W ratio of caudal ramus is 2.3; lateral caudal seta inserted at 57 % of the total caudal ramus length *M. mediasetosus*
- Length ratio of the innermost (VI): outermost (III) caudal setae is 1.6 to 2.0; L: W ratio of caudal ramus is 2.7 to 6.0; lateral caudal seta inserted at 60 to 75 % of the total caudal ramus length 7
- 7 Inner basis of P1 with hair-like setae, medial spine absent (Fig. 14B); inner basis of P4 hairy (Fig. 14D); intercoxal sclerite of P1 nude; intercoxal sclerite of P4 armed *M. finitimus*
- P1 basis with medial spine (Figs 4A, B; 10B, C; 13A); inner basis of P4 hairy (Fig. 4F), or with strong spine-like setae (Figs 10I, K), or with a combination of both (Fig. 13C); intercoxal sclerite of P1 nude (Fig. 4A, B) or armed (Fig. 13A); intercoxal sclerite of P4 nude (Fig. 2D) or armed (Fig. 10K) 8
- 8 Inner basis of P1 naked, medial spine reaching the proximal half of Enp2P1 and with homonomous ornamentation; L: W ratio of caudal ramus is 5 to 6, with a row of spines at the base of the lateral caudal seta (II) that extends dorsally; and no spines at the base of the outermost caudal seta (III) *M. elongatus*
- Inner basis of P1 hairy, medial spine reaching the distal half of Enp2P1 and with heteronomous ornamentation (Figs 2C; 4A, B; 13A); L: W ratio of caudal ramus is 2.5 to 6, with or without spines at the base of both the lateral (II) and the outermost caudal seta (III) 9
- 9 Anal somite with a row of spines on ventral margin; no spines at the bases of the caudal setae II and III (Figs 5A, C–E); intercoxal sclerites of P1 to P4 unarmed (Fig. 4A–D); basipodite of P4 with long hair-like setae on inner margin (Fig. 2D); one group of spines on caudal view of antennal basis (Fig. 3B); L: W ratio of caudal ramus 2.5 ± 0.4 *M. inarmatus* sp. n.
- Anal somite with a row of spines along both ventral and dorsal margins; with spines at the bases of the caudal setae II and III (Fig. 11C); intercoxal sclerites of P1 to P4 armed (Fig. 10C–H, K); basipodite of P4 with strong spine-like

- setae (Figs 10I, K), or with a combination of hair-like setae and spinules (Fig. 13C); more than one group of spines on caudal view of antennal basis (Figs 9F; 12B); L: W ratio of caudal ramus between 3.2 to 6.3.....10
- 10 Fifth pediger with spines on ventral and lateral surfaces (Fig. 13D); caudal ramus is 5.9 ± 0.4 times longer than wide (Fig. 13E); inner basis of P4 with heteronomous ornamentation: short spine-like plus long hair-like setae (Fig. 13C); caudal surface of antennal basis with two rows of long spines next to exopodal seta (Fig. 12B) *M. echinatus*
- Fifth pediger nude ventrally and laterally; caudal ramus is 3.6 ± 0.4 times longer than wide (Fig. 11C); inner basis of P4 with homonomous ornamentation: strong spine-like setae (Figs 10I, K); caudal surface of antennal basis without spines next to exopodal seta (Fig. 9F) *M. ceibaensis*
- 11 No spines on the base of the caudal setae II and III; inner basis of P4 naked, unarmed *M. anceps* var. *minor*
- Spines on the base of caudal seta III, no spines on the base of the caudal seta II (Fig. 17B); inner basis of P4 ornamented (Fig. 16D) 12
- 12 W ratio of caudal ramus is 3.7 ± 0.3 (Fig. 17B); inner basis of P1 hairy (Fig. 16A); inner basis of P4 with long spine-like setae (Fig. 16D) *M. anceps anceps*
- L: W ratio of caudal ramus is 2.4; inner basis of P1 naked; inner basis of P4 with short hair-like setae *M. anceps pauxensis*

Acknowledgements

Danielle Defaye and Paula Rodríguez Moreno-Martin Lefèvre, Muséum National d'Histoire Naturelle, Paris; Hubert Höfer and Hans Walter Mittmann, Staatliches Museum für Naturkunde, Karlsruhe; Chad Walter, Smithsonian Institution National Museum of Natural History, kindly allowed us to review MNHN, SMNK, and NMNH specimens, respectively. The authors received grants from the National Council of Science and Technology (CONACYT) through the Mexican Barcode of Life (MEXBOL). We are grateful to Danielle Defaye, Maria Hołyńska, and Iskandar M. Mirabdullayev who made valuable comments and suggestions to improve the manuscript. We are grateful with the academic edition of two native English-spoken reviewers.

References

- Alekseev VR (2002) Copepoda. In: Fernando CH (Ed.) A Guide to Tropical Freshwater Zoo-plankton. Backhuys Publishers, Leiden, 123–187.
- Alekseev V, Dumont HJ, Pensaert J, Baribwegure D, Vanfleteren JR (2006) A redescription of *Eucyclops serrulatus* (Fischer, 1851) (Crustacea: Copepoda: Cyclopoida) and some related taxa, with a phylogeny of the *E. serrulatus*-group. Zoologica Scripta 35(2): 123–147. doi: 10.1111/j.1463-6409.2006.00223.x

- Dussart BH (1984) Some Crustacea Copepoda from Venezuela. *Hydrobiologia* 113(1): 25–67. doi: 10.1007/BF00026592
- Dussart BH, Frutos S-M (1985) Sur quelques Copépodes d'Argentine. *Revue d'hydrobiologie Tropicale* 18(4): 305–314.
- Dussart BH, Frutos S-M (1986) Sur quelques Copépodes d'Argentine. 2. Copépodes du Pará Medio. *Revue d'hydrobiologie Tropicale* 19(3-4): 241–262.
- Einsle U (1993) Crustacea: Copepoda: Calanoida und Cyclopoida. *Süßwasserfauna von Mitteleuropa*: 8/4-1. Gustav Fischer Verlag, Stuttgart, 209 pp.
- Elías-Gutiérrez M, Suárez-Morales E, Gutiérrez-Aguirre MA, Silva-Briano M, Granados-Ramírez JG, Garfias-Espejo T (2008) Cladocera y Copepoda de las aguas continentales de México, guía ilustrada. Universidad Nacional Autónoma de México FES Iztacala, Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, El Colegio de la Frontera Sur, Consejo Nacional de Ciencia y Tecnología, Secretaría de Medio Ambiente y Recursos Naturales, México, 322 pp.
- Fiers F, Ghenne V, Suárez-Morales E (2000) New species of Continental cyclopoid copepods (Crustacea, Cyclopoida) from the Yucatán Peninsula, Mexico. *Studies on Neotropical Fauna & Environment* 35(3): 209–251. doi: 10.1076/snfe.35.3.209.8862
- Franke U (1989) Katalog zur Sammlung limnischer Copepoden von Prof. Dr. Friedrich Kiefer. Carolinea. Staatliches Museum für Naturkunde, Karlsruhe, 433 pp.
- Herbst HV (1962) Crustacea aus dem Amazonasgebiet, gesammelt von professor Dr. H Sioli und Dr. R Braun. 1. Litorale und substratgebundene Cyclopoida Gnathostoma (Copepoda). *Crustaceana* 3(4): 259–278. doi: 10.1163/156854062X00508
- Holyńska M (2000) Revision of the Australasian species of the genus *Mesocyclops* Sars, 1914 (Copepoda: Cyclopidae). *Annales Zoologici* 50(3): 363–447.
- Huys R, Boxshall GA (1991) Copepod Evolution. The Ray Society, London, 468 pp.
- Karanovic T, Krajicek M (2012) When anthropogenic translocation meets cryptic speciation globalized bouillon originates; molecular variability of the cosmopolitan freshwater cyclopoid *Macrocyclops albidus* (Crustacea: Copepoda). *Annales de Limnologie-International Journal of Limnology* 48(1): 63–80. doi: 10.1051/limn/2011061
- Karaytug S (1999) Genera *Paracyclops*, *Ochridacyclops* and Key to the Eucyclopinae. Guides to the Identification of the Macroinvertebrates of the Continental Waters of the World, Vol. 14. Backhuys Publishers, Leiden, 217 pp.
- Kiefer F (1935) Neue Süßwassercyclopiden (Crustacea Copepoda) aus Uruguay. *Zoologischer Anzeiger* 109(7/8): 181–188.
- Lilljeborg W (1901) Synopsis specierum huc usque in Suecia observatarum generis *Cyclops*, sive Bidrag till en Översigt af de inom Sverige iakttagna arterna af släktet *Cyclops*. K. Svenska Vetensakad. Forhhandl., Stockholm 35(4): 1–118.
- Lindberg K (1942) Cyclopoïdes nouveaux du continent Indo-Iranien. III-IV. Records of the Indian Museum Calcutta 44(1): 15–27.
- Mercado-Salas NF, Suárez-Morales E, Silva-Briano M (2015) Taxonomic revision of the Mexican *Eucyclops* (Copepoda: Cyclopoida) with comments on the biogeography of the genus. *Journal of Natural History* 2015: 1–123. doi: 10.1080/00222933.2015.1061715

- Mirabdullayev IM (1998) Redescription of *Microcyclops rechtyae* Lindberg, 1960 (Crustacea, Copepoda). *Hydrobiologia* 362(1–3): 219–223. doi: 10.1023/A:1003178400061
- Mirabdullayev IM (2007) Redescription of *Microcyclops cunningtoni* (GO Sars, 1909) (Copepoda, Cyclopoida). *Munis Entomology and Zoology* 2(1): 79–85.
- Mirabdullayev IM, Urazova RS (2006) Redescription of the female and first description of the male of *Microcyclops karvei* (Kiefer & Moorthy, 1935) (Copepoda, Cyclopoida). *Zoology in the Middle East* 38(1): 57–62. doi: 10.1080/09397140.2006.10638165
- Mirabdullayev IM, Van Damme K, Dumont H (2002) Freshwater cyclopoids (Crustacea: Copepoda) from the Socotra Archipelago, Yemen, with description of a new species of *Bryocyclops*. *Fauna of Arabia* 19: 261–271.
- Pennak RW, Ward JV (1985) New cyclopoid copepods from interstitial habitats of a Colorado Mountain stream. *Transactions of the American Microscopical Society* 104(3): 216–222. doi: 10.2307/3226433
- Reid JW (1985) Chave de Identificação e lista de referências bibliográficas para as espécies continentais Sulamericanas de vida livre da Orden Cyclopoida (Crustacea, Copepoda). Bolm. Instituto de Biociências, Universidade de São Paulo 9: 17–143.
- Reid JW (1986) A redescription of *Microcyclops ceibaensis* (Marsh, 1919) (Copepoda: Cyclopoida) from Marsh's specimens in the National Muesum of Natural History. *Proceedings of the Biological Society of Washington* 99(1): 71–78.
- Reid JW (1990) Continental and coastal free-living Copepoda (Crustacea) of Mexico, Central America and the Caribbean region. In: Navarro D, Robinson JG (Eds) *Centro de Investigaciones de Quintana Roo, Program of Studies in Tropical Conservation*. University of Florida, Chetumal, Quintana Roo, México, 175–213.
- Reid JW (1992) Copepoda (Crustacea) from fresh waters of the Florida Everglades, U.S.A. with a description of *Eucyclops conrowae* n. sp. *Transactions of the American Microscopial Society* 111(3): 229–254. doi: 10.2307/3226612
- Rocha CEF, Da (1998) New morphological characters useful for the taxonomy of the genus *Microcyclops* (Copepoda, Cyclopoida). *Journal of Marine Systems* 15(1–4): 425–431. doi: 10.1016/S0924-7963(97)00091-2
- Rylov VM (1948) Fauna of U.S.S.R. Crustacea. Freshwater Cyclopoida. Translated from Russian, 3(3). Zoological Institute of the Academy of Sciences of the U.S.S.R., Leningrad, 314 pp.
- Sars GO (1863) *Oversigt af de indenlandske Ferskvands copepoder*. Forh Vidensk Selsk Christiana 29: 212–262.
- Van de Velde I (1984a) Introduction of new characters in *Mesocyclops* with African species as example. *Crustaceana Supplement* 2: 404–419.
- Van de Velde I (1984b) Revision of the African species of the genus *Mesocyclops* Sars, 1914 (Copepoda: Cyclopidae). *Hydrobiologia* 109(1): 3–66. doi: 10.1007/BF00006297
- Williamson CE, Reid JW (2001) Copepod. In: Thorp JH, Covich AP (Eds) *Ecology and Classification of North American Freshwater Invertebrates*. Academic Press, San Diego California, 915–954. doi: 10.1016/B978-012690647-9/50023-5
- Wyngaard GA, Hołyńska M, Schulte JA II (2010) Phylogeny of the freshwater copepod *Mesocyclops* (Crustacea: Cyclopidae) based on combined molecular and morphological data, with notes on biogeography. *Molecular Phylogenetics and Evolution* 55(3): 753–764. doi: 10.1016/j.ympev.2010.02.029

Supplementary material I

Table 2

Authors: Martha Angélica Gutiérrez-Aguirre, Adrián Cervantes-Martínez

Data type: RTF file

Explanation note: Biological material examined.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.