



Published in final edited form as:

Pancreat Disord Ther. 2016 ; 6(1): 1–2. doi:10.4172/2165-7092.1000167.

Developing Effective Salmonella-based Approaches to Treat Pancreatic Cancer

Jeremy Chen, Don J. Diamond, and Edwin R. Manuel*

Department of Experimental Therapeutics, City of Hope, Duarte CA, USA

Keywords

Salmonella; Pancreatic cancer; Treatment; Pancreatic ductal adenocarcinoma; Tumor malignancies

Commentary

Pancreatic ductal adenocarcinoma (PDAC) is predicted to have the second highest incidence of fatalities among all solid tumor malignancies worldwide. Despite three decades of research focused on treating PDAC, the five-year survival rate is less than 6%, thus leaving much room for improvement. Historically, PDAC has been difficult to treat due to the absence of early detection methods resulting in clinical disease that is significantly advanced and characterized by immunosuppression, an extreme form of fibrosis known as desmoplasia, and metastasis to vital organs [1]. Current chemotherapeutic combinations, such as gemcitabine with Abraxane, have been shown to extend patient survival; however, they only do so by two to three months [2]. The newly released irinotecan liposome injection tested in pancreatic cancer patients is, at best, equal in efficacy to gemcitabine with Abraxane [3-5]. The low efficacy and high toxicity of chemotherapy has led to innovative new strategies using immunotherapeutic approaches for treatment of PDAC. For example, the whole pancreatic cancer cell vaccine expressing human macrophage-colony stimulating factor (GM-CSF), known as GVAX, combined with a Listeriased vaccine expressing the PDAC antigen mesothelium (CRS-207) shows promise in extending survival evidenced by both pre-clinical and preliminary clinical data [6]. Furthermore, antibody therapies targeting CTLA-4 and PD-1 have shown great benefit toward enhancing antitumor immunity resulting in tumor regression and extension of survival in other solid tumor models [7]. Despite these successes, there is a growing consensus that a “multi-pronged” approach to induce anti-tumor immunity and simultaneously targeting immune suppression and desmoplasia will have the greatest effect in eliminating PDAC. However, balancing such aggressive approaches with minimal toxicity to the patient will prove to be an incredibly daunting task.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Corresponding author: Assistant Professor, City of Hope, 1500 East Duarte Road Fox South 2000A, Duarte, CA 91010, USA, Tel: +6262564673,60452; emanuel@coh.org.

Conflicts of Interest

ERM and DJD are consultants for, and have received royalty and SRA payments from, Tara Immuno-Oncology LLC who have licensed SHIDO-ST for clinical development.

We first tested the hypothesis of targeting immune suppression to enhance tumor-specific responses in an aggressive, highly immunogenic murine model of melanoma [8]. We found that therapeutic vaccination alone using the tumor antigen survivin, which is over-expressed and enhances survival in melanoma cells, was unable to generate tumor-specific responses to control melanoma growth. However, when we administered, prior to vaccination, an attenuated *Salmonella typhimurium* (ST) targeting Signal Transducer and Activator of Transcription 3 (STAT3), which is overexpressed in many cancers and induces which contributes to tumor-derived immune suppression, prior to vaccination, we observed enhancement of the survivin-specific response resulting in significant control of the primary tumor, reduction in lung metastases, and extension in survival. Thus, modulating tumor-derived immune suppression prior to therapeutic vaccination was sufficient to rescue the immune response and cause tumor regression. We then sought to apply the same approach to PDAC. We postulated, however, that although while PDAC is a prime candidate for *Salmonella*-based treatment due to its hypoxic nature [9,10], it is poorly immunogenic and many of the physical barriers, such as interstitial pressure generated by surrounding fibrosis, would prevent ST from penetrating into the hypoxic regions, thus requiring alternative intervention methods. In the PDAC stromal compartment, hyaluronic acid (HA) can be found at extremely high levels in the extracellular matrix (ECM) [11]. This results in a biophysical barrier that significantly reduces delivery of therapeutics to tumor cells. Our recent studies have shown that using PEGPH20, a PEGylated human recombinant PH20 hyaluronidase that depletes the HA found in the ECM of PDAC, helps to decrease interstitial and increase the permeability of the tumor to biological vectors such as attenuated ST [12]. Following PEGPH20 treatment in aggressive murine models of PDAC, we have found that administration of ST transformed with an shRNA plasmid specific to the immunosuppressive protein indoleamine 2,3-dioxygenase [13,14], results in enhanced ST colonization and intratumoral recruitment of PMN which are involved in the direct killing of surrounding tumor cells. The advantage of this therapeutic approach is that it induces innate immunity specifically within tumor tissue to suppress growth. This is beneficial in patients who are immune compromised; especially those who have been the recipients of chemotherapy treatments, as PMN have extremely rapid turnover rates. There is also evidence that the combination of PEGPH20/shIDO-ST treatment induces late anti-tumor CD8+ T cell responses critical for long-term tumor control. We have initiated studies to further characterize this response in hopes of determining its specific antigenic target and the role of PMN in its generation. While we have observed induction of potent innate and adaptive anti-tumor responses, we have not observed significant toxicity in pre-clinical cancer models, thus providing some evidence of safety. Further studies of efficacy and toxicity in patient-derived xenograft and humanized-IDO transgenic models are planned. In conclusion, we believe that PEGPH20/shIDO-ST represents a multi-pronged strategy that will potentially have greater efficacy than present chemotherapeutic options for PDAC and other solid tumors and is currently being developed for clinical application.

Acknowledgements

We would like to acknowledge that work relating to pancreatic cancer discussed in this commentary was supported by NIH grants R21-CA174306 and done in collaboration with Halozyme Therapeutics. Research reported in this publication was supported by the National Cancer Institute of the National Institutes of Health under award number

P30CA33572. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Erkan MS, Hausmann CW, Michalski AA, Fingerle M, Dobritz J, et al. The role of stroma in pancreatic cancer: diagnostic and therapeutic implications. *Nature reviews. gastroenterology & hepatology*. 2012; 9:454–467.
2. Han H, Von Hoff DD. SnapShot: pancreatic cancer. *Cancer Cell*. 2013; 23:424–424. [PubMed: 23518352]
3. Schober M, Javed MA, Beyer G, Le N, Vinci A, et al. New Advances in the Treatment of Metastatic Pancreatic Cancer. *Digestion*. 2015; 92:175–184. [PubMed: 26372949]
4. Wang-Gillam A, Li CP, Bodoky G, Dean A, Shan YX, et al. Nanoliposomal irinotecan with fluorouracil and folinic acid in metastatic pancreatic cancer after previous gemcitabine-based therapy (NAPOLI-1): a global, randomised, open-label, phase 3 trial. *Lancet*. 2015
5. Yi SY, Park YS, Kim HS, Jun HJ, Kim KH, et al. Irinotecan monotherapy as second-line treatment in advanced pancreatic cancer. *Cancer Chemother Pharmacol*. 2009; 63:1141–1145. [PubMed: 18839175]
6. Lutz ER, Wu AA, Bigelow E, Sharma R, Mo G, et al. Immunotherapy converts nonimmunogenic pancreatic tumors into immunogenic foci of immune regulation. *Cancer immunology research*. 2014; 2:16–631.
7. Arslan C, Yalcin S. Current and future systemic treatment options in metastatic pancreatic cancer. See comment in PubMed Commons below *J Gastrointest Oncol*. 2014; 5:280–295.
8. Manuel ER, Blache CA, Paquette R, Kaltcheva TI, Ishizaki I, et al. Enhancement of cancer vaccine therapy by systemic delivery of a tumor-targeting Salmonella-based STAT3 shRNA suppresses the growth of established melanoma tumors. *Cancer research*. 2011; 71:4183–4191. [PubMed: 21527558]
9. Cheng X, Zhang X, Cheng W, Chen J, Ma C, et al. Tumorspecific delivery of histidine-rich glycoprotein suppresses tumor growth and metastasis by antiangiogenesis and vessel normalization. *Current gene therapy*. 2014; 14:75–85. [PubMed: 24606115]
10. Yu B, Yang M, Shi L, Yao Y, Jiang Q, et al. Explicit hypoxia targeting with tumor suppression by creating an “obligate” anaerobic Salmonella Typhimurium strain. *Scientific reports*. 2012; 2:436. [PubMed: 22666539]
11. Jacobetz MA, Chan DS, Nesses A, Bapiro TE, Cook N, et al. Hyaluronan impairs vascular function and drug delivery in a mouse model of pancreatic cancer. *Gut*. 2013; 62:112–120. [PubMed: 22466618]
12. Manuel ER, Chen J, D’Apuzzo M, Lampa MG, Kaltcheva TI, et al. Salmonella-Based Therapy Targeting Indoleamine 2,3-Dioxygenase Coupled with Enzymatic Depletion of Tumor Hyaluronan Induces Complete Regression of Aggressive Pancreatic Tumors. *Cancer immunology research*. 2015; 3:1096–1107. [PubMed: 26134178]
13. Blache CA, Manuel ER, Kaltcheva TI, Wong AN, Ellenhorn JD, et al. Systemic delivery of Salmonella typhimurium transformed with IDO shRNA enhances intratumoral vector colonization and suppresses tumor growth. *Cancer research*. 2012; 72:6447–6456. [PubMed: 23090116]
14. Uyttenhove C, Pilotte L, Theate I, Stroobant V, Colau D, et al. Evidence for a tumoral immune resistance mechanism based on tryptophan degradation by indoleamine 2,3-dioxygenase. *Nat Med*. 2003; 9:1269–1274. [PubMed: 14502282]