

Improvements to the FIGO staging for ovarian cancer: reconsideration of lymphatic spread and intraoperative tumor rupture

Dong Hoon Suh¹, Tae Hun Kim², Jae-Weon Kim², Sun Young Kim³, Hee Seung Kim², Taek Sang Lee⁴, Hyun Hoon Chung², Yong-Beom Kim¹, Noh Hyun Park², Yong Sang Song^{2,5,6}

¹Department of Obstetrics and Gynecology, Seoul National University Bundang Hospital, Seongnam; ²Department of Obstetrics and Gynecology, Seoul National University College of Medicine, Seoul; ³Department of Obstetrics and Gynecology, Hanvit Women's Hospital, Ansan; ⁴Department of Obstetrics and Gynecology, SMG-SNU Boramae Medical Center, Seoul; ⁵Cancer Research Institute, Seoul National University College of Medicine, Seoul; ⁶WCU Biomodulation Major, Department of Agricultural Biotechnology, Seoul National University, Seoul, Korea

See accompanying editorial by Denny on page 293.

Objective: To evaluate the improvement in prognosis prediction with reassignment of International Federation of Gynecology and Obstetrics (FIGO) stages for ovarian carcinoma.

Methods: This was a retrospective study of patients with epithelial ovarian, fallopian tube, and primary peritoneal cancers. Sub-staging criteria used in stage reassignment were defined as follows: surgical spillage (IC1), capsule rupture before surgery or tumor on the surface (IC2), and positive cytology results (IC3); microscopic (IIB1) and macroscopic (IIB2) pelvic spread; microscopic extrapelvic spread (IIIA1) and retroperitoneal lymph node (LN) metastasis without extrapelvic spread (IIIA2); and supraclavicular LN metastasis (IVA) and other distant metastasis (IVB). Survival outcomes associated with the current and reassigned stages were compared.

Results: Overall, 870 patients were eligible for analysis. The median follow-up period was 45 months (range, 0 to 263 months). The 5-year overall survival rates (5YSRs) according to the current staging were 93.5% (IA), 82.5% (IC), 75.0% (IIB), 74.5% (IIC), 57.5% (IIIA), 54.0% (IIBB), 38.5% (IIIC), and 33.0% (IV). The 5YSRs of patients with IC1, IC2, and IC3 after sub-staging were 92.0%, 85.0%, and 71.0%, respectively ($p=0.004$). Patients who were reassigned to stage IIIA2 had a better 5YSR than those with extrapelvic tumors >2 cm (66.3% vs. 35.8%; $p=0.005$). Additionally, patients with newly assigned stage IVA disease had a significantly better 5YSR than those with stage IVB disease (52.0% vs. 28.0%; $p=0.015$).

Conclusion: The modified FIGO staging for ovarian carcinoma appears superior to the current staging for discriminating survival outcomes of patients with surgical spillage, retroperitoneal LN metastasis without extrapelvic peritoneal involvement, or distant metastasis to supraclavicular LNs.

Keywords: Lymph node metastasis, Ovarian cancer, Stage reassignment, Supraclavicular lymph node metastasis, Surgical spillage

Received Jul 27, 2013, Revised Aug 10, 2013, Accepted Aug 12, 2013

Correspondence to Jae-Weon Kim
Department of Obstetrics and Gynecology, Seoul National University College of Medicine, 101 Daehak-ro, Jongno-gu, Seoul 110-744, Korea.
E-mail: kjwksh@snu.ac.kr

INTRODUCTION

Among gynecologic malignancies, ovarian cancer is the leading cause of cancer-related deaths [1]. The International Federation of Gynecology and Obstetrics (FIGO) staging for ovarian cancer reflects the prognosis of affected patients. The

Copyright © 2013. Asian Society of Gynecologic Oncology, Korean Society of Gynecologic Oncology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

current staging was accepted by FIGO in 1988. However, many studies have indicated the necessity of a revision of the current FIGO staging guidelines for better prognostic discrimination. For stage IC disease, a growing body of evidence suggests that intraoperative rupture might not increase the risk of tumor recurrence [2,3]. For stage IIB disease, macroscopic pelvic peritoneal tumor masses might be associated with poor survival outcomes compared with microscopic tumor infiltration or adherence [4]. Many studies have reported better prognosis for stage IIIC ovarian cancer with lymph node (LN) involvement alone (without peritoneal carcinomatosis) than with LN involvement and concomitant peritoneal carcinomatosis [5-7]. The revision process is currently underway by the Gynecology Oncology Committee of FIGO in collaboration with various international societies and agencies, and the revision would be addressed about the stage IIIC classification, at least [8].

Since the revision process should be finalized only after a consensus is reached by all relevant international organizations through extensive consultations, we performed this timely study to evaluate whether the revision of FIGO sub-staging for ovarian carcinoma could improve prognosis prediction. Furthermore, the favorable prognosis of FIGO stage IIIC ovarian cancer characterized by LN-positive disease only, prompted us to compare the survival outcomes of stage IV disease so-assigned based on supraclavicular LN metastasis with those of stage IV disease so-assigned based on the other metastatic sites.

MATERIALS AND METHODS

Data were obtained from the review of the medical charts of all patients with epithelial ovarian, fallopian tube, and primary

peritoneal cancers at three Seoul National University-affiliated hospitals between 1990 and 2011. Each of the Institutional Review Boards of the three hospitals approved this study. Data on patient demographics, primary cancer sites, stages, grades, histology, first courses of treatment, intraoperative findings regarding tumor spread, operation procedures performed, residual disease, adjuvant chemotherapy, recurrence, and survival were collected. Optimal cytoreduction was defined as a residual tumor size of <1 cm. A total of 870 patients were eligible for analysis, as shown in **Fig. 1**.

Stages were reassigned according to the following criteria when indicated: surgical spillage with intraoperative tumor rupture (IC1), capsule rupture before surgery or presence of a tumor on the surface (IC2), and presence of malignant cells on ascites or peritoneal washing cytology (IC3); microscopic (IIB1) and macroscopic (IIB2) pelvic spread; microscopic extrapelvic spread (IIIA1) and retroperitoneal LN metastasis without extrapelvic spread (IIIA2); and supraclavicular LN metastasis (IVA) and other distant metastasis (IVB).

Kaplan-Meier survival estimates were derived for overall survival (OS) both before and after stage reassignment. Univariate and multivariate analyses were performed using Cox proportional hazards models in order to evaluate the association between the site of distant metastasis and OS outcomes. Various prognostic factors for the sub-stages were also compared using the chi-square test, Student's t-test, and Kruskal-Wallis test, as appropriate, to assess the association of these factors with survival outcomes. The OS rates were defined as duration from the date of diagnosis to either the recorded date of death from ovarian cancer or the date of death provided by Statistics Korea. Statistical tests were 2-sided, with $p < 0.05$ indicating significance. Statistical analyses were performed using SPSS ver. 19.0 (IBM Co., Armonk, NY, USA).




Fig. 1. Patient enrollment. SNUH, Seoul National University Hospital; SNUBH, Seoul National University Bundang Hospital; SMG-SNU BMC, Seoul Metropolitan Government-Seoul National University Boramae Medical Center; MMMT, malignant mixed müllerian tumor.

sub-stage IC1 (n=39), IC2 (n=27), and IC3 (n=53) disease were 92.0%, 85.0%, and 71.0%, respectively (p=0.004) (**Fig. 2**). Age, initial serum CA-125 level, tumor grade, lymphadenectomy rate, use of taxane-platinum-based adjuvant chemotherapy, and completion of chemotherapy ≥6 cycles did not differ according to sub-stage. Although patients who were assigned an advanced sub-stage had a higher frequency of serous cancer than those with early sub-stage disease did (p=0.02), multivariate analysis using Cox proportional hazards models revealed that serous-type cancer was not an independent prognostic factor for OS (p=0.766).

We performed additional analysis in the patients (n=64) who underwent staging operation along with lymphadenectomy, considering the low rate of lymphadenectomy in the patients with early stage disease. There were 20 IC1, 15 IC2, and 29 IC3 patients after stage reassignment. We found a significant OS difference among the patients according to the three sub-stages in this subset (p=0.021).

2. Stage IIB

The 5YSR for patients with stage IIB disease (n=32) was 75.0%. After stage reassignment, the 5YSRs of patients with sub-stage IIB1 (n=11) and IIB2 (n=21) disease were 70.0% and 76.0%, respectively (p=0.461) (**Table 2**).

3. Stage III

The 5YSR for patients with stage IIIC disease (n=410) was 38.5%. However, after stage reassignment, the 5YSR of patients with sub-stage IIIA2 disease (n=33) was significantly higher than that of the remaining patients with stage IIIC disease (n=377) who had extrapelvic tumors >2 cm (66.3% vs. 35.8%; p=0.005) (**Fig. 3**), and even higher than that of

patients with sub-stage IIIA1 disease (57.5%) (**Table 2**). The rate of optimal cytoreduction was significantly higher in stage IIIA2 disease than in stage IIIC disease (p=0.006). Lymphadenectomy was performed more frequently in the patients with stage IIIA2 disease than in those with stage IIIC disease (p<0.001). However, patients with stage IIIA2 and IIIC disease did not show any difference in age, initial serum CA-125 level, serous histology, tumor grade, neoadjuvant chemotherapy rate, use of taxane-platinum based adjuvant chemotherapy, and receipt of ≥ 6 chemotherapy cycles. Multivariate analysis using Cox proportional hazards models revealed that optimal cytoreduction was an independent prognostic factor for OS in stage III disease (hazard ratio [HR], 2.1; 95% confidence interval [CI], 1.6 to 2.7; p<0.001).




Fig. 3. Overall survival of patients with stage III ovarian cancer according to reassigned stages. Retroperitoneal lymph node metastasis without extrapelvic involvement was downstaged from stage IIIC to IIIA2.




Fig. 2. Overall survival of patients with stage IC ovarian cancer according to reassigned stages. IC1, intraoperative tumor rupture; IC2, capsule ruptured before surgery or tumor on surface; IC3, malignant cells in the ascites or peritoneal washings.




Fig. 4. Overall survival of patients with stage IV ovarian cancer according to sub-staging. Supraclavicular lymph node metastasis (stage IVA) vs. other sites metastasis (stage IVB).

Table 3. Univariate and multivariate analyses of the site of distant metastasis as a function of overall survival in stage IV ovarian cancer (n=77)

Site of distant metastasis	No. (%)	Multiple sites ≥ 2, no. (%)	Overall survival (mo), median (range)	Univariate analysis, HR (95% CI)	Multivariate analysis	
					HR (95% CI)	p-value
Pleural effusion	36 (46.8)*	8 (22.2)	31 (1–198)	1.01 (0.58–1.77)	0.54 (0.26–1.10)	0.090
Liver parenchyma	24 (31.2)	13 (54.2)	38 (1–91)	1.19 (0.66–2.14)	0.64 (0.30–1.34)	0.237
Supraclavicular LN	17 (22.1)	7 (41.2)	46 (6–162)	0.34 (0.13–0.85)	0.22 (0.08–0.63)	0.005
Lung	14 (18.2)	13 (92.9)	22 (1–54)	1.65 (0.79–3.45)	1.44 (0.61–3.36)	0.404
Bone	13 (16.9)	12 (92.3)	20 (1–57)	1.93 (0.89–4.17)	3.49 (1.10–11.08)	0.034
Brain	11 (14.3)	10 (90.9)	57 (1–143)	1.02 (0.42–2.40)	0.33 (0.11–1.02)	0.053

HR, hazard ratio; CI, confidence interval; LN, lymph node.

*Three were not pathologically confirmed.

4. Stage IV

The 5YSR for the patients with stage IV disease (n=77) was 33.0%. After stage reassignment, the 5YSRs of the patients with sub-stage IVA (n=17) and IVB (n=60) disease were 52.0% and 28.0%, respectively (p=0.015) (**Fig. 4**). Age, initial serum CA-125 level, serous histology, tumor grade, neoadjuvant chemotherapy rate, rate of optimal cytoreduction, lymphadenectomy rate, use of taxane-platinum-based adjuvant chemotherapy, receipt of ≥6 chemotherapy cycles, and the rate of chemoresistance did not differ between patients with the 2 sub-stages. Nevertheless, chemoresistance (HR 7.4; 95% CI, 2.8 to 19.3; p<0.001), no use of taxane-platinum based adjuvant chemotherapy (HR, 35.4; 95% CI, 1.7 to 733.4; p=0.021), and receipt of <6 chemotherapy cycles (HR, 17.7; 95% CI, 4.3 to 72.3; p<0.001) were independent poor prognostic factors for OS in stage IV ovarian cancer (data not shown). **Table 3** shows the results of univariate and multivariate survival analyses for stage IV ovarian cancer, according to the site of distant metastasis. There was no case of spleen parenchymal metastasis in our study population. Although patients with brain metastasis had the longest median OS, multivariate analysis using Cox proportional hazards models revealed that supraclavicular LN metastasis was an independent favorable prognostic factor for OS with regard to the site of metastasis in ovarian cancer (HR, 0.22; 95% CI, 0.08 to 0.63; p=0.005). In contrast, bone metastasis was an independent prognostic factor for poor OS (HR, 3.49; 95% CI, 1.10 to 11.08; p=0.034). Compared with the lung, bone, and brain metastases, which frequently showed multiple metastatic lesions, supraclavicular LNs often showed single one.

DISCUSSION

Our study demonstrates the necessity for revision of the

current FIGO staging for ovarian cancer with regard to the 3 sub-staging issues for stages IC, III, and IV. The sub-staging process has 2 main implications. First, stage IC was sub-staged as stages IC1, IC2, and IC3, to account for the distinct prognoses associated with the etiology of tumor rupture. Second, considering the relatively favorable prognosis associated with lymphatic tumor spread compared with peritoneal tumor spread, stage IIIC, which was classified solely on the basis of LN metastasis, was down-staged to stage IIIA2. Similarly, stage IV was sub-staged as stage IVA, with distant metastasis to the supraclavicular LNs, and stage IVB, with other distant metastasis.

There has been much controversy regarding the cause of tumor rupture that upstages cancers to stage IC with regard to prognosis. Vergote et al. [9] demonstrated that tumor rupture during surgery (HR, 2.65; 95% CI, 1.53 to 4.56; p<0.001) and before surgery (HR, 1.64; 95% CI, 1.07 to 2.51; p=0.022) had an independent unfavorable impact on disease-free survival. However, other researchers failed to show that capsular rupture caused by the surgeon affected the prognosis of patients with early-stage ovarian cancer [3,10,11]. A recently published meta-analysis supported this finding and concluded that intraoperative rupture might not decrease progression-free survival compared to no rupture in early-stage ovarian cancer with complete surgical staging and adjuvant platinum-based chemotherapy [2]. This finding is consistent with our findings that the 5YSR of patients with sub-stage IC1 disease was similar to that of patients with stage IA disease and was clearly higher than that of patients with sub-stage IC3 disease.

Many studies have supported Berek's suggestion of a separate entity for LN-positive stage IIIC ovarian cancer [5-7]. Berek [5] argued that FIGO should consider modifying the ovarian cancer staging by further stratifying stage III disease on the basis of the better OS in patients with retroperitoneal LN metastasis without peritoneal carcinomatosis than in

patients with macroscopic peritoneal carcinomatosis. One plausible explanation for the favorable prognosis of those patients with sub-stage IIIA2 disease might be the higher optimal cytoreduction rate compared to the patients with stage IIIC disease showing intraperitoneal tumor implants >2 cm. Optimal cytoreduction was a well-known and important prognostic factor for advanced-stage ovarian cancer [12]. Our study also showed that optimal cytoreduction was an independent prognostic factor for OS in patients with stage III disease. Bachmann et al. [13] reported that the influence of LN metastasis on prognosis decreases with the increase in residual tumor volume. The authors also reported that the nodal status seemed to be the next most important prognostic factor for advanced-stage ovarian cancer. Additionally, successful retreatment of recurrence in patients upstaged to stage IIIC on the basis of LN metastasis alone with second surgery and chemotherapy could partly account for the favorable prognosis of these patients [6,14,15].

The primary routes of ovarian cancer metastasis include intra-peritoneal implantation of exfoliated cells at distant sites and spreading through retroperitoneal lymphatic channels [16]. Spreading through the lymphatic channels of the diaphragm and the retroperitoneal LNs can lead to dissemination above the diaphragm, especially to the supraclavicular LNs [5]. In accordance with the favorable prognosis of LN-positive-only stage IIIC disease, stage IV disease, assigned on the basis of supraclavicular LN metastasis, might be associated with better survival outcomes than stage IV disease. Nevertheless, no study has investigated the favorable prognosis of stage IV ovarian cancer patients with supraclavicular LN metastasis. To the best of our knowledge, this study is the first one to demonstrate a distinctly better OS for stage IV ovarian cancer patients with supraclavicular LN metastasis than for patients with other forms of distant metastasis. Since ovarian cancer is known to spread both intraperitoneally and retroperitoneally almost simultaneously [17], the presence of tumor spreading mainly through lymphatic channels without intra-peritoneal dissemination suggests that such tumors might be associated with a favorable biologic behavior. In support of this idea, our study shows that sub-stage IVA disease was associated with a relatively limited number of simultaneous metastatic sites compared with sub-stage IVB disease. However, in this study, we failed to show any significant positive or negative associations between the prognosis of patients with sub-stage IVA disease and the prognostic factors that were already proven important for advanced-stage ovarian cancer, such as optimal cytoreduction and chemoresistance.

This study has some limitations. First, we did not show a significant difference in OS between sub-stages IIB1 and IIB2.

This might be partly due to the small number of patients with stage II disease. FIGO stage II ovarian cancer accounts for 8% of all ovarian cancers [4] and 6.4% in our study. Further studies with larger population numbers are needed in order to achieve sufficient statistical power. Second, the follow-up period for the study population was relatively short. This might result in an overestimation of the favorable prognosis of patients with stage IVA disease (5YSR, 52.0%), which was even better than that of patients with stage IIIC disease. A longer follow-up period could ensure that the survival estimates are more accurate. Third, the low rate of lymphadenectomy in early stage disease implicated that true pathologic stage IIIC disease could be included in the analysis of stage IC disease. However, the same result of a significant OS difference in stage IC patients who underwent staging lymphadenectomy was observed. Finally, mortality was not disease-specific but overall, without considering the cause of death. This might cause an underestimation of prognosis in this study. Nevertheless, the potential impact of this underestimation on the study results could be minimal because the age difference between the compared sub-stages was not significant.

In conclusion, modification of the current FIGO staging for ovarian carcinoma appears to improve the discrimination of the survival outcomes of patients with surgical spillage, retroperitoneal LN metastasis without extrapelvic peritoneal involvement, or distant metastasis to the supraclavicular LN. Further studies are warranted to explore the biologic mechanisms that underlie the favorable prognosis of patients with stage IV disease, so-assigned on the basis of supraclavicular LN metastasis.

CONFLICT OF INTEREST

Dong Hoon Suh and Jae-Weon Kim serve as Editors of the JGO but have no role in the decision to publish this article. No other conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

This study was presented at the 46th Semi-Annual Meeting of Korean Society of Gynecologic Oncology and awarded Shin Poong Howol Scholarship prize.

REFERENCES

1. Oh HK, Sin JI, Choi J, Park SH, Lee TS, Choi YS. Overexpression of CD73 in epithelial ovarian carcinoma is associated with better

- prognosis, lower stage, better differentiation and lower regulatory T cell infiltration. *J Gynecol Oncol* 2012;23:274-81.
2. Kim HS, Ahn JH, Chung HH, Kim JW, Park NH, Song YS, et al. Impact of intraoperative rupture of the ovarian capsule on prognosis in patients with early-stage epithelial ovarian cancer: a meta-analysis. *Eur J Surg Oncol* 2013;39:279-89.
 3. Goudge CS, Li Z, Downs LS Jr. The influence of intraoperative tumor rupture on recurrence risk in Stage Ic epithelial ovarian cancer. *Eur J Gynaecol Oncol* 2009;30:25-8.
 4. Seidman JD, Cosin JA, Wang BG, Alsop S, Yemelyanova A, Fields A, et al. Upstaging pathologic stage I ovarian carcinoma based on dense adhesions is not warranted: a clinicopathologic study of 84 patients originally classified as FIGO stage II. *Gynecol Oncol* 2010;119:250-4.
 5. Berek JS. Lymph node-positive stage IIIC ovarian cancer: a separate entity? *Int J Gynecol Cancer* 2009;19 Suppl 2:S18-20.
 6. Cliby WA, Aletti GD, Wilson TO, Podratz KC. Is it justified to classify patients to stage IIIC epithelial ovarian cancer based on nodal involvement only? *Gynecol Oncol* 2006;103:797-801.
 7. Baek SJ, Park JY, Kim DY, Kim JH, Kim YM, Kim YT, et al. Stage IIIC epithelial ovarian cancer classified solely by lymph node metastasis has a more favorable prognosis than other types of stage IIIC epithelial ovarian cancer. *J Gynecol Oncol* 2008;19:223-8.
 8. Denny L, Quinn M, Hacker N. FIGO cancer report 2012. *Int J Gynaecol Obstet* 2012;119 Suppl 2:S89.
 9. Vergote I, De Brabanter J, Fyles A, Bertelsen K, Einhorn N, Sevelde P, et al. Prognostic importance of degree of differentiation and cyst rupture in stage I invasive epithelial ovarian carcinoma. *Lancet* 2001;357:176-82.
 10. Kodama S, Kase H, Tanaka K, Matsui K. Multivariate analysis of prognostic factors in patients with endometrial cancer. *Int J Gynaecol Obstet* 1996;53:23-30.
 11. Bakkum-Gamez JN, Richardson DL, Seamon LG, Aletti GD, Powless CA, Keeney GL, et al. Influence of intraoperative capsule rupture on outcomes in stage I epithelial ovarian cancer. *Obstet Gynecol* 2009;113:11-7.
 12. Raungkaewmanee S, Tangjitgamol S, Manusirivithaya S, Srijaipracharoen S, Thavaramara T. Platelet to lymphocyte ratio as a prognostic factor for epithelial ovarian cancer. *J Gynecol Oncol* 2012;23:265-73.
 13. Bachmann C, Bachmann S, Fehm T, Staebler A, Becker S, Rothmund R, et al. Nodal status: its impact on prognosis in advanced ovarian cancer. *J Cancer Res Clin Oncol* 2012;138:261-7.
 14. Santillan A, Karam AK, Li AJ, Giuntoli R 2nd, Gardner GJ, Cass I, et al. Secondary cytoreductive surgery for isolated nodal recurrence in patients with epithelial ovarian cancer. *Gynecol Oncol* 2007;104:686-90.
 15. Benedetti Panici P, Perniola G, Angioli R, Zullo MA, Manci N, Palaia I, et al. Bulky lymph node resection in patients with recurrent epithelial ovarian cancer: impact of surgery. *Int J Gynecol Cancer* 2007;17:1245-51.
 16. Powless CA, Aletti GD, Bakkum-Gamez JN, Cliby WA. Risk factors for lymph node metastasis in apparent early-stage epithelial ovarian cancer: implications for surgical staging. *Gynecol Oncol* 2011;122:536-40.
 17. Pickel H, Lahousen M, Stettner H, Girardi F. The spread of ovarian cancer. *Baillieres Clin Obstet Gynaecol* 1989;3:3-12.