

Utilizing Nanobody Technology to Target Non-Immunodominant Domains of VAR2CSA

Sisse B. Ditlev^{1*}, Raluca Florea^{2,3}, Morten A. Nielsen¹, Thor G. Theander¹, Stefan Magez^{2,3}, Philippe Boeuf^{4,5}, Ali Salanti¹

1 Centre for Medical Parasitology at Department of International Health, Immunology, and Microbiology, University of Copenhagen and at Department of Infectious Diseases, Copenhagen University Hospital (Rigshospitalet), Copenhagen, Denmark, **2** Cellular and Molecular Immunology Research Unit, Vrije Universiteit Brussel, Brussels, Belgium, **3** Department of Structural Biology, VIB, Brussels, Belgium, **4** The University of Melbourne, Department of Medicine, Parkville, Victoria, Australia, **5** Victorian Infectious Diseases Service, Royal Melbourne Hospital, Parkville, Victoria, Australia

Abstract

Placental malaria is a major health problem for both pregnant women and their fetuses in malaria endemic regions. It is triggered by the accumulation of *Plasmodium falciparum*-infected erythrocytes (IE) in the intervillous spaces of the placenta and is associated with foetal growth restriction and maternal anemia. IE accumulation is supported by the binding of the parasite-expressed protein VAR2CSA to placental chondroitin sulfate A (CSA). Defining specific CSA-binding epitopes of VAR2CSA, against which to target the immune response, is essential for the development of a vaccine aimed at blocking IE adhesion. However, the development of a VAR2CSA adhesion-blocking vaccine remains challenging due to (i) the large size of VAR2CSA and (ii) the extensive immune selection for polymorphisms and thereby non-neutralizing B-cell epitopes. Camelid heavy-chain-only antibodies (HcAbs) are known to target epitopes that are less immunogenic to classical IgG and, due to their small size and protruding antigen-binding loop, able to reach and recognize cryptic, conformational epitopes which are inaccessible to conventional antibodies. The variable heavy chain (VHH) domain is the antigen-binding site of camelid HcAbs, the so called Nanobody, which represents the smallest known (15 kDa) intact, native antigen-binding fragment. In this study, we have used the Nanobody technology, an approach new to malaria research, to generate small and functional antibody fragments recognizing unique epitopes broadly distributed on VAR2CSA.

Citation: Ditlev SB, Florea R, Nielsen MA, Theander TG, Magez S, et al. (2014) Utilizing Nanobody Technology to Target Non-Immunodominant Domains of VAR2CSA. PLoS ONE 9(1): e84981. doi:10.1371/journal.pone.0084981

Editor: Gordon Langsley, Institut national de la santé et de la recherche médicale - Institut Cochin, France

Received: October 7, 2013; **Accepted:** November 28, 2013; **Published:** January 21, 2014

Copyright: © 2014 Ditlev et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: SBD is supported by a PhD studentship from the Danish Research Council for Development Research (RUF). RF is supported by IWT (Agentschap voor Innovatie door Wetenschap en Technologie) Belgium. This study received funding from Danish National Advanced Technology Foundation and Proof of Concept Foundation DTU. The project also was supported by Novo Nordisk Fonden, Project number: 1015627 and also supported by a Project grant from the Australian NHMRC to Philippe Boeuf Grant # 1003384. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: sditlev@sund.ku.dk

These authors contributed equally to this work.

These authors also contributed equally to this work.

Introduction

Placental malaria is caused by the protozoan *Plasmodium falciparum* transmitted by the female Anopheles mosquito and can lead to maternal anemia, low birth weight, preterm delivery and increased infant and maternal mortality. *P. falciparum*-infected erythrocytes (IE) accumulate in the placenta by adhering to chondroitin sulfate A (CSA) chains on chondroitin sulfate proteoglycans (CSPG) in the intervillous spaces and on the microvillous membrane of the placental syncytiotrophoblast [1]. IE adhesion is mediated by VAR2CSA, a pregnancy-specific member of the *P. falciparum* erythrocyte membrane protein 1 (PfEMP1) family expressed on the surface of IE [2]. In malaria endemic areas, children develop clinical immunity through the acquisition of a broad repertoire of anti-PfEMP1 antibodies [3]. Pregnant women become susceptible to malaria, as they have not previously acquired antibodies to the pregnancy-specific PfEMP1 variant VAR2CSA. IE adhesion to the placenta triggers the recruitment and activation of maternal mononuclear cells

secreting pro-inflammatory cytokines, leading to further inflammation and negative effects on placental function [4] and fetal development [5]. During subsequent pregnancies, women build up protective immunity to placental malaria by acquiring anti-VAR2CSA antibodies that prevent IE binding to CSA in the placenta [6,7]. VAR2CSA is therefore an attractive candidate for a vaccine against placental malaria. VAR2CSA is a large protein (~350 kDa) consisting of six Duffy-Binding-Like domains and several inter domains [8,9]. Even though VAR2CSA is conserved relative to other PfEMP1 proteins, there is a substantial sequence variation [10]. Thus, a major challenge for vaccine development is to define VAR2CSA epitopes that can induce a broad anti-adhesive antibody response. Several single domains of VAR2CSA have been shown to be able to induce functional adhesion-blocking antibodies by immunization in laboratory animals, even though these domains do not directly take part in VAR2CSA binding to CSA [11–17]. Recent studies have highlighted the importance of the N-terminal part of VAR2CSA in CSA-binding and antibodies targeting this region effectively prevent VAR2CSA




Figure 1. Sequence alignment of the 17 VAR2CSA-specific Nbs. The clones used for Nanobody production were sequenced, converted to amino acid sequences and aligned. The complementarity determining regions (CDRs) 1–3 make up the binding paratope and the framework regions (FRs) 1–4 are indicated.

doi:10.1371/journal.pone.0084981.g001

binding to CSA [18–20]. However, identification of smaller VAR2CSA regions responsible for CSA binding is a major challenge since VAR2CSA is a large and complex antigen. The identification of such epitopes could pave the way towards designing an effective multivalent VAR2CSA vaccine. We have extensively explored the naturally-acquired response to VAR2CSA in order to differentiate the protective adhesion-blocking response from the immuno-dominant, non-functional response focused towards the DBL3X, DBL5ε and DBL6ε domains of VAR2CSA [21,22]. Indeed, the majority of the naturally-acquired response targets the C-terminal part of VAR2CSA that does not mediate binding to CSA [22]. The majority of hybridomas cloned from mice and rats immunized with full-length VAR2CSA produced IgG against DBL3X and DBL5ε domains and these antibodies did not block IE adhesion to CSA (unpublished data).

In this study, we introduce an approach new to malaria research to produce versatile and functional monoclonal reagents against VAR2CSA circumventing IgG immuno-dominant epitopes, based on camelid heavy-chain-only antibodies (HcAbs). The variable heavy chain (VHH) domain is the antigen-binding site of camelid HcAbs and represents the smallest (15 kDa), intact, native antigen-binding fragment [23]. Recombinantly-produced VHHs are termed Nanobodies (Nbs). Nbs are easily expressed in large quantities, are soluble, have high thermal stability, and bind the

target antigen with the *high affinity and specificity typical of conventional antibodies* [24]. Due to their small size and protruding antigen-binding complementarity determining region-3 (CDR3) loop, Nbs have the capacity to reach and recognize cryptic, conformational epitopes that are inaccessible to conventional antibodies [25–27]. Moreover, Nbs generally interact with epitopes that are less antigenic as compared to conventional antibodies [28]. These properties make them potent alternatives to conventional antibodies for non-immuno-dominant epitopes.

To investigate the potential of Nbs as a tool for targeting VAR2CSA, an alpaca was immunized with full-length VAR2CSA and the Nbs generated were screened for VAR2CSA-specificity and functionality. Using this approach, we produced a large panel of VAR2CSA-specific Nbs targeting epitopes broadly distributed over VAR2CSA, including against the poorly immunogenic CSA-binding regions. This study highlights the advantages of using the Nanobody technology for production of monoclonal reagents to pathogenic antigens that have evolved immuno-dominant regions.

Results

Library screening and selection of VAR2CSA-specific Nanobodies (Nbs)

A Nanobody phagemid library with a size of 1.7×10^8 colonies was generated by cloning HcAbs from peripheral blood mononu-




Figure 2. Recognition of immobilized full-length VAR2CSA by each of the 17 Nbs. Microtiter plates were coated with VAR2CSA protein (50 nM) and incubated with individual Nbs (50 nM). Binding was detected with rabbit anti-camel antibody and HRP-conjugated goat anti-rabbit antibody. Optical density was measured at 490 nm after 20 min. A non-VAR2CSA-PfEMP1 (50 nM) was used as negative control. Data are represented as mean and standard deviations of three independent experiments.

doi:10.1371/journal.pone.0084981.g002




Figure 3. Recognition of immobilized VAR2CSA domains by Nbs (Nb01–Nb17). Baculovirus-produced domains of VAR2CSA (50 nM) were coated on microtiter plates and incubated with each of the Nbs (50 nM). The binding was detected with rabbit anti-camel and goat anti-rabbit HRP-conjugated antibodies and optical density measured at 490 nm after 20 min. Non-VAR2CSA PfEMP1 (50 nM) was used as negative control. The assay was performed several times with similar result. doi:10.1371/journal.pone.0084981.g003

clear cells of the VAR2CSA immunized alpaca. An aliquot of the library was infected with M13K07 helper phages and phages expressing VAR2CSA-specific VHs were enriched by three consecutive rounds of bio-panning on VAR2CSA. From the

second and third rounds of panning, 100 colonies randomly chosen were screened for VAR2CSA recognition by periplasmic extraction, followed by ELISA. The majority (70%) of these clones were found to specifically bind VAR2CSA but not the control

Table 1. Summary of the reactivity of the individual Nbs against VAR2CSA (full length and domains).

	FV2-FCR3	ID1-ID2a	DBL1	DBL2	DBL3	DBL4	DBL5	DBL6
Nb01	X	X						
Nb02	X						X	
Nb03	X					X		
Nb04	X							X
Nb05	X						X	
Nb06	X							X
Nb07	X	X						
Nb08	X							X
Nb09	X	X					X	
Nb10	X	X						
Nb11	X						X	
Nb12	X	X						
Nb13	X					X		
Nb14	X						X	
Nb15	X							X
Nb16	X					X		
Nb17	X					X		

A cross in the box indicates specific binding.

doi:10.1371/journal.pone.0084981.t001

protein BSA. Nucleotide sequence analysis of these clones revealed 17 genetically distinct VAR2CSA binders. Their paratope (CDR-3 region) amino acid sequences differed by several amino acids (Figure 1).

Expression and purification of anti-VAR2CSA Nbs

The VHH vectors encoding the 17 VAR2CSA-specific Nbs were sub-cloned into the pHEN6c expression vector containing a C-terminal His₆ tag. Nbs were expressed in WK6 *E. coli* cells and purified using HisTrap columns. The production yields of each Nb varied from 4 mg to 11 mg per litre culture. The SDS PAGE analysis of the purified Nbs showed no impurities after the purification steps and only showed formation of dimers in the Nb03 production. (Figure S1).

Nanobody reactivity to recombinant VAR2CSA protein

To verify the specificity of the purified Nbs (Nb01–Nb17), direct binding to different domains of recombinant VAR2CSA was analyzed by ELISA (Figure 2). All 17 Nbs specifically recognized full-length VAR2CSA (FV2) and not a non-pregnancy specific PfEMP1 used as negative control. The Nbs were subsequently screened against individual VAR2CSA domains (DBL1-6) and against the ID1–ID2a region, which represents the minimal-CSA binding region of VAR2CSA [18,20] (Figure 3). These domains were produced using a baculo-virus expression system, as described in [11,20]. Table 1 shows an overview of the VAR2CSA domain recognition by the individual Nbs. The three C-terminal domains (DBL4-6) were recognized by twelve of the Nbs (71%). None of the Nbs recognized the individual N-terminal domains (DBL1-3) of VAR2CSA whereas five Nbs (Nb01, Nb07, Nb09, Nb10 and Nb12) recognized the ID1–ID2a minimal CSA-binding domain expressed in the baculovirus expression system.

These five Nbs recognized to a similar degree ID1–ID2a minimal CSA-binding domains expressed in *E. coli* and S2 cells (Figure 4A). When tested against the minimal CSA-binding region

of a heterologous parasite strain (3D7), three of these five Nbs (Nb09 and to some degree Nb10 and Nb12) were found to be cross-reactive (Figure 4B).

We tested whether the epitopes recognized by the 17 Nbs were discontinuous using Western Blotting of reduced or non-reduced recombinant VAR2CSA protein. The Nbs specific for single domains showed similar binding to both the reduced and the non-reduced recombinant protein (Figure 5A–C), whereas the minimal CSA-binding region-specific Nbs showed no or very limited reactivity to the reduced protein (Figure 5D).

Nanobody reactivity to native VAR2CSA protein expressed on the surface of IE

Epitopes exposed on recombinant proteins may not be surface-exposed on the native VAR2CSA protein expressed by IE. Therefore, we used flow cytometry to test the reactivity of the Nbs to VAR2CSA-expressing parasite lines (Figure 6). All Nbs showed (except Nb06 and Nb13) some degree of reactivity to VAR2CSA-expressing IE, including the homologous parasite line FCR3 and two heterologous parasite lines.

Nanobody-mediated inhibition of IE binding to CSA

We evaluated the capacity of the Nbs to inhibit the adhesion of VAR2CSA-expressing IE to the placental receptor chondroitin sulfate A (Decorin) (Figure 7). Most Nbs increased IE adhesion to CSA but Nb01, Nb09 and Nb10, specific for VAR2CSA minimal CSA-binding region reproducibly inhibited CSA adhesion of the homologous FCR3 parasite line (15%–49% mean adhesion inhibition). The cross-inhibitory activity of the Nbs specific for VAR2CSA minimal CSA-binding region was assessed using two heterologous parasite lines (NF54 and 7201). All the Nbs specific for VAR2CSA minimal CSA-binding region reduced 7201 IE adhesion to CSA by at least 42% whereas NF54 IE adhesion to CSA was only inhibited by Nb09 (57% adhesion inhibition).




Figure 4. Recognition of the minimal CSA-binding region of VAR2CSA. (A) Reactivity of Nbs to specific minimal CSA-binding regions produced in different expression systems. ID1-ID2a proteins produced either in a baculo-virus expression system (BV), or in *E. coli* (coli) or in Schneider 2 (S2) cells were coated (50 nM) on microtiter plates and incubated with 50 nM Nb01, Nb07, Nb09, Nb10 or Nb12 (Nbs specific for VAR2CSA minimal CSA-binding domain). Nb02 (DBL5-specific) was used as a negative control. (B) Cross-reactivity of Nbs specific for the minimal CSA-binding region of FCR3 to recombinant proteins covering the minimal CSA-binding region (DBL1-ID2a and ID1-ID2a) of the heterologous 3D7 parasite line produced in the baculovirus expression system. In both assays, the binding was detected with rabbit-anti-camel antibody and HRP-labeled goat anti-rabbit antibody and the optical density was measured at 490 nm after 20 min. A non-VAR2CSA protein was used as a negative control.

doi:10.1371/journal.pone.0084981.g004

Discussion

Identification of VAR2CSA epitopes that are target of protective antibodies is key to the development of multivalent vaccines that can protect pregnant women against placental malaria. However, the mapping of such epitopes has been hampered by the large and complex nature of VAR2CSA and the poor understanding of its interaction with the placental receptor CSA. Production and isolation of monoclonal antibodies to VAR2CSA from malaria-exposed women or VAR2CSA-immunized animals has been limited to the immuno-dominant DBL3X and DBL5ε domains [22,29]. Because HcAbs can recognize poorly immunogenic epitopes [30] we hypothesized that HcAbs generated against VAR2CSA could circumvent the

immuno-dominance of epitopes of the DBL3X and DBL5ε domains and induce a response against other VAR2CSA domains.

We immunized an alpaca with full-length VAR2CSA (FV2) and selected seventeen VHHs that specifically recognized FV2. This approach avoided a focused response towards the DBL3X and DBL5ε immuno-dominant domains since none of the Nbs targeted DBL3X and some Nbs recognized the less immunogenic CSA-binding N-terminal region of VAR2CSA. The twelve Nbs specific for the three C-terminal domains (DBL4ε, DBL5ε and DBL6ε) recognized these as single domains, whereas the five Nbs recognizing the N-terminal region did not react with single domains but was dependent on the entire minimal-CSA binding region, ID1-ID2a, encompassing DBL2X with the flanking parts ID1 and ID2. These results suggest that Nbs specific for the




Figure 5. Nbs recognize both continuous and discontinuous epitopes. Reduced and non-reduced baculovirus-produced VAR2CSA domains were transferred onto membranes by Western blotting and probed with the corresponding domain-specific Nbs; (A) DBL4 ϵ protein probed with Nb03, Nb13, Nb16 and Nb17. (B) DBL5 ϵ protein probed with Nb02, Nb05, Nb11, Nb14. (C) DBL6 ϵ protein probed with Nb15, Nb08, Nb06 and Nb04. (D) DBL1-ID2a protein (containing minimal CSA-binding region) probed with Nb12, Nb10, Nb09, Nb07 and Nb01. Binding was detected with rabbit-anti-camel and HRP-labeled goat-anti-rabbit antibodies. Bands larger than monomer size correspond to multimers formed by intermolecular disulfide bonds (monomers are marked with arrows).
doi:10.1371/journal.pone.0084981.g005

minimal CSA-binding region target discontinuous epitopes. This is in line with Western blot data showing that C-terminal-specific Nbs recognized linear epitopes whereas the N-terminal-specific Nbs recognize discontinuous epitopes. Taken together, these results support the proposed overall fold of VAR2CSA, in which the C-terminal domains are presented as single and accessible

domains whereas the N-terminal domains containing the minimal CSA-binding region have a more complex conformation.

Nb01, Nb09, Nb10, and Nb12, which specifically recognize the minimal-CSA binding region of VAR2CSA, were found to partially block binding of VAR2CSA to CSA. Importantly, Nb09 was both cross-inhibitory and cross-reactive since it both




Figure 6. Nanobody recognition of native VAR2CSA expressed on the surface of *P. falciparum*-infected erythrocytes (IE). Binding of VAR2CSA-specific Nbs to VAR2CSA-expressing IE was measured by flow cytometry. Three different *P. falciparum* strains (FCR3, 7201 and NF54) were tested. Each bar represents reactivity of 25 μ l nanobody (0.1 mg/ml) to 50 μ l IE (2×10^5 parasites/ml). Values are normalized to the mean fluorescence intensity (MFI) of the negative control (neg ctr; IE incubated only with the secondary and tertiary antibodies). Specific recognition was defined by an MFI ratio >1.2 . Error bars in A) represents standard deviation of three independent experiments. doi:10.1371/journal.pone.0084981.g006

recognized the NF54 parasite and the recombinant VAR2CSA from the 3D7 strain (which has identical VAR2CSA sequence as the NF54 parasite) and reduced the NF54 parasite binding. The level of recognition was generally lower for the minimal CSA-binding region-specific Nbs than for Nbs recognizing the C-terminal domains. This could indicate that the epitopes recognized by these Nbs specific for the minimal CSA-binding region are more hidden in the structure, making it physically difficult for the secondary antibodies to bind the Nbs.

The CDR regions of Nbs are often longer and more flexible than corresponding regions of conventional antibodies and thus can bind epitopes that physically cannot be targeted by conventional immunoglobulins [28,31,32]. This could explain the concordance between reactivity to recombinant protein and to native VAR2CSA. The fact that Nbs can “penetrate” and bind hidden epitopes could stabilize the minimal CSA-binding region and facilitate crystallization, which has proven to be very challenging.

Natural selection of pathogen-derived antigens is associated with epitopes of varying immunogenicity, and it is likely that epitopes of functional importance will have evolved to avoid host antibody response. We demonstrated that the nanobody technology, through its capacity to recognize non-immuno-dominant and hidden epitopes, produces versatile monoclonal reagents to such antigens. Besides being used for quality control of VAR2CSA vaccine construct, VAR2CSA-specific Nbs could be novel diagnostic or therapeutic tools and could provide novel insights

into structure/function of this complex antigen. This is essential to the design of a multivalent VAR2CSA vaccine.

Materials and Methods

Immunization

An alpaca was immunized with purified full-length VAR2CSA (FV2) recombinant protein expressed as described in [33]. Immunization of the alpaca and a bleed of 50 ml were done by Alpa-Vet (www.alpa-vet.be) and the procedure was approved by the ethics committee of the Free University Brussels (file number 12-601-1). The peripheral blood lymphocytes were isolated from the 50 ml of blood of the immunized alpaca using Lymphoprep (Nycomed).

Construction of the Nanobody library and selection of VAR2CSA-specific Nanobodies

The nanobody library was constructed as previously described by Conrath et al, 2001. Briefly, total mRNA was extracted from 2×10^7 lymphocytes from which 50 μ g mRNA was used for the synthesis of cDNA with oligodT primer. Using the cDNA as template, the fragments encoding for both the VH and VHH domains of camelid IgGs were amplified by PCR with the CALL001 and CALL002 primers [34]. The VHH gene fragment (600 bp) was separated from the VH genes (900 bp) by agarose gel extraction and re-amplified with the framework-1 and framework-4 primers [35] in order to introduce PstI and NotI restriction sites. The PCR fragments were cloned into the phage-display phagemid




Figure 7. Adhesion-inhibitory capacity of VAR2CSA-specific nanobodies. (A) Ability of the individual Nbs to inhibit the adhesion of VAR2CSA-expressing IE (FCR3 line) to Decorin. Nbs specific for the minimal CSA-binding-region of VAR2CSA (Nb01, Nb07, Nb09, Nb10 and Nb12) were tested three times on the homologous parasite FCR3-CSA. (B) Adhesion inhibition capacity of the minimal CSA-binding-specific Nbs to VAR2CSA-expressing heterologous parasite lines NF54 and (C) 7201. Parasite binding to CSA ligand without Nbs was set to 100. doi:10.1371/journal.pone.0084981.g007

vector pHEN4 and electro-transformed in electro-competent *E. coli* TG1 cells. The VHH library was expressed on phage after infection with M13K07 helper phages (Invitrogen). Phages expressing VAR2CSA-specific VHHs on their coat proteins were enriched via three consecutive rounds of panning on microtiter plates (Nunc MaxiSorp®) coated with FV2 (10 µg/well). From the second and third round of panning, individual colonies were selected, grown in Terrific Broth (TB) media and VHH expression was induced with 1 mM isopropyl β-d-thiogalactopyranoside (IPTG). Identification of *E. coli* TG1 clones expressing anti-VAR2CSA Nbs was performed by periplasmic extract ELISA.

Expression and purification of selected nanobodies

Genes encoding VAR2CSA-specific VHH were sub-cloned into pHEN6c expression vector using BstEII and PstI restriction sites, between a *pelB* leader signal sequence (for VHH periplasmic transport) and a C-terminal hexa-histidine (His₆) tag and transformed into WK6 *E. coli* cells. Production of the recombinant Nbs was done in shaker flasks by growing the cells in TB media supplemented with ampicillin (0.1 mg/ml), MgCl₂ (2 mM) and glucose (0.1%). When optical density at 600 nm was between 0.6–0.8, Nbs expression was induced with 1 mM IPTG for 16 h at 28°C under agitation (200 rpm) overnight. Periplasmic extract proteins were released by osmotic shock. The recombinant Nbs were purified from the periplasmic extract using HisTrap HP columns (17-5247-01, GE Healthcare). The bound protein was eluted with 10 mM NaH₂PO₄ (pH 7.4), 500 mM NaCl, and

500 mM imidazole. Nbs purity and formation of disulfide bonds were verified by SDS-PAGE using NuPage Novex Bis-Tris mini gels (Invitrogen).

ELISA mapping of VAR2CSA-specific nanobodies

Nanobody recognition of recombinant VAR2CSA protein and domains was measured by ELISA. Recombinant VAR2CSA (50 nM in PBS) was coated on Nunc MaxiSorp® plates overnight at 4°C. Non-specific binding sites were blocked by incubating the plates with 5% skim milk in PBS for one hour at room temperature (RT). After three washes in PBS+0.05% Tween-20, Nbs diluted in 1% skim milk in PBS were added to the wells (50 nM) and incubated for 90 min at RT. The plates were washed three times in PBS+0.05% Tween-20 before polyclonal rabbit anti-camel Ab (kindly provided by B. Stijlemans [36]) was added for 1 h (non-commercial; diluted 1:2000 in 1% skim milk in PBS.) After three washes in PBS+0.05% Tween-20, horseradish peroxidase-conjugated (HRP) anti-rabbit-IgG (DAKO, P0448) diluted 1:2000 in 1% skim milk in PBS was added for 1 h. The plates were washed three times in PBS+0.05% Tween-20 and binding of Nb was visualized by adding o-phenylenediamine substrate. After 20 min, the HRP enzymatic reaction was stopped by adding 2.5 M H₂SO₄ and the optical density at 490 nm measured using an ELISA plate reader (VersaMax Molecular Devices).

Western Blot Analysis

Western blot was performed to determine if the individual Nbs recognized continuous or discontinuous VAR2CSA epitopes. Recombinant VAR2CSA protein (0.4 µg/lane) was loaded on an 8–12% Bis-Tris SDS gel (Invitrogen) under reduced or non-reduced conditions. The separated proteins were transferred to nitrocellulose membranes by wet blotting. After blocking non-specific sites with 5% skim milk in TBS+0.05% Tween 20 (TBST), the membranes were incubated for 1½ hour at RT sequentially with VAR2CSA-specific nanobody (diluted to 50 nM in TBST), rabbit polyclonal anti-camel Ab (diluted 1:2000 in TBST) and HRP-conjugated goat anti-rabbit antibody (DAKO; P0448 diluted 1:3000 in TBST). Between incubations, the membranes were washed three times with TBST and Nb recognition of VAR2CSA was detected by chemiluminescence.

Nanobody reactivity to IE using flow cytometry

Nbs reactivity to VAR2CSA-expressing IE was measured by flow cytometry. Approximately 100,000 late stage IE (50 µl at 2×10^6 IE/ml) labelled with ethidium bromide were incubated with 50 µl Nbs (0.1 mg/ml). Nanobody binding was detected by 50 µl mouse-anti-penta-His Ab (Qiagen; 34660 diluted 1:100 in PBS) and a FITC-labelled anti-mouse Ab (Vector; FI-2000; diluted 1:200 in PBS). Each labelling step was conducted for 30 min at 4°C. As a negative control, IE were incubated with the secondary and tertiary antibody only. Data from 5,000 IE were collected on a FC500 flow cytometer (Beckman Coulter). The mean FITC fluorescence intensity was determined using Winlist Software (Verity Software House).

Adhesion-inhibition capacity of nanobodies

The adhesion-inhibition capacity of various Nbs was measured in a high throughput assay as previously described [37]. Briefly,

References

- Achur RN, Valiyaveetil M, Alkhalil A, Ockenhouse CF, Gowda DC (2000) Characterization of proteoglycans of human placenta and identification of unique chondroitin sulfate proteoglycans of the intervillous spaces that mediate the adherence of *Plasmodium falciparum*-infected erythrocytes to the placenta. *J Biol Chem* 275: 40344–40356. 10.1074/jbc.M006398200 [doi];M006398200 [pii].
- Salanti A, Staaloe T, Lavstsen T, Jensen AT, Sowa MP et al. (2003) Selective upregulation of a single distinctly structured var gene in chondroitin sulphate A-adhering *Plasmodium falciparum* involved in pregnancy-associated malaria. *Mol Microbiol* 49: 179–191. 3570 [pii].
- Cham GK, Turner L, Kurtis JD, Mutabingwa T, Fried M et al. (2010) Hierarchical, domain type-specific acquisition of antibodies to *Plasmodium falciparum* erythrocyte membrane protein 1 in Tanzanian children 10. *Infect Immun* 78: 4653–4659. IAL00593-10 [pii];10.1128/IAI.00593-10 [doi].
- Boeuf P, Aitken EH, Chandrasiri U, Chua CL, McInerney B et al. (2013) *Plasmodium falciparum* malaria elicits inflammatory responses that dysregulate placental amino acid transport 7. *PLoS Pathog* 9: e1003153. 10.1371/journal.ppat.1003153 [doi];PPATHOGENS-D-12-01766 [pii].
- Brabin BJ, Romagosa C, Abdelgalil S, Menendez C, Verhoeff FH et al. (2004) The sick placenta—the role of malaria. *Placenta* 25: 359–378. 10.1016/j.placenta.2003.10.019 [doi];S0143400403003072 [pii].
- Salanti A, Dahlback M, Turner L, Nielsen MA, Barfod L et al. (2004) Evidence for the involvement of VAR2CSA in pregnancy-associated malaria. *J Exp Med* 200: 1197–1203.
- Duffy PE, Fried M (2003) Antibodies that inhibit *Plasmodium falciparum* adhesion to chondroitin sulfate A are associated with increased birth weight and the gestational age of newborns. *Infect Immun* 71: 6620–6623.
- Trimnell AR, Kraemer SM, Mukherjee S, Phippard DJ, Janes JH et al. (2006) Global genetic diversity and evolution of var genes associated with placental and severe childhood malaria. *Mol Biochem Parasitol* 148: 169–180. S0166-6851(06)00110-1 [pii];10.1016/j.molbiopara.2006.03.012 [doi].
- Andersen P, Nielsen MA, Resende M, Rask TS, Dahlback M et al. (2008) Structural insight into epitopes in the pregnancy-associated malaria protein VAR2CSA. *PLoS Pathog* 4: e42. 07-PLPA-RA-0432 [pii];10.1371/journal.ppat.0040042 [doi].
- Bockhorst J, Lu F, Janes JH, Keebler J, Gamain B et al. (2007) Structural polymorphism and diversifying selection on the pregnancy malaria vaccine

candidate VAR2CSA. *Mol Biochem Parasitol* 155: 103–112. S0166-6851(07)00163-6 [pii];10.1016/j.molbiopara.2007.06.007 [doi].

2×10^5 tritium-labelled late stage IE, pre-incubated or not with Nbs, were added to a decorin-coated plate (Sigma; D8428: 2 µg/ml) for 90 min at 37°C. Unbound IE were washed away by re-suspension performed by a pipetting robot (Beckman Coulter) and the detection of adhering IE was determined by liquid scintillation counting on a Topcount NXT (Perkin-Elmer).

Supporting Information

Figure S1 Coomassie-stained SDS-PAGE of purified Nbs Nb01–Nb17. A: example of expression and purification of two different Nbs (Nb04 in lanes 2–6 and Nb07 in lanes 8–12). Lanes 2 and 8: Total protein in periplasmic lysate as loaded onto a HIS-column non-reduced. Lanes 3 and 9: Run through after purification on a HIS-column non-reduced. Lanes 4 and 10: Column wash non-reduced, Lanes 5, 6, 11 and 12: HIS-purified nanobody with (Lanes 5 and 11) or without (Lanes 6 and 12) reducing agent DTT. Lanes 1 and 7 are molecular markers (ProSieve™ Color Protein Marker, Lonza). Figure B and C show the 17 produced and purified nanobodies under non-reduced conditions.

(TIF)

Acknowledgments

The authors would like to thank Elham Alijazaeri and Christina Holm for excellent technical assistance. The protein produced in Schneider2 (S2) cells was kindly provided by ExpreS²ion Biotechnologies

Author Contributions

Conceived and designed the experiments: SBD RF MAN TGT SM PB AS. Performed the experiments: SBD RF MAN PB. Analyzed the data: SBD RF MAN PB AS. Wrote the paper: SBD RF MAN TGT SM PB AS.

- Nielsen MA, Pinto VV, Resende M, Dahlback M, Ditlev SB et al. (2009) Induction of adhesion-inhibitory antibodies against placental *Plasmodium falciparum* parasites by using single domains of VAR2CSA. *Infect Immun* 77: 2482–2487. IAL00159-09 [pii];10.1128/IAI.00159-09 [doi].
- Obiakor H, Avril M, Macdonald NJ, Srinivasan P, Reiter K et al. (2013) Identification of VAR2CSA domain-specific inhibitory antibodies of the *Plasmodium falciparum* erythrocyte membrane protein 1 using a novel flow cytometry assay 4. *Clin Vaccine Immunol* 20: 433–442. CVI.00638-12 [pii];10.1128/CVI.00638-12 [doi].
- Avril M, Cartwright MM, Hathaway MJ, Smith JD (2011) Induction of strain-transcendent antibodies to placental-type isolates with VAR2CSA DBL3 or DBL5 recombinant proteins. *Malar J* 10: 36. 1475-2875-10-36 [pii];10.1186/1475-2875-10-36 [doi].
- Fried M, Avril M, Chaturvedi R, Fernandez P, Lograsso J et al. (2013) Multilaboratory approach to preclinical evaluation of vaccine immunogens for placental malaria 12. *Infect Immun* 81: 487–495. IAL01106-12 [pii];10.1128/IAI.01106-12 [doi].
- Magistrado PA, Minja D, Doritchamou J, Ndam NT, John D et al. (2011) High efficacy of anti DBL4varepsilon-VAR2CSA antibodies in inhibition of CSA-binding *Plasmodium falciparum*-infected erythrocytes from pregnant women. *Vaccine* 29: 437–443. S0264-410X(10)01593-8 [pii];10.1016/j.vaccine.2010.10.080 [doi].
- Fernandez P, Petres S, Mecheri S, Gysin J, Scherf A (2010) Strain-transcendent immune response to recombinant Var2CSA DBL5-epsilon domain block P. *falciparum* adhesion to placenta-derived BeWo cells under flow conditions 15. *PLoS One* 5: e12558. 10.1371/journal.pone.0012558 [doi].
- Fernandez P, Viebig NK, Dechavanne S, Lepolard C, Gysin J et al. (2008) Var2CSA DBL6-epsilon domain expressed in HEK293 induces limited cross-reactive and blocking antibodies to CSA binding parasites. *Malar J* 7: 170. 1475-2875-7-170 [pii];10.1186/1475-2875-7-170 [doi].
- Dahlback M, Jorgensen LM, Nielsen MA, Clausen TM, Ditlev SB et al. (2011) The chondroitin sulfate A-binding site of the VAR2CSA protein involves multiple N-terminal domains. *J Biol Chem* 286: 15908–15917. M110.191510 [pii];10.1074/jbc.M110.191510 [doi].

19. Srivastava A, Gangnard S, Dechavanne S, Amirat F, Lewit BA et al. (2011) Var2CSA Minimal CSA Binding Region Is Located within the N-Terminal Region. *PLoS One* 6: e20270. 10.1371/journal.pone.0020270 [doi];PONE-D-11-03862 [pii].
20. Clausen TM, Christoffersen S, Dahlback M, Langkilde AE, Jensen KE et al. (2012) Structural and Functional Insight Into How the Plasmodium falciparum VAR2CSA Protein Mediates Binding to Chondroitin Sulfate A in Placental Malaria 18. *J Biol Chem*. M112.348839 [pii];10.1074/jbc.M112.348839 [doi].
21. Barfod L, Bernasconi NL, Dahlback M, Jarrossay D, Andersen PH et al. (2007) Human pregnancy-associated malaria-specific B cells target polymorphic, conformational epitopes in VAR2CSA. *Mol Microbiol* 63: 335–347. MMI5503 [pii];10.1111/j.1365-2958.2006.05503.x [doi].
22. Tutterow YL, Avril M, Singh K, Long CA, Leke RJ et al. (2012) High levels of antibodies to multiple domains and strains of VAR2CSA correlate with the absence of placental malaria in Cameroonian women living in an area of high Plasmodium falciparum transmission 19. *Infect Immun* 80: 1479–1490. IAI00071-12 [pii];10.1128/IAI.00071-12 [doi].
23. Muyldermans S (2001) Single domain camel antibodies: current status. *J Biotechnol* 74: 277–302. S1389-0352(01)00021-6 [pii].
24. De Genst EJ, Guillemins T, Wellens J, O'Day EM, Waudby CA et al. (2010) Structure and properties of a complex of alpha-synuclein and a single-domain camelid antibody 26. *J Mol Biol* 402: 326–343. S0022-2836(10)00742-4 [pii];10.1016/j.jmb.2010.07.001 [doi].
25. Stijlemans B, Conrath K, Cortez-Retamozo V, Van XH, Wyns L, et al. (2004) Efficient targeting of conserved cryptic epitopes of infectious agents by single domain antibodies. African trypanosomes as paradigm. *J Biol Chem* 279: 1256–1261. 10.1074/jbc.M307341200 [doi];M307341200 [pii].
26. Muyldermans S, Baral TN, Retamozzo VC, De BP, De GE, et al. (2009) Camelid immunoglobulins and nanobody technology 106. *Vet Immunol Immunopathol* 128: 178–183. S0165-2427(08)00401-7 [pii];10.1016/j.vetimm.2008.10.299 [doi].
27. Saerens D, Conrath K, Govaert J, Muyldermans S (2008) Disulfide bond introduction for general stabilization of immunoglobulin heavy-chain variable domains 59. *J Mol Biol* 377: 478–488. S0022-2836(08)00040-5 [pii];10.1016/j.jmb.2008.01.022 [doi].
28. Harmsen MM, de Haard HJ (2007) Properties, production, and applications of camelid single-domain antibody fragments. *Appl Microbiol Biotechnol* 77: 13–22. 10.1007/s00253-007-1142-2 [doi].
29. Barfod L, Dobrilovic T, Magistrado P, Khunrae P, Viwami F et al. (2010) Chondroitin sulfate A-adhering Plasmodium falciparum-infected erythrocytes express functionally important antibody epitopes shared by multiple variants. *J Immunol* 185: 7553–7561. jimmunol.1002390 [pii];10.4049/jimmunol.1002390 [doi].
30. Minaeian S, Rahbarizadeh F, Zarkesh-Esfahani SH, Ahmadvand D, Broom OJ (2012) Neutralization of human papillomavirus by specific nanobodies against major capsid protein L1 98. *J Microbiol Biotechnol* 22: 721–728. JMB022-05-20 [pii].
31. Muyldermans S, Atarhouch T, Saldanha J, Barbosa JA, Hamers R (1994) Sequence and structure of VH domain from naturally occurring camel heavy chain immunoglobulins lacking light chains. *Protein Eng* 7: 1129–1135.
32. Wesolowski J, Alzogaray V, Reyelt J, Unger M, Juarez K et al. (2009) Single domain antibodies: promising experimental and therapeutic tools in infection and immunity 6. *Med Microbiol Immunol* 198: 157–174. 10.1007/s00430-009-0116-7 [doi].
33. Khunrae P, Dahlback M, Nielsen MA, Andersen G, Ditlev SB et al. (2010) Full-length recombinant Plasmodium falciparum VAR2CSA binds specifically to CSPG and induces potent parasite adhesion-blocking antibodies. *J Mol Biol* 397: 826–834. S0022-2836(10)00086-0 [pii];10.1016/j.jmb.2010.01.040 [doi].
34. Conrath KE, Lauwereys M, Galleni M, Matagne A, Frere JM et al. (2001) Beta-lactamase inhibitors derived from single-domain antibody fragments elicited in the camelidae 73. *Antimicrob Agents Chemother* 45: 2807–2812. 10.1128/AAC.45.10.2807-2812.2001 [doi].
35. Hmila I, Abdallah RB, Saerens D, Benlasar Z, Conrath K et al. (2008) VHH, bivalent domains and chimeric Heavy chain-only antibodies with high neutralizing efficacy for scorpion toxin AahI^r 15. *Mol Immunol* 45: 3847–3856. S0161-5890(08)00161-2 [pii];10.1016/j.molimm.2008.04.011 [doi].
36. Stijlemans B, Caljon G, Natesan SK, Saerens D, Conrath K et al. (2011) High affinity nanobodies against the Trypanosome brucei VSG are potent trypanolytic agents that block endocytosis 30. *PLoS Pathog* 7: e1002072. 10.1371/journal.ppat.1002072 [doi];10-PLPA-RA-4101 [pii].
37. Nielsen MA, Resende M, Alifrangis M, Turner L, Hviid L et al. (2007) Plasmodium falciparum: VAR2CSA expressed during pregnancy-associated malaria is partially resistant to proteolytic cleavage by trypsin 103. *Exp Parasitol* 117: 1–8. S0014-4894(07)00076-8 [pii];10.1016/j.exppara.2007.03.002 [doi].