

extraluminal extension of the tumor^(6,12). The prognosis is poor, with mean survival time of approximately one year and a half after symptoms onset⁽⁸⁾. Due to pulmonary artery occlusion and acute symptoms, surgical resection is generally the treatment of choice⁽⁸⁾.

In conclusion, the present case reinforces the important role of the imaging methods in the differentiation between pulmonary artery intimal sarcoma and chronic PTE. The relevant aspects for this differentiation, such as contrast enhancement, distention of the affect vessels and extraluminal extension, allow for a correct diagnosis, avoiding delay in the required surgical approach.

REFERENCES

1. Yamanari MGI, Mansur MCD, Kay FU, et al. Bullet embolism of pulmonary artery: a case report. *Radiol Bras.* 2014;47:128–30.
2. Agnollitto PM, Barreto ARF, Barbieri RFP, et al. Rendu-Osler-Weber syndrome: what radiologists should know. Literature review and three cases report. *Radiol Bras.* 2013;46:168–72.
3. Yamada AM, Melo ALKO, Lopes GP, et al. Bilateral breast swelling secondary to superior vena cava obstruction and subclavian vein thrombosis. *Radiol Bras.* 2013;46:252–4.
4. Daud DF, Campos MMF, Fleury Neto LAP. Cardiac tamponade in an infant during contrast infusion through central venous catheter for chest computed tomography. *Radiol Bras.* 2013;46:385–6.
5. Eifer DA, Arsego FV, Torres FS. Unilateral pulmonary veins atresia: evaluation by computed tomography. *Radiol Bras.* 2013;46:376–8.

6. Chong S, Kim TS, Kim BT, et al. Pulmonary artery sarcoma mimicking pulmonary thromboembolism: integrated FDG PET/CT. *AJR Am J Roentgenol.* 2007;188:1691–3.
7. Grosse C, Grosse A. CT findings in diseases associated with pulmonary hypertension: a current review. *Radiographics.* 2010;30:1753–77.
8. Wong HH, Gounaris I, McCormack A, et al. Presentation and management of pulmonary artery sarcoma. *Clin Sarcoma Res.* 2015;5:3.
9. Dornas AP, Campos FT, Rezende CJ, et al. Intimal sarcoma of the pulmonary artery: a differential diagnosis of chronic pulmonary thromboembolism. *J Bras Pneumol.* 2009;35:814–8.
10. Cheng HM, Chou ASB, Chiang KH, et al. Serial CT findings of pulmonary artery intimal sarcoma in 4 months: a case report. *Chin J Radiol.* 2009;34:35–8.
11. Wittram C, Maher MM, Yoo AJ, et al. CT angiography of pulmonary embolism: diagnostic criteria and causes of misdiagnosis. *Radiographics.* 2004;24:1219–38.
12. Yi CA, Lee KS, Choe YH, et al. Computed tomography in pulmonary artery sarcoma: distinguishing features from pulmonary embolic disease. *J Comput Assist Tomogr.* 2004;28:34–9.

Marianna Nunes Batista¹, Miriam Menna Barreto¹, Renata Fukamati Cavaguti¹, Gláucia Zanetti¹, Edson Marchiori¹

1. Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, RJ, Brazil. Mailing Address: Dr. Edson Marchiori. Rua Thomaz Cameron, 438, Valparaíso. Petrópolis, RJ, Brazil, 25685-120. E-mail: edmarchiori@gmail.com.

<http://dx.doi.org/10.1590/0100-3984.2015.0046>

Central nervous system involvement in sarcoidosis

Envolvimento do sistema nervoso central na sarcoidose

Dear Editor,

A 51-year-old female patient complained of mild frontotemporal headache of insidious-onset for two years. One year ago, she had an episode of focal, tonic-clonic seizures (with right lower limb paresthesia) and was prescribed carbamazepine. Cerebrospinal fluid demonstrated increased protein levels and intrathecal immunoglobulin (IgG) synthesis, suggesting an inflammatory component. Magnetic resonance imaging was performed (Figure 1).

Sarcoidosis is a multisystem disease of unknown etiology characterized by noncaseating granulomatous inflammation⁽¹⁾. There is a genetic predisposition, with T-lymphocyte receptor

activation by some unknown antigen. The disease affects preferentially the respiratory system⁽¹⁾. In the lungs, granulomas are observed in the interstitial compartment, showing a perilymphatic distribution along the peribronchovascular sheaths, interlobular septa and pleural surface⁽¹⁾.

It is estimated that in about 5% to 15% of cases sarcoidosis affects the central nervous system. Rarely the patient presents with exclusively neurological manifestations like in the present case. Most commonly, neurosarcoidosis is observed in cases of disseminated disease⁽²⁾.

The clinical manifestations of neurosarcoidosis are pleomorphic. Cranial nerve compromise, visual alterations, headache, weakness, paresis, paresthesia, psychiatric alterations and signs of meningeal irritation may be observed. Although rare, symp-




Figure 1. A: Coronal magnetic resonance imaging – T2-weighted sequence demonstrating diffuse pachymeningeal thickening most prominent at the high convexity and extending bilaterally toward the falx, with predominance of hypointensity in association with reduction in volume and hypersignal of the left hippocampus (mesial sclerosis). **B:** Paramagnetic contrast-enhanced axial T1-weighted sequence showing diffuse and homogeneous pachymeningeal enhancement.




Figure 2. A: Biopsy of thickened area of the dura mater showing numerous noncaseating epithelioid and giant cell granulomas and predominantly lymphocytic inflammatory infiltrate intermingled with dense collagenous fibrosis. **B:** Masson's trichrome stain: granulomas (pink) against dense collagenous connective tissue (blue).

toms of diabetes insipidus such as polydipsia and polyuria may also occur due to the involvement of the hypothalamus and hypophysis. In cases of spinal cord involvement, weakness of lower limbs and other nonspecific signs of myelopathy are observed⁽³⁾.

Although sarcoidosis may manifest in all the regions of the central nervous system, it is most commonly seen in the skull base, hypothalamus, pituitary and optic chiasm⁽⁴⁾. At magnetic resonance imaging, a common finding is intraparenchymal lesions with hypersignal on T2-weighted and FLAIR sequences, generally multifocal, periventricular, subcortical or in the deep white matter. Such findings can hardly be differentiated from vasculitis or demyelinating diseases. Intraparenchymal lesions are generally located near the areas with leptomeningeal involvement (with enhancement by paramagnetic contrast medium), and may be either single or multiple, possibly also involving cranial nerves⁽⁴⁾.

Like in the present case, diffuse pachymeningeal thickening may be observed, with hyposignal on T2-weighted, isosignal on T1-weighted sequences and contrast enhancement. Thus, differential diagnoses such as neurotuberculosis, dural lymphoma, meningioma en plaque, IgG4 deposition disease, pseudotumor, adenocarcinoma metastasis, Wegener's granulomatosis, idiopathic hypertrophic pachymeningitis might be considered, requiring biopsy to define the etiology. Simultaneous dural and leptomeningeal involvement is rarely observed⁽⁴⁾. In the present case, the anatomopathological findings corresponded to typical noncaseating granulomas in the pachymeninges (Figure 2). Intracranial hypotension is another differential diagnosis to be considered, gen-

erally presenting with diffuse pachymeningeal thickening, but with hypersignal on T2-weighted sequences (in the present case, hyposignal was observed on T2-weighted sequences).

A consensus is still to be reached on the treatment for sarcoidosis. In cases where the patient is symptomatic the treatment is initiated with high doses of corticosteroids, gradually reduced along the treatment up to complete withdrawal⁽³⁾.

REFERENCES

1. Melo ASA, Marchiori E, Capone D. Tomographic and pathological findings in pulmonary sarcoidosis. *Radiol Bras.* 2011;44:220–4.
2. Spencer TS, Campellone JV, Maldonado I, et al. Clinical and magnetic resonance imaging manifestations of neurosarcoidosis. *Semin Arthritis Rheum.* 2005;34:649–61.
3. Nozaki K, Judson MA. Neurosarcoidosis: clinical manifestations, diagnosis and treatment. *Presse Med.* 2012;41(6 Pt 2):e331–48.
4. Christoforidis GA, Spickler EM, Recio MV, et al. MR of CNS sarcoidosis: correlation of imaging features to clinical symptoms and response to treatment. *AJNR Am J Neuroradiol.* 1999;20:655–69.

Vinicius Silles Machado¹, Nivaldo Adolfo Silva Junior¹, Luciano Souza Queiroz¹, Fabiano Reis¹, Danilo dos Santos Silva¹, Flavia Fagundes Bueno¹, Ana Carolina Coan¹

1. Universidade Estadual de Campinas (Unicamp), Campinas, SP, Brazil. Mailing Address: Dr. Fabiano Reis. Faculdade de Ciências Médicas – Universidade Estadual de Campinas, Departamento de Radiologia. Rua Tessália Vieira de Camargo, 126, Cidade Universitária Zeferino Vaz. Caixa Postal: 6111. Campinas, SP, Brazil, 13083-887. E-mail: fabianoreis2@gmail.com.

<http://dx.doi.org/10.1590/0100-3984.2014.0113>

Femoral artery injury during aneurysm coiling

Lesão da artéria femoral durante embolização de um aneurisma

Dear Editor,

Endovascular artery reconstruction with low-profile stents, flow-diverters and flow-disrupting devices represent a significant progress in the endovascular therapy of intracranial aneurysms. Despite the improvement in technical expertise and developments in device technology, endovascular treatment still has inherent risks⁽¹⁾. In the literature, most reports are focused on neurological complications during procedures⁽²⁾, however, reports on access vessel complications are scarce. Some of the well known access-related complications include: arterial pseudoaneurysms, arterio-

venous fistulae, hematomas, arterial dissection leading to acute vessel occlusion^(3,4), intracavitary bleeding, and retroperitoneal hematoma following femoral artery puncture⁽⁵⁾. The authors report the case of a large groin hematoma caused by a hypodermic needle connected with the black cable of the detachable coil power supply (Boston Scientific; Natick, MA, USA) and its endovascular management.

Local compression is the first line treatment for femoral access complications⁽⁶⁾, but such strategy may fail when indicated for patients under combined antiplatelet and anticoagulation regimens. Open surgery is effective in the treatment of groin complications⁽⁷⁾. However, the endovascular approach is a safe and effective minimally invasive alternative to surgery in the management