



جامعة نايف العربية للعلوم الأمنية  
Naif Arab University for Security Sciences

Naif Arab University for Security Sciences  
Arab Journal of Forensic Sciences & Forensic Medicine

www.nauss.edu.sa  
http://ajfsfm.nauss.edu.sa



الجمعية العربية للعلوم الجنائية والطب الشرعي  
Arab Society for forensic Sciences and forensic Medicine

# Lead and Cadmium Toxicity in Tile Manufacturing Workers in Assiut, Egypt

Ragaa M Abd Elmaaboud<sup>1\*</sup>, Zaghoul T. Mohamed<sup>1</sup>

Safaa M. George<sup>1</sup>, Azaa M. Ez El-dine<sup>2</sup>, Doaa M. El Shehaby<sup>1</sup>.

<sup>1</sup> Forensic Medicine and Clinical Toxicology Department, Faculty of Medicine, Assiut University, Egypt.

<sup>2</sup> Clinical Pathology Department, Faculty of Medicine, Assiut University, Egypt.

Open Access



Original Article

## Abstract

Occupational lead and cadmium exposure are important health issues in developing countries. This study aimed to detect toxic metal contents in raw materials used to make tiles and to assess exposure health impacts on workers. The study sample consisted of 74 tile workers, having a mean age of 35.2 years, in the Industrial City of Arab El Awamer, Assiut (Egypt). Elemental analysis of the raw materials was performed by using scanning electron microscopy. The data collection questionnaire was divided into two parts; the first included demographic data, symptoms attributed to toxic elements and possible sources of exposure to metals. The second part was designated to assess heavy metal exposure health impacts through clinical examination and biological

investigations. Many toxic elements were identified in the raw materials used to make tiles, and the most abundant were lead and cadmium. Analysis of the clinical data revealed that 66% of the workers suffered from headache, constipation (8%), abdominal colic (33.8%) and 30% suffered from a variety of respiratory problems such as dyspnea (60%), cough (13%) and chest tightness (27%). Fifty percent of the workers complained of weak grip, 33.8% of foot drop, and 54% had tremors. Burton's line in gums was present in 28% of workers and 28.2% were diagnosed with constrictive lung diseases. Of the 74 workers, 90.5% showed toxic lead levels and 80% had toxic cadmium levels. 10.8% had abnormal alpha glutathione levels with a positive strong linear correlation between lead and cadmium levels and years of work. It is mandatory to develop and implement measures to prevent these hazardous exposure effects among tile industry workers.

**Keywords:** Lead, Cadmium, Occupational hazard, Toxicity, Tiles Manufacturing, Assiut.

\* Corresponding Author: Ragaa M. Abd Elmaaboud  
Email: ragaamohabd@yahoo.com

1658-6794© 2016 AJFSFM. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial License.

doi: 10.12816/0026462

التسمم بالرصاص والكاديوم لدى عمال صناعة الأجر

(البلاط) بمدينة أسيوط في مصر

المستخلص

يُعد التعرض المهني للرصاص والكاديوم قضية صحية هامة وخاصة في البلدان النامية. وحيث تعد صناعة الأجر إحدى مصادر التعرض لهذه العناصر، فقد هدفت هذه الدراسة إلى تحديد ما



Production and hosting by NAUSS

تحتويه المواد المستخدمة في صناعة الأجر (البلاط) من العناصر السامة ومدى تأثيرها على صحة العاملين. شملت الدراسة أربعة وسبعين عاملاً في ستة أماكن لتصنيع الأجر بالمدينة الصناعية بعرب العوامر بأسسوط، حيث أجريت التحاليل لتحديد العناصر في عينات المواد الخام المستخدمة في التصنيع بواسطة مجهر المسح الإلكتروني، وتم دراسة استبيان تضمن أسئلة حول المعلومات الديموغرافية للعاملين وأسئلة عن الأعراض التي يمكن أن تُعزى إلى التعرض للعناصر السامة، كما أجريت بعض الفحوص السريرية وأشعة سينية على الصدر وبعض التحاليل على عينات دم وبول مأخوذة من العاملين. وقد أثبتت نتائج تحليل المواد الخام المستخدمة في صناعة الأجر أنها تحتوي على العديد من العناصر الثقيلة السامة وأن النسبة الأكبر كانت لعنصري الرصاص والكاديوم. وأظهرت الفحوص السريرية للعاملين أن (66%) منهم يشكون من الصداع، و(8%) يعانون من الإمساك مع تغير في لون البراز، وحوالي ثلثهم (33.8%) يعانون من أعراض المغص المتكرر. وحوالي (30%) من العاملين لديهم مشاكل في الجهاز التنفسي؛ عانى (60%) منهم من ضيق تنفس. كما بين الاستبيان أن (50%) من العاملين يشكون من ضعف في قبضة اليد، وثلثهم (33%) كانوا يشكون من هبوط القدم. وأن (60%) من العاملين لديهم ظلال داكنة تحت العينين، كما وجد الخط الأزرق في اللثة لدى (28%) من العمال، وظهرت أعراض التليف الرئوي لدى (27.8%) من الذين أجروا اختبار وظائف التنفس. وأنه من أصل (39) حالة وافقت على فحص الصدر بالأشعة السينية (11) حالة ما يمثل (28.2%) كان لديهم علامات التليف الرئوي. وأن (90.5%) من العاملين لديهم مستوى رصاص سام ومرتفع بالدم، و(80%) من العمال كانوا يعانون من مستوى سام ومرتفع للكاديوم. و(10.8%) كان لديهم مستوى غير طبيعي من ألفا-جلوتاثيون-اس-ترانسفيريز (alphaGST) في البول، وأن مستوى الرصاص السام في الدم يزيد بزيادة عدد سنوات التعرض، ووجود علاقة ذات دلالة إحصائية بين مستوى الرصاص في الدم والأعراض العصبية عند العاملين. وإن نتائج هذا البحث تستوجب وضع وتفعيل ضوابط لحماية العاملين بصناعة الأجر لمنع ما يتعرضون له من مخاطر صحية.

الكلمات المفتاحية: الرصاص، الكاديوم، السموم المهنية، صناعة الأجر (البلاط)، أسسوط.

## 1. Introduction

Tile are one of the most widely used materials in construction [1]. Toxic products of the tile production process are mixtures of predominantly clays, dust, vapor and other natural occurring minerals that have been mixed with water and fired in a high temperature [2]. In the tile industry, a certain amount of coloring agents are added to the clay and mixed in a ball mill. Exposure to these heavy metals in the tile production process occurs mainly during mixing, storing, handling and other activities where large amounts of respirable dust particles are generated [3].

Occupational lead exposure in many developing countries is entirely unregulated, often with no monitoring of exposure. Although there are numerous small-scale industries which use lead-based raw materials that may pose health risks to workers, there are no workplace regulations for monitoring lead exposure among these workers [4] and no such data are available with the labour departments or any other related authority.

Cadmium is a naturally occurring heavy metal and is an occupational hazard within a variety of industrial settings. The workplace is the most common source of cadmium exposure, occurring mainly through inhalation of dust and fumes [5]. Cadmium principally affects the lung, kidney, liver, and testes following acute intoxication, which leads to nephrotoxicity due to prolonged exposure [6]. Cadmium is a potent cell poison and which causes oxidative stress by increasing lipid peroxidation and/or by changing intracellular glutathione levels. Inhalation exposure can result in acute hepatic and renal injury, while intake of cadmium-contaminated food causes acute gastrointestinal effects, such as nausea, vomiting and diarrhea [7].

This study aimed to do an elemental analysis of raw materials used in the manufacturing of tiles, detect different heavy metals present in their raw materials and to determine the toxicological health hazards among tile manufacturing workers that may have resulted from heavy metal exposure during the tile manufacturing process.

## 2. Subjects and Methods

This is a community based, cross sectional study including all workers (total coverage technique, 74 workers





**Figure 1-** Direct contact with raw materials used in tile manufacture process



**Figure 2-** Direct contact during cutting and glazing in tile manufacture process



**Figure 3-** Glazing waste

working in about 6 tile manufacturing factories) fulfilling the predetermined inclusion criteria. A written informed consent was obtained from each participating worker.

### 2.1 Inclusion criteria

- Adult male (more than 18 years old).
- Working in direct contact with raw materials used in the tile manufacture process (cutting, glazing as shown in Figure 1-3).
- Working at least 8 hours daily for not less than one year in this industry.
- Not known to be diseased at the time of joining the tile factory.

A structured questionnaire was used for data collection and was divided into two parts. The first part included questions about demographic data of workers, all possible symptoms that may be attributed to exposure to the previously mentioned toxic elements, general health status and all possible sources of exposure to heavy metals. The second part of the questionnaire consisted of a clinical data sheet which includes signs of poor health and a clinical examination for suspected toxicological health hazards such as pallor, blue line in the gums, chest findings, abdominal signs, and peripheral neuropathy.

Blood samples were collected from all workers (5 mL) and tested for complete blood count, blood lead and cadmium levels, blood urea, serum creatinine, and urine Alpha Glutathione S- transferase level by Glory Science Co. ELISA kits (2400 Veterans Blvd. Suite 16-101, Del Rio, TX, United

**Table 1-** Elemental analysis of raw materials used in tile manufacture

Element	Material 1(%)	Material 2 (%)
Mn	0.42	0.46
Fe	1.44	5.95
Ni	10.75	12.35
Cu	16.86	20.28
Zn	2.73	0.62
Cd	34.39	25.09
Pb	36.58	35.24

**Table 2-** Neurological complaints among tile manufacturing workers

Complaints	No. (%)	
Weak grip strength	37 (50.0)	
Degree of grip weakness:	Mild	14 (37.8)
	Moderate	20 (54.1)
	Severe	3 (8.1)
Trend of grip weakness:	Increase with time	35 (94.6)
	Fixed	2 (5.4)
Foot drop	25 (33.8)	
Tremors	40(54.0)	
Numbness	17 (23.1)	
Headache	49 (66.2)	

State). Chest X-rays and pulmonary function tests were also performed.

## 2.2 Method of detection of lead and cadmium

1 mL of whole blood was collected in a vacutainer prothrombin tube and then acid digested using nitric acid (5 mL of HNO<sub>3</sub> to 1 mL blood) then put on a hot plate for approximately 30 min until the volume of the sample reduced to 1 mL [8]. Analysis of blood lead and cadmium levels was performed using Buck Model 210 VGP Atomic Absorption Spectrophotometer in the Central Laboratory of the Faculty of Science, Assiut University.

Samples of two different raw materials consisting of waste product resulting from glazing process (Material 1) and raw material (Material 2) were collected in 500 g quantities and subjected to elemental analysis by Dual

JSM-5400 Scanning Electron Microscopy equipped with Oxford Link ISIS 300 EDX system at Assiut University, Egypt.

## 2.3 Statistical analysis

SPSS software package version 16 was used to analyse the data. Descriptive statistics were done in the form of frequencies, mean and SD. Analytic statistics were done as chi-square, independent sample t-test and pearson correlation. ANOVA Test was used and values were considered significant when *p* value was <0.05.

## 3. Results

Samples of different raw materials were analyzed by scanning electron microscopy (Table-1). The SEM analysis showed that there were many toxic elements in these

**Table 3-** Clinical signs among tile manufacturing workers

Clinical Signs	No.	%
Pallor of face and hands	27	36.5
Darkness under the eyes	45	60.8
Blue line on the gums	46	62.2
Chest wheezes and rhonchi	19	25.7
Abdominal tenderness	4	5.4



**Table 4-** Blood and urine analysis of the studied cases

Parameter	No.	%	(Mean $\pm$ SD)
Blood lead level ( $\mu\text{g/dL}$ )	7	9.5	75.1 $\pm$ 46.1
Accepted (up to 20)	67	90.5	
Toxic (more than 20)			
Blood cadmium level ( $\mu\text{g/dL}$ )	15	20.3	18.2 $\pm$ 24.4
Accepted (less than 5)	59	79.7	
Toxic (5 or more)			
Hb(mg/dL)	12	25.5	14.8 $\pm$ 2.1
Normal (14 and more )	35	74.5	
Anaemic (less than 14)			
Urinary alpha-glutathione S- Transferase(IU/L)	66	89.2	57.3 $\pm$ 3.3
Normal (22-50)	8	10.8	
Abnormal (more than 50)			
Blood urea (mg/dL)	47	100.0	26.2 $\pm$ 3.7
Normal (20 – 40)	0	00.0	
Elevated (more than 40)			
Serum creatinine (mg/dL)	47	100.0	0.78 $\pm$ 0.2
Normal (0.6 - 1.2)	0	00.0	
Elevated (more than 1.2)			

materials such as cadmium, lead, copper, arsenic, iron, zinc and traces of other elements. The most abundant toxic elements were lead and cadmium; lead represented more than one third of the raw material (36.58% and 35.24%, respectively) followed by cadmium (34.39% and 25.09%, respectively), while other elements (Mn, Cu, Ni, and Zn) collectively represented less than one third of these raw materials.

Toxic effects of lead on workers' CNS are shown in Table-2. Our results revealed that 66.2% of workers complained of headache and 50% of workers complained of weak grip that was moderate in 54.1%, mild in 37.8% and severe in 8.1%. More than half of workers (54%) had tremors and about one quarter of them (23.1%) had numbness in upper and lower extremities.

Table-3 shows clinical signs observed among tile man-

ufacturing workers. A majority of the workers (62.2%) reported blue line on the gums (Burton line) followed by darkness under the eyes (60.8%), pallor on face and hands (36.5%), chest wheezes and rhonchi (25.7%) and abdominal tenderness (5.4%) respectively.

Results of laboratory investigations are shown in Table-4 which indicates that 90.5% of the workers had toxic blood lead levels (BLL) with a mean  $\pm$  SD concentration of 75.1  $\mu\text{g/dL}$ . Regarding urine alpha-glutathione S-transferase levels, it was found that 8 cases (10.8%) had abnormal levels while the majority of workers (89.2%) had normal levels. Blood urea and serum creatinine levels were determined in 47/74 workers and all of them were within the normal reference range of 20-40 mg/dL and 0.6-1.2 mg/dL, respectively. After testing the urine Alpha glutathione in workers, it was found that 8 cases (10.8%) had abnormal



**Table 5-** Relationship between blood lead level and systolic/diastolic blood pressure in studied cases

Grade of blood pressure (mmHg)	No.	Mean blood Lead level ( $\mu\text{g/dL}$ )	<i>p</i> value
Systolic blood pressure			
• Low	3	69.1	0.007**
• Normal	47	73.9	
• High	24	78.1	
Diastolic blood pressure			
• Normal	27	70.6	0.034*
• High	47	82.8	

\*Statistically significant, \*\*statistically highly significant

levels while a great majority of workers (89.2%) had normal levels.

Table-5 shows the relationship between blood pressure and blood lead levels (BLL) among 74 workers. Our results revealed a statistically significant relation between BLL and both systolic and diastolic blood pressure as there is a linear trend of increase in BLL from low to normal to higher grade of blood pressure (Low 90/60 or lower, normal between 90/60 and 139/89, and high 140/90 or higher) with a statistically significant difference between the three grades ( $p = 0.007$ ). The same findings were observed in diastolic blood pressure ( $p = 0.034$ ), as those suffering from high blood pressure reported higher blood lead levels than

normotensive workers.

Results show a positive, highly statistically significant moderate correlation between BLL and years of work ( $p = 0.002$ ,  $r = 0.33$ ) (Table-6). There was a negative, moderately significant correlation between weight ( $p = 0.001$ ,  $r = -0.53$ ) and BLL. Hemoglobin levels showed a negative, moderate statistically significant correlation with blood lead level ( $r = -0.41$  and  $p = 0.023$ ). No statistically significant correlation was seen between blood cadmium level and years of work, blood urea, serum creatinine and alpha-glutathione S-transferase level.

The relation between BLL and cadmium levels and years of work is shown in Table-7, which indicates a linear

**Table 6-** Correlation between blood lead and cadmium with some clinical parameters

Parameters	<i>r</i>	<i>p</i> value	Sig.	
Years of work	0.33	0.002	**	
Weight	-0.53	0.001	**	
Body mass index	-0.66	0.008	**	
Lead ( $\mu\text{g/dL}$ ) *	Hemoglobin	-0.41	0.023	*
	Blood urea	0.12	0.28	NS
	Serum creatinine	0.07	0.63	NS
	Alpha-glutathione S-transferase (IU/L)	0.07	0.583	NS
Years of work	0.16	0.91	NS	
Cadmium ( $\mu\text{g/dL}$ )	Lead ( $\mu\text{g/dL}$ )	0.63	0.003	**
	Blood urea	0.04	0.78	NS
	Serum creatinine	0.05	0.54	NS
	Alpha-glutathione S-transferase (IU/L)	0.08	0.16	NS

\* Statistically significant \*\* Statistically highly significant, NS: Statistically not significant



**Table 7-** Relationship between blood lead and cadmium levels with years of work among tile manufacture workers

Years of Work	No. of workers	Mean $\pm$ SD Blood lead ( $\mu\text{g/dL}$ )	Mean $\pm$ SD Blood cadmium ( $\mu\text{g/dL}$ )
1 to 5 years	14	51.1 $\pm$ 6.3	9.2 $\pm$ 1.3
6 to 10 years	13	52.9 $\pm$ 7.2	16.7 $\pm$ 4.2
11 to 15 years	17	68.7 $\pm$ 4.4	18.5 $\pm$ 3.4
16 to 20 years	23	86.3 $\pm$ 5.4	20.6 $\pm$ 2.4
21 to 25 years	7	91.1 $\pm$ 8.6	38.1 $\pm$ 4.6
F value		6.3	3.8
p value		0.009**	0.029*

\*Statistically significant, \*\*statistically highly significant

**Table 8-** Relationship between blood cadmium level and chest X- ray findings in our studied cases

Chest X- rays	Toxic cadmium level $\geq 5 \mu\text{g/dL}$		Normal cadmium level $< 5 \mu\text{g/dL}$		Total	
	No.	%	No.	%	No.	%
Normal	4	33.3	24	88.9	28	71.8
Abnormal Findings	8	66.7	3	11.1	11	28.2
Total	12	30.8	27	69.2	39	100.0
p value				0.02*		
Chi-square				5.12		

\*Statistically significant

association between intoxication and years of work in the tile manufacture industry. The highest mean $\pm$ SD concentration of blood lead and cadmium levels were present in the workers who had worked from 16 to 20 years.

Regarding the toxic effects of heavy metal (cadmium) exposure on the respiratory system, a respiratory function test was performed. Table-8 shows a statistically significant relationship between blood cadmium level and chest x-rays in the studied workers ( $p = 0.02$ ). It was found that 12 out of 39 workers who underwent chest x rays were suffering from toxic blood cadmium levels (equal to or more than  $5 \mu\text{g/dL}$ ). Among these workers, two thirds had abnormal

findings in their x-rays compared to one third who showed normal x-ray findings.

Table-9 shows logistic multivariate regression analysis model that was highly predictive of major symptoms that may have resulted from exposure to high blood lead levels in the studied sample. The symptoms in order were: tremors (odds ratio = 14.03), Weak grip (odds ratio = 11.1), presence of foot drop (odds ratio = 10.4), occurrence of numbness (odds ratio = 9.68), presence of pallor (odds ratio = 7.52), presence of blue line in the gums (odds ratio = 5.42), high systolic blood pressure (odds ratio = 3.07), high diastolic blood pressure (odds ratio = 2.2) and occurrence



**Table 9-** *Multivariate regression analysis for the most important risk factors of high blood lead levels among tile manufacturing workers*

Variables	B	S.E	Wald	df	Sig.	Odds ratio
<b>Tremors</b>	3.334	1.120	8.854	1	0.003	14.03
<b>Weak grip strength</b>	7.79	60.45	0.017	1	0.002	11.132
<b>Foot drop</b>	2.56	0.989	6.89	1	0.009	10.43
<b>Numbness</b>	2.04	0.97	4.423	1	0.003	9.684
<b>Presence of pallor in face and hands</b>	1.709	0.984	3.12	1	0.008	7.522
<b>Blue line on the gums</b>	1.69	1.43	1.39	1	0.023	5.421
<b>High systolic blood pressure</b>	1.62	0.894	3.304	1	0.06	3.07
<b>High diastolic blood pressure</b>	0.79	0.95	0.69	1	0.04	2.20
<b>Headache</b>	0.79	0.95	0.69	1	0.08	1.20
<b>Constant</b>	-3.17	1.943	2.66	1	0.01	0.042

of headache (odds ratio = 1.2). Foot drop reflects the affection of lead toxicity on the motor function of the workers and may interfere with the performance and productivity of such workers as well as the psychological disturbance that may have resulted from this complaint.

#### 4. Discussion

Occupational heavy metal toxicity is still a health problem in many countries such as Egypt and Turkey [9]. Although there are numerous industries which use lead-based raw materials that may pose health risks to workers, there are no standardized workplace regulations to avoid lead and cadmium exposure. Moreover, there are no sufficient studies carried out on blood lead and cadmium levels in workers or on the contribution of common workplace practices to lead poisoning [10]. Occupational lead and cadmium exposure with or without symptoms has not been sufficiently investigated in Egyptian workers, in particular, workers who are involved in glazing traditional tiles [11].

In the present study, samples of different raw materials used in tile manufacturing were analyzed. As shown in Table-1, there were many toxic elements in these materials

such as cadmium, lead, copper, arsenic, iron zinc and traces of other elements. The two most abundant toxic elements were lead and cadmium. This study revealed the same low standard work conditions as described by Hisham et al. Tile workers did not wear any kind of protective clothes or any protective respiratory devices during their job [12]. Regarding toxic effects of lead on CNS, results in Table-2 revealed that 66% of workers complained from headache. This may be explained by the fact that lead increases the permeability of the blood-brain barrier and causes disruption, allowing large molecules such as albumin to enter the brain which increase osmotic pressure and causes ions and water to follow, resulting in edema of the brain and an increase in intracranial pressure [13].

Neurological complaints of workers that may be related to lead exposure are described in Table-2. Once lead enters the body, it interferes with normal cell function and physiological processes. The nervous system seems to be the most sensitive system to lead poisoning [13].

Weak grip strength increased with time in about 95% of the workers, which may point to the cumulative effect of lead toxicity in those workers; with an increase in work-



ing time there is a concomitant increase in their complaint of weakness. Regarding other neurological complaints, our results showed that about one third of workers (33.8%) suffered from foot drop. In addition, more than half of workers (54%) suffered from tremors and about one quarter of them (23%) had numbness. These findings are consistent with many other studies that documented the effects of lead on the peripheral nervous system, especially motor nerves [14].

Bellinger (2004) investigated the behavioral and neurobiological mechanisms of the functional deficits observed in lead-exposed humans [15]. He stated that lead causes the axons of nerve cells to degenerate and lose their myelin coats. It was found that in adults, manifestations of lead toxicity may appear at blood lead levels greater than 80  $\mu\text{g}/\text{dL}$  [16]. Signs that occur in adults at blood lead levels exceeding 100  $\mu\text{g}/\text{dL}$  include wrist and foot drop, signs of encephalopathy, such as delirium, coma, seizures, and headache. It is rare to be asymptomatic if blood lead levels exceed 100  $\mu\text{g}/\text{dL}$  [17].

Pallor on face and hands, a clinical sign that may reflect lead and cadmium exposure, was reported in approximately one third of cases (Table-3). This may point to anemia and peripheral vasoconstriction. Blue line in the gum was found in 62.2% of workers. This bluish to black edging of the teeth, known as Burton line, is an indication of chronic lead poisoning [18]. This line is caused by a reaction between circulating lead with sulphur ions released by oral bacterial activity, which deposits lead sulphide at the junction of the teeth and gums [19]. This symptom is an important sign in the detection and diagnosis of chronic lead toxicity; however, it is not always present in all cases and usually related to chronic oral lead exposure. Our results are in agreement with Mood et al. who studied lead toxicity in traditional tile workers in Mashhad in Iran, 2010, and reported that blue line was present in 64.8% of the studied cases [20].

Table-3 also shows that one quarter of the workers (25%) had chest related complaints in the form of chest wheezes and ronchi. Only 4 workers (5%) showed abdominal symptoms in the form of abdominal tenderness. This was lower than reported by Mood et al. who documented

abdominal tenderness in 15.1% of their studied sample [20]. Five of our workers (27.8%) showed symptoms that suggested obstructive lung disease while the remaining workers were normal. The table also shows that out of 39 cases that agreed to undergo chest X-rays, 11 cases showed symptoms suggestive of obstructive lung disease, which may be attributed to occupational exposure to cadmium and partially to occupational silicosis. These findings are constant with Sorahan et al. who studied 624 workers employed in a cadmium alloy factory and reported that exposure to cadmium oxide fumes increases risks of chronic non-malignant diseases of the respiratory system [21].

Lead poisoning may be acute (from intense exposure of short duration) or chronic (from repetitive low-level exposure over a prolonged period), but the latter is much more common [22]. Classically, "adult lead poisoning" or "lead intoxication" has been defined as exposure to high levels of lead typically associated with severe health effects when BLL is more than 20  $\mu\text{g}/\text{dL}$  [14]. Our results showed that 90.5% of the workers had toxic BLL, exceeding 20  $\mu\text{g}/\text{dL}$  with a mean of 75.10  $\mu\text{g}/\text{dL}$  (Table-4). This value is greater than reported in the study by Mood et al. in which workers had a mean toxic BLL of 57.80  $\mu\text{g}/\text{dL}$ , while only 7 cases (9.5%) showed acceptable BLL levels [9]. Our results are partially consistent with Rashid et al. who studied the effect of lead on hematological parameters in 51 occupationally exposed individuals comprising 27 lead furnace workers, 24 lead pellet handlers and 20 healthy and service-matched controls [18]. The lead workers had the highest blood lead levels (71.20  $\mu\text{g}/\text{dL}$ ) compared to healthy subjects (29.80  $\mu\text{g}/\text{dL}$ ), while workers handling pellets had moderately increased blood lead levels (45.50  $\mu\text{g}/\text{dL}$ ). The furnace exposed workers had higher blood lead levels compared to the lead pellet handlers [21-22].

Table-4 also shows that more than three quarters of the workers suffered from anemia (74.5%), which is in accordance with the findings of Rashid et al who concluded that chronic lead exposure causes normocytic normochromic (NC/NC) anemia and demonstrated a dose-response relationship between lead levels and severity of anemia [22]. Anemia may appear at blood lead levels higher than 50  $\mu\text{g}/\text{dL}$ . This depends upon the amount of lead in blood and tis-



sues, as well as the time course of exposure.

After testing the urine Alpha glutathione of workers, it was found that 8 cases (10.8%) had abnormal levels, while a great majority of workers (89.2%) had normal levels (Table- 4). Our results are consistent with a cross-sectional study done by Garcon et al. (2007) that aimed to evaluate the usefulness of a set of early biological markers of oxidative stress of nephrotoxicity for the bio-monitoring of workers occupationally exposed to lead [25]. They found that mean levels of Pb in blood and urine were also  $387.10 \pm 99.10 \mu\text{g/L}$  and  $217.7 \pm 117.7 \mu\text{g/g}$  creatinine, respectively. The mean levels of Cd in blood and urine were  $3.26 \pm 2.11 \mu\text{g/L}$  and  $2.51 \pm 1.89 \mu\text{g/g}$  creatinine, respectively, suggesting relatively low occupational exposure levels [25]. Alpha-glutathione S-transferase and urinary protein levels were reported between the two groups and were closely correlated with Pb and/or Cd exposure levels. These results suggest the use of alpha-glutathione S-transferase excretion in urine as a hallmark of early changes in the proximal tubular integrity. Hence, successful prevention of renal diseases induced by occupational exposure to lead and/or cadmium largely relies on the capability to detect nephrotoxic effects at a stage when they are still reversible or at least not yet compromising renal function.

Table-4 also shows that out of the 74 workers, blood urea & serum creatinine tests were done in 47 cases and levels were normal in all workers. This is explained by the fact that kidney damage occurs with exposure to high levels of lead, which were not reported in the study (40-60  $\mu\text{g/dL}$ ). Long-term exposure to toxic levels of lead cause nephropathy that may lead to Fanconi syndrome, in which the proximal tubular function of the kidney is impaired [24]. Long-term exposure to lead levels lower than those which cause lead nephropathy have also been reported as nephrotoxic in patients from developed countries that had chronic kidney disease or were at risk because of hypertension [25].

Numerous studies have indicated a relationship between moderate or heavy lead exposure and high blood pressure [26-27]. The present results have shown that low-levels of lead exposure are related to changes in blood pressure (Table-5). The same findings were also noted in dia-

stolic blood pressure ( $p = 0.034$ ), as those suffering from high blood pressure reported higher blood lead levels than normotensive workers. These results are in agreement with Pocock et al. (1988) who studied the relationship between blood lead concentration and blood pressure in a survey of 7371 men from 24 British towns and found a statistically significant positive association [29].

The present results also support findings of an earlier report by Harlan (1988), who studied the American population and analyzed data from the National Health and Nutrition Examination Survey and revealed significant correlations between blood lead levels and blood pressure [27]. He also stated that blood lead levels were significantly higher in groups with high diastolic blood pressure (greater than 90 mm Hg). These findings and those from other studies [28] confirm the relationship between blood lead levels and blood pressure at relatively low levels that are commonly observed in the general population; however, the strength and importance of this relationship require further studies through epidemiologic and metabolic investigations.

Regarding cadmium levels, about 80% of the workers reported toxic levels of cadmium compared to one fifth of workers who showed normal levels. As shown in table - 4, the mean level of Cd was  $18.2 \pm 24.4 \mu\text{g/dL}$ , which was higher than that reported by Garcon et al (2007) [25].

The correlation studies in table-6 showed a positive, highly significant moderate correlation between blood lead level and years of work ( $p = 0.002$ ,  $r = 0.33$ ). This important finding explains the cumulative toxicity of occupational lead exposure in workers that may have a negative impact on their health status and a bad impact on their work performance. The weight of workers is usually an indication of their health status, and malnutrition and being underweight is usually associated with many health problems. Table-6 shows a negative, statistically moderate, significant correlation between workers weight and blood lead levels ( $p = 0.001$ ,  $r = -0.53$ ). This negative correlation may be attributed to the toxic effects of lead on the gastro-intestinal tract in the form of nausea and loss of appetite, together with reduction in the protein content in workers' meals. In the same table, hemoglobin level shows a negative, moderate, statistically significant correlation with blood lead level ( $r$



= - 0.41 and  $p = 0.023$ ). Also, there was a strong positive, highly significant linear correlation between blood lead level and blood cadmium ( $r = 0.63$  &  $p = 0.003$ ) in our studied workers. This finding was expected in our study because both lead and cadmium were the most abundant elements in the raw materials that the workers were exposed to (Table-1). Also, these are the elements which are usually associated with many health hazards among workers in the tile industry.

In the present study, no statistically significant correlations were found between blood cadmium and years of work, blood urea, serum creatinine and alpha-glutathione levels. These findings may be related to the fact that kidney function tests that were used in the present study are usually elevated in late phases of kidney infection and are usually normal in early renal diseases [29]. Despite this, alpha-glutathione was more sensitive than serum creatinine and blood urea to detect early kidney problems [30]. Alpha-glutathione was elevated in about 11% of workers as shown in table 4. It is, therefore, highly recommended to do screening for nephrotoxic effects of cadmium by alpha-glutathione routinely and regularly rather than to perform tests for usual kidney functions (blood urea and serum creatinine) [31].

Table-8 shows a statistically significant relationship between blood cadmium level and chest x-rays in the studied workers ( $p = 0.02$ ). It was found that 12 out of 39 workers who underwent chest x rays were suffering from toxic blood cadmium levels (equal to or more than  $5 \mu\text{g/dL}$ ). Among these workers, two thirds had abnormal findings in their x-rays compared to one third who showed normal x-ray findings. The present results agree with other reports which showed that glazing and enameling of tiles can lead to metal poisoning and lung diseases, which can lower workers' productivity [31-32].

Table-9 shows logistic multivariate regression analysis model that was highly predictive of major symptoms that may have resulted from exposure to high blood lead levels in the studied sample. The symptoms in order were: tremors (odds ratio = 14.03), weak grip (odds ratio = 11.1), presence of foot drop (odds ratio = 10.4), occurrence of numbness (odds ratio = 9.68), presence of pallor (odds ra-

tio = 7.52), presence of blue line in the gums (odds ratio = 5.42), high systolic blood pressure (odds ratio = 3.07), high diastolic blood pressure (odds ratio = 2.2) and occurrence of headache (odds ratio = 1.2). Foot drop reflects the affection of lead toxicity on the motor function of the workers and may interfere with the performance and productivity of such workers beside the psychological upset that may have resulted from this complaint [33].

## 5. Conclusion

Occupational exposure to heavy metals, especially lead and cadmium, should be viewed as a very serious health risk to the workers engaged in tile manufacturing. Workers should be informed about the multiple health hazards and complications that may occur from their persistent exposure to toxic raw materials used in the manufacturing of tiles. In order to reduce the exposure to lead, cadmium and other toxic substances, simple protective gear should be adopted and strictly observed to safeguard the health of workers.

## References

1. Maria D B, Elena D, Antonio G, Francisco JC, Julio. Environmental Performance of Ceramic Tiles: Improvement Proposals. *Materials and Designs* 2010; 31: 35-41.
2. Richard R. Environmental Health and Safety Department. Emergency Assistance: Material Safety Data Sheet. 2012; 214: 309-429.
3. Fateme K, Seyed J Shaht A, Farideh G, Abolfazle B, Abbas R. Measurement of Urinary Cadmium in Glazers Using Solid Phase Extraction Followed by Inductively Coupled Plasma Atomic Emission Spectroscopy. *Int J Occup Hyg* 2012; 2: 63-68.
4. Yalemsew A, Argaw A, Dejene AT. Occupational lead exposure among automotive garage workers – a case study for Jimma town, Ethiopia. *Occup Med Toxicology* 2012; 7: 15.
5. Agency for toxic substances and disease registry. Toxicological profile for cadmium. U.S Department of health And Human Services Public Health Service Agency for Toxic Substances and Disease Registry



- 2008
6. Patra R, Rautray A, Swarup D. Oxidative Stress in Lead and Cadmium Toxicity and Its Amelioration. *Vet Med Int* 2011; 47-52.
  7. Nordberg G. *Handbook on the Toxicology of Metals*, Academic Pr., Burlington, 2007.
  8. Noish.: *Manual of Analytical Method (NMAM)*. Fourth Edition(1994).
  9. Kayihan P, Alpaslan T, Erdinc O, Hamdi A. Occupational lead exposure: Blood lead levels of apprentices in Bursa, Turkey. *Industrial Health* 2009; 47: 97–102.
  10. World Health Organization (WHO) Childhood lead poisoning. WHO Library Cataloguing-in-Publication Data, 2010.
  11. Ceramic Factory Outlet Copyright (2010): CANBIS (Canadian Building & Interior Services) Newmarket, Ontario, Canada (905) 898-899.
  12. Hisham M. A., Safia B. A., Inas A. S. Respiratory Hazards Among Egyptian Ceramics Workers. *Environmental and Occupational Researcher* 2010; 2: 65-73.
  13. Patel A. How Does Lead affects the Nervous System? *Biology* 202. 2000 Second Web Report 07-Jan-2002; 14:24.
  14. Karri S K, Saper RB, Kales S N. Lead encephalopathy due to traditional medicines. *Curr Drug Saf* 2008; 3:54–59.
  15. Bellinger D C Supplement Article: Lead. *J Paed* 2004; 113: 1016-1022.
  16. Grant L D. "Lead and compounds". In Lippmann, M. *Environmental Toxicants: Human Exposures and Their Health Effects*, 3rd edition. Wiley-Inter science 2009.
  17. Merrill J, Morton J, Soileau S, Hayes A. *Principles and Methods of Toxicology* (5th ed.), CRC Press, 2007; 4:102–253.
  18. Rambousek A J. The symptoms and treatment of industrial poisoning". *Industrial Poisoning from Fumes, Gases, and Poisons of Manufacturing Processes* 2008.
  19. Schwartz S, Stephenson B, Sarkar D, Freyholtz H, Ruth G. Quantitative assay of erythrocyte "free" and zinc-protoporphyrin: clinical and genetic studies. *Int J Biochem* 1980; 12: 1053.
  20. Mood B, Shademanfar S, Rastegar M, Afshari R, Namaei G, Allah N H. Occupational lead poisoning in workers of traditional tile factories in Mashhad, North-east of Iran. *Int J Occup Environ Med* 2010; 1: 29-38.
  21. Sorahan T, Lister A, Gilthorpe M, Harrington J. Mortality of copper cadmium alloy workers with special reference to lung cancer and non-malignant diseases of the respiratory system, 1946-92. *Occup Environ Med* 1995; 52: 804-812
  22. Rashid B, Khurshid U, Dilshad A, Muhammad S, Iftikhar A. Blood Lead Levels and Anemia in Lead Exposed Workers. *JPMA* 1995; 65: 66-68.
  23. Guidotti T and Ragain L. Protecting children from toxic exposure: three strategies. *Pediatr Clin North Am* 2007; 54: 227-235.
  24. Agency for Toxic Substances and Disease Registry (ATSDR). *Case Studies in Environmental Medicine (CSEM): Lead Toxicity*. ATSDR Publication No: ATSDR-HE-CS-2001-2002. Atlanta, 2007GA: Agency for Toxic Substances and Disease Registry.
  25. Garçon G, Leleu B, Marez T, Zerimech F, Haguenoer JM, Furon D, Shirali P. Biomonitoring of the adverse effects induced by the chronic exposure to lead and cadmium on kidney function: usefulness of alpha-glutathione S-transferase. *Sci Total Environ* 2007; 377: 165-172.
  26. Missouni F, Slimani M, Aoues A. Toxic effect of lead on kidney function in rat Wistar. *African J Biochem Res* 2010; 4: 21-27.
  27. Harlan W R. The relationship of blood lead levels to blood pressure in the U.S. population. *Environ Health Perspect* 1988; 78: 9–13.
  28. Kopp S J, Barron J T, Tow J P. Cardiovascular actions of lead and relationship to hypertension: a review. *Environ Health Perspect* 1988;78: 91-99.
  29. Pocock S, Shaper A, Ashby D, Delves H, Clayton. The relationship between blood lead, blood pressure, stroke, and heart attacks in middle-aged British men. *Environ Health Perspect* 1988; 78: 23–30.
  30. Tiwari S, Tripathi I, Tiwari H. Lead Poisoning - A Review. *Res J Chem Sci* 2013; 3: 86-88.
  31. Kosnett J M. Health effects of low dose lead exposure in adults and children and preventable risk posed by



- consumption of game meat harvested with lead ammunition. University of Colorado, Denver/o 1630 Welton, Suite 2005.
32. Shadick NA, Kim R, Weiss S, Liang MH, Sparrow D, Hu H. Effect of low level lead exposure on hyperuricemia and gout among middle aged and elderly men: the Normative Aging Study. *J Rheumatol* 2000; 27:1708–1712.
33. Environmental Guidelines for Small-Scale Activities in Africa (EGSSAA): Cleaner Production Fact Sheet and Resource Guide. Chapter 4.1: Brick and Tile Production, 2009, [www.encapafrika.org](http://www.encapafrika.org).

