

Case report

A Pediatric case of Bickerstaff's Brainstem Encephalitis

Ju Yi Park, MD, Kyong Ok Ko, MD, Jae Woo Lim, MD, Eun Jeong Cheon, MD, Jung Min Yoon,
MD, Hyo Jeong Kim, MD

Department of Pediatrics, Konyang University College of Medicine, Daejeon, Korea

Corresponding author

Hyo Jeong Kim, M.D.

Department of Pediatrics, Konyang University College of Medicine,

685 Gasuwon-dong, Seo-gu, Daejeon, Korea

Tel: 82-42-600-9230, Fax: 82-42-600-9090

E-mail: greatelena@naver.com

Abstract

Bickerstaff's brainstem encephalitis is characterized by ophthalmoplegia, ataxia, and disturbance of consciousness. It is similar to Miller Fisher syndrome, a variant of Guillain-Barre syndrome, because they share some features such as ophthalmoplegia and ataxia. The difference is that patients with Bickerstaff's brainstem encephalitis have impaired consciousness and patients with Miller Fisher syndrome have alert consciousness and areflexia. Here, we report the case of a 3-year-old child who was diagnosed with Bickerstaff's brainstem encephalitis presenting typical clinical features and interesting radiological findings. The patient showed ophthalmoplegia, ataxia, and subsequent stuporous mentality. Brain magnetic resonance imaging revealed high signal intensity in the pons and cerebellum around the 4th ventricle on a T2-weighted image. He was successfully treated with intravenous immunoglobulin. Differentiation of Bickerstaff's brainstem encephalitis and Miller Fisher syndrome is often difficult because they share many overlapping features. Brain magnetic resonance imaging could help diagnose Bickerstaff's brainstem encephalitis, especially when definite lesions are found.

Key words: Encephalitis, Miller Fisher syndrome, ophthalmoplegia, ataxia

Introduction

Bickerstaff's brainstem encephalitis was defined as ophthalmoplegia, ataxia, and impaired consciousness¹⁾. Bickerstaff's brainstem encephalitis is in the same spectrum with Miller Fisher syndrome and Guillain-Barre syndrome. They are autoimmune, postinfectious diseases and present common clinical features such as ophthalmoplegia and ataxia²⁾. Furthermore, the etiology is regarded as similar, because they commonly show prodromal upper respiratory infection, CSF albuminocytological dissociation, and serum IgG antibody to ganglioside GQ1b^{3),4)}.

While Guillain-Barre syndrome and Miller Fisher syndrome are peripheral nervous system (PNS) diseases, Bickerstaff's brainstem encephalitis is described as a central nervous system (CNS) disease. However, the nosologic classification of Bickerstaff's brainstem encephalitis, Guillain-Barre syndrome and Miller Fisher syndrome remains uncertain. Many patients with overlap between these conditions have been reported and there is variable CNS and PNS involvement in the spectrum of these diseases⁴⁻⁶⁾.

Here, we present the case of a 3-year-old child with typical features of Bickerstaff's brainstem encephalitis, and magnetic resonance evidence of pons and cerebellar involvement therein.

Case Report

A 3-year-old boy was hospitalized because of high fever, headache, vomiting, and unsteady gait for 1 day. He had flu-like symptoms for 3 days. There was no history of recent travelling, exposure to toxic materials, or head injury. He had scheduled vaccination, and had previously been a healthy child. On examination, he showed an alert mental state and, except for a high fever, vital signs within normal limits. His pharynx was injected but tonsils were not enlarged.

He showed alert consciousness. His both pupils were normal and responsive to light but he showed bilateral ptosis. Except cranial nerve III, other cranial nerve function test was founded normal. On motor examination, muscle power and muscle tone of both upper and lower extremities showed normal findings. Sensory function was also normal. But, he couldn't stand upright and showed staggered gait besides, finger-to-nose did comparatively well. He represented hyperreflexia, and revealed bilateral Babinski sign. Neck stiffness or other meningeal irritation sign was not observed.

White blood cell (WBC) count in blood was 13,000 cells/ μ L. Cerebrospinal fluid (CSF) study revealed a WBC count of 68 cells/ μ L, a protein level of 75 mg/dL, and a glucose level of 87 mg/dL (blood glucose was 83 mg/dL). Enterovirus was not isolated in either stool or CSF samples. Serological tests of herpes simplex virus (HSV), Epstein-Barr virus (EBV), varicella-zoster virus (VZV), and cytomegalovirus (CMV) showed negative results. A CSF culture was also negative.

We empirically administered ceftriaxone, acyclovir, dexamethasone, and mannitol to him. On the third hospital day, he showed stuporous mentality, hyperventilation and persistent fever. Furthermore, he represented sustained ptosis, upward gaze disability, slurred speech and truncal ataxia. He began to be treated in the intensive care unit (ICU).

Non-contrast brain magnetic resonance imaging (MRI) showed high signal intensity in the pons and cerebellum around the 4th ventricle on a T2-weighted image (Fig. 1). Electroencephalography

(EEG) showed high amplitude slow wave activities without epileptiform discharges, suggesting encephalopathy. He was administrated immunoglobulin 1 gm/kg/day for 2 days and methylprednisolone pulse therapy. After these 2 days of treatment with immunoglobulin, he showed alert mentality and improvement of persistent downward gaze and truncal ataxia. On the 9th hospital day, he was transferred to the general ward under maintaining steroid regimen. Two weeks later, we confirmed improvement of the previously detected lesions on brain MRI (Fig. 2) and clinical symptoms including decreased mentality, ataxia and ptosis were fully recovered.

On the basis of the ophthalmoplegia-ataxia-disturbance of consciousness triad with radiological evidence of CNS involvement, the diagnosis of Bickerstaff's brainstem encephalitis was made.

Korean JJP Accepted diatrics

Discussion

Diagnostic criteria of Bickerstaff's brainstem encephalitis were described as follows^{4,7)}. "Progressive, relatively symmetric external ophthalmoplegia and ataxia by 4 weeks" and "disturbance of consciousness or hyperreflexia" are both required¹¹⁾. The following diseases must also be excluded: vascular disease involving the brain stem, Wernicke's encephalopathy, botulism, myasthenia gravis, brain stem tumor, pituitary apoplexy, acute disseminated encephalomyelitis, multiple sclerosis, neuro-Behcet's disease, vasculitis, lymphoma, and Creutzfeldt-Jakob disease⁴⁾. In our case, the initial presentation of ataxia and ophthalmoplegia suggested the diagnosis of Miller Fisher syndrome, but brisk deep tendon reflexes and subsequent presentation of a disturbance of consciousness suggested the diagnosis of Bickerstaff's brainstem encephalitis rather than Miller Fisher syndrome. Furthermore, definite lesions on brain MRI could explain CNS involvement. In our case, the diagnosis of Bickerstaff's brainstem encephalitis was possible because of the neurological manifestations.

However, in many cases, differentiation of Bickerstaff's brainstem encephalitis and Miller Fisher syndrome is difficult because not all patients with Bickerstaff's brainstem encephalitis present abnormal brain MRI, abnormal EEG, or hyperreflexia. According to a previous report, 11% of 47 patients with Bickerstaff's brainstem encephalitis had abnormalities on brain MRI, whereas 57% of 30 patients with Bickerstaff's brainstem encephalitis had abnormalities on EEG⁴⁾. Another study on 37 patients with Bickerstaff's brainstem encephalitis reported abnormalities at 23% and 50% for brain MRI and EEG, respectively⁸⁾. Thus, brain MRI could not detect abnormalities in more than two-third of the patients with Bickerstaff's brainstem encephalitis. There was, however, a case report of Bickerstaff's brainstem encephalitis with normal brain MRI in which the evidence of cerebellar involvement was noted by statistical parametric mapping (SPM) analysis using positron emission tomography (PET)⁹⁾. As most cases of Bickerstaff's brainstem encephalitis show no

abnormal lesions on brain MRI, functional imaging tools such as PET could be useful to document CNS involvement.

In fact, brain MRI findings in Bickerstaff's brainstem encephalitis have rarely been reported. Usually the findings are hyperintensity on T2-weighted images of the pons, medulla, thalamus, or cerebellum^{4,7)}. Typical lesions are noted in the brainstem and a few cases showed cerebellar lesions^{4,10)}. In our presented case, pons and cerebellar lesions could explain ophthalmoplegia, hyperreflexia, altered consciousness, and ataxia.

As mentioned above, Miller Fisher syndrome and Bickerstaff's brainstem encephalitis are supposed to have a common pathogenesis. Anti-GQ1b antibodies are commonly found in both, but more frequently in Miller Fisher syndrome⁴⁾. In Miller Fisher syndrome, the anti-GQ1b antibody binds to GQ1b, expressed on the ocular nerve and primary sensory neuron; conversely, in Bickerstaff's brainstem encephalitis, it binds to GQ1b expressed on the brainstem reticular formation^{4,11)}. Therefore, the current consensus is that the two diseases are in fact part of a spectrum of autoimmune disease variably involving the PNS and CNS.

Nerve conduction study was not performed because of poor compliance; however, the patient showed no sensory loss or limb weakness despite showing ataxia. No limb weakness, but the presence of hyperreflexia and definite brain MRI lesions in the pons and cerebellum, suggested CNS rather than PNS involvement.

Unfortunately, in this case, a study of anti-GQ1b antibodies was not performed. In fact, according to the epidemiological study in Japan, only 56% of the patients with Bickerstaff's brainstem encephalitis showed less than 5/mm³ of CSF cell count and 20% showed more than 50/mm³ of CSF cell count⁸⁾. Moreover, dramatic response to the immunoglobulin treatment supported the evidence of Fisher-Bickerstaff spectrum disease than infective disease.

Bickerstaff's brainstem encephalitis is a rare disorder. Epidemiologic study in Korea has not been

done until now. Only a few cases including a pediatric case who presented typical clinical features with positive anti-GM1 antibodies and several cases in adults with typical clinical symptoms but variety of laboratory findings were reported¹²⁻¹⁵⁾. However, according to the Japanese epidemiological study, Bickerstaff's brainstem encephalitis accounted for 43% of brainstem encephalitis and anti-GQ1b antibodies were present in 75% of the patients with Bickerstaff's brainstem encephalitis⁸⁾.

In conclusion, we report the case of Bickerstaff's brainstem encephalitis which is rare in childhood. The patient presented typical clinical manifestations of ataxia, ophthalmoplegia and impaired consciousness and MRI findings of definite lesions in the pons and cerebellum. Among the brain stem encephalitis, Bickerstaff's brainstem encephalitis accounts for major proportion. In cases with typical clinical manifestations, Bickerstaff's brainstem encephalitis should be considered and Anti-GQ1b antibodies may be useful in diagnosis of Bickerstaff's brainstem encephalitis.

Conflict of interest

No potential conflict of interest relevant to this article was reported.


Korean JJP Accepted diatrics

References

1. Bickerstaff E, Cloake P. Mesencephalitis and rhombencephalitis. *Br Med J* 1951;2:77-81.
2. Fargas A, Roig M, Vazquez E, Fito A. Brainstem involvement in a child with ophthalmoplegia, ataxia, areflexia syndrome. *Pediatr neurol* 1998;18:73-5.
3. Rebecca C, Gologorsky BA, Jerome A, Barakos, Farhad S. Rhomb- and Bickerstaff Encephalitis: Two Clinical Phenotypes? *Pediatr neurol* 2013;244-8.
4. Ito M, Kuwabara S, Odaka M, Misawa S, Koga M, Hirata K et al. Bickerstaff's brainstem encephalitis and Fisher syndrome form a continuous spectrum: clinical analysis of 581 cases. *J Neurol* 2008;255:674-82.
5. Yuki N. Fisher syndrome and Bickerstaff brainstem encephalitis (Fisher-Bickerstaff syndrome). *J Neuroimmunol* 2009;215:1-9.
6. Tsapis M, Laugel V, Koob M, de Saint Martin A, Fischbach M. A pediatric case of Fisher-Bickerstaff spectrum. *Pediatr Neurol* 2010;42:147-50.
7. Odaka M, Yuki N, Yamada M, Koga M, Takemi T, Hirata K et al. Bickerstaff's brainstem encephalitis: clinical features of 62 cases and a subgroup associated with Guillain-Barre syndrome. *Brain* 2003;126:2279-90.
8. Koga M, Kusunoki S, Kaida K, Uehara R, Nakamura Y, Kohriyama T et al. Nationwide survey of patients in Japan with Bickerstaff's brainstem encephalitis: epidemiological and clinical characteristics. *J Neurol Neurosurg Psychiatry* 2012;83:1210-5.
9. Kwon HM, Hong YH, Sung JJ, Paeng JC, Lee DS, Lee KW. A case of Bickerstaff's brainstem encephalitis; the evidence of cerebellum involvement by SPM analysis using PET. *Clin Neurol Neurosurg* 2006;108:418-20.
10. Mondéjar RR, Santos JM, Villalba EF. MRI findings in a remitting-relapsing case of Bickerstaff encephalitis. *Neuroradiology* 2002;44:411-4.

11. Chiba A, Kusunoki S, Obata H, Machinami R, Kanazawa I. Serum anti-GQ1b IgG antibody is associated with ophthalmoplegia in Miller Fisher syndrome and Guillain-Barre syndrome: clinical and immunohistochemical studies. *Neurology* 1993;43:1911-7.
12. Kim JY, Kim YO, Son YG, Woo YJ. A case of Bickerstaff's brainstem encephalitis in childhood. *Korean J Pediatr* 2010;53:607-11.
13. Kang BG, Lim JG, Kim KD, Sohn SI, Jung DK, Cho YW et al. A Case of Recurrent Bickerstaff's Brainstem Encephalitis with an Overlap of Guillain-Barre Syndrome. *J Korean Neurol Assoc* 2002;20:561-3.
14. Lee JY, Park KC, Lee IH, Park CI, Lee MS, Lee SS. Bickerstaff's brain stem encephalitis after a motor vehicle crash. *J Trauma*. 2004;56:179-81.
15. Kwon HM, Hong YH, Sung JJ, Paeng JC, Lee DS, Lee KW. A case of Bickerstaff's brainstem encephalitis; the evidence of cerebellum involvement by SPM analysis using PET. *Clin Neurol Neurosurg*. 2006;108:418-20.

Fig.1. T2-weighted brain MRI showed hyperintensity in the pons (A) and cerebellum (B).



Korean JJP Accepted diatriCS

Fig.2. Brain MRI performed after 2 weeks showed only a focal T2 hyperintense lesion in the pons (A), with improvement of the previous lesion (B).






Fig.1



KJP Accepted

Fig.2



KJP Accepted