

The Effect of Lavender Aromatherapy on the Pain Severity of Primary Dysmenorrhea: A Triple-blind Randomized Clinical Trial

Nikjou R^{1,2}, Kazemzadeh R¹, Rostamnegad M¹, Moshfegi S³, Karimollahi M³, Salehi H^{1,2}

¹Midwifery Department, School of Nursing and Midwifery, Ardabil University of Medical Sciences, ²Student Research Centre, Ardabil University of Medical Sciences, ³Nursing Department, School of Nursing and Midwifery, Ardabil University of Medical Sciences, Ardabil, Iran

Address for correspondence:

Dr. Karimollahi M,
Nursing Department, School of
Nursing and Midwifery, Ardabil
University of Medical Sciences,
Shohad Headway, Ardabil, Iran.
E-mail: m.karimollahi@arums.ac.ir

Abstract

Background: Primary dysmenorrhea is the most common complaint in adolescents and adult young women that disturbs their daily life performance. **Aim:** The current study investigated the effect of lavender aromatherapy on pain severity in primary dysmenorrhea. **Subjects and Methods:** This triple-blind randomized clinical trial was conducted on 200 students of Ardabil University of Medical Sciences, Iran. Subjects were allocated randomly into intervention (lavender) and control (placebo) groups. The researcher assistant asked the participants to smell the lavender in the first 3 days of menstruation, 30 min in a day in two menstrual cycles. Control group was also administered placebo (diluted milk) to be used as lavender in treatment group. Pain severity was scored by visual analog scale in the first 3 days of menstruation before intervention and 2 months after intervention. Data were analyzed through descriptive statistics and independent and paired samples *t*-tests. **Results:** There was a significant difference in average pain severity between treatment and control groups after intervention. However, students in the treatment group reported significantly less pain severity 2 months after intervention ($P < 0.01$). **Conclusion:** Using lavender aromatherapy for 2 months may be effective in decreasing the pain severity of primary dysmenorrhea.

Keywords: Aromatherapy, Lavender, Menstruation, Pain, Primary dysmenorrhea

Introduction

Primary dysmenorrhea is the most common gynecologic complaint in adolescent and young women, and it is linked to the biological cycle of ovulation.^[1,2] About 75% of women experience primary dysmenorrhea during their fertility period.^[3] Berk reported the prevalence of primary dysmenorrhea as 60%,^[4]

It has been estimated that 7–15% of women suffer from severe pain which causes disturbances in their performance for 1–3 days in a month,^[5] and 14–52% of female students

miss school because of complaining dysmenorrhea.^[6] This issue is associated with decline in career and educational function and results in disturbed quality of life and emerging economic problems.^[7] Various reasons have been proposed for primary dysmenorrhea such as progesterone reduction in the final stages of luteal phase that causes lissome rupture and subsequent release of A2 phospholipase of the endometrium. Origin of increasing phospholipase is prostaglandin that causes retraction of uterine muscles vessels which finally result in ischemia and pain.^[8] However, mild physical

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Nikjou R, Kazemzadeh R, Rostamnegad M, Moshfegi S, Karimollahi M, Salehi H. The effect of lavender aromatherapy on the pain severity of primary dysmenorrhea: A triple-blind randomized clinical trial. *Ann Med Health Sci Res* 2016;6:211-5.

Access this article online

Quick Response Code:



Website: www.amhsr.org

DOI:
10.4103/amhsr.amhsr_527_14

activities, proper diet, relaxation, massage, biofeedback techniques, some Yoga exercises, contraceptive pills, and nonsteroidal anti-inflammatory drugs such as mefenamic acid and ibuprofen are recommended for decreasing the pain of primary dysmenorrhea.^[9,10] Since most of the drugs utilized for controlling the pain of dysmenorrhea have various side-effects and also are expensive, most of the nonpharmacological methods are in the category of complementary medicine for this disorder. The use of nonpharmacological methods such as aromatherapy, massage, acupuncture, and acupressure are also having been popular in the recent years.^[11]

Traditionally, aromatherapy has been used as one of the nonpharmacological methods for reducing the symptoms of dysmenorrhea, releasing uterine cramps, and decreasing the pain and anxiety after childbirth.^[12]

However, a common complementary drug which has been used in aromatherapy is lavender that belongs to the Mediterranean region and it is found in Africa and India.^[13] Lavender is just a little herbaceous plant with narrow and long leaves, covered by white cottony fluff and purple flowers and is in the form of a spike. Its essence is made from distilling flowers from blossoming plants, and its muscle relaxant and antispasmodic effects have been confirmed by many studies.^[14,15] Some studies suggest lavender and its main extract such as linalool acetate and linalool had topical analgesic effects on laboratory animals,^[16] which is made by increasing regional perfusion.^[17] Results of studies supporting aromatherapy suggest that essential oils stimulate the receptors in the olfactory bulb and transfer the message of olfaction to limbic system and cause releasing endorphin, enkephalin, and serotonin, which results in the sense of relaxation and stress reduction.^[18]

About the prevalence of the primary dysmenorrhea and the side effects of the current treatments and because of lack of similar studies in this field, we decided to perform a study on the female students in dormitories of Ardabil University of Medical Sciences with the aim of determining the effect of lavender aromatherapy on the pain severity in primary dysmenorrhea.

Subjects and Methods

This study, a triple-blinded randomized clinical trial, was done on 200 female students who had primary dysmenorrhea settled in the dormitories of Ardabil University of Medical Sciences in 2013. According to the population of female students at the medical university dormitories that 420 of them had dysmenorrhea, The number of samples is determined using a formula sample size was estimated 200 people. The inclusion criteria were being single, age range 20–30-year-old, onset of menstrual pain before the age of 20, having regular menstruation cycles with 21–35 days interval, menstrual bleeding without clots disposal (low and medium bleeding), onset and duration of menstrual pain several hours before

bleeding to maximum 5th day of menstruation, and having dysmenorrhea.

Whereas exclusion criteria were very severe and intolerable dysmenorrhea that limits activities and not controllable by medicine, onset of menstruation pain after 20-years-old, having chronic disease (diabetes, hyperthyroidism, and hypothyroidism), previous allergy to medicinal plants, pelvic inflammatory disease and pelvic mass, gynecologic surgery (cyst, tumor, and pelvic adhesions), unwillingness to participate in the study, and occurrence of any allergies during study.

Sampling has been conducted by the cluster sampling method in female dormitories of Ardabil University of Medical Sciences, but placement in groups was done by a simple random method through randomized blocks of four and six. After listing all possible four and six blockings and assigning their numbers, using True Random Number Generator software, Randomness and Integrity Services Limited, Ireland. Several blocks were selected till reaching to 200 subjects. There was no significant difference between two groups according to their demographic properties.

After approving this project by the research deputy of the Ardabil University of Medical Sciences, the researcher assistant started sampling in the dormitories and explaining the goals and research method and stressing on the autonomy of subjects for surrender from research in every stage of the study and confidentiality of obtained information, the students interested in taking part in the study were invited to sign the consent form to show their agreement for taking part in this study.

Afterward, subjects were embedded in two lavender (A) and diluted milk (B) groups. The researcher assistant expressed instructive guidelines on filling the questionnaire and quantifying the pain severity by using visual analog scale (VAS) to all the subjects. Both groups completed the questionnaire which included demographic characteristics and pain severity in the first cycle of menstruation (nonintervention cycle). After filling questionnaires, Groups A and B were offered a glass of 10 cc lavender essences and 10 cc diluted milk, respectively. Then, the subjects were asked for strew three drops of the solution to a piece of cotton and smell that for the first 3 days of menstruation, daily 1 time for 30 min in two continual cycles and complete the VAS in the first and second menstruation. None of the students had the need for more analgesics. Glasses with a same appearance in number from 1 to 200 were prepared by the pharmacological firm in Tabriz, Iran; and the researcher assistant and the subjects had no information about the content; also the statistical analyzer had no information about study groups. At the end of the sampling, the questionnaires were gathered and descriptive statistics done for calculating frequency, mean, and standard deviation. In addition, independent and paired *t*-tests and Chi-square were utilized to analyze data in SPSS version 21 (IBM, USA).

Results

The age range of subjects was 19–29-year-old, and 80% (160/200) of them were single. The average age of subjects at menarche was 13.0 (1.3) years. Eighty-five percent (170/200) of participants had dysmenorrhea for more than 4 years, 60% (120/200) had never used analgesics, and 76% (152/200) had intermediate bleeding according to their own disclosure.

The mean (and standard deviation) of pain severity in the treatment group has decreased after 1st and 2nd month, which was statistically significant ($P < 0.01$) [Table 1]. As well in the control group, the mean of pain severity has not been reduced in cycle-1, 7.06 (1.43) and cycle-2, 7.04 (1.33) after intervention [Table 1].

The mean difference in pain severity was not statistically significant between the two groups before the intervention (cycle-0), while it was significantly different in the 1st and 2nd month after the intervention ($P = 0.001$ and $P < 0.01$) [Table 2].

Discussion

The findings showed that lavender had a significant effect on primary dysmenorrhea and pain.

Aroma inhalation is effective for mental and physical stability, refreshing, and concentration. In this method, the essential oil is absorbed into the lung and blood through the nose.^[18,19] Lavender is used as a sedative, anti-depressant,^[20] anti-spasmodic,^[19] anti-flatulent,^[21] and also to treat infertility, infection, anxiety, fever, stress,^[21,22] restlessness or colic in infants,^[23] varicose ulcers,^[24] and carpal tunnel syndrome.^[25]

Amiri-Farahani *et al.* referred to the effect of massage with lavender oil and mint essence on decreasing the pain intensity of menstruation which showed a significant difference in pain intensity between groups.^[26] Furthermore, Ou *et al.* in a research in 2012 reported aromatherapy with lavender, clary sage, and marjoram oils in a 2:1:1 ratio reduced pain intensity of dysmenorrhea, which was statistically significant ($P < 0.001$),^[18] whereas aromatherapy with lavender did not decrease the pain of intrauterine device (IUD) placement in the study of Shahnazi *et al.* in 2012.^[27] These controversial findings maybe because of the duration of lavender essence inhalation, which was inhaled for 30 min in

our research, while in the study of Shahnazi *et al.* the time was about 20 min. In addition, the reason of difference between the results may be due to different etiology and mechanism of pain in two studies (IUD and dysmenorrhea).

The study by Kane *et al.* also suggested that lavender essence can decrease the pain scores in changing wound bandage ($P < 0/05$).^[28] Similarly, Han *et al.* compared the pain in the 1st and 2nd days just in one cycle of menstruation, in which the mean pain intensity in the group of aromatherapy with lavender, rose, and salvia was significantly lower than the group undergoing massaging with almond oil and the group without the intervention.^[29] In addition, the study of Kim *et al.* on 100 patients in the Intensive Care Unit suggested massaging with lavender decreases the pain intensity about 50%.^[30]

A study showed that lavender and linalool have antispasmodic effect and increase the local blood flow.^[31] Moreover, Shokohi *et al.* studied the effect of lavender aroma in the pain of cesarean section; the results showed that treatment with lavender aroma has a significant effect on the pain of cesarean section compared to placebo.^[32] In 2007, Kim *et al.* has done study with the title, “effect of treatment with lavender aromatherapy in the postanesthesia care unit on opioid requirements of morbidly obese patients undergoing laparoscopic adjustable gastric banding” that showed significantly more patients in the placebo group required analgesics for postoperative pain (82%) than patients in the lavender group (46%) ($P = 0.007$).^[33]

This study had several limitations, including that there was a possible treatment bias (exceeding the recommended treatment duration) because the participants performed the self-inhalation freely following instructions. We should also acknowledge the limitation in use of the VAS for measuring pain severity. It has been reported that the VAS is difficult to use in clinical practice and has a higher failure rate than other scales including the verbal rating scale and the numerical rating scale.^[34]

Conclusion

Findings showed the positive effect of lavender inhalation in reducing pain severity during menstruation. Because there were no side-effects reported by subjects, aromatherapy can be regarded as a safe and effective treatment for menstrual pain. Further randomized studies, using more objective measures, larger samples, and long-term follow-up are needed to verify

Table 1: The mean (standard deviation) of pain severity in two groups, before and after 1st and 2nd month of treatment

Pain intensity group	Cycle-0	1 st month later	2 nd month later	P^* The difference cycle-0 and other months
Lavender	7.27 (1.24)	5.09 (1.02)	5.04 (1.01)	<0.001
Diluted milk	7.03 (1.04)	7.06 (1.43)	7.04 (1.33)	0.67

*Paired t-test analysis

Table 2: Difference of pain severity between lavender and placebo groups over the study

The time of study	Means difference (SE)	Df	t	P*
Cycle-0	0.06 (0.22)	198	1.923	0.72
1 st month later (cycle-1)	1.57 (1.53)	198	6.23	0.001
2 nd month later (cycle-2)	1.93 (1.87)	198	8.17	<0.001

*Independent t-test. SE: Standard error

self-aromatherapy massage's effects on menstrual pain and anxiety.

Acknowledgment

This project was registered with ethics code number "Arums. ERC.93.49" from the Ethics Committee of Ardabil University of Medical Sciences and RCT code "IRCT201470616252N2" in the clinical trial registering center. We appreciate the research deputy of Ardabil University of Medical Sciences, the members of research council, the ethics committee of the university, and all the colleagues, dormitory workers, and all the participants in this research.

Financial support and sponsorship

This project was funded by the research deputy of Ardabil University of Medical Sciences.

Conflicts of interest

There are no conflicts of interest.

References

- Harel Z. Dysmenorrhea in adolescents and young adults: From pathophysiology to pharmacological treatments and management strategies. *Expert Opin Pharmacother* 2008;9:2661-72.
- Smith CA, Crowther CA, Petrucco O, Beilby J, Dent H. Acupuncture to treat primary dysmenorrhea in women: A randomized controlled trial. *Evid Based Complement Alternat Med* 2011;2011:612464.
- Stoppard M. *Family Health Guide*. 1st ed. London: Penguin Books Ltd.; 2002. p. 247-80.
- Berek JS, Novak E. *Berek and Novak's Gynecology*. Philadelphia Lippincott Williams & Wilkins; 2012. p. 607-11.
- El-Gilany AH, Badawi K, El-Fedawy S. Epidemiology of dysmenorrhoea among adolescent students in Mansoura, Egypt. *East Mediterr Health J* 2005;11:155-63.
- Doty E, Attaran M. Managing primary dysmenorrhea. *J Pediatr Adolesc Gynecol* 2006;19:341-4.
- Dawood MY. Primary dysmenorrhea: Advances in pathogenesis and management. *Obstet Gynecol* 2006;108:428-41.
- Daley A. The role of exercise in the treatment of menstrual disorders: The evidence. *Br J Gen Pract* 2009;59:241-2.
- Decherney AH, Prenoll ML. *Current Obstetrics*. Philadelphia: Lange; 2000. p. 238-9.
- Marjoribanks J, Proctor M, Farquhar C, Derks RS. Nonsteroidal anti-inflammatory drugs for dysmenorrhoea. *Cochrane Database Syst Rev* 2010; Jan 20;(1):CD001751.
- Zahradnik HP, Hanjalic-Beck A, Groth K. Nonsteroidal anti-inflammatory drugs and hormonal contraceptives for pain relief from dysmenorrhea: A review. *Contraception* 2010;81:185-96.
- Tillett J, Ames D. The uses of aromatherapy in women's health. *J Perinat Neonatal Nurs* 2010;24:238-45.
- Mirzaei F, Keshtgar S, Kaviani M, Rajaeifard AR. The effect of lavender essence smelling during labor on cortisol and serotonin plasma levels and anxiety reduction in nulliparous women. *J Kerman Univ Med Sci* 2009;16:245-54. [Persian]
- Nikfarjam M, Parvin N, Asaradegan N. The effect of *Lavandula angustifolia* in the treatment of mild to moderate depression. *Shahrekord Univ Med Sci J* 2010;11:66-73. [Persian]
- Hadi N, Hanid AA. Lavender essence for post-cesarean pain. *Pak J Biol Sci* 2011;14:664-7.
- Efferth T, Greten HJ. Medicinal and aromatic plant research in the 21st century. *J Med Aromatic Plants Res* 2012;1:100-10.
- Sheikhan F, Jahdi F, Khoei EM, Shamsalizadeh N, Sheikhan M, Haghani H. Episiotomy pain relief: Use of lavender oil essence in primiparous Iranian women. *Complement Ther Clin Pract* 2012;18:66-70.
- Ou MC, Hsu TF, Lai AC, Lin YT, Lin CC. Pain relief assessment by aromatic essential oil massage on outpatients with primary dysmenorrhea: A randomized, double-blind clinical trial. *J Obstet Gynaecol Res* 2012;38:817-22.
- Ghelardini C, Galeotti N, Salvatore G, Mazzanti G. Local anaesthetic activity of the essential oil of *Lavandula angustifolia*. *Planta Med* 1999;65:700-3.
- Cornwell S, Dale A. Lavender oil and perineal repair. *Mod Midwife* 1995;5:31-3.
- Seo JY. The effects of aromatherapy on stress and stress responses in adolescents. *J Korean Acad Nurs* 2009;39:357-65.
- Lehrner J, Marwinski G, Lehr S, Jöhren P, Deecke L. Ambient odors of orange and lavender reduce anxiety and improve mood in a dental office. *Physiol Behav* 2005;86:92-5.
- Kenner D. Using aromatics in clinical practice. *Cal J Orient Med* 1998;9:30-2.
- Schulz V, Hänsel R, Tyler VE. *Rational Phytotherapy: A Physician's Guide to Herbal Medicine*. 4th ed. Berlin: Rutledge; 2001.
- Combest WL. *Alternative therapies: Lavender*. US Pharm 1999;24:24-33.
- Amiri-Farahani L, Heidari T, Roozbahani N, Attarha M, Akbari-Torkestani N, Bekhradi R, *et al.* Effect of aromatherapy on pain severity in primary dysmenorrhea. *Arak Med Univ J* 2012;15:26-33. [Persian]
- Shahnazi M, Nikjoo R, Yavarikia P, Mohammad-Alizadeh-Charandabi S. Inhaled lavender effect on anxiety and pain caused from intrauterine device insertion. *J Caring Sci* 2012;1:255-61.
- Han SH, Hur MH, Buckle J, Choi J, Lee MS. Effect of aromatherapy on symptoms of dysmenorrhea in college students: A randomized placebo-controlled clinical trial. *J Altern Complement Med* 2006;12:535-41.
- Kane FM, Brodie EE, Coull A, Coyne L, Howd A, Milne A, *et al.* The analgesic effect of odour and music upon dressing change. *Br J Nurs* 2004;13:S4-12.
- Kim JT, Wajda M, Cuff G, Serota D, Schlame M, Axelrod DM, *et al.* Evaluation of aromatherapy in treating postoperative

- pain: Pilot study. *Pain Pract* 2006;6:273-7.
31. Auld B, Sinha P. Dysmenorrhoea: Features and current treatment options. *Prescriber* 2006;5:33-40.
 32. Shokohi F. Lavender Aromatherapy Effect on Pain Relief After Caesarean Section Patients Alzahra Hospital in Rasht. Sixth International Congress of Obstetrics and Gynecology, Rasht; 2003. p. 62-4.
 33. Kim JT, Ren CJ, Fielding GA, Pitti A, Kasumi T, Wajda M, *et al.* Treatment with lavender aromatherapy in the post-anesthesia care unit reduces opioid requirements of morbidly obese patients undergoing laparoscopic adjustable gastric banding. *Obes Surg* 2007;17:920-5.
 34. Williamson A, Hoggart B. Pain: A review of three commonly used pain rating scales. *J Clin Nurs* 2005;14:798-804.