

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dicaesium tetrachloridodioxido-plutonate(VI)

 Marianne P. Wilkerson^{a*} and Brian L. Scott^b
^aC-ADI, MS J565, Los Alamos National Laboratory, Los Alamos, New Mexico 87545 USA, and ^bMPA-MC, MS J514, Los Alamos National Laboratory, Los Alamos, New Mexico 87545 USA

Correspondence e-mail: mpw@lanl.gov


Received 4 October 2007; accepted 10 December 2007

 Key indicators: single-crystal X-ray study; $T = 141$ K; mean $\sigma(\text{Pu}-\text{O}) = 0.004$ Å; R factor = 0.015; wR factor = 0.037; data-to-parameter ratio = 26.3.

The anion of the title complex, $\text{Cs}_2[\text{PuCl}_4\text{O}_2]$, adopts a pseudo-octahedral geometry ($2/m$ crystallographic site symmetry) with two plutonyl oxide ligands in axial sites and four chloride ligands occupying the equatorial plane. Charge balance is maintained by two caesium cations per tetrachloridodioxido-plutonate(VI) anion. Principal bond lengths include $\text{Pu}-\text{O} = 1.752$ (3) Å and $\text{Pu}-\text{Cl} = 2.6648$ (8) Å.

Related literature

For related literature, see: Hall *et al.* (1966); Watkin *et al.* (1991); Wilkerson *et al.* (2004); Wilkerson *et al.* (2007); Bean *et al.* (2004, 2005); Grenthe *et al.* (2006); Grigoriev *et al.* (2004); Runde *et al.* (2003); Sessler *et al.* (2002).



Experimental

Crystal data

 $\text{Cs}_2[\text{PuCl}_4\text{O}_2]$
 $M_r = 678.67$
 Monoclinic, $C2/m$
 $a = 11.9489$ (7) Å
 $b = 7.7286$ (5) Å
 $c = 5.7855$ (4) Å
 $\beta = 96.439$ (1)°

 $V = 530.91$ (6) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 13.92$ mm⁻¹
 $T = 141$ (2) K
 $0.22 \times 0.18 \times 0.12$ mm

Data collection

 Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2001)
 $T_{\min} = 0.150$, $T_{\max} = 0.286$
 (expected range = 0.099–0.188)

 3212 measured reflections
 711 independent reflections
 659 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.015$
 $wR(F^2) = 0.037$
 $S = 1.26$
 711 reflections

 27 parameters
 $\Delta\rho_{\text{max}} = 0.78$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.64$ e Å⁻³

Data collection: APEX2 (Bruker, 2003); cell refinement: SAINT-Plus (Bruker, 2001); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: SHELXTL (Bruker, 2000); software used to prepare material for publication: SHELXTL.

We thank Mr Phillip Palmer and Mr Sean D. Reilly for technical assistance. This research at Los Alamos National Laboratory was supported by NA-22, Office of Nonproliferation Research and Engineering, US Department of Energy, National Nuclear Security Administration (NNSA), and Office of Basic Energy Sciences, US Department of Energy. Los Alamos National Laboratory is operated by Los Alamos National Security, LLC for the US Department of Energy, NNSA under contract number DE-AC52-06NA25396.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FI2048).

References

- Bean, A. C., Abney, K., Scott, B. L. & Runde, W. (2005). *Inorg. Chem.* **44**, 5209–5211.
- Bean, A. C., Albrecht-Schmitt, T. E. & Runde, W. (2004). *J. Solid State Chem.* **177**, 1346–1351.
- Bruker (2000). SHELXTL. Version 6.10. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2001). SAINT-Plus. Version 7.06. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2003). APEX2. Version 1.08. Bruker AXS Inc., Madison, Wisconsin, USA.
- Grenthe, I., Drozdzyński, J., Fujino, T., Buck, E. C., Albrecht-Schmitt, T. E. & Wolf, S. F. (2006). *The Chemistry of the Actinide and Transactinide Elements*, Vol. 1, ch. 5, edited by L. R. Morrs, N. M. Edelstein, J. Fuger & J. J. Katz, 3rd ed., p. 578. Dordrecht: Springer.
- Grigoriev, M. S., Antipin, M. Yu., Krot, N. N. & Bessonov, A. A. (2004). *Radiochim. Acta*, **92**, 405–409.
- Hall, D., Rae, A. D. & Waters, T. N. (1966). *Acta Cryst.* **20**, 160–162.
- Runde, W., Bean, A. C., Albrecht-Schmitt, T. E. & Scott, B. L. (2003). *Chem. Commun.* pp. 478–479.
- Sessler, J. L., Gorden, A. E. V., Seidel, D., Hannah, S., Lynch, V., Gordon, P. L., Donohoe, R. J., Tait, C. D. & Keogh, D. W. (2002). *Inorg. Chim. Acta*, **341**, 54–70.
- Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.
- Sheldrick, G. M. (2001). SADABS. University of Göttingen, Germany.
- Watkin, D. J., Denning, R. G. & Prout, K. (1991). *Acta Cryst.* **C47**, 2517–2519.
- Wilkerson, M. P., Arrington, C. A., Berg, J. M. & Scott, B. L. (2007). *J. Alloys Compd.* **444–445**, 634–639.
- Wilkerson, M. P., Dewey, H. J., Gordon, P. L. & Scott, B. L. (2004). *J. Chem. Crystallogr.* **34**, 807–811.

supporting information

Acta Cryst. (2008). E64, i5 [https://doi.org/10.1107/S1600536807066354]

Dicaesium tetrachloridodioxidoplutonate(VI)**Marianne P. Wilkerson and Brian L. Scott****S1. Comment**

The title compound, (1), is isostructural with the uranium analogue $\text{Cs}_2\text{U(VI)O}_2\text{Cl}_4$ and the neptunium analogue $\text{Cs}_2\text{Np(VI)O}_2\text{Cl}_4$, which crystallize in the space group $C2/m$ (Hall *et al.*, 1966; Watkin *et al.*, 1991; Wilkerson *et al.*, 2007). The Pu atom sits on a site of $2/m$ crystallographic symmetry; the twofold rotation axis bisects the Cl—U—Cl angle, and the plutonyl axis lies in the mirror plane. The plutonium metal is coordinated in a pseudo-octahedral fashion by two oxo groups and four chloride ligands. The oxo groups of the plutonyl ion lie *trans* to one another with a bond angle of $180.0(3)^\circ$. This angle is equivalent within 3σ to those reported for the limited number of plutonyl structures published (range $178.0(4)$ – $179.4(2)^\circ$) (Bean *et al.*, 2004; Bean *et al.*, 2005; Grigoriev *et al.*, 2004; Runde *et al.*, 2003). This value is within the range reported for the majority of actinyl compounds (Grenthe *et al.*, 2006). The Pu—O(oxo) distances are $1.752(3)$ Å. Although there are no reported six coordinate plutonyl structures with which to compare, this value is within range of Pu—O(oxo) bond lengths reported for seven coordinate plutonyl structures ($1.727(4)$ – $1.771(11)$ Å) (Bean *et al.*, 2004; Bean *et al.*, 2005; Grigoriev *et al.*, 2004; Runde *et al.*, 2003). The chloride ligands lie in the equatorial plane of the plutonyl ion, and the Pu—Cl bond distances are $2.6648(8)$ Å. Although there are no plutonyl chloride structures reported, these values are longer than Np—Cl bond lengths reported for six-coordinate $\text{Cs}_2\text{NpO}_2\text{Cl}_4$ ($2.653(3)$ Å) (Wilkerson *et al.*, 2004).

S2. Experimental

Caesium chloride (0.021 g, 0.12 mmol; Aldrich, 99.999%) was dissolved in 2M HCl (0.5 ml; Aldrich, ACS reagent, 37%), and this solution was added to a stock solution of 0.063 M Pu(VI)O_2^{2+} in 2M HCl (1 ml, 0.063 mmol; Fisher, Certified ACS Plus) (Sessler *et al.*, 2002). The vial containing the solution was covered loosely with parafilm and allowed to stand for 2 weeks at room temperature. Following evaporation of the solvent, dark golden-brown blocks formed. A dark golden-brown block of $0.22 \times 0.18 \times 0.12$ mm was cut from a larger crystal, and then thinly coated with epoxy and placed in a capillary. The capillary was coated with a thin film of acrylic dissolved in ethyl acetate. (Note: this triple containment was necessitated by the health hazards of transuranic materials.)




Figure 1

Molecular structure (70% probability displacement ellipsoids) of $\text{Cs}_2\text{PuO}_2\text{Cl}_4$ and the atom-numbering scheme used in Table 1. Symmetry codes as in Table 1.

Dicaesium tetrachloridodioxidoplutonium(VI)

Crystal data

$\text{Cs}_2[\text{PuCl}_4\text{O}_2]$

$M_r = 678.67$

Monoclinic, $C2/m$

Hall symbol: $-C 2y$

$a = 11.9489(7) \text{ \AA}$

$b = 7.7286(5) \text{ \AA}$

$c = 5.7855(4) \text{ \AA}$

$\beta = 96.439(1)^\circ$

$V = 530.91(6) \text{ \AA}^3$

$Z = 2$

$F(000) = 576$

$D_x = 4.245 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

$\theta = 3.2\text{--}28.7^\circ$

$\mu = 13.92 \text{ mm}^{-1}$

$T = 141 \text{ K}$

Irregular, golden brown

$0.22 \times 0.18 \times 0.12 \text{ mm}$

Data collection

Bruker D8 with APEXII CCD diffractometer	3212 measured reflections
Radiation source: fine-focus sealed tube	711 independent reflections
Graphite monochromator	659 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.025$
Absorption correction: multi-scan (SADABS; Sheldrick, 2001)	$\theta_{\text{max}} = 28.7^\circ$, $\theta_{\text{min}} = 3.2^\circ$
$T_{\text{min}} = 0.150$, $T_{\text{max}} = 0.286$	$h = -15 \rightarrow 15$
	$k = -10 \rightarrow 10$
	$l = -7 \rightarrow 7$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.0151P)^2]$
$R[F^2 > 2\sigma(F^2)] = 0.015$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.037$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.26$	$\Delta\rho_{\text{max}} = 0.78 \text{ e } \text{\AA}^{-3}$
711 reflections	$\Delta\rho_{\text{min}} = -0.64 \text{ e } \text{\AA}^{-3}$
27 parameters	Extinction correction: SHELXL97 (Sheldrick, 1997), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
0 restraints	Extinction coefficient: 0.0050 (2)
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Pu1	0.0000	0.0000	0.0000	0.02605 (11)
Cs1	0.34147 (2)	0.0000	0.69601 (6)	0.03801 (11)
Cl1	-0.10351 (7)	-0.24889 (10)	0.21367 (16)	0.0390 (2)
O1	0.1124 (3)	0.0000	0.2222 (6)	0.0395 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pu1	0.02849 (15)	0.02078 (14)	0.02946 (15)	0.000	0.00583 (9)	0.000
Cs1	0.03790 (19)	0.03608 (19)	0.0419 (2)	0.000	0.01250 (15)	0.000
Cl1	0.0436 (4)	0.0305 (4)	0.0454 (5)	-0.0015 (3)	0.0162 (4)	0.0058 (3)
O1	0.0381 (19)	0.0424 (19)	0.0371 (19)	0.000	0.0002 (16)	0.000

Geometric parameters (\AA , $^\circ$)

Pu1—O1 ⁱ	1.752 (3)	Cs1—Cl1 ^{xii}	3.5201 (8)
Pu1—O1	1.752 (3)	Cs1—Cl1 ^{viii}	3.5201 (8)

Pu1—Cl1 ⁱⁱ	2.6648 (8)	Cs1—Cl1 ^{xiii}	3.5674 (10)
Pu1—Cl1	2.6648 (8)	Cs1—Cl1 ^{xiv}	3.5674 (10)
Pu1—Cl1 ⁱⁱⁱ	2.6648 (8)	Cs1—O1	3.650 (4)
Pu1—Cl1 ⁱ	2.6648 (8)	Cs1—Cl1 ^{xv}	3.6710 (9)
Pu1—Cs1 ^{iv}	4.5680 (3)	Cs1—Cl1 ^{vi}	3.6710 (9)
Pu1—Cs1 ^v	4.5680 (3)	Cs1—Pu1 ^{xiii}	4.5680 (3)
Pu1—Cs1 ^{vi}	4.5680 (3)	Cs1—Pu1 ^{xvi}	4.5680 (3)
Pu1—Cs1 ^{vii}	4.5680 (3)	Cs1—Pu1 ^{xvii}	4.6207 (3)
Pu1—Cs1 ^{viii}	4.6207 (3)	Cl1—Cs1 ^{xviii}	3.5209 (9)
Pu1—Cs1 ^{ix}	4.6207 (3)	Cl1—Cs1 ^{viii}	3.5201 (8)
Cs1—Cl1 ^x	3.5209 (9)	Cl1—Cs1 ^{iv}	3.5674 (10)
Cs1—Cl1 ^{xi}	3.5209 (9)	Cl1—Cs1 ^{vi}	3.6710 (9)
O1 ⁱ —Pu1—O1	180.0 (3)	Cl1 ^x —Cs1—Cl1 ^{xiii}	158.75 (3)
O1 ⁱ —Pu1—Cl1 ⁱⁱ	90.98 (8)	Cl1 ^{xi} —Cs1—Cl1 ^{xiii}	109.41 (2)
O1—Pu1—Cl1 ⁱⁱ	89.02 (8)	Cl1 ^{xii} —Cs1—Cl1 ^{xiii}	104.390 (15)
O1 ⁱ —Pu1—Cl1	89.02 (8)	Cl1 ^{viii} —Cs1—Cl1 ^{xiii}	69.76 (2)
O1—Pu1—Cl1	90.98 (8)	Cl1 ^x —Cs1—Cl1 ^{xiv}	109.41 (2)
Cl1 ⁱⁱ —Pu1—Cl1	87.58 (3)	Cl1 ^{xi} —Cs1—Cl1 ^{xiv}	158.75 (3)
O1 ⁱ —Pu1—Cl1 ⁱⁱⁱ	89.02 (8)	Cl1 ^{xii} —Cs1—Cl1 ^{xiv}	69.76 (2)
O1—Pu1—Cl1 ⁱⁱⁱ	90.98 (8)	Cl1 ^{viii} —Cs1—Cl1 ^{xiv}	104.390 (15)
Cl1 ⁱⁱ —Pu1—Cl1 ⁱⁱⁱ	180.00 (5)	Cl1 ^{xiii} —Cs1—Cl1 ^{xiv}	65.91 (3)
Cl1—Pu1—Cl1 ⁱⁱⁱ	92.42 (3)	Cl1 ^x —Cs1—O1	66.15 (5)
O1 ⁱ —Pu1—Cl1 ⁱ	90.98 (8)	Cl1 ^{xi} —Cs1—O1	66.15 (5)
O1—Pu1—Cl1 ⁱ	89.02 (8)	Cl1 ^{xii} —Cs1—O1	63.98 (5)
Cl1 ⁱⁱ —Pu1—Cl1 ⁱ	92.42 (3)	Cl1 ^{viii} —Cs1—O1	63.98 (5)
Cl1—Pu1—Cl1 ⁱ	180.00 (4)	Cl1 ^{xiii} —Cs1—O1	133.04 (4)
Cl1 ⁱⁱⁱ —Pu1—Cl1 ⁱ	87.58 (3)	Cl1 ^{xiv} —Cs1—O1	133.04 (4)
O1 ⁱ —Pu1—Cs1 ^{iv}	57.778 (4)	Cl1 ^x —Cs1—Cl1 ^{xv}	99.912 (18)
O1—Pu1—Cs1 ^{iv}	122.222 (4)	Cl1 ^{xi} —Cs1—Cl1 ^{xv}	65.75 (3)
Cl1 ⁱⁱ —Pu1—Cs1 ^{iv}	53.469 (18)	Cl1 ^{xii} —Cs1—Cl1 ^{xv}	163.34 (3)
Cl1—Pu1—Cs1 ^{iv}	51.24 (2)	Cl1 ^{viii} —Cs1—Cl1 ^{xv}	112.348 (8)
Cl1 ⁱⁱⁱ —Pu1—Cs1 ^{iv}	126.531 (18)	Cl1 ^{xiii} —Cs1—Cl1 ^{xv}	61.25 (2)
Cl1 ⁱ —Pu1—Cs1 ^{iv}	128.76 (2)	Cl1 ^{xiv} —Cs1—Cl1 ^{xv}	95.403 (19)
O1 ⁱ —Pu1—Cs1 ^v	122.222 (4)	O1—Cs1—Cl1 ^{xv}	131.50 (4)
O1—Pu1—Cs1 ^v	57.778 (4)	Cl1 ^x —Cs1—Cl1 ^{vi}	65.75 (3)
Cl1 ⁱⁱ —Pu1—Cs1 ^v	126.531 (18)	Cl1 ^{xi} —Cs1—Cl1 ^{vi}	99.912 (18)
Cl1—Pu1—Cs1 ^v	128.76 (2)	Cl1 ^{xii} —Cs1—Cl1 ^{vi}	112.348 (8)
Cl1 ⁱⁱⁱ —Pu1—Cs1 ^v	53.469 (18)	Cl1 ^{viii} —Cs1—Cl1 ^{vi}	163.34 (3)
Cl1 ⁱ —Pu1—Cs1 ^v	51.24 (2)	Cl1 ^{xiii} —Cs1—Cl1 ^{vi}	95.403 (19)
Cs1 ^{iv} —Pu1—Cs1 ^v	180.000 (8)	Cl1 ^{xiv} —Cs1—Cl1 ^{vi}	61.25 (2)
O1 ⁱ —Pu1—Cs1 ^{vi}	122.222 (4)	O1—Cs1—Cl1 ^{vi}	131.50 (4)
O1—Pu1—Cs1 ^{vi}	57.778 (4)	Cl1 ^{xv} —Cs1—Cl1 ^{vi}	63.83 (3)
Cl1 ⁱⁱ —Pu1—Cs1 ^{vi}	51.24 (2)	Cl1 ^x —Cs1—Pu1 ^{xiii}	131.698 (14)
Cl1—Pu1—Cs1 ^{vi}	53.469 (18)	Cl1 ^{xi} —Cs1—Pu1 ^{xiii}	74.494 (15)
Cl1 ⁱⁱⁱ —Pu1—Cs1 ^{vi}	128.76 (2)	Cl1 ^{xii} —Cs1—Pu1 ^{xiii}	135.406 (14)
Cl1 ⁱ —Pu1—Cs1 ^{vi}	126.531 (18)	Cl1 ^{viii} —Cs1—Pu1 ^{xiii}	77.734 (14)
Cs1 ^{iv} —Pu1—Cs1 ^{vi}	64.454 (8)	Cl1 ^{xiii} —Cs1—Pu1 ^{xiii}	35.626 (12)

Cs1 ^v —Pu1—Cs1 ^{vi}	115.546 (8)	Cl1 ^{xiv} —Cs1—Pu1 ^{xiii}	96.174 (16)
O1 ⁱ —Pu1—Cs1 ^{vii}	57.778 (4)	O1—Cs1—Pu1 ^{xiii}	122.194 (5)
O1—Pu1—Cs1 ^{vii}	122.222 (4)	Cl1 ^{xv} —Cs1—Pu1 ^{xiii}	35.683 (12)
Cl1 ⁱⁱ —Pu1—Cs1 ^{vii}	128.76 (2)	Cl1 ^{vi} —Cs1—Pu1 ^{xiii}	94.713 (15)
Cl1—Pu1—Cs1 ^{vii}	126.531 (18)	Cl1 ^x —Cs1—Pu1 ^{xvi}	74.494 (15)
Cl1 ⁱⁱⁱ —Pu1—Cs1 ^{vii}	51.24 (2)	Cl1 ^{xi} —Cs1—Pu1 ^{xvi}	131.698 (14)
Cl1 ⁱ —Pu1—Cs1 ^{vii}	53.469 (18)	Cl1 ^{xii} —Cs1—Pu1 ^{xvi}	77.734 (14)
Cs1 ^{iv} —Pu1—Cs1 ^{vii}	115.546 (8)	Cl1 ^{viii} —Cs1—Pu1 ^{xvi}	135.406 (14)
Cs1 ^v —Pu1—Cs1 ^{vii}	64.454 (8)	Cl1 ^{xiii} —Cs1—Pu1 ^{xvi}	96.174 (16)
Cs1 ^{vi} —Pu1—Cs1 ^{vii}	180.000 (7)	Cl1 ^{xiv} —Cs1—Pu1 ^{xvi}	35.626 (12)
O1 ⁱ —Pu1—Cs1 ^{viii}	69.04 (11)	O1—Cs1—Pu1 ^{xvi}	122.194 (5)
O1—Pu1—Cs1 ^{viii}	110.96 (11)	Cl1 ^{xv} —Cs1—Pu1 ^{xvi}	94.713 (15)
Cl1 ⁱⁱ —Pu1—Cs1 ^{viii}	130.706 (17)	Cl1 ^{vi} —Cs1—Pu1 ^{xvi}	35.683 (12)
Cl1—Pu1—Cs1 ^{viii}	49.294 (17)	Pu1 ^{xiii} —Cs1—Pu1 ^{xvi}	115.546 (8)
Cl1 ⁱⁱⁱ —Pu1—Cs1 ^{viii}	49.294 (17)	Cl1 ^x —Cs1—Pu1 ^{xvii}	123.572 (15)
Cl1 ⁱ —Pu1—Cs1 ^{viii}	130.706 (17)	Cl1 ^{xi} —Cs1—Pu1 ^{xvii}	123.572 (15)
Cs1 ^{iv} —Pu1—Cs1 ^{viii}	78.510 (5)	Cl1 ^{xii} —Cs1—Pu1 ^{xvii}	35.021 (13)
Cs1 ^v —Pu1—Cs1 ^{viii}	101.490 (5)	Cl1 ^{viii} —Cs1—Pu1 ^{xvii}	35.021 (13)
Cs1 ^{vi} —Pu1—Cs1 ^{viii}	101.490 (5)	Cl1 ^{xiii} —Cs1—Pu1 ^{xvii}	76.579 (13)
Cs1 ^{vii} —Pu1—Cs1 ^{viii}	78.510 (5)	Cl1 ^{xiv} —Cs1—Pu1 ^{xvii}	76.579 (13)
O1 ⁱ —Pu1—Cs1 ^{ix}	110.96 (11)	O1—Cs1—Pu1 ^{xvii}	70.49 (5)
O1—Pu1—Cs1 ^{ix}	69.04 (11)	Cl1 ^{xv} —Cs1—Pu1 ^{xvii}	136.161 (13)
Cl1 ⁱⁱ —Pu1—Cs1 ^{ix}	49.294 (17)	Cl1 ^{vi} —Cs1—Pu1 ^{xvii}	136.161 (13)
Cl1—Pu1—Cs1 ^{ix}	130.706 (17)	Pu1 ^{xiii} —Cs1—Pu1 ^{xvii}	101.490 (5)
Cl1 ⁱⁱⁱ —Pu1—Cs1 ^{ix}	130.706 (17)	Pu1 ^{xvi} —Cs1—Pu1 ^{xvii}	101.490 (5)
Cl1 ⁱ —Pu1—Cs1 ^{ix}	49.294 (17)	Pu1—Cl1—Cs1 ^{xviii}	154.58 (3)
Cs1 ^{iv} —Pu1—Cs1 ^{ix}	101.490 (5)	Pu1—Cl1—Cs1 ^{viii}	95.68 (2)
Cs1 ^v —Pu1—Cs1 ^{ix}	78.510 (5)	Cs1 ^{xviii} —Cl1—Cs1 ^{viii}	87.583 (18)
Cs1 ^{vi} —Pu1—Cs1 ^{ix}	78.510 (5)	Pu1—Cl1—Cs1 ^{iv}	93.13 (2)
Cs1 ^{vii} —Pu1—Cs1 ^{ix}	101.490 (5)	Cs1 ^{xviii} —Cl1—Cs1 ^{iv}	109.41 (2)
Cs1 ^{viii} —Pu1—Cs1 ^{ix}	180.000 (3)	Cs1 ^{viii} —Cl1—Cs1 ^{iv}	110.24 (2)
Cl1 ^x —Cs1—Cl1 ^{xi}	66.90 (3)	Pu1—Cl1—Cs1 ^{vi}	90.85 (2)
Cl1 ^x —Cs1—Cl1 ^{xii}	92.417 (18)	Cs1 ^{xviii} —Cl1—Cs1 ^{vi}	80.088 (18)
Cl1 ^{xi} —Cs1—Cl1 ^{xii}	130.133 (13)	Cs1 ^{viii} —Cl1—Cs1 ^{vi}	163.34 (3)
Cl1 ^x —Cs1—Cl1 ^{viii}	130.133 (13)	Cs1 ^{iv} —Cl1—Cs1 ^{vi}	84.597 (19)
Cl1 ^{xi} —Cs1—Cl1 ^{viii}	92.417 (18)	Pu1—O1—Cs1	178.55 (16)
Cl1 ^{xii} —Cs1—Cl1 ^{viii}	66.25 (3)		

Symmetry codes: (i) $-x, -y, -z$; (ii) $-x, y, -z$; (iii) $x, -y, z$; (iv) $x-1/2, y-1/2, z-1$; (v) $-x+1/2, -y+1/2, -z+1$; (vi) $-x+1/2, -y-1/2, -z+1$; (vii) $x-1/2, y+1/2, z-1$; (viii) $-x, -y, -z+1$; (ix) $x, y, z-1$; (x) $x+1/2, -y-1/2, z$; (xi) $x+1/2, y+1/2, z$; (xii) $-x, y, -z+1$; (xiii) $x+1/2, y+1/2, z+1$; (xiv) $x+1/2, -y-1/2, z+1$; (xv) $-x+1/2, y+1/2, -z+1$; (xvi) $x+1/2, y-1/2, z+1$; (xvii) $x, y, z+1$; (xviii) $x-1/2, y-1/2, z$.