



# Spotted fever group rickettsiae transmitted by *Dermacentor* ticks and determinants of their spread in Europe

Weronika Buczek<sup>1,B,D,F</sup>, Agnieszka Koman-Iżko<sup>1,B-D,F</sup>, Alicja M. Buczek<sup>1,B,D</sup>, Alicja Buczek<sup>1,A,D-F</sup>, Katarzyna Bartosik<sup>1,E-F</sup>, Dorota Kulina<sup>2,D,F</sup>, Dariusz Ciura<sup>3,D,F</sup>

<sup>1</sup> Chair and Department of Biology and Parasitology, Medical University, Lublin, Poland

<sup>2</sup> Department of Basic Nursing and Medical Teaching, Medical University, Lublin, Poland

<sup>3</sup> Department of Health Promotion and Treatment of Obesity, Medical University of Silesia, Katowice, Poland

A – Research concept and design, B – Collection and/or assembly of data, C – Data analysis and interpretation, D – Writing the article, E – Critical revision of the article, F – Final approval of article

Buczek W, Koman-Iżko A, Buczek AM, Buczek A, Bartosik K, Kulina D, Ciura D. Spotted fever group rickettsiae transmitted by *Dermacentor* ticks and determinants of their spread in Europe. *Ann Agric Environ Med*. DOI: 10.26444/aaem/120602

## Abstract

Rickettsiae from the spotted fever group, i.e. the etiological agents of tick-borne lymphadenopathy/*Dermacentor*-borne necrotic erythema and lymphadenopathy (TIBOLA /DEBONEL) syndrome, are associated with ticks, including *Dermacentor marginatus* and *Dermacentor reticulatus*. The expansion of these ticks into new areas increases the risk of infection of their hosts with tick-borne pathogens. The study summarises the importance of 2 species from the genus *Dermacentor*, i.e. *D. marginatus* and *D. reticulatus*, in the spread of spotted fever group rickettsiae in various regions of Europe. The study also focuses on the determinants of the presence of vectors and transmission of rickettsiae, as well as the effects of human infections with these pathogens. The climate changes observed nowadays affect vectors and increase the incidence and spread of tick-borne diseases worldwide. Due to the existing risk of exposure to an increasing number of people, knowledge about the course of these serious diseases and their etiological factors should be disseminated among healthcare professionals as well as in society. There is a great challenge for epidemiological services to provide access to medical and veterinary facilities in order to diagnose and treat rickettsioses. Therefore, the development of a strategy for tick control and the popularisation of knowledge concerning prophylaxis of tick-borne diseases is indispensable.

## Key words

*Rickettsia*; spotted fever group rickettsiae; TIBOLA/DEBONEL; tick-borne diseases; *Dermacentor reticulatus*; *Dermacentor marginatus*

## INTRODUCTION

Tick-borne rickettsioses are caused by various bacterial species from the genera *Rickettsia*, *Ehrlichia*, *Neoehrlichia*, and *Anaplasma* [1, 2, 3, 4]. The greatest medical importance in this bacterial group is ascribed to spotted fever group rickettsiae (SFGR) (genus *Rickettsia*). The association of different rickettsia species with their hosts is determined by many factors, mainly the pathogen traits, the vector competence of tick species, and the presence of zoonotic reservoirs indispensable for the persistence of pathogens in a specific environment. The presented study analyses the role of *Dermacentor* ticks in the transmission of SFGR rickettsiae in Europe, especially in its central and south parts, and the effects of infection with these pathogens in humans. The increase is emphasised in the risk of attacks by *Dermacentor reticulatus* and *Dermacentor marginatus* ticks on humans and animals related to climate and weather change, and human activity affecting the landscape and the structure of plants and animals, the latter of which are potential hosts of various developmental stages of these ticks.

**Spread of spotted fever group rickettsiae.** SFGR are rod-shaped bacteria with a size of 0.3–0.5 × 0.8–2 µm with trilaminar cell walls surrounded by a thick electron-lucent layer (halo zone) without flagella. As reported by Socolovschi et al. [5], as many as 16 out of the 25 *Rickettsia* species identified to-date may infect humans, and 13 species have been detected in the inhabitants of Europe. Species with confirmed pathogenicity and remarkable epidemiological importance include *Rickettsia conorii* subsp. *conorii* (agent of Mediterranean spotted fever, MSF), *Rickettsia conorii* subsp. *israelensis* (Israeli spotted fever, ISF), *Rickettsia conorii* subsp. *caspia*, *Rickettsia conorii* subsp. *indica* (Indian tick typhus), *Rickettsia massiliae*, *Rickettsia monacensis* (MSF-like illness), *R. sibirica* subsp. *mongolitimonae* (lymphangitis-associated rickettsiosis, LAR), *Rickettsia slovaca* and *Rickettsia raoultii* (SENLAT- scalp eschars and neck lymphadenopathy, formerly referred to as TIBOLA- tickborne lymphadenopathy or DEBONEL- *Dermacentor*-borne necrotic erythema and lymphadenopathy), *Rickettsia aeschlimannii* (symptoms similar to those of MSF), *Rickettsia helvetica*, and *R. sibirica* subsp. *sibirica* [6, 7, 8, 9, 10, 11]. In central Europe, 5 *Rickettsia* species have been detected in ticks: *R. slovaca*, *R. raoultii*, *R. helvetica*, *R. monacensis*, and *Candidatus Rickettsia mendelii* [12, 13]. With the exception of *R. mendelii*, all these species have been identified in *D. marginatus* and *D. reticulatus* (Fig. 1).

Address for correspondence: Alicja Buczek, Chair and Department of Biology and Parasitology, Medical University, Poland  
E-mail: alicja.buczek@umlub.pl

Received: 20.02.2020; accepted: 15.04.2020; first published: 07.05.2020



Figure 1. *Dermacentor reticulatus*: (a) female, (b) male

SFGR have been detected in ticks from the genus *Dermacentor* across Europe, from the UK [14], Denmark [15], The Netherlands [16], and the Baltic countries [17] in the north, through countries in the western [8, 18, 19], central [20–30], and eastern parts [8, 31–34] and Spain [35], Portugal [36–38], Italy [39–43], Greece [44], and Croatia [45] in the south. In Poland, the presence of SFG rickettsiae in *D. reticulatus* has been documented in many regions of the country [46–53].

Ticks are infected with rickettsiae while feeding on a host with rickettsiosis. Within the tick population, rickettsiae are disseminated via transovarial [5, 33, 54, 55], transstadial [5, 33] and, probably less frequently, transspermal [56] transmission. The possibility of transmission of SFGR *R. massiliae* and *R. rickettsii* during the co-feeding of ixodid ticks has also been suggested [5].

In natural conditions, rickettsiae may be transferred in another way. The phenomenon of oral-anal contact between two *D. reticulatus* and *Ixodes ricinus* ticks, described recently for the first time, may indicate an alternative route of transmission of some pathogenic and non-pathogenic microorganisms from one population of sympatric tick species to another [57]. The scale of this type of interspecific tick contacts in natural conditions and their epidemiological importance are as yet unknown and require further investigations. The possibility of transmission of rickettsiae during the feeding of hungry or not fully engorged specimens on engorged ticks from the same species, also needs elucidation. Such research is even more advisable as hyperparasitism has been described in various ticks, including those with great epidemiological importance [58]. It has been experimentally shown that some pathogens, such as *Dipetalonema viteae*, *Borrelia crociduriae* and *Borrelia burgdorferi* s.l., can be transmitted during parasitization of some ticks by others [59–61].

As in the case of other tick-borne pathogens, the prevalence of rickettsiae in ticks differs between regions and between different years in the regions. In the northern regions, *D. reticulatus* occurrence varies, with the prevalence of *D. raoultii* in adult ticks varying from 0% – 36.9% (mean 4.9%) in different parts of Lithuania and Latvia [17]. In Belarus, as many as 41.9% of *D. reticulatus* specimens collected from animals and 44.5% of these ticks found on vegetation were infected with *Rickettsia* spp., with the dominance of *R. raoultii* (35.5% and 22.6% in specimens collected from

animals and vegetation, respectively) [34]. Higher SFGR infection rates in *Dermacentor* ticks have been reported from many localities in central and southern Europe.

In recreational areas in Białowieża (Podlaskie Province, north-eastern Poland), *R. raoultii* DNA was detected in 56.7% of *D. reticulatus* specimens collected from vegetation [47]. Later, Stańczak et al. [53] identified rickettsial DNA in 41% of adult *D. reticulatus* in this province. In Lublin Province (eastern Poland), the SFGR prevalence in this species during the study period was 53.0% [49]. *Dermacentor reticulatus* ticks from Łęczyńsko-Włodawskie Lakeland (Lublin Province) have been most frequently diagnosed with *R. raoultii* infection (43.8%) [52]. In the endemic regions of north-eastern (Warmińsko-Mazurskie Province) and central Poland (Mazowieckie Province), *Rickettsia* spp. were identified in 44.1% of *D. reticulatus* [50]. A higher percentage of SFGR (60%) was detected in adult stages of the species collected from dogs and cats in the Wrocław agglomeration [51].

Infection with SFG rickettsiae has been detected in 85.15% and 97.0% of *D. marginatus* ticks in southern and northern Spain, respectively [62, 63]. In Slovakia during 2004 – 2010, the infection rate ranged from 25.49% – 36.17% depending on the tick collection. The prevalence of *R. raoultii* was 8.1 – 8.6% and 22.3 – 27% in *D. marginatus* and *D. reticulatus*, respectively. In turn, the prevalence of *R. slovacca* was higher in *D. marginatus* (20.6 – 24.3%) than in *D. reticulatus* (1.7 – 3.4%) [26]. In hunting and recreational forest areas in southern Hungary, *R. raoultii* infection has been confirmed in 58% of examined *D. reticulatus* specimens [30]. A high prevalence of *Rickettsia* spp. (70.5%) in questing *D. reticulatus* and only 11.4% cases of *I. ricinus* have been reported from Saxony (Germany) [64]. As in the case of *D. reticulatus*, the infection rate in *D. marginatus* varied widely from 0.7% – 13.3% in Germany [24, 25], 32.1% in Italy [40] and 41.5% in Portugal [37].

Particularly noteworthy is the fact that SGFR are transferred into new areas together with the spread of *Dermacentor* ticks into urban and suburban areas. In a Kiev city park (Ukraine), 10.1% of *D. reticulatus* ticks were infected by *R. raoultii* [65]. As many as 36% of ticks collected in a park in Rome (Italy), including 14% of *D. marginatus*, 26% of *Rhipicephalus turanicus*, and 70% of *I. ricinus*, exhibited the presence of rickettsial DNA. *D. marginatus* were also found to be infected by *R. monacensis* [43]. SFGR may co-infect *D. reticulatus* ticks with other pathogens, i.e. TBE virus, *Anaplasma phagocytophilum*, *B. burgdorferi*, *Babesia* spp., or *Toxoplasma gondii* [52]. Ticks can also be infected by 2 SFG *Rickettsia* species. In Croatia, the coexistence of *R. helvetica* and *R. slovacca* was noted in 1% of *D. reticulatus* ticks [45]. Co-infections of SFGR (*R. slovacca* and *R. helvetica*) with *Borrelia lusitaniae* were also found in ticks, including *D. marginatus* (23.3%), collected in a Safari Park in Alentejo (Portugal) [37].

The circulation of *Rickettsia* spp. in nature proceeds with the participation of vectors – ticks and hosts of their juvenile and adult stages [66–70]. Larvae and nymphs of *D. reticulatus* and *D. marginatus* ticks parasitize small rodents and insectivores, while adult stages feed on medium-sized and large animals. *R. raoultii* has also been detected in blood collected 24 h after *D. reticulatus* was placed in feeding units [71]. In Slovakia, 11% of small mammals are infected by SFG rickettsiae [70]. The dominant species is *R. helvetica*, which has been detected most frequently in the striped field mouse



(*Apodemus agrarius*), the yellow-necked mouse (*Apodemus flavicollis*), the bank vole (*Myodes glareolus*), and the common vole (*Microtus arvalis*) in natural and suburban habitats. In turn, *R. felis* was identified in *A. flavicollis* captured in a suburban habitat. *R. slovaca* was detected in one *A. flavicollis* captured in a natural habitat. The highest species diversity of rickettsiae was observed in *A. flavicollis*, and the highest prevalence of the bacteria was recorded in *M. glareolus* [66–69]. The prevalence of *Rickettsia* spp. in the investigated small mammals (*A. agrarius*, *A. flavicollis*, *M. arvalis*, *M. agrestis*, *Mustela nivalis*, *M. glareolus*, *Sorex araneus*, and *Talpa europaea*) from Germany (Saxony) was 25.3% [64].

The strong relationship between rickettsiae and host species is supported by the 7-fold higher prevalence of *Rickettsia* spp. in *M. glareolus* infested with *D. reticulatus* than that in non-infested animals. In south-western Poland, DNA of SFG rickettsiae was detected in tissues of 17.6% of wild-living rodents – *A. agrarius*, *A. flavicollis*, and *M. glareolus* [72]. Some of the hosts of *D. reticulatus* and *D. marginatus* adult stages, i.e. wild and domestic large ungulates, carnivores, horses, or wild boars, can also be reservoirs of rickettsiae in nature [40, 51, 73, 74].

Analyses of mammalian biological material for the presence of the DNA of SFG rickettsiae and/or antibodies against pathogens have confirmed the widespread distribution and high prevalence of rickettsiae in *Dermacentor* spp. hosts in the European occurrence range of these ticks [51, 74–76]. For instance, high seroprevalence of *R. slovaca* in Spain was recorded in domestic ruminants – 65% in bullfighting cattle, 21% in goats, and 16% in sheep [77].

Ticks infected by SFG rickettsiae can invade humans and feed on the new host, which poses a high risk of transmission of the pathogens during blood ingestion. Cases of tick-borne lymphadenopathy (TIBOLA) caused by *R. slovaca* have been reported in patients who had contact with horses [78]. Humans can be infected not only by a feeding *Dermacentor* female tick, but also by a foraging male specimen [78, 79].

**Impact of environmental factors on *Dermacentor* ticks and transmission of SFG rickettsiae.** Climate change leading to global warming alters tick fauna in various habitats and the structure of animals, i.e. potential hosts of various tick development stages and animal reservoirs of pathogens. Examples of species with high dynamics of expansion in the last 10 years include *D. reticulatus* [80–83] and *D. marginatus* [84–86]. These ticks migrate with their hosts via ecological routes along rivers. The transfer of *D. reticulatus* ticks with hosts is the quickest route of extension of the occurrence range of this species. *D. reticulatus* females and males alone are able to move at a distance of  $66.35 \pm 100$  cm and  $54.85 \pm 45$  cm over 7 weeks, respectively [87]. Studies conducted on other tick species suggest that pathogens may influence the physical activity of ticks. It has been demonstrated that *Rickettsia* increase and *Arsenophonus* limit the locomotor activity of larvae of 3 tick species, i.e. *Amblyomma americanum*, *Dermacentor variabilis*, and *Ixodes scapularis*, which are pathogen vectors in eastern North America [88]. The presence of *Rickettsia* bacteria in *D. variabilis* ticks affect their behaviour [89].

Twenty years ago in Poland, *D. reticulatus* ticks were found only in the eastern and north-eastern parts of the country [90–92]. Currently, tick localities are reported from many other places in the eastern [92, 93], central and north-

eastern [94–97] western [98–100], and in south-western [101] provinces. In Europe, a similar trend towards expansion of the occurrence range is observed in other species with great epidemiological importance [102].

Due to the climate and weather changes, adult *D. reticulatus* stages in central Europe can attack hosts even during winter [103]. Active specimens were found in January in the habitat of this species in eastern Poland, which is usually a diapause period for these ticks. Similarly, *D. reticulatus* activity in winter has been reported from western Poland [104]. Besides the expansion of the occurrence range of *Dermacentor* ticks in Europe, their population size is also increasing. Over 10 years, the number of *D. reticulatus* adults in a meadow ecosystem used for grazing cattle in eastern Poland has increased two- and three-fold in spring and autumn, respectively [105]. The dynamics of the seasonal activity of this species with 2 peaks, in spring and autumn, and a period of decline in the activity in July persists in central Europe. However, the beginning of spring activity and the end of autumn activity shift in time together with the climate change [106]. Depending on the temperature, ticks can attack hosts already at the end of the calendar winter and in early spring. Favourable weather conditions contribute to the prolongation of the autumn activity of *D. reticulatus*. The risk of pathogen infection of hosts is increased at the peak of tick diurnal activity. In eastern Poland, adult *D. reticulatus* stages exhibit the highest activity from 12.00 midday – 16.25 in spring and from 10.00 a.m. – 18.25 in autumn, with a peak at 14.00 – 14.25 in both these periods [107].

The behaviour of ticks during the parasitic phase can be modified when 2 species coexist on the same host. During *D. reticulatus* and *D. marginatus* co-feeding on the same host, 2 inter-specific systems with different physiological features are formed, which may influence the transmission of tick-borne pathogens. In the inter-specific group, *D. marginatus* females attach to the host for a longer time, and although their feeding time is shorter than in a mono-specific group, feeding efficiency does not decrease [108]. The current study demonstrates that the co-feeding of *I. ricinus* and *D. reticulatus* ticks on the same host influences their non-parasitic stage of the life cycle by increasing significantly the egg amount and hatching success in *I. ricinus* feeding in inter-specific groups [109]. Additionally, the dynamics of tick feeding on hosts changes with the intensity of invasion [110] and tick gender structure [111].

**Human infection with SFG rickettsiae.** Human infections with SFG rickettsiae have been reported from various European countries, in particular from those located in the southern and central parts of the continent. As many as 8 species of different spotted fever group (SFG) rickettsiae, i.e. *R. slovaca*, *Rickettsia* sp. IRS3/IRS4, *R. massiliae*/Bar29, *R. aeschlimannii*, *Rickettsia* sp. RpA4/DnS14, *R. helvetica*, *Rickettsia* sp. DmS1, and *R. conorii*, were identified in 10 species of ticks infesting humans in north-western Spain [35]. *R. slovaca*, *R. raoultii*, and *R. roji* transmitted by *D. marginatus* and/or *D. reticulatus* are the agents of TIBOLA (tickborne lymphadenopathy) or DEBONEL (*Dermacentor*-borne necrotic erythema and lymphadenopathy) syndromes, which are also referred to as SENLAT, and reflects their clinical symptoms. The first case of *R. slovaca* infection following a tick bite was described in 1980. The patient was diagnosed with acute meningoencephalitis, febrile

relapses, and prolonged persistence of neurasthenic disorders [112]. The next case of *R. slovaca* infection in humans was documented in 1997 in a patient in France [113]. Human infection with *R. slovaca* was also diagnosed in, e.g. Slovakia, Bulgaria, Hungary, Germany, Italy, Spain and Portugal [9, 38, 39, 114–118]. Cases of human *R. raoultii* infections have also been reported from several other countries, e.g. France, Slovakia and Poland [116, 118, 119]. In Poland, *Rickettsia* spp. were detected for the first time in a patient with TIBOLA/DEBONEL symptoms in Warsaw [120].

SFG rickettsiae introduced with tick saliva during feeding enter human blood. Via interaction between the adhesive protein (OmpB), which is an abundant surface protein in *Rickettsia*, and Ku70 receptors in host cells, rickettsiae penetrate the vascular endothelial cells and are divided in the cell cytosol. Since rickettsiae penetrate other cells without passing through the intracellular space, they avoid the immune response. This promotes the spread of the infection in the organism. Most of the clinical symptoms caused by SFG rickettsiae are attributed to disseminated endothelial infection. *Rickettsia* spp. and other intracellular microorganisms have been identified in pathological aortic valves in humans [121]. Rickettsial infection results in increased microvascular permeability, generalised vascular inflammation, oedema, and release of vasoactive mediators, which exert an effect on coagulation and promote the release of pro-inflammatory cytokines.

The presence of antibodies against tick-borne SFG rickettsiae was detected in 36.0% of inhabitants of areas with a high risk of tick attacks in eastern Poland. In groups exposed to contact with ticks occupationally, antibodies against tick-borne SFG rickettsiae were confirmed in 50.7% of forestry workers and 21.3% of agriculture workers [122]. In Catalonia (north-eastern Spain), the seroprevalence of *R. slovaca* in the general population was reported to range from 3.7% – 5.5% [123], whereas in a group exposed to tick bites seroprevalence was as high as 16.9% [124].

Tick-borne rickettsioses (TBR) in humans may be asymptomatic [125], but usually with moderate symptoms [3, 9, 125]. However, deaths of patients have also been reported [9, 38].

The length of the TIBOLA/DEBONEL incubation time varies from 1 – 15 days (typically between 5 – 10 days). The most common symptoms of *R. slovaca* infection include asthenia, headache, painful adenopathies, painful scalp eschar, and oedema of lymph nodes close to the tick bite. Low fever, and rarely rash, as well as face oedema may develop in patients. Symptoms of SENLAT syndrome may also include inoculation eschar, cervical lymphadenopathy, high fever, malaise, and headaches [6, 38, 115–117, 126]. Alopecia around the eschar persisting for a long time (several months), even after antibiotic therapy, has been reported [116]. Clinical symptoms of *R. raoultii*-induced TIBOLA syndrome are similar to those of *R. slovaca* infection, except for the absence of alopecia.

There are known cases of an emerging syndrome of scalp eschar and neck lymphadenopathy after a tick bite [127]. SFGs can cause chronic symptoms in patients. Nillson et al. [128] reported the occurrence of *R. helvetica* in a patient with chronic perimyocarditis and suggested that the pathogen may play an important role in its etiology.

Due to the colonisation of new areas by *D. marginatus* and *D. reticulatus*, which are the main SFG vectors in Europe, the risk of human and animal infections (including

companion animals such as dogs and cats) with these rickettsiae and other tick-borne pathogens is increasing. The challenge for medical and epidemiological services is therefore to provide access to medical and veterinary facilities in order to diagnose and treat rickettsioses. Development of a strategy for tick control and popularisation of prophylaxis of tick-borne diseases are also indispensable.

## REFERENCES

1. Raoult D, Roux V. Rickettsioses as paradigms of new or emerging infectious diseases. *Clin Microbiol Rev.* 1997; 10(4): 694–719.
2. Raoult D, Lakos A, Fenollar F, Beytout J, Brouqui P, Fournier PE. Spotless rickettsiosis caused by *Rickettsia slovaca* and associated with *Dermacentor* ticks. *Clin Infect Dis.* 2002; 34(10): 1331–1336. <https://doi.org/10.1086/340100>
3. Parola P, Paddock CD, Raoult D. Tick-borne rickettsioses around the world: emerging diseases challenging old concepts. *Clin Microbiol Rev.* 2005; 18(4): 719–756. <https://doi.org/10.1128/CMR.18.4>
4. Brites-Neto J, Duarte KM, Martins TF. Tickborne infections in human and animal population worldwide. *Vet World.* 2015; 8(3): 301–315.
5. Socolovschi C, Mediannikov O, Raoult D, Parola P. The relationship between spotted fever group Rickettsiae and ixodid ticks. *Vet Res.* 2009; 40(2): 34.
6. Lakos A. Tick-borne lymphadenopathy- a new rickettsial disease? *Lancet* 1997; 350(9083): 1006. [https://doi.org/10.1016/S0140-6736\(05\)64072-X](https://doi.org/10.1016/S0140-6736(05)64072-X).
7. Raoult D, Berbis P, Roux V, Xu W, Maurin M. A new tick-transmitted disease due to *Rickettsia slovaca*. *Lancet* 1997; 350(9071): 112–113.
8. Mediannikov O, Matsumoto K, Samoylenko I, Drancourt M, Roux V, Rydkina E, Davoust B, Tarasevich I, Brouqui P, Fournier PE. *Rickettsia raoultii* sp. nov., a spotted fever group rickettsia associated with *Dermacentor* ticks in Europe and Russia. *Int J Syst Evol Microbiol.* 2008; 58(7): 1635–1639. <https://doi.org/10.1099/ijs.0.64952-0>
9. Oteo JA, Portillo A. Tick-borne rickettsioses in Europe. *Ticks Tick Borne Dis.* 2012; 3(5–6): 271–278. <https://doi.org/10.1016/j.ttbdis.2012.10.035>
10. Madeddu G, Mancini F, Caddeo A, Ciervo A, Babudieri S, Maida I, Fiori ML, Rezza G, Mura MS. *Rickettsia monacensis* as cause of Mediterranean spotted fever-like illness, Italy. *Emerg Infect Dis.* 2012; 18(4): 702–704. <https://doi.org/10.3201/eid1804.111583>
11. Parola P, Paddock CD, Socolovschi C, Labruna MB, Mediannikov O, Kernif T, Abdad MY, Stenos J, Bitam I, Fournier P-E, Raoult D. Update on Tick-Borne Rickettsioses around the World: a Geographic Approach. *Clin Microbiol Rev.* 2013; 26(4): 657–702.
12. Hajduskova E, Literak I, Papousek I, Costa FB, Novakova M, Labruna MB, Zdrzilova-Dubská L. *Candidatus Rickettsia mendelii*, a novel basal group rickettsia detected in *Ixodes ricinus* ticks in the Czech Republic. *Ticks Tick Borne Dis.* 2016; 7(3): 482–486. <https://doi.org/10.1016/j.ttbdis.2016.02.004>
13. Kowalec M, Szewczyk T, Welc-Falęciak R, Siński E, Karbowski G, Bajer A. Rickettsiales Occurrence and Co-occurrence in *Ixodes ricinus* Ticks in Natural and Urban Areas. *Microb. Ecol.* 2019; 77(4): 890–904. <http://doi.org/10.1007/s00248-018-1269-y>
14. Tijssse-Klasen E, Jameson LJ, Fonville M, Leach S, Sprong H, Medlock JM. First detection of spotted fever group rickettsiae in *Ixodes ricinus* and *Dermacentor reticulatus* ticks in the UK. *Epidemiol Infect.* 2011; 139(4): 524–529.
15. Klitgaard K, Chriél M, Isbrand A, Jensen TK, Bødker R. Identification of *Dermacentor reticulatus* ticks carrying *Rickettsia raoultii* on migrating jackal, Denmark. *Emerg Infect Dis.* 2017; 23(12): 2072–2074. <https://dx.doi.org/10.3201/eid2312.170919>
16. Nijhof AM, Bodaan C, Postigo M, Nieuwenhuijs H, Opsteegh M, Franssen L, Jebbink F, Jongejan F. Ticks and associated pathogens collected from domestic animals in the Netherlands. *Vector Borne Zoonotic Dis.* 2007; 7(4): 585–595.
17. Radzijeuskaja J, Paulauskas A, Aleksandraviciene A, Jonauskaitė I, Stanko M, Karbowski G, Petko B. New records of spotted fever group rickettsiae in Baltic region. *Microbes Infect.* 2015; 17(11–12): 874–878. <https://doi.org/10.1016/j.micinf.2015.09.006>
18. Beati L, Finidori JP, Raoult D. First isolation of *Rickettsia slovaca* from *Dermacentor marginatus* in France. *Am J Trop Med Hyg.* 1993; 48(2): 257–268.
19. Bonnet S, de la Fuente J, Nicollet P, Liu X, Madani N, Blanchard B, Maingourd C, Alongi A, Torina A, Fernández de Mera IG, Vicente J,



- George JC, Vayssier-Taussat M, Joncour G. Prevalence of tick-borne pathogens in adult *Dermacentor* spp. ticks from nine collection sites in France. *Vector Borne Zoonotic Dis.* 2013; 13(4): 226–236. <https://doi.org/10.1089/vbz.2011.0933>
20. Řeháček J. *Rickettsia slovaca*, the organism and its ecology. *Acta Scient Nat Acad Sci Bohemoslovaca Brno.* 1984; 18: 1–50.
  21. Řeháček J. Ecological relationships between ticks and rickettsiae. *Eur J Epidemiol.* 1989; 5(4): 407–413.
  22. Dautel H, Dippel C, Oehme R, Hartelt K, Schettler E. Evidence for an increased geographical distribution of *Dermacentor reticulatus* in Germany and detection of *Rickettsia* sp. RpA4. *Int J Med Microbiol.* 2006; 296(Suppl. 1): 149–156.
  23. Smetanová K, Schvarzová K, Kocianová E. Detection of *Anaplasma phagocytophilum*, *Coxiella burnetii*, *Rickettsia* spp., and *Borrelia burgdorferi* s.l. in ticks and wild-living animals in western and middle Slovakia. *Ann NY Acad Sci.* 2006; 1078: 312–315.
  24. Pluta S, Teward F, Hartelt K, Oehme R, Kimmig P, Mackenstedt U. *Rickettsia slovaca* in *Dermacentor marginatus* ticks, Germany. *Emerg Infect Dis.* 2009; 15(12): 2077–2078. <https://doi.org/10.3201/eid1512.090843>
  25. Silaghi C, Hamel D, Thiel C, Pfister K, Pfeiffer M. Spotted fever group rickettsiae in ticks, Germany. *Emerg Infect Dis.* 2011; 17(5): 890–892.
  26. Špitalská E, Štefanidesová K, Kocianová E, Boldiš V. *Rickettsia slovaca* and *Rickettsia raoultii* in *Dermacentor marginatus* and *Dermacentor reticulatus* ticks from Slovak Republic. *Exp Appl Acarol.* 2012; 57(2): 189–197. <https://doi.org/10.1007/s10493-012-9539-8>
  27. Duscher GG, Hodžić A, Weiler M, Vaux AGC, Rudolf I, Sixl W, Medlock JM, Versteirt V, Hubálek Z. First report of *Rickettsia raoultii* in field collected *Dermacentor reticulatus* ticks from Austria. *Ticks Tick Borne Dis.* 2016; 7(5): 720–722. <http://dx.doi.org/10.1016/j.ttbdis.2016.02.022>
  28. Ionita M, Silaghi C, Mitrea IL, Edouard S, Parola P, Pfister K. Molecular detection of *Rickettsia conorii* and other zoonotic spotted fever group rickettsiae in ticks, Romania. *Ticks Tick Borne Dis.* 2016; 7(1): 150–153. <https://doi.org/10.1016/j.ttbdis.2015.10.006>
  29. Rudolf I, Vencliková K, Blažejová H, Betášová L, Mendel J, Hubálek Z, Parola P. First report of *Rickettsia raoultii* and *Rickettsia helvetica* in *Dermacentor reticulatus* ticks from the Czech Republic. *Ticks Tick Borne Dis.* 2016; 7(6): 1222–1224. <https://doi.org/10.1016/j.ttbdis.2016.07.011>
  30. Szekeres S, van Leeuwen AD, Rigó K, Jablonszky M, Majoros G, Sprong H, Földvári G. Prevalence and diversity of human pathogenic rickettsiae in urban versus rural habitats, Hungary. *Exp Appl Acarol.* 2016; 68(2): 223–226.
  31. Rydkina E, Roux V, Rudakov N, Gafarova M, Tarasevich I, Raoult D. New Rickettsiae in ticks collected in territories of the Former Soviet Union. *Emerg Infect Dis.* 1999; 5(6): 811–814.
  32. Brouqui P, Parola P, Fournier PE, Raoult D. Spotted fever rickettsioses in southern and eastern Europe. *FEMS Immunol Med Microbiol.* 2007; 49(1): 2–12.
  33. Samoylenko I, Shpynov S, Raoult D, Rudakov N, Fournier P-E. Evaluation of *Dermacentor* species naturally infected with *Rickettsia raoultii*. *Clin Microbiol Infect.* 2009; 15 (Suppl 2): 305–306. <https://doi.org/10.1111/j.1469-0691.2008.02249.x>
  34. Reye AL, Stegny V, Mishaeva NP, Velhin S, Hübschen JM, Ignatyev G, Muller CP. Prevalence of tick-borne pathogens in *Ixodes ricinus* and *Dermacentor reticulatus* ticks from different geographical locations in Belarus. *PLoS One* 2013; 8(1): e54476. <https://doi.org/10.1371/journal.pone.0054476>
  35. Fernandez-Soto P, Perez-Sanchez R, Alamo-Sanz R, Encinas-Grandes A. Spotted fever group rickettsiae in ticks feeding on humans in northwestern Spain: is *Rickettsia conorii* vanishing? *Ann NY Acad Sci.* 2006; 1078: 331–333.
  36. Márquez FJ. Spotted fever group *Rickettsia* in ticks from southeastern Spain natural parks. *Exp Appl Acarol.* 2008; 45(3–4): 185–194.
  37. Milhano N, de Carvalho IL, Alves AS, Arroubé S, Soares J, Rodriguez P, Carolino M, Nuncio MS, Piesman J, de Sousa R. Coinfections of *Rickettsia slovaca* and *Rickettsia helvetica* with *Borrelia lusitaniae* in ticks collected in a Safari Park, Portugal. *Ticks Tick Borne Dis.* 2010; 1(4): 172–177. <https://doi.org/10.1016/j.ttbdis.2010.09.003>
  38. de Sousa R, Pereira BI, Nazareth C, Cabral S, Ventura C, Crespo P, Marques N, da Cunha S. *Rickettsia slovaca* Infection in Humans, Portugal. *Emerg Infect Dis.* 2013; 19(10): 1627–1629. <https://dx.doi.org/10.3201/eid1910.130376>
  39. Selmi M, Bertolotti L, Tomassone L, Mannelli A. *Rickettsia slovaca* in *Dermacentor marginatus* and tick-borne lymphadenopathy, Tuscany, Italy. *Emerg Infect Dis.* 2008; 14(5): 817–820. <https://doi.org/10.3201/eid1405.070976>
  40. Selmi M, Martello E, Bertolotti L, Bisanzio D, Tomassone L. *Rickettsia slovaca* and *Rickettsia raoultii* in *Dermacentor marginatus* ticks collected on wild boars in Tuscany, Italy. *J Med Entomol.* 2009; 46(6): 1490–1493.
  41. Masala G, Chisu V, Satta G, Socolovschi C, Raoult D, Parola P. *Rickettsia slovaca* from *Dermacentor marginatus* ticks in Sardinia, Italy. *Ticks Tick Borne Dis.* 2012; 3(5–6): 393–395.
  42. Martello E, Selmi M, Ragagli C, Ambrogi C, Stella MC, Mannelli A, Tomassone L. *Rickettsia slovaca* in immature *Dermacentor marginatus* and tissues from Apodemus spp. in the northern Apennines, Italy. *Ticks Tick Borne Dis.* 2013; 4(6): 518–521. <https://doi.org/10.1016/j.ttbdis.2013.07.002>
  43. Mancini F, Ciccozzi M, Lo Presti A, Cella E, Giovanetti M, Di Luca M, Toma L, Bianchi R, Khoury C, Rezza G, Ciervo A. Characterization of spotted fever group Rickettsiae in ticks from a city park of Rome, Italy. *Ann Ist Super Sanita.* 2015; 51(4): 284–290. [https://doi.org/10.4415/ANN\\_15\\_04\\_07](https://doi.org/10.4415/ANN_15_04_07)
  44. Psaroulaki A, Ragiadakou D, Kouris G, Papadopoulos B, Chaniotis B, Tselentis Y. Ticks, tick-borne rickettsiae, and *Coxiella burnetii* in the Greek Island of Cephalonia. *Ann NY Acad Sci.* 2006; 1078: 389–399.
  45. Dobec M, Golubic D, Punda-Polic V, Kaeppli F, Sievers M. *Rickettsia helvetica* in *Dermacentor reticulatus* ticks. *Emerg Infect Dis.* 2009; 15(1): 98–100. <https://doi.org/10.3201/eid1501.080815>
  46. Stańczak J. Detection of spotted fever group (SFG) rickettsiae in *Dermacentor reticulatus* (Acari: Ixodidae) in Poland. *Int J Med Microbiol.* 2006; 296(Suppl 1): 144–148. <https://doi.org/10.1016/j.ijmm.2006.01.014>
  47. Chmielewski T, Podsiadly E, Karbowski G, Tylewska-Wierzbianańska S. *Rickettsia* spp. in ticks, Poland. *Emerg Infect Dis.* 2009; 15: 486–488. <https://doi.org/10.3201/eid1503.080711>
  48. Matsumoto K, Grzeszczuk A, Brouqui P, Raoult D. *Rickettsia raoultii* and *Anaplasma phagocytophilum* in *Dermacentor reticulatus* ticks collected from Białowieża Primeval Forest European bison (*Bison bonasus bonasus*), Poland. *Clin Microbiol Infect.* 2009; 15(Suppl 2): 286–287.
  49. Wójcik-Fatla A, Cisak E, Zając V, Sroka J, Sawczyn A, Dutkiewicz J. Study on tick-borne rickettsiae in eastern Poland. I. Prevalence in *Dermacentor reticulatus* (Acari: Amblyomidae). *Ann Agric Environ Med.* 2013; 20(2): 276–279.
  50. Mierzejewska EJ, Pawełczyk A, Radkowski M, Welc-Fałęciak R, Bajera A. Pathogens vectored by the tick, *Dermacentor reticulatus*, in endemic regions and zones of expansion in Poland. *Parasit Vectors.* 2015; 8: 490. <https://doi.org/10.1186/s13071-015-1099-4>
  51. Król N, Obiegala A, Pfeiffer M, Lonc E, Kiewra D. Detection of selected pathogens in ticks collected from cats and dogs in the Wrocław Agglomeration, South-West Poland. *Parasit Vectors.* 2016; 9: 351. <https://doi.org/10.1186/s13071-016-1632-0>
  52. Zając V, Wójcik-Fatla A, Sawczyn A, Cisak E, Sroka J, Kloc A, Zając Z, Buczek A, Dutkiewicz J, Bartosik K. Prevalence of infections and coinfections with 6 pathogens in *Dermacentor reticulatus* ticks collected in eastern Poland. *Ann Agric Environ Med.* 2017; 24(1): 26–32.
  53. Stańczak J, Biernat B, Racewicz M, Zalewska M, Matyjasek A. Prevalence of different *Rickettsia* spp. in *Ixodes ricinus* and *Dermacentor reticulatus* ticks (Acari: Ixodidae) in north-eastern Poland. *Ticks Tick Borne Dis.* 2018; 9(2): 427–434. [doi.org/10.1016/j.ttbdis.2017.12.010](https://doi.org/10.1016/j.ttbdis.2017.12.010)
  54. Macaluso KR, Sonenshine DE, Ceraul SM, Azad AF. Infection and transovarial transmission of rickettsiae in *Dermacentor variabilis* ticks acquired by artificial feeding. *Vector Borne Zoonotic Dis.* 2004; 1(1): 45–53. <https://doi.org/10.1089/153036601750137660>
  55. Moore TC, Pulscher LA, Caddell L, von Fricken ME, Anderson BD, Gonchigoo B, Gray GC. Evidence for transovarial transmission of tick-borne rickettsiae circulating in Northern Mongolia. *PLoS Negl Trop Dis.* 2018; 12(8): e0006696. <https://doi.org/10.1371/journal.pntd.0006696>
  56. Hayes SF, Burgdorfer W, Aeschlimann A. Sexual transmission of spotted fever group rickettsiae by infected male ticks: detection of rickettsiae in immature spermatozoa of *Ixodes ricinus*. *Infect Immun.* 1980; 27(2): 638–642.
  57. Buczek A, Bartosik K, Buczek W, Buczek AM, Kulina D, Kulisz J, Tomasiewicz K. A unique phenomenon of oral-anal contact between ticks observed in two tick species *Ixodes ricinus* and *Dermacentor reticulatus*. *Ann Agric Environ Med.* 2018; 25(4): 686–689. <https://doi.org/10.26444/aaem/99054>
  58. Buczek A, Bartosik K, Buczek AM, Buczek W, Stanko M. Conspecific hyperparasitism in the *Hyalomma excavatum* tick and considerations on the biological and epidemiological implications of this phenomenon. *Ann Agric Environ Med.* 2019; 26(4): 548–554. <https://doi.org/10.26444/aaem/110128>

59. Votava CL, Rabalais FC, Ashley DC. Transmission of *Dipetalonema viteae* by hyperparasitism in *Ornithodoros tartakovskyi*. J Parasitol. 1974; 60(3): 479.
60. Helmy N, Khalil GM, Hoogstraal H. Hyperparasitism in *Ornithodoros erraticus*. J Parasitol. 1983; 69(1): 229–233.
61. Alekseev AN, Dubinina HV. Exchange of *Borrelia burgdorferi* between *Ixodes persulcatus* (Ixodidae: Acarina) sexual partners. J Med Entomol. 1996; 33(3): 351–354.
62. Márquez FJ, Rojas A, Ibarra V, Cantero A, Rojas J, Oteo JA, Muniain MA. Prevalence data of *Rickettsia slovaca* and other SFG Rickettsiae species in *Dermacentor marginatus* in the southeastern Iberian peninsula. Ann NY Acad Sci. 2006; 1078: 328–330.
63. Oteo JA, Portillo A, Santibáñez S, Pérez-Martínez L, Blanco JR, Jiménez S, Ibarra V, Pérez-Palacios A, Sanz M. Prevalence of spotted fever group *Rickettsia* species detected in ticks in La Rioja, Spain. Ann NY Acad Sci. 2006; 1078: 320–323.
64. Obiegala A, Król N, Oltersdorf C, Nader J, Pfeffer M. The enzootic life-cycle of *Borrelia burgdorferi* (sensu lato) and tick-borne rickettsiae: an epidemiological study on wild-living small mammals and their ticks from Saxony, Germany. Parasit Vectors. 2017; 10: 115. <https://doi.org/10.1186/s13071-017-2053-4>
65. Didyk YM, Blaňárová L, Pogrebnyak S, Akimov I, Petko B, Vichová B. Emergence of tick-borne pathogens (*Borrelia burgdorferi* sensu lato, *Anaplasma phagocytophilum*, *Rickettsia raoultii* and *Babesia microti*) in the Kyiv urban parks, Ukraine. Ticks Tick Borne Dis. 2017; 8(2): 219–225. <https://doi.org/10.1016/j.ttbdis.2016.10.002>
66. Pluta S, Hartelt K, Oehme R, Mackenstedt U, Kimmig P. Prevalence of *Coxiella burnetii* and *Rickettsia* spp. in ticks and rodents in southern Germany. Ticks Tick Borne Dis. 2010; 1(3): 145–147.
67. Burri C, Schumann O, Schumann C, Gern L. Are *Apodemus* spp. mice and *Myodes glareolus* reservoirs for *Borrelia miyamotoi*, *Candidatus Neoehrlichia mikurensis*, *Rickettsia helvetica*, *R. monacensis* and *Anaplasma phagocytophilum*? Ticks Tick Borne Dis. 2014; 5(3): 245–251.
68. Minichová L, Hamšíková Z, Mahriková L, Slovák M, Kocianová E, Kazimírová M, Škultéty L, Štefanidesová K, Špitalská E. Molecular evidence of *Rickettsia* spp. in ixodid ticks and rodents in suburban, natural and rural habitats in Slovakia. Parasit Vectors. 2017; 10: 158. <https://doi.org/10.1186/s13071-017-2094-8>
69. Baráková I, Derdákova M, Selyemová D, Chvostáč M, Špitalská E, Rosso F, Collini M, Rosà R, Tagliapietra V, Girardi M, Ramponi C, Hauffe HC, Rizzoli A. Tick-borne pathogens and their reservoir hosts in northern Italy. Ticks Tick Borne Dis. 2018; 9(2): 164–170. <https://doi.org/10.1016/j.ttbdis.2017.08.012>
70. Heglasová I, Vichová B, Kraljik J, Mošanský L, Miklisová D, Stanko M. Molecular evidence and diversity of the spotted-fever group *Rickettsia* spp. in small mammals from natural, suburban and urban areas of Eastern Slovakia. Ticks Tick Borne Dis. 2018; 9(6): 1400–1406. <https://doi.org/10.1016/j.ttbdis.2018.06.011>
71. Olivieri E, Wijnveld M, Bonga M, Berger L, Manfredi MT, Veronesi F, Jongejan F. Transmission of *Rickettsia raoultii* and *Rickettsia massiliae* DNA by *Dermacentor reticulatus* and *Rhipicephalus sanguineus* (s.l.) ticks during artificial feeding. Parasit Vectors. 2018; 11: 494. <https://doi.org/10.1186/s13071-018-3075-2>
72. Gajda E, Hildebrand J, Sprong H, Buńkowska-Gawlik K, Perec-Matysiak A, Coipan EC. Spotted fever rickettsiae in wild-living rodents from south-western Poland. Parasit Vectors. 2017; 10: 413. <https://doi.org/10.1186/s13071-017-2356-5>
73. Ortuño A, Quesada M, López-Claessens S, Castellà J, Sanfeliu I, Antón E, Segura-Porta F. The role of wild boar (*Sus scrofa*) in the eco-epidemiology of *R. slovaca* in Northeastern Spain. Vector Borne Zoonotic Dis. 2007; 7(1): 59–64. <https://doi.org/10.1089/vbz.2006.0576>
74. Chisu V, Foxi C, Masala G. First molecular detection of the human pathogen *Rickettsia raoultii* and other spotted fever group rickettsiae in Ixodid ticks from wild and domestic mammals. Parasitol Res. 2018; 117(11): 3421–3429. <https://doi.org/10.1007/s00436-018-6036-y>
75. Schreiber C, Krücken J, Beck S, Maaz D, Pachnicke S, Krieger K, Gross M, Kohn B, von Samson-Himmelstjerna G. Pathogens in ticks collected from dogs in Berlin/Brandenburg, Germany. Parasit Vectors. 2014; 7: 535. <https://doi.org/10.1186/s13071-014-0535-1>
76. Wächter M, Wölfel S, Pfeffer M, Dobler G, Kohn B, Moritz A, Pachnicke S, Silaghi C. Serological differentiation of antibodies against *Rickettsia helvetica*, *R. raoultii*, *R. slovaca*, *R. monacensis* and *R. felis* in dogs from Germany by a micro-immunofluorescent antibody test. Parasit Vectors. 2015; 8: 126. <https://doi.org/10.1186/s13071-015-0745-1>
77. Ortuño A, Pons I, Quesada M, Lario S, Anton E, Gil A, Castellà J, Segura F. Evaluation of the presence of *Rickettsia slovaca* infection in domestic ruminants in Catalonia, Northeastern Spain. Vector Borne Zoonotic Dis. 2012; 12(12): 1019–1022. <https://doi.org/10.1089/vbz.2012.0972>
78. Lakos A, Kőrösi A, Földvári G. Contact with horses is a risk factor for tick-borne lymphadenopathy (TIBOLA): a case control study. Wien Klin Wochenschr. 2012; 124(17–18): 611–617. <https://doi.org/10.1007/s00508-012-0217-y>
79. Földvári G, Rigó K, Lakos A. Transmission of *Rickettsia slovaca* and *Rickettsia raoultii* by male *Dermacentor marginatus* and *Dermacentor reticulatus* ticks to humans. Diagn Microbiol Infect Dis. 2013; 76(3): 387–389. <https://doi.org/10.1016/j.diagmicrobio.2013.03.005>
80. Bullová E, Lukán M, Stanko M, Petko B. Spatial distribution of *Dermacentor reticulatus* tick in Slovakia in the beginning of the 21st century. Vet Parasitol. 2009; 165(3–4): 357–360. <https://doi.org/10.1016/j.vetpar.2009.07.023>
81. Chitimia-Dobler L. Spatial distribution of *Dermacentor reticulatus* in Romania. Vet Parasitol. 2015; 214(1–2): 219–223. <https://doi.org/10.1016/j.vetpar.2015.09.018>
82. Földvári G, Siroký P, Szekeres S, Majoros G, Sprong H. *Dermacentor reticulatus*: a vector on the rise. Parasit Vectors. 2016; 9: 314. <http://dx.doi.org/10.1186/s13071-016-1599-x>
83. Hofmeester TR, van der Lei PB, Docters van Leeuwen A, Sprong H, van Wieren SE. New foci of *Haemaphysalis punctata* and *Dermacentor reticulatus* in the Netherlands. Ticks Tick Borne Dis. 2016; 7(2): 367–370. <https://doi.org/10.1016/j.ttbdis.2015.12.009>
84. Rubel F, Brugger K, Pfeffer M, Chitimia-Dobler L, Didyk YM, Leverenz S, Dautel H, Kahl O. Geographical distribution of *Dermacentor marginatus* and *Dermacentor reticulatus* in Europe. Ticks Tick Borne Dis. 2016; 7(1): 224–233. <https://doi.org/10.1016/j.ttbdis.2015.10.015>
85. Walter M, Brugger K, Rubel F. The ecological niche of *Dermacentor marginatus* in Germany. Parasitol Res. 2016; 115(6): 2165–2174. <https://doi.org/10.1007/s00436-016-4958-9>
86. Selmi M, Tomassone L, Ceballos LA, Crisci A, Ragagli C, Pintore MD, Mignone W, Pautasso A, Ballardini M, Casalone C, Mannelli A. Analysis of the environmental and host-related factors affecting the distribution of the tick *Dermacentor marginatus*. Exp Appl Acarol. 2018; 75(2): 209–225. <https://doi.org/10.1007/s10493-018-0257-8>
87. Buczek A, Zając Z, Woźniak A, Kulina D, Bartosik K. Locomotor activity of adult *Dermacentor reticulatus* ticks (Ixodida: Ixodidae) in natural conditions. Ann Agric Environ Med. 2017; 24(2): 271–275. <https://doi.org/10.5604/12321966.1230736>
88. Kagemann J, Clay K. Effects of infection by *Arsenophonus* and *Rickettsia bacteria* on the locomotive ability of the ticks *Amblyomma americanum*, *Dermacentor variabilis*, and *Ixodes scapularis*. J Med Entomol. 2013; 50(1): 155–162.
89. El Nababout A, Kho J, Lloyd V, Rossolimo T. *Rickettsia* spp. and its Effects on the physiology and behaviour of *Dermacentor variabilis*. Int J Biol. 2017; 9(2): 39–49. <https://doi.org/10.5539/ijb.v9n2p39>
90. Siuda K. 1993. Ticks (Acari: Ixodida) of Poland. Part II Taxonomy and Distribution. PTP, Warszawa (in Polish).
91. Buczek A, Kuśmierz A, Olszewski K, Buczek L, Czerny K, Łañcut M. Comparison of rabbit skin changes after feeding of *Ixodes ricinus* (L.) and *Dermacentor reticulatus* (Fabr.), In: Bernini F, Nannelli R, Nuzzaci G, de Lillo E. (Eds.), Acarid phylogeny and evolution: adaptation in mites and ticks. Proceedings of the IV Symposium of the European Association of Acarologists. Kluwer Academic Publ., Dordrecht, 2002, pp. 419–424.
92. Buczek A, Bartosik K. Ticks (Ixodida: Ixodidae, Amblyomidae) in south-eastern Poland and their medical and epidemiological importance. Zdr Publ. 2011; 121(4): 392–397.
93. Zając Z, Bartosik K, Buczek A. Factors influencing the distribution and activity of *Dermacentor reticulatus* (F.) ticks in an anthropopressure unaffected area in central-eastern Poland. Ann Agric Environ Med. 2016; 23(2): 270–275. <https://doi.org/10.5604/12321966.1203889>
94. Zygnier W, Wędrychowicz H. Occurrence of hard ticks in dogs from Warsaw area. Ann Agric Environ Med. 2006; 13(2): 355–359.
95. Zygnier W, Górski P, Wędrychowicz H. New localities of *Dermacentor reticulatus* tick (vector of *Babesia canis canis*) in central and eastern Poland. Pol J Vet Sci. 2009; 12(4): 549–555.
96. Kadulski S, Izdebska JN. New data on distribution of *Dermacentor reticulatus* (Fabr.) (Acari, Ixodidae) in Poland, In: Buczek A, Błaszak C. (Eds.), Arthropods: Invasions and their control. Akapit, Lublin, 2009, pp. 53–58.
97. Mierzejewska EJ, Estrada-Peña A, Alsarraf M, Kowalec M, Bajer A. Mapping of tick *Dermacentor reticulatus* expansion in Poland in 2012–2014. Ticks Tick Borne Dis. 2016; 7(1): 94–106. <https://doi.org/10.1016/j.ttbdis.2015.09.003>



98. Karbowski G, Kiewra D. New locations of *Dermacentor reticulatus* ticks in Western Poland: the first evidence of the merge in *D. reticulatus* occurrence areas? *Wiad Parazytol.* 2010; 56(4): 333–336.
99. Nowak M. Discovery of *Dermacentor reticulatus* (Acari: Amblyomidae) populations in the Lubuskie Province (Western Poland). *Exp Appl Acarol.* 2011; 54(2): 191–197. <https://doi.org/10.1007/s10493-010-9422-4>
100. Kiewra D, Czulowska A. Evidence for an increased distribution range of *Dermacentor reticulatus* in south-west Poland. *Exp Appl Acarol.* 2013; 59(4): 501–506. <https://doi.org/10.1007/s10493-012-9612-3>
101. Cuber P, Solarz K, Mosialek A, Jakubiec-Spanier M, Spanier A. The first record and occurrence of the ornate cow tick *Dermacentor reticulatus* (Fabricius, 1794) in south-western Poland. *Ann Parasitol.* 2013; 59(1): 49–51.
102. Gray JS, Dautel H, Estrada-Peña A, Kahl O, Lindgren E. Effects of climate change on ticks and tick-borne diseases in Europe. *Interdiscip Perspect Infect Dis.* 2009; 593232: 1–12. <https://doi.org/10.1155/2009/593232>
103. Buczek A, Bartosik K, Zajac Z. Changes in the activity of adult stages of *Dermacentor reticulatus* (Ixodida: Amblyomidae) induced by weather factors in eastern Poland. *Parasit Vectors.* 2014; 7: 245. <https://doi.org/10.1186/1756-3305-7-245>
104. Kiewra D, Czulowska A, Lonc E. Winter activity of *Dermacentor reticulatus* (Fabricius, 1794) in the newly emerging population of Lower Silesia, south-west Poland. *Ticks Tick Borne Dis.* 2016; 7(6): 1124–1127. <https://doi.org/10.1016/j.ttbdis.2016.08.012>
105. Buczek A, Bartosik K, Wiśniowski Ł, Tomaszewicz K. Changes in population abundance of adult *Dermacentor reticulatus* (Acari: Amblyomidae) in long-term investigations in eastern Poland. *Ann Agric Environ Med.* 2013; 20(2): 269–272.
106. Bartosik K, Wiśniowski Ł, Buczek A. Abundance and seasonal activity of adult *Dermacentor reticulatus* (Acari: Amblyomidae) in eastern Poland in relation to meteorological conditions and the photoperiod. *Ann Agric Environ Med.* 2011; 18(2): 340–344.
107. Bartosik K, Wiśniowski Ł, Buczek A. Questing behavior of *Dermacentor reticulatus* adults (Acari: Amblyomidae) during diurnal activity periods in eastern Poland. *J Med Entomol.* 2012; 49(4): 859–864.
108. Buczek A, Bartosik K, Zajac Z, Stanko M. Host-feeding behaviour of *Dermacentor reticulatus* and *Dermacentor marginatus* in mono-specific and inter-specific infestations. *Parasit Vectors.* 2015; 8: 470. <https://doi.org/10.1186/s13071-015-1078-9>
109. Bartosik K, Buczek A, Borzęcki A, Kulina D. Study of the non-parasitic stage in *Ixodes ricinus* after co-feeding with *Dermacentor reticulatus* in three infestations. *Ann Agric Environ Med.* 2017; 24(1): 90–95. <https://doi.org/10.5604/12321966.1234005>
110. Bartosik K, Buczek A. The impact of intensity of invasion of *Ixodes ricinus* and *Dermacentor reticulatus* on the course of the parasitic phase. *Ann Agric Environ Med.* 2012; 19(4): 651–655.
111. Bartosik K, Buczek A, Buczek W, Buczek AM, Kulina D, Koman-Iżko A. Host feeding behaviour of *Dermacentor reticulatus* males in relation to the transmission of pathogens. *Ann Agric Environ Med.* 2019; 26(2): 227–230. <https://doi.org/10.26444/aaem/105402>
112. Mittermayer T, Brezina R, Urvölgyi J. First report of an infection with *Rickettsia slovaca*. *Folia Parasitol.* 1980; 27(4): 373–376.
113. Raoult D, Berbis P, Roux V, Xu W, Maurin M. A new tick-transmitted disease due to *Rickettsia slovaca*. *Lancet* 1997; 350(9071): 112–113. [https://doi.org/10.1016/S0140-6736\(05\)61814-4](https://doi.org/10.1016/S0140-6736(05)61814-4)
114. Bacellar F, Nuncio MS, Alves MJ, Filipe AR. *Rickettsia slovaca*: un agente del grupo de las fiebres exantemáticas, en Portugal. *Enferm Infecc Microbiol Clin.* 1995; 13(4): 218–223.
115. Komitova R, Lakos A, Aleksandrov A, Christova I, Murdjeva M. A case of tick-transmitted lymphadenopathy in Bulgaria associated with *Rickettsia slovaca*. *Scand J Infect Dis.* 2003; 35(3): 213. <https://doi.org/10.1080/0036554021000027016>
116. Parola P, Roveery C, Rolain JM, Brouqui P, Davoust B, Raoult D. *Rickettsia slovaca* and *R. raoultii* in tick-borne rickettsioses. *Emerg Infect Dis.* 2009; 15(7): 1105–1108. <https://doi.org/10.3201/eid1507.081449>
117. Rieg S, Schmoldt S, Theilacker C, de With K, Wölfel S, Kern WV, Dobler G. Tick-borne lymphadenopathy (TIBOLA) acquired in Southwestern Germany. *BMC Infect Dis.* 2011; 11: 167. <https://doi.org/10.1186/1471-2334-11-167>
118. Sekeyova Z, Subramanian G, Mediannikov O, Diaz MQ, Nyitray A, Blaskovicova H, Raoult D. Evaluation of clinical specimens for *Rickettsia*, *Bartonella*, *Borrelia*, *Coxiella*, *Anaplasma*, *Francisella* and *Diplorickettsia* positivity using serological and molecular biology methods. *FEMS Immunol Med Microbiol.* 2012; 64(1): 82–91. <https://doi.org/10.1111/j.1574-695X.2011.00907.x>
119. Świtaj K, Chmielewski T, Borkowski P, Tylewska-Wierzbanowska S, Olszyna-Krowicka M. Spotted fever rickettsiosis caused by *Rickettsia raoultii*—case report. *Przegl Epidemiol.* 2012; 66(2): 347–350.
120. Chmielewski T, Rudzka D, Fiecek B, Mączka I, Tylewska-Wierzbanowska S. Case of TIBOLA/DEBONEL (tick-borne lymphadenopathy/*Dermacentor*-spp. borne necrosis-erythema lymphadenopathy) in Poland. *Przegl Epidemiol.* 2011; 65(4): 583–586.
121. Nilsson K, Liu A, Pahlson C, Lindquist O. Demonstration of intracellular microorganisms (*Rickettsia* spp., *Chlamydia pneumoniae*, *Bartonella* spp.) in pathological human aortic valves by PCR. *J Infect.* 2005; 50(1): 46–52.
122. Zajac W, Wojcik-Fatla A, Cisak E, Sroka J, Sawczyn A, Dutkiewicz J. Study on tick-borne rickettsiae in eastern Poland. II. Serological response of occupationally exposed populations. *Ann Agric Environ Med.* 2013; 20(2): 280–282.
123. Antón E, Nogueras MM, Pons I, Font B, Muñoz T, Sanfeliu I, Segura F. *Rickettsia slovaca* infection in humans in the northeast of Spain: seroprevalence study. *Vector Borne Zoonotic Dis.* 2008; 8(5): 689–694. <https://doi.org/10.1089/vbz.2007.0246>
124. Lledó L, Gegúndez MI, Fernandez N, Sousa R, Vicente J, Alamo R, Fernández-Soto P, Pérez-Sánchez R, Bacellar F. The seroprevalence of human infection with *Rickettsia slovaca*, in an area of northern Spain. *Ann Trop Med Parasitol.* 2006; 100(4): 337–343.
125. Fournier PE, Xeridat B, Raoult D. Isolation of a rickettsia related to Astrakhan fever rickettsia from a patient in Chad. *Ann NY Acad Sci.* 2003; 990: 152–157.
126. Portillo A, Santibáñez S, García-Álvarez L, Palomar AM. Rickettsioses in Europe. *Microbes Infect.* 2015; 17(11–12): 834–838. <https://doi.org/10.1111/j.1749-6632.2003.tb07356.x>
127. Dubourg G, Socolovschi C, Del Giudice P, Fournier PE, Raoult D. Scalp eschar and neck lymphadenopathy after tick bite: an emerging syndrome with multiple causes. *Eur J Clin Microbiol Infect Dis.* 2014; 33(8): 1449–1456. <http://dx.doi.org/10.1007/s10096-014-2090-2>
128. Nilsson K, Lindquist O, Pahlso C. Association of *Rickettsia helvetica* with chronic perimyocarditis in sudden cardiac death. *Lancet* 1999; 354(9185): 1169–1173.