

Case Report

Cystic Fibrosis of Pancreas and Nephrotic Syndrome: A Rare Association

Author 1 (Corresponding author): Selvi Kelekçi, Assistant Professor. Dicle University, Medicine School, Department of Pediatric Pulmonology.

E- mail: selvikelekci@gmail.com.

Mobile Phone: 090 505 953 23 01, Fax: 090 412 248 80 01

Kıtlıbil, 21280/ Diyarbakır, Turkey.

Author 2: Müsemma Karabel. Assistant Professor. Dicle University, Medicine School, Department of Pediatric Pulmonology. Diyarbakır, Turkey. E-mail: musemma.alagoz@gmail.com

Author 3: Aydın Ece, Professor. Dicle University, Medicine School, Director of Pediatric Nephrology Department. Diyarbakır, Turkey. E-mail: drece@gmail.com

Author 4: Velat Şen, Assistant Professor. Dicle University, Medicine School, Department of Pediatric Pulmonology, Diyarbakır, Turkey. E-mail: drvelatsen@hotmail.com

Author 5: Ali Güneş, Assistant Professor. Dicle University, Medicine School, Department of Pediatrics. Diyarbakır, Turkey. E-mail: dragunes@gmail.com

Author 6: İlyas Yolbaş, Assistant Professor. Dicle University, Medicine School, Department of Pediatrics. Diyarbakır, Turkey. E-mail: dryolbas@yahoo.com

Author 7: Cahit Şahin, Research Assistant. Dicle University, Medicine School, Department of Pediatrics. Diyarbakır, Turkey. E-mail: csahin@gmail.com

Running title: Cystic fibrosis and nephrotic syndrome in children

*There is no conflict of interest between authors.

Abstract

Cystic fibrosis (CF) is a genetic disease with autosomal recessive inheritance and is more common in Caucasian people. The prevalence of the disease is between 1/2000 and 1/3500 live births, and the incidence varies between populations. Although the cystic fibrosis transmembrane conductance regulator gene is expressed in the kidneys, renal involvement is not common. With advances in the treatment of cystic fibrosis, life expectancy has increased, and some previously unobserved disease associations are now seen in patients. It is important to follow patients for possible abnormalities that may accompany CF. In this paper, we present two rare cases of cystic fibrosis accompanied by nephrotic syndrome.

Keywords: Cystic fibrosis, nephrotic syndrome, child.

Korean JJP Accepted diatrics

Introduction

Cystic fibrosis (CF) is a recessively inherited genetic disease that decreases quality of life and is more commonly seen in Caucasian people. CF occurs due to defective epithelial cell chloride channels caused by mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene (1). Nephrotic syndrome (NS) is more common in children than adults. NS is a clinical syndrome characterized by massive proteinuria and hypoalbuminemia due to increased permeability of the glomerular capillary membrane.

Because several systems are involved in CF, clinical symptoms show a heterogeneous distribution (2). The gastrointestinal system, reproductive system, and sweat glands are affected in patients with CF, and CF also causes progressive lung disease, which is still the most important cause of morbidity and mortality (3). Early diagnosis and appropriate treatment are very effective in reducing long-term morbidity from CF.

The lungs are generally affected by CF, while pancreatic involvement is seen in 85-90% of cases (4, 5). The **CFTR** gene is usually expressed in the kidneys; however, renal involvement is not very common in patients with CF (6). Patients that experience urinary system involvement most commonly have nephrolithiasis and cough related mechanical urological problems (7, 8). Urolithiasis in CF may result from enteric hyperoxaluria caused by fat malabsorption and the lack of intestinal oxalate degrading bacteria as *Oxalobacter formigenes*. Additional renal manifestations occasionally reported in children and teenagers include AA **type** amyloidosis, membranoproliferative glomerulonephritis, and postinfectious glomerulonephritis (9, 10,11). The number of CF-NS associated cases is limited in the literature (12,13). In this report, we present two rare cases of both NS and CF in children.

Case 1:

A nine-year-old male patient diagnosed with **CF**, lung disease, and pancreatic insufficiency had been monitored for three years. A gradual increase in the ratio of decline in the urine and

complaints of swelling of the face and both ankles had begun two years previous. Physical examination revealed the following: temperature 36.8°C, pulse 92/minute, blood pressure 109/71 mmHg (95th percentile), edemas in both lower extremities and the sacral region, and more prominent edemas around the face. Other system examinations were normal. Liver function tests, viral hepatitis panel results, alpha-1 antitrypsin levels, and abdominal ultrasonography were normal but sweat test was 98mmol/l (chloride concentration).

At that time he was detected hypoalbuminemia and proteinuria on a random urine sample. Complement factors 3 and 4 were normal, anti-nuclear antibody was negative, serum urea was 11 mg/dl, triglyceride was 175 mg/dl, total cholesterol 230 mg/dl, creatinine was 0.6 mg/dl, and albumin was 0.5 g/dl in diagnosis time and 24 hour urine collection revealed nephrotic range proteinuria (>960 mg/m²/day).

The patient began steroid therapy following NS diagnosis. At follow-up, the patient was found to have steroid-resistant NS, and a renal biopsy was performed under light microscopy (LM) and immunofluorescence microscopy (IF). The renal biopsy showed mesangial proliferation without AA. Partial remission was achieved using low-dose steroids (15 mg/day prednisolone, for a period 1 year) and cyclosporine A therapy (3 mg/kg/day; for a period 4 months). Serum albumin and total protein levels in serum were 3.2 g/dl, 5.5 g/dl after treatment, respectively. Renal biopsy was the second time in the last 6 months. Serum Amyloid A protein was negative. At the moment, our patient has treated with 0.5mg/kg/every other day prednisone and his proteinuria (30g/day) has continued. The patient is still in partial remission.

The patient's radiological imaging showed destruction around the peripheral airways, and bronchiectasis was detected. Pulmonary function tests provided the following values: FEV1 78% and FVC 85%; FEV1/FVC values were low. The patient was treated with

recombinant human DNase, followed by treatment with inhaled bronchodilators and pulmonary rehabilitation.

Case 2:

A two-year-old male patient was referred to our center. In history, he had diagnosed as CF due to chronic diarrhea, while he was two months of age. During follow-up, no pulmonary involvement was observed; however, eight months prior to this study a decrease in urine volume was detected, as well as, edemas in the patient's face and bilateral lower extremities.

The patient's medical history included an upper respiratory tract infection two weeks prior. Physical examination revealed the following: temperature 36.8°C, pulse 84/min, blood pressure 94/50mm Hg (50th percentile), and edemas in the patient's face and bilateral lower extremities. Other system examinations were normal.

Laboratory examinations results revealed the following in diagnosis time; sweat test was repeated 122 mmol/l, liver function tests and complement factor 3 and 4 were normal; anti-nuclear antibody was negative; albumin was 0.6 mg/dl; total cholesterol was 220 mg/dl, urea was 16 mg/dl; and creatinine was 0.4 mg/dl. 24 hour urine collection was found nephrotic proteinuria (>960 mg/m²/day) and the patient was diagnosed with NS and started oral corticosteroids 2 mg/kg/ day, 6weeks therapy and achieved remission. The patient was regarded as MCD. The patient is still in remission and being followed for NS.

Discussion

Early diagnosis, intensive antibiotic therapy and developments in supportive treatment have extended the life expectancy for CF patients by reducing mortality and morbidity. Due to the increase in patient lifespans, previously unobserved disease associations have been detected. Renal involvement in CF patients is rare, but recently, renal pathologies such as glomerulosclerosis, mesangial proliferation, nephrocalcinosis and microscopic hematuria, tubular damage, fibrillary glomerulonephritis (14), and amyloidosis (15) have been reported.

The first case in this report was monitored for three years. The patient's biopsy revealed mesangial proliferation. The patient was diagnosed with steroid-resistant **NS**, and cyclosporine treatment was initiated in addition to prednisolone treatment. As a result of treatment, the patient is in partial remission. The second patient was diagnosed with **NS** when he was 1.5 years old, and since he did not have hematuria-hypertension. Systemic amyloidosis with renal involvement has been reported in some patients with **CF** (16). Secondary amyloidosis is in fact a complication of chronic inflammatory processes such as tuberculosis and bronchiectasis, and it is conceivable that this condition may develop in patients with **CF** which is characterised by chronic and recurrent respiratory infections (10). Renal amyloidosis is a cause of **NS**. The prognosis of patients with **CF** and AA amyloidosis is generally poor, with nearly all patients dying within 1 year of clinical presentation (17). In second patient; there was no need for a renal biopsy because of responded to steroid therapy and renal functions returned to normal.

There is a widespread belief that kidneys are not affected by **CF**. **NS** may be randomly associated with **CF**, or it may be a complication of **CF**. **CF** patients may develop kidney disease due to nephrotoxic agent exposure, immune complexes circulating in the serum due to recurrent pulmonary infections, or nephrotoxic antibiotics, such as aminoglycosides. Regular patient follow-up will allow full systemic evaluation and early diagnosis of accompanying complications, especially renal disorders. **CF** patients must undergo urine tests and evaluations for possible renal disorders, including proteinuria.

Conflicts of Interest Statement

There is no conflict of interest between authors.

References

1-Gadsby DC, Vergani P, Csanády L. The ABC protein turned chloride channel whose failure causes cystic fibrosis. *Nature*. 2006;440:477-483.

- 2-Wallis C. Diagnosis of cystic fibrosis. In: Hodson M, Geddes D, Bush A, editors. Cystic Fibrosis. London: Hodder Arnold, 2007: 99-108.
- 3-Ratjen F, Döring G.Cystic fibrosis. Lancet. 2003;361:681-689
- 4-O'Sullivan BP, Freedman SD.Cystic fibrosis. Lancet. 2009;373:1891-1904.
- 5-McCormick J, Green MW, Mehta G, Culross F, Mehta A. Demographics of the UK cystic fibrosis population: implications for neonatal screening. Eur J Hum Genet. 2002;10:583-590.
- 6- Jouret F, Devuyst O.CFTR and defective endocytosis: new insights in the renal phenotype of cystic fibrosis. Pflugers Arch. 2009;457:1227-1236.
- 7-Perez-Brayfield MR, Caplan D, Gatti JM, Smith EA, Kirsch AJ.Metabolic risk factors for stone formation in patients with cystic fibrosis. J Urol. 2002;167:480-484.
- 8-Orr A, McVean RJ, Webb AK, Dodd ME. Questionnaire survey of urinary incontinence in women with cystic fibrosis. BMJ. 2001;322:1521.
- 9-Mc Laughlin AM, Crotty TB, Egan JJ, Watson AJ, Gallagher CGAmyloidosis in cystic fibrosis: a case series. J Cyst Fibros. 2006;5:59-61.
- 10- Melzi ML, Costantini D, Giani M, Appiani AC, Giunta AM.Severe nephropathy in three adolescents with cystic fibrosis. Arch Dis Child. 1991;66:1444-1447.
- 11-Soriano E, Fischman D, Cheriya P. Membranoproliferative glomerulonephritis in patients with cystic fibrosis: coincidence or comorbidity? A case series. South Med J. 2008;101:641-655.
- 12- Al-Shawwa BA, Rao AR.Cystic fibrosis and renal disease: a case report. J Med Case Rep. 2007 ;1:24.
- 13-Melzi ML, Costantini D, Giani M, Appiani AC, Giunta AM.Severe nephropathy in three adolescents with cystic fibrosis. Arch Dis Child. 1991;66:1444-1447.

14-Laufer J, Augarten A, Szeinberg A, Rapoport J, Katzenelson D, Yahav Y. Nephrotic syndrome and fibrillary glomerulonephritis. *J Intern Med.* 1997;242:83-86.

15-Mc Laughlin AM, Crotty TB, Egan JJ, Watson AJ, Gallagher CG. Amyloidosis in cystic fibrosis: a case series. *J Cyst Fibros.* 2006;5:59-61.

16-Canciani M, Pederzini F, Mastella G, Boccato P. Systemic amyloidosis in cystic fibrosis. *Acta Paediatr Scand.* 1985;74:613-614.

17-Yahiaoui Y, Jablonski M, Hubert D, Mosnier-Pudar H, Noël LH, Stern M et al. Renal involvement in cystic fibrosis: diseases spectrum and clinical relevance. *Clin J Am Soc Nephrol.* 2009;4:921-928.

Korean JJP Accepted diatrics

Korean KJP Accepted diacritics