



RESEARCH ARTICLE

A bioactive compound isolated from Duku (*Lansium domesticum* Corr) fruit peels exhibits cytotoxicity against T47D cell line [version 1; peer review: 1 approved with reservations]

Khusnul Fadhilah ¹, Subagus Wahyuono², Puji Astuti²

¹Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, 55281, Indonesia

²Departement of Biology, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, 55281, Indonesia

V1 First published: 06 Jan 2020, 9:3
<https://doi.org/10.12688/f1000research.21072.1>
Latest published: 06 Jan 2020, 9:3
<https://doi.org/10.12688/f1000research.21072.1>

Abstract

Background: Breast cancer is a major health problem for women globally. Many attempts have been promoted to cure cancer by finding new anticancer medicines from natural resources. Despite the richness of biodiversity discovered, there are some natural resources that remain unexplored. Fruit peels of Duku (*Lansium domesticum* Corr.) are rich with compounds that may have the potential to be developed as anticancer drugs. This study aimed to isolate cytotoxic compounds from the fruit peels of *L. domesticum* and assess their cytotoxic nature against T47D cells.

Methods: Powdered peels were macerated with ethyl acetate and the filtrate was evaporated to give EtOAc extract A. Dried extract A was triturated with n-hexane to give n-hexane soluble fraction B and insoluble fraction C. The cytotoxic nature of these three samples were assessed using MTT assay using T47D cells and doxorubicin as a control.

Results: Fraction C that showed the smallest IC₅₀ (25.56 ± 0.64 µg/mL) value compared to extract A and fraction B. Fraction C was further fractionated by vacuum liquid chromatography to give 6 subfractions. Subfraction 2 showed a single compound based on thin layer chromatography, and this compound was identified as Lamesticum A on the basis of its spectroscopic data. Lamesticum A demonstrated cytotoxic activity against T47D cell lines with an IC₅₀ value of 15.68 ± 0.30 µg/mL.

Conclusions: Further research is needed to investigate the potential of the natural compound Lamesticum A derived from *L. domesticum* fruit peel as an anticancer therapy.

Keywords

Lansium domesticum Corr., cytotoxic, T47D cell line, Lamesticum A

Open Peer Review

Reviewer Status ?


Invited Reviewers

1

version 1

06 Jan 2020

?
report

1. **Ratana Banjerdpongchai** , Chiang Mai University, Chiang Mai, Thailand

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Subagus Wahyuono (subagusw_fa@ugm.ac.id)

Author roles: **Fadhilah K:** Investigation, Writing – Original Draft Preparation; **Wahyuono S:** Conceptualization, Data Curation, Methodology, Supervision, Writing – Review & Editing; **Astuti P:** Conceptualization, Data Curation, Methodology, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Ministry of Higher Education of the Republic of Indonesia (grant No: 5832/UN1.DITLIT/DIT-LIT/LT/2018). *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2020 Fadhilah K *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Fadhilah K, Wahyuono S and Astuti P. **A bioactive compound isolated from Duku (*Lansium domesticum* Corr) fruit peels exhibits cytotoxicity against T47D cell line [version 1; peer review: 1 approved with reservations]** F1000Research 2020, **9:3** <https://doi.org/10.12688/f1000research.21072.1>

First published: 06 Jan 2020, **9:3** <https://doi.org/10.12688/f1000research.21072.1>

Introduction

The most frequent cancer in women and that which causes the highest mortality is breast cancer. In Indonesia, it was reported that approximately 21% of cancer deaths among women were due to breast cancer¹. Therefore, new medicines to eradicate this type of cancer is required. Duku (*Lansium domesticum* Correa) widely grows in Indonesia. Traditionally, *L. domesticum* bark and seeds have been used to treat dysentery and fever². Based on previous studies, chloroform and methanol extracts of *L. domesticum* displayed cytotoxic activity on murine melanoma (B₁₆F₁₀) and colon cancer (HT29) cells³. In addition, it has been shown that ethanol and ethyl acetate fractions of the peel have a deterrent activity on DNA damage in lymphoblast cells induced by H₂O₂ exposure⁴. Onoceranoid-type of triterpenoids have been isolated from twigs and leaves of *L. domesticum*, and these compounds showed antibacterial and antimutagenic activities^{5,6}. In this study, the cytotoxic effects of compound extracted from the peels of *L. domesticum* are assayed against breast cancer T47D cells.

Methods

Plant material

The fruits of *L. domesticum* were collected on March 2018 from Bantul, Yogyakarta (GPS : -7.871098, 110.394854) and identified at the Department of Pharmaceutical Biology, Faculty of Pharmacy, Universitas Gadjah Mada.

Chemicals and equipment

Organic solvents (methanol, ethyl acetate, chloroform, n-hexane) used were pro analytical grades obtained from Merck. Silica gel F₂₅₄, Silica gel PF₂₅₄ (Merck), RPMI 1640, Fetal Bovine Serum, Penicillin-Streptomycin, Fungizon, Sodium bicarbonate (Gibco), HEPES (Invitrogen), Phosphate Buffered Saline, MTT (Sigma Aldrich cat. M5655), Doxorubicin (Sigma Aldrich). Infrared (KBr) spectrum was obtained from spectrophotometer (Shimadzu) using a method previously described by Ashokkumar and Ramaswamy⁷. Ultraviolet spectrum (CHCl₃) was obtained from UV spectrophotometer (Hitachi UH 5300). Sample (1 mg) were diluted in 1 mL CHCl₃ and was run between 200–400 nm. Spectra of ¹H- and ¹³C- NMR in CDCl₃ solvent were measured using JEOL JNM-ECZ 500R/S1 at 500 MHz.

Extraction and fractionation

The peel was separated from the fruit, dried in oven 50°C for 24 hours and powdered using a blender. Powdered *L. domesticum* fruit peel (200 gr) was macerated with ethyl acetate (EtOAc; 2 L) overnight. This solution was filtrated (0,45µm) and the filtrate was evaporated to dryness with a rotary evaporator set at 50°C, to give dried EtOAc extract (A, 50.13 g). In order to separate the extract into non-polar and polar fractions, 6 gr of extract A was diluted 5 times using n-hexane (20 mL) to give soluble n-hexane fraction B (supernatant; 3.03 g) and insoluble n-hexane fraction C (residue; 2.83 g).

Fraction C was the most active among other fractions (see *Results*), and was therefore further fractionated using vacuum liquid chromatography as described by Mae Sri Hartati *et al.*⁸. In brief, using silica gel preparation grade (15 gr) as stationary

phase this was eluted with n-hexane and increasing amounts of ethyl acetate. Six subfractions were obtained and subfraction 2 contained a major compound which appeared as white crystals (referred to as compound 1). Compound 1 was obtained as a single compound from subfraction 2, while the other subfractions still contained various compounds. Compound 1 (142.6 mg), had a melting point at 140–150°C (Figure 1). Compound 1 was identified using spectroscopy data such as ultraviolet (UV), infrared (IR), ¹³C-NMR and ¹H-NMR (see section *Chemicals and equipment*).

Cytotoxic assay

The bioassay followed the methodology described by Bahuguna *et al.*⁹ with modifications. In brief, 100 µl T47D cells (in RPMI media Faculty of Medicine, Universitas Gadjah Mada) were placed in each well of a 96 well microplates, resulting in 1 × 10⁴ cells/well. The cells were incubated for 24 hours at 37°C in a CO₂ incubator.

Extract A, fractions B and C and compound 1 (5mg) were dissolved in DMSO (50 µL). Serial concentrations of extract and fractions (50, 25, 12.5, 6.25, 3.125 µg/mL), compound 1 (25, 12.5, 6.25, 3.125 µg/mL) and doxorubicin (positive control; 0.5, 0.25, 0.125, 0.0625, 0.0312 µg/mL) were obtained. Cells were treated with the dose dependent samples and incubated for 24 hours at 37°C. The culture medium was removed by pipette, and MTT solution (100µL) was added to each well and incubated for 4 hours at 37°C. After incubation, stop solution (10% SDS, 100 µL) was added to each well and let stand at room temperature for 24 hours.

Absorbance was measured by microplate reader (Bio Rad) at 595 nm. positive control The data generated were used to plot a dose-response curve and IC₅₀ of the samples was determined.

Statistical analysis

The IC₅₀ values were analyzed by one-way ANOVA with statistical significance P < 0.05 using IBM SPSS ver.23.

Results

Compound 1 characterization

Identified as Lamesticum A.

White crystal. IR (KBr) ν_{max} cm⁻¹: 3074, 2960, 1712; UV (MeOH) λ_{max} 236,5; ¹H,¹³C-NMR: see Table 1; m/z 502; (Calculated for C₃₁H₅₀O₅)

The infrared spectroscopy (KBr) spectrum of 1 showed a broad band at 3400–2800 cm⁻¹, which indicated the presence of –OH group, specifically –COOH due to intermolecular bonding. This data is supported by the appearance of –C=O at 1712 cm⁻¹. Compound 1 displayed UV absorption at 236,5 nm. The ¹³C-NMR spectrum (500 MHz, CDCl₃) of compound 1 showed 30 carbons (Table 1). There were two down field carbon signals (δ , 147.6 and 148.1 ppm) identified as C=O signal carbons. Two characteristic terminals =CH₂ signals (δ , 107.4 and 114.2 ppm) were observed, and this identity was confirmed by 2D (Het-Cor) NMR technique. Based on ¹³C-NMR




Figure 1. Isolation process of cytotoxic compound 1 (subfraction 2) from *Lansium domesticum* fruit peels.

Table 1. ^{13}C -NMR spectrum (500 MHz, CDCl_3) of compound 1, Lamesticum A.

Position	^1H -NMR (J, Hz)		^{13}C -NMR δ (ppm)
	δ (ppm)	Multiplicity	
1	1.2	2H, triplet (7.0)	27.98
2	2.1	2H, triplet (6.8)	29.18
3	-	-	147.60
4	-	-	51.78
5	0.9	1H, triplet (7.1)	50.78
6	1.4	2H, multiplet	27.51
7	1.9	2H, triplet (6.8)	28.75
8	-	-	122.10
9	1.1	1H, triplet (7.0)	48.91
10	-	-	47.68
11	1.4	2H, multiplet	30.59
12	1.7	2H, multiplet	29.67
13	1.13	1H, triplet (7.0)	41.71
14	-	-	135.91
15	5.4	1H, triplet (6.8)	113.93

Position	^1H -NMR (J, Hz)		^{13}C -NMR δ (ppm)
	δ (ppm)	Multiplicity	
16	1.7	2H, dd (7.0, 6.8)	30.59
17	1.1	1H, triplet	31.81
18	-	-	38.85
19	1.2	2H, multiplet	32.09
20	2.1	2H, multiplet	33.15
21	-	-	148.10
22	-	-	147.20
23	1.7	3H, singlet	23.37
24	1.9	3H, singlet	23.76
25	0.8	3H, singlet	16.16
26	4.8	2H, doublet (9.1)	107.4
27	1.7	3H, Singlet	17.91
28	0.7	3H, singlet	14.37
29	4.6	2H, doublet (9.2)	114.23
30	1.6	3H, singlet	22.88

^1H - and ^{13}C -NMR (CDCl_3) spectra were obtained from JEOL JNM-ECZ 500R/S1, 500 MHz

and $^1\text{H-NMR}$ data, compound 1 (Figure 2) was identified as Lamesticum A ($\text{C}_{31}\text{H}_{50}\text{O}_5$, m/z , 502) which was previously isolated from *L. domesticum* twigs⁵.

Cytotoxicity of extract A, fractions B and C, and compound 1

The cytotoxicity of extract A, fractions B and C, compound 1 and doxorubicin (positive control) is shown in Table 2. Fraction C was the most cytotoxic (IC_{50} 25.57 $\mu\text{g/mL}$) compared with extract A (29.41 $\mu\text{g/mL}$) and fraction B (43.51 $\mu\text{g/mL}$). The IC_{50} of the isolated compound from fraction C, compound 1/Lamesticum A was 15.68 $\mu\text{g/mL}$. All samples inhibited T47D cell growth in a dose dependent behavior (Figure 3).

Discussion

In this study, the cytotoxic activity of Lamesticum A, derived from the peel of *L. domesticum*, was demonstrated in the T47D cell line with IC_{50} 15.68 ($\mu\text{g/ml}$). The T47D cell line is an epithelial breast cancer cell subtype luminal A cell line that express estrogen and progesterone receptors¹⁰. Based on National Cancer Institute guidelines, a natural compound has potent anticancer activity if it has $\text{IC}_{50} < 4 \mu\text{g/ml}$ or $10 \mu\text{M}$ ¹¹.

Many triterpenoid compounds have been previously isolated from *L. domesticum*. Most of these compounds are UV inactive or have no strong UV absorbance because triterpenoid's lack of a conjugated functional group¹². Lansiosida A and Dukunolida A has been isolated from n-hexane extract of *L. domesticum* fruit peel^{13,14}. Lamesticum A is an onoceranoïd-type triterpenoid, isolated previously from *L. domesticum* twigs, that has antibacterial activity against *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Micrococcus luteus*, *Bacillus subtilis*, *Micrococcus pyogenes* and *Bacillus cereus* with minimum inhibitory concentration of $< 15 \mu\text{g/ml}$ ⁵. Another onoceranoïd-type triterpenoid Lansium acid I-IX were isolated from *L. domesticum* leaves, which was reported to have antimutagenic activity⁶.

Based on several *in vitro* tests, some terpenoid compounds had anticancer activity. Sesquiterpene lactone compounds are known to inhibit NF- κB , thereby inducing apoptosis¹⁵. Celastrol has anticancer properties by regulating various transcription factors, angiogenesis processes, cell cycle arrest and induction of apoptosis¹⁶. Betulinic acid can induce apoptosis in HT-29 colon cancer cells and acts as a chemosensitizer for chemotherapeutic agents in wildtype adenocarcinoma cancer cells (SNU-C5/WT)¹⁷.




Figure 2. Isolated compound 1, Lamesticum A.

Table 2. IC_{50} values of extract, fractions and isolated compound 1 against T47D breast cancer cell line.

	IC_{50} ($\mu\text{g/ml}$), mean \pm SD
Extract A	29.41 \pm 0.67
Fraction B	43.51 \pm 1.77
Fraction C	25.56 \pm 0.64
Lamesticum A	15.68 \pm 0.30
Doxorubicin	0.18 \pm 0.01

T47D Cell viability




Figure 3. T47D breast cancer cell line viability after treatments with extracts from *Lansium domesticum* fruit peels. Error bars shows standard deviation.

Clematangoticosides D and F from *Clematis tangutica* are known to have cytotoxic activity against human gastric cancer cell line (SGC-7901) with IC_{50} 24.22 and 21.35 μ M, respectively¹⁸. Cycloartane-type and oleanane-type triterpenoids from *Ligularia przewalskii* show cytotoxicity in Hela, HEPG2, SGC7901, MDA231, HL-60, and Lewis cell lines with IC_{50} 8.40–24.39 μ M¹⁹.

It has been reported that natural compounds combined with low doses of antineoplastics can increase effectiveness and reduce toxic effects²⁰. Betulinic acid can induce apoptosis when combined with 5-fluorouracil, irinotecan and oxaliplatin⁴. Ursolic acid (UA), a pentacyclic triterpenoid, is known to have anticancer activity through interfering with multiple signaling pathways. Furthermore, UA has been shown to act as a chemosensitizing agent to increase the effect of conventional anticancer drugs²¹, and to increase the effect of doxorubicin by increasing the cellular amount of the drug in the MCF cell line²². Further study is needed to investigate the possibility of Lamesticum A to be combined with doxorubicin for its potential to have synergistic effect.

Conclusions

Extract, fractions and Lamesticum A derived from the peel of *L. domesticum* showed cytotoxic activity against the T47D breast cancer cell line. Further research is needed to investigate the potential of the natural compound Lamesticum A derived from *L. domesticum* fruit peel as an anticancer therapy.

Data availability

Underlying data

Zenodo: A bioactive compound isolated from Duku (*Lansium domesticum* Corr) fruit peels exhibits cytotoxicity against T47D cell line, <http://doi.org/10.5281/zenodo.3539670>²³.

This project contains the following underlying data:

- UV, infrared, ¹³C-NMR and ¹H-NMR spectra of compound 1.
- Cell viability and IC_{50} values of extract A, fractions B and C, compound 1 and doxorubicin in T47D cell line.

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/) (CC-BY 4.0).

References

1. WHO: **Cancer country profiles 2014**. 2014; (Accessed: 20th April 2018). [Reference Source](#)
2. Heyne K: **Tumbuhan berguna Indonesia**. Badan Litbang Kehutanan, 1988. [Reference Source](#)
3. Manosroi A, Jantrawut P, Sainakham M, *et al.*: **Anticancer activities of the extract from Longkong (*Lansium domesticum*) young fruits**. *Pharm Biol.* 2012; 50(11): 1397–407. [PubMed Abstract](#) | [Publisher Full Text](#)
4. Klungsunya P, Suthepakul N, Muangman T, *et al.*: **Determination of Free Radical Scavenging, Antioxidative DNA Damage Activities and Phytochemical**

- Components of Active Fractions from *Lansium domesticum* Corr. Fruit.**
Nutrients. 2015; **7**(8): 6852–73.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Dong SH, Zhang CR, Dong L, *et al.*: **Onoceranoid-type triterpenoids from *Lansium domesticum*.** *J Nat Prod*. 2011; **74**(5): 1042–1048.
[PubMed Abstract](#) | [Publisher Full Text](#)
 6. Matsumoto T, Kitagawa T, Teo S, *et al.*: **Structures and Antimutagenic Effects of Onoceranoid-Type Triterpenoids from the Leaves of *Lansium domesticum*.** *J Nat Prod*. 2018; **81**(10): 2187–2194.
[PubMed Abstract](#) | [Publisher Full Text](#)
 7. Ashok kumar R, Ramaswamy M: **Phytochemical screening by FTIR spectroscopic analysis of leaf extracts of selected Indian Medicinal plants.** *Int J Curr Microbiol App Sci*. 2014; **3**(1): 395–406.
[Reference Source](#)
 8. Mae Sri Hartati W, Mubarika S, Gandjar IG, *et al.*: **Phalerin, a new benzophenoid glucoside isolated from the methanolic extract of Mahkota Dewa [*Phaleria macrocarpa* (Scheff.) Boerl] leaves.** *Maj Farm Indones*. 2005; **16** (1): 51–57.
[Reference Source](#)
 9. Bahuguna A, Khan I, Bajpai VK, *et al.*: **MTT assay to evaluate the cytotoxic potential of a drug.** *Bangladesh J Pharmacol*. 2017; **12**: 115–118.
[Reference Source](#)
 10. Yu S, Kim T, Yoo KH, *et al.*: **The T47D cell line is an ideal experimental model to elucidate the progesterone-specific effects of a luminal A subtype of breast cancer.** *Biochem Biophys Res Commun*. 2017; **486**(3): 752–758.
[PubMed Abstract](#) | [Publisher Full Text](#)
 11. Kuete V, Ndontsa BL, Nguokeu YMM, *et al.*: **Ardisinol III, a naturally occurring alkenylmethylresorcinol displayed cytotoxic effects in carcinoma cells.** *Investig Med Chem Pharmacol*. 2018; **1**: 1–6.
[Reference Source](#)
 12. Xu R, Ye Y, Zhao W: **Introduction to natural products chemistry.** 2011.
[Reference Source](#)
 13. Nishizawa M, Nademoto Y, Sastrapradja S, *et al.*: **Structure of dukunolide A: A tetranortriterpenoid with a new carbon skeleton from *Lansium domesticum*.** *J Chem Soc Chem Commun*. 1985; 395–396.
[Publisher Full Text](#)
 14. Nishizawa M, Nishide H, Hayashi Y, *et al.*: **The structure of Lansioside A : A novel triterpene glycoside with amino-sugar from *Lansium domesticum*.** *Tetrahedron Lett*. 1982; **23**(13): 1349–1350.
[Publisher Full Text](#)
 15. Sharma SH, Thulasigam S, Nagarajan S: **Terpenoids as anti-colon cancer agents - A comprehensive review on its mechanistic perspectives.** *Eur J Pharmacol*. 2017; **795**: 169–178.
[PubMed Abstract](#) | [Publisher Full Text](#)
 16. Kashyap D, Sharma A, Tuli HS, *et al.*: **Molecular targets of celestrol in cancer: Recent trends and advancements.** *Crit Rev Oncol Hematol*. 2018; **128**: 70–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
 17. Rzeski W, Stepulak A, Szymański M, *et al.*: **Betulinic acid decreases expression of bcl-2 and cyclin D1, inhibits proliferation, migration and induces apoptosis in cancer cells.** *Naunyn Schmiedebergs Arch Pharmacol*. 2006; **374**(1): 11–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
 18. Zhao M, Da-Wa ZM, Guo DL, *et al.*: **Cytotoxic triterpenoid saponins from *Clematis tangutica*.** *Phytochemistry*. 2016; **130**: 228–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
 19. Shi ZN, Wang YD, Gong Y, *et al.*: **New triterpenoid saponins with cytotoxic activities from *Ligularia przewalskii*.** *Phytochem Lett*. 2019; **30**: 215–219.
[Publisher Full Text](#)
 20. Lagoa R, Silva J, Rodrigues JR, *et al.*: **Advances in phytochemical delivery systems for improved anticancer activity.** *Biotechnol Adv*. 2019; pii: S0734-9750(19)30063-1.
[PubMed Abstract](#) | [Publisher Full Text](#)
 21. Prasad S, Tyagi AK, Aggarwal BB: **Chemosensitization by Ursolic Acid: A New Avenue for Cancer Therapy.** In *Role of Nutraceuticals in Chemoresistance to Cancer*. 2018; **2**: 99–109.
[Publisher Full Text](#)
 22. Zong L, Cheng G, Liu S, *et al.*: **Reversal of multidrug resistance in breast cancer cells by a combination of ursolic acid with doxorubicin.** *J Pharm Biomed Anal*. 2019; **165**: 268–275.
[PubMed Abstract](#) | [Publisher Full Text](#)
 23. Fadhilah K, Wahyuono S, Astuti P: **A bioactive compound isolated from Duku (*Lansium domesticum* Corr) fruit peels exhibits cytotoxicity against T47D cell line [Data set].** *Zenodo*. 2019.
<http://www.doi.org/10.5281/zenodo.3539670>

Open Peer Review

Current Peer Review Status: ?

Version 1

Reviewer Report 26 August 2020

<https://doi.org/10.5256/f1000research.23191.r70005>

© 2020 Banjerdpongchai R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

? **Ratana Banjerdpongchai** 

Department of Biochemistry, Chiang Mai University, Chiang Mai, Thailand

Major points:

The extracts and purified bioactive compound, Lamesticum A, from this plant, is not novel, since it was isolated from the twigs from this same herb in the previous report. The anticancer activity of the active compound is not promising since it is more than 4 microgram/mL or 10 micromolar, the level of which the National Cancer Institution defines is 4 microgram/mL or less than this. However, the authors discussed of using this compound as a chemosensitizing agent, which various steps still are needed before launching it as a therapeutic anti-cancer drug. The authors should explore the cytotoxic effects of the extracts, EtOAc extract (A), N-hexane fraction (B), N-hexane insoluble fraction (C) and compound 1 (Lamesticum A) on normal cells.

In Figure 3, the percent cell viabilities of EtOAc extract (A), N-hexane fraction (B), N-hexane insoluble fraction (C) and compound 1 (Lamesticum A) should be presented with the statistical significance by using asterisks, such as * $p < 0.05$, ** $p < 0.01$, and so on, above the dots.

Minor points:

The sources of chemicals and instruments should indicate the city and country that they are produced from. Sometimes the authors used gr and sometimes g, it should be consistent with each other. In subtopic "Cytotoxic assay", in 3rd – 4th lines, the authors should rewrite the sentences. There are many grammar and various typological errors, such as the number of g it should be 43.4 rather than 43,4, etc. For the previous findings, the sentences should be in present tense, but for the authors' research data in the Results and Discussion parts, the sentences would be in past tense. For MDA231, it should be MDA-MB-231 breast cancer cells. For MCF, it should be clarified as MCF-7 or not? In Ref. No. 12, there is no information of company, city, country of publication, is it a book? Ref. No. 21, the name of journal is full name, whereas others are in abbreviated forms. It should be consistent with each other.

References

1. Manosroi A, Jantrawut P, Sainakham M, Manosroi W, et al.: Anticancer activities of the extract from Longkong (*Lansium domesticum*) young fruits. *Pharm Biol.* 2012; **50** (11): 1397-407 [PubMed](#)

[Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Programmed cell death, Natural products, Antioxidants, Cancer.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research