

QUASI-CONVEX UNIVALENT FUNCTIONS

K. INAYAT NOOR and D.K. THOMAS

Kerman University
P. O. Box 182
Kerman, Iran
University College of Swansea
SWANSEA SA2 8PP, Wales

(Received May 9, 1979 and in Revised form in June 25, 1979)

ABSTRACT. In this paper, a new class of normalized univalent functions is introduced. The properties of this class and its relationship with some other subclasses of univalent functions are studied. The functions in this class are close-to-convex.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES: Primary 30A32; Secondary 30A34

KEYWORDS AND PHRASES. Univalent functions, Quasi-convex, Close-to-convex.

1. INTRODUCTION.

Denote by S the class of functions f which are regular and univalent in the unit disc E and satisfy $f(0) = 0$ and $f'(0) = 1$. The subclasses S^* and C of star-like and convex functions respectively are well known and have been extensively studied. S^* and C are connected by the basic property

$$f \in C \text{ if and only if } zf' \in S^*. \quad (1.1)$$

The subclass K of S consisting of close-to-convex function is also well known and many of the properties of S^* can be extended to the wider class K .

The purpose of this paper is to introduce a natural analogue of the class C in terms of the property defined in (1.1).

2. MAIN RESULTS.

Def. Let f be regular in E with $f(0) = 0$ and $f'(0) = 1$. Then f is said to be *quasi-convex* in E if there exists a convex function g with $g(0) = 0$, $g'(0) = 1$ such that for $z \in E$,

$$\operatorname{Re} \frac{(zf'(z))'}{g'(z)} > 0. \quad (2.2)$$

Denote the class of quasi-convex functions by Q .

It is clear that, when $f(z) \equiv g(z)$, $C = Q$ so that $C \subset Q$. We show first that $Q \subset K$, so that every quasi-convex functions is univalent.

THEOREM 1. Let $f \in Q$. Then, for $z \in E$,

$$\operatorname{Re} \frac{zf'(z)}{g(z)} > 0,$$

and so $Q \subset K \subset S$; thus, every quasi-convex function is close-to-convex and hence univalent in E .

PROOF: A result of Libera [4] shows that, if s and t are functions regular in E with $s(0) = t(0) = 0$ and $t \in S^*$, then for $z \in E$,

$$\operatorname{Re} \frac{s'(z)}{t'(z)} > 0 \rightarrow \operatorname{Re} \frac{s(z)}{t(z)} > 0.$$


An immediate application of this with $s(z) = zf'(z)$ and $t(z) = g(z)$ proves

Theorem 1.

It follows at once from the definition that

$$f \in Q \text{ if and only if } zf' \in K. \tag{2.2}$$

We can thus write



where the direction of the arrow indicates set inclusion.

Theorem 1 shows that the image domain for all $f \in Q$ is close-to-convex. However a specific characterisation of the image domain for $f \in Q$ remains an open question.

We state now some basic properties of quasi-convex functions which can easily be extended from the class of convex functions. We omit the proofs as they are simple extensions from the convex case.

THEOREM 2. Let $f \in Q$ with $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$. Then for $|z| = r < 1$,

$$(i) |a_n| \leq 1, \quad n = 2, 3, \dots,$$

$$(ii) \frac{1}{(1+r)^2} \leq |f'(z)| \leq \frac{1}{(1-r)^2},$$

$$(iii) \frac{r}{1+r} \leq |f(z)| \leq \frac{r}{1-r},$$

$$(iv) |w| \geq \frac{1}{2}, \text{ where } f(z) \neq w \text{ in } E.$$

All inequalities are sharp, equality being attained for $f_0(z) = \frac{z}{1-z}$.

We now give an example of a function in Q which is not convex.

Example 1.3.1 Let f_1 be the Koebe function; i.e., $f_1(z) = \frac{z}{(1-z)^2}$. Then f_1 maps E 1-1 conformally onto the w -plane cut from $-\frac{1}{4}$ to $-\infty$ along the negative real axis. Let $f_2: f_2(z) = \frac{x+z}{1+\bar{x}z}$, $x \in E$. Then f_2 maps E onto itself and takes the origin onto the point x . Define f_3 by

$$f_3(z) = f_1[f_2(z)] - f_1(x); \quad f_3(0) = 0 \quad \text{and}$$

$f_3'(0) \neq 0$ (since f_3 is univalent), and let f_4 :

$$f_4(z) = \frac{f_3(z)}{f_3'(0)}, \quad z \in E.$$

Combining all these transformations, we can write

$$F(z) = \frac{f_1\left(\frac{x+z}{1+\bar{x}z}\right) - f_1(x)}{f_1'(x)(1-|x|^2)}; \quad f_1(z) = \frac{z}{(1-z)^2}.$$

The function F is close-to-convex. In fact,

$$F'(z) = \frac{f_1'\left(\frac{x+z}{1+\bar{x}z}\right)}{f_1'(x)(1+\bar{x}z)^2}$$

$$\frac{F''(z)}{F'(z)} = \frac{f_1''\left(\frac{x+z}{1+\bar{x}z}\right)(1-|x|^2)}{f_1'\left(\frac{x+z}{1+\bar{x}z}\right)(1+\bar{x}z)^2} - \frac{2\bar{x}}{(1+\bar{x}z)}$$

and

$$\begin{aligned}
 1 + \frac{zF''(z)}{F'(z)} &= \left\{ 1 + \left(\frac{x+z}{1+\bar{x}z} \right) \frac{f_1''\left(\frac{x+z}{1+\bar{x}z}\right)}{f_1'\left(\frac{x+z}{1+\bar{x}z}\right)} \right\} \frac{z(1-|x|^2)}{(x+z)(1+\bar{x}z)} \\
 &\quad - \frac{2\bar{x}z}{1+\bar{x}z} - \frac{z(1-|x|^2)}{(x+z)(1+\bar{x}z)} + 1 \\
 &= \left\{ 1 + \frac{x+z}{1+\bar{x}z} \frac{f_1''\left(\frac{x+z}{1+\bar{x}z}\right)}{f_1'\left(\frac{x+z}{1+\bar{x}z}\right)} \right\} \frac{z(1-|x|^2)}{(x+z)(1+\bar{x}z)} + \frac{x-\bar{x}z^2}{(x+z)(1+\bar{x}z)} \\
 &= \left\{ 1 + \frac{\xi f_1''(\xi)}{f_1'(\xi)} \right\} \frac{(1-|x|^2)z}{(x+z)(1+\bar{x}z)} + \frac{x-\bar{x}z^2}{(x+z)(1+\bar{x}z)}, \quad \xi = \frac{x+z}{1+\bar{x}z}
 \end{aligned}$$

Let $z = re^{i\theta}$, $\frac{x+re^{i\theta}}{1+\bar{x}re^{i\theta}} = r_1 e^{i\varphi}$; $\frac{re^{i\theta}(1-|x|^2)}{(x+re^{i\theta})(1+\bar{x}re^{i\theta})} d\theta = d\varphi$.

For φ_1 and φ_2 with $(\varphi_1 < \varphi_2)$, we have corresponding θ_1 and θ_2 with $(\theta_1 < \theta_2)$.

Now,

$$\begin{aligned}
 \operatorname{Re} \left\{ 1 + re^{i\theta} \frac{F''(re^{i\theta})}{F'(re^{i\theta})} \right\} d\theta &= \operatorname{Re} \left\{ 1 + \frac{\xi f_1''(\xi)}{f_1'(\xi)} \right\} d\varphi; \quad \xi = r_1 e^{i\varphi} \\
 &= \operatorname{Re} \left\{ 1 + r_1 e^{i\varphi} \frac{f_1''(r_1 e^{i\varphi})}{f_1'(r_1 e^{i\varphi})} \right\} d\varphi.
 \end{aligned}$$

Hence, for $\phi_1 < \phi_2$ and $\theta_1 < \theta_2$,

$$\int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ 1 + r e^{i\theta} \frac{F''(r e^{i\theta})}{F'(r e^{i\theta})} \right\} d\theta = \int_{\phi_1}^{\phi_2} \operatorname{Re} \left\{ 1 + r_1 e^{i\phi} \frac{f_1''(r_1 e^{i\phi})}{f_1'(r_1 e^{i\phi})} \right\} d\phi > -\pi,$$

which shows that $f \in K$.

Now,

$$\begin{aligned} F'(z) &= \left(\frac{1 + \frac{x+z}{1+xz}}{1 - \frac{x+z}{1+xz}} \right)^3 \cdot \frac{(1-x)^3}{(1+x)} \cdot \frac{1}{(1+xz)^2} \\ &= \frac{1 + \frac{1+x}{1+x}z}{\left(1 - \frac{1-x}{1-x}z \right)^3} \\ &= \frac{(1+\beta z)}{(1-\alpha z)^3}; \quad \beta = \frac{1+x}{1+x}, \quad \alpha = \frac{1-x}{1-x}; \quad |\alpha| = 1 = |\beta|. \end{aligned}$$

Integrating, we have for $z \in E$,

$$F(z) = \frac{z(1 + \frac{\beta-\alpha}{2}z)}{(1-\alpha z)^2}.$$

We notice that F maps E onto the w -plane cut along a half-line ℓ . Since the choice of the point x in E is arbitrary, we can select x in such a way that the half-line ℓ does not pass through the origin in $F(E)$, which means F is not, in general, starlike. Because of relationships (3) and (1) between the classes Q and $K - C$ and S^* , we conclude that, in general,

the function f_* defined in E by $f_*(z) = \int_0^2 \frac{F(\xi)}{\xi} d\xi$ belongs to Q but not to C .

3. SOME GROWTH PROBLEMS

Clunie and Keogh [1] showed that, if $f \in C$ with $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ and $f(E)$ has finite area. Then $n a_n = o(1)$ as $n \rightarrow \infty$ and the exponent is best possible. We extend this result to quasi-convex functions.

THEOREM 3. Let $f \in Q$ with $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$. If $f(E)$ has finite area, then $n a_n = o(1)$ as $n \rightarrow \infty$, the index of n being best possible.

PROOF: We use a modified version of the method of Clunie and Pommerenke [2].

By (2), we can write

$$(zf'(z))' = g'(z)h(z), \tag{3.1}$$

where $\text{Re } h(z) > 0$ for $z \in E$ and $h(0) = 1$. Thus,

$$z(zf'(z))' = 2zg'(z) \text{Re } h(z) - zg'(z) \overline{h(z)},$$

and so with $z = re^{i\theta}$, $0 \leq r < 1$, Cauchy's formula gives for $n \geq 1$

$$\begin{aligned} n^2 a_n &= \frac{1}{2\pi r^n} \int_0^{2\pi} z(zf'(z))' e^{-in\theta} d\theta \\ &= \frac{1}{\pi r^n} \int_0^{2\pi} zg'(z) \text{Re } h(z) e^{-in\theta} d\theta - \frac{1}{2\pi r^n} \int_0^{2\pi} zg'(z) \overline{h(z)} e^{-in\theta} d\theta. \end{aligned}$$

Since $\text{Re } h(z) > 0$ for $z \in E$,

$$n^2 |a_n| \leq \frac{1}{\pi r^n} \int_0^{2\pi} |zg'(z)| \text{Re } h(z) d\theta + \frac{1}{2\pi r^n} \left| \int_0^{2\pi} zg'(z) h(z) e^{in\theta} d\theta \right|. \tag{3.2}$$

$$\text{Now } |zg'(z)| \operatorname{Re} h(z) = \operatorname{Re} [|zg'(z)| h(z)]$$

= $\operatorname{Re} [z(zf'(z))' e^{-i \arg zg'(z)}]$ from (3.1), and so integrating the first of the above two integrals by parts we have

$$\begin{aligned} \frac{1}{\pi r^n} \int_0^{2\pi} |zg'(z)| \operatorname{Re} h(z) d\theta &= \operatorname{Re} \frac{1}{\pi r^n} \left[\int_0^{2\pi} z(zf'(z))' e^{-i \arg zg'(z)} d\theta \right] \\ &= \operatorname{Re} \frac{1}{\pi r^n} \int_0^{2\pi} z f'(z) e^{-i \arg zg'(z)} d_\theta(\arg zg'(z)) \end{aligned} \quad (3.3)$$

Also, $\overline{zg'(z)} h(z) = z(zf'(z))' e^{-2i \arg zg'(z)}$ and so (3.2) and (3.3) give

$$\begin{aligned} n^2 |a_n| &\leq \frac{1}{\pi r^n} \int_0^{2\pi} \operatorname{Re} \left[z f'(z) e^{-i \arg zg'(z)} \right] d_\theta(\arg zg'(z)) \\ &\quad + \frac{1}{2\pi r^{2n}} \left| \int_0^{2\pi} z^{n+1} (zf'(z))' e^{-2i \arg zg'(z)} d\theta \right| \\ &= \frac{1}{\pi r^n} I_1 + \frac{1}{2\pi r^{2n}} I_2, \text{ say.} \end{aligned} \quad (3.4)$$

To estimate I_1 , we note that, since f is regular in E and the area of $f(E)$ is finite, $M(r, f') = \frac{o(1)}{1-r}$ as $r \rightarrow 1$, where $M(r, f) = \max_{\theta} |f(re^{i\theta})|$. Since $\int_0^{2\pi} d_\theta(\arg zg'(z)) = 2\pi$, we have $I_1 = \frac{o(1)}{1-r}$ as $r \rightarrow 1$.

Integrating I_2 by parts gives

$$I_2 = 2 \left| \int_0^{2\pi} F_n(z) e^{-2i \arg z} g'(z) \operatorname{Re} \frac{(zg'(z))'}{g'(z)} d\theta \right|,$$

where

$$F_n(z) = \int_0^z t^n (tf'(t))' dt = z^{n+1} f'(z) - n f_n(z),$$

and

$$f_n(z) = \int_0^z t^n f'(t) dt.$$

Now $|F_n(z)| \leq r^{n+1} M(r, f') + n \int_0^r t^n dt$

$$\leq 2r^{n+1} M(r, f').$$

Since $\int_0^{2\pi} \operatorname{Re} \frac{(zg'(z))'}{g'(z)} d\theta = 2\pi$, we have

$$I_2 \leq 8\pi r^{n+1} M(r, f') = \frac{o(1)}{1-r} \text{ as } r \rightarrow 1 \text{ as before.}$$

Finally, choosing $r = 1 - \frac{1}{n}$ in (3.4), the estimates for I_1 and I_2 give $na_n = o(1)$ as $n \rightarrow \infty$, and Theorem 3 is proved.

An examination of the proof of Theorem 3 gives

COROLLARY: Let $f \in Q$, $E_r = \{z: |z| = r < 1\}$ and $A(r)$ be the area of $f(E_r)$.

Then, for $n \geq 2$,

$$n|a_n| = o(1) A(1 - \frac{1}{n})^{\frac{1}{2}}. \quad (3.5)$$

We remark that (3.5) holds for the class S^* , but appears still to be an open problem for the class K .

Denote by $C(r)$ the closed curve which is the image of $f(E_r)$ and by $L(r)$ the length of $C(r)$. We prove

THEOREM 4. Let $f \in Q$. Then, for $0 \leq r \leq 1$,

$$2\sqrt{(\pi A(r))} \leq L(r) \leq 2\sqrt{(\pi A(\sqrt{r}))} (\log \frac{1}{1-r})^{\frac{1}{2}}. \quad (3.6)$$

Further, if $A(r) < \infty$ for $0 \leq r < 1$, then

$$L(r) = o(1) (\log \frac{1}{1-r})^{\frac{1}{2}} \text{ as } r \rightarrow 1.$$

PROOF: The left hand inequality follows at once from the isoperimetric inequality. Since $f \in Q$, $F(z) = zf'(z)$ is close-to-convex. Thus from [3,p.45]

$$\begin{aligned} L(r) &= \int_0^{2\pi} |zf'(z)| d\theta = \int_0^{2\pi} |B(z)| d\theta \\ &\leq 2\pi \int_0^r M(\rho, zf') \frac{d\rho}{\rho} \\ &\leq 2\pi \sum_{n=1}^{\infty} |a_n| r^n \\ &\leq 2\pi \left(\sum_{n=1}^{\infty} n|a_n|^2 r^n \right)^{\frac{1}{2}} \left(\sum_{n=1}^{\infty} \frac{r^n}{n} \right)^{\frac{1}{2}} \\ &= 2\sqrt{(\pi A(\sqrt{r}))} \log \frac{1}{1-r}^{\frac{1}{2}}. \end{aligned}$$

If $a(r) < \infty$ for $0 \leq r < 1$, then from (3.7)

$$\begin{aligned} L(r) &\leq 2\pi \sum_{n=1}^{\infty} |a_n| r^n \\ &\leq 2\pi \sum_{n=1}^N |a_n| r^n + \sum_{n=N}^{\infty} n |a_n|^2 \frac{1}{2} \sum_{n=1}^{\infty} \frac{r^{2n}}{n} \\ &= 2\pi \sum_{n=1}^N |a_n| r^n + \delta_N \log \frac{1}{1-r^2} \end{aligned}$$

where $\delta_N \rightarrow 0$ as $N \rightarrow \infty$. Thus $L(r) = o(1) \log \frac{1}{1-r}$ as $r \rightarrow 1$.

The convex function $f_1(z) = \log \frac{1}{1-z}$ shows that the factor $\log \frac{1}{1-r}$ in

(3.6) is best possible.

For $f \in C$ it is well known that $L(r) \leq 2\pi M(r)$. It follows from (3.6) that for $f \in Q$, $L(r) = o(1) M(r) \log \frac{1}{1-r}$ as $r \rightarrow 1$. The question of whether the factor $\log \frac{1}{1-r}$ can be removed remains open.

In conclusion, we remark that other results for the class C can be extended to quasi-convex functions, often with only minor alterations in the proof. The objective of this paper has been to introduce the class Q , exhibit its basic properties and give some results whose proofs are not trivial extensions from the class C .

REFERENCES

1. CLUNIE, J.G. and KEOGH, F.R., "On starlike and convex schlicht functions", J. London Math.Soc., 35 (1960), 229-233.
2. CLUNIE, J.G. and POMMERENKE, Ch., "On the coefficients of close-to-convex univalent functions", J. London Math.Soc., 41 (1966), 161-165.
3. HAYMAN, W.K., 'Multivalent functions', Cambridge, 1967.
4. LIBERA, R.J., "Some classes of regular univalent functions", Proc.Amer. Math Soc., 16 (1965), 755-758.

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mssklai@cityu.edu.hk