

CORRIGENDUM

Clinical Implication of Energy Loss Coefficient in Patients With Severe Aortic Stenosis Diagnosed by Doppler Echocardiography

Teruyoshi Kume, MD; Hiroyuki Okura, MD; Takahiro Kawamoto, MD; Nozomi Watanabe, MD;
Akihiro Hayashida, MD; Yoji Neishi, MD; Yoshinori Miyamoto, MD;
Koichiro Imai, MD; Ryotaro Yamada, MD; Kiyoshi Yoshida, MD

(*Circ J* 2008; **72**: 1265–1269)

The author (T. Kume) apologizes for errors which appeared in the Table 2, page 1266, Vol 72. Aortic cross-sectional area was $5.8 \pm 1.5 \text{ cm}^2$ in group A and $5.5 \pm 0.8 \text{ cm}^2$ in group B. The corrected Table 2 is given below.

Table 2 Hemodynamic and Echocardiographic Data

	Group A (n=26)	Group B (n=7)	p value
<i>Hemodynamic data</i>			
LV maximum pressure (mmHg)	197±34	191±25	0.659
LV end-diastolic pressure (mmHg)	20±6	19±5	0.698
Ascending aorta maximum pressure (mmHg)	140±25	161±31	0.061
Ascending aorta minimum pressure (mmHg)	65±12	66±13	0.836
Cardiac index ($\text{L} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$)	3.0±0.9	3.8±1.3	0.107
Pulmonary capillary wedge pressure (mmHg)	13±6	12±3	0.804
Pulmonary artery maximum pressure (mmHg)	32±12	33±9	0.915
Pulmonary artery minimum pressure (mmHg)	15±7	14±3	0.815
<i>Echocardiographic data</i>			
LV diastolic dimension (mm)	44±6	42±6	0.628
LV systolic dimension (mm)	28±7	27±6	0.888
LV mass index (g/m^2)	195±73	180±51	0.626
LV ejection fraction (%)	64±9	61±8	0.445
Aortic cross-sectional area (cm^2)	5.8±1.5	5.5±0.8	0.631

LV, left ventricular.