

Dielectric Boundary Force in Molecular Solvation with the Poisson–Boltzmann Free Energy: A Shape Derivative Approach

Bo Li* Xiaoliang Cheng[†] Zhengfang Zhang[‡]

November 11, 2011

Abstract

In an implicit-solvent description of molecular solvation, the electrostatic free energy is given through the electrostatic potential. This potential solves a boundary-value problem of the Poisson–Boltzmann equation in which the dielectric coefficient changes across the solute-solvent interface—the dielectric boundary. The dielectric boundary force acting on such a boundary is the negative first variation of the electrostatic free energy with respect to the location change of the boundary. In this work, the concept of shape derivative is used to define such variations and formulas of the dielectric boundary force are derived. It is shown that such a force is always in the direction toward the charged solute molecules.

Keywords: implicit solvent, the Poisson–Boltzmann equation, dielectric boundary force, shape derivative.

1 Introduction

We consider electrostatic interactions in the solvation of molecules within the framework of widely used implicit-solvent or continuum-solvent modeling [9, 19, 24, 45, 53]. In such a model, the solvent molecules and ions are treated implicitly and their effects are coarse grained. Most of the existing implicit-solvent models are based on various kinds of fixed solute-solvent interfaces, such as the van der Waals surface, solvent-excluded surface, or

*Department of Mathematics and NSF Center for Theoretical Biological Physics, University of California, San Diego, 9500 Gilman Drive, Mail code: 0112, La Jolla, CA 92093-0112, U.S.A. Email: bli@math.ucsd.edu.

[†]Department of Mathematics, Zhejiang University, Hangzhou, Zhejiang, 310027, P. R. China. Email: xiaoliangcheng@zju.edu.cn.

[‡]Department of Mathematics, Hangzhou Dianzi University, Hangzhou, Zhejiang, 310018, P. R. China. Email: zhengfangzhang@hdu.edu.cn.

solvent-accessible surface [17, 18, 35, 43, 44]. Such a predefined interface is used to compute the solvation free energy as the sum of two separate parts. One is the surface energy, proportional to the area of interface. The other is the electrostatic contribution determined by the Poisson–Boltzmann (PB) [1, 7, 20, 25–27, 30, 31, 36, 37, 39, 49, 58] or generalized Born (GB) [2, 3, 52] approach in which the solute-solvent interface is used as the dielectric boundary.

In recent years, a new class of implicit-solvent models, termed variational implicit-solvent models (VISM), have emerged [22, 23]. Coupled with the robust level-set numerical method [41, 42, 46], such models allow an efficient and quantitative description of molecular solvation [12, 13, 15]. Central in the VISM is a free-energy functional of all possible solute-solvent interfaces, or dielectric boundaries, that separate the continuum solvent from all solute atoms. In a simple setting, such a free-energy functional consists of surface energy of solute molecules, solute-solvent van der Waals interaction energy, and continuum electrostatic free energy, all coupled together and depending solely on a given solute-solvent interface. Minimizing the functional determines the solvation free energy and stable equilibrium solute-solvent interfaces. Initial applications of the level-set VISM to nonpolar molecular systems have demonstrated its success in capturing the hydrophobic interaction, multiple equilibrium states of hydration, and fluctuation between such states [12, 14, 15, 47, 57], all of which are difficult to be captured by a fixed-surface implicit-solvent model. See [4, 9, 10, 24, 56] for other related models and methods.

In this work, we study the dielectric boundary force—the normal component of such a force, to be more precisely—acting on a dielectric boundary or solute-solvent interface. Such a force is the negative variation of the electrostatic free energy with respect to the location change of the dielectric boundary. It is the electrostatic part of the total boundary force associated with the VISM free-energy functional, determining the conformation and dynamics of an underlying molecular system with an implicit solvent. Practically, it is also the electrostatic part of the “normal velocity” in the level-set relaxation of the free-energy functional.

For a given solvation system with a fixed charge density and dielectric coefficient, any given, possible dielectric boundary Γ determines the electrostatic potential $\psi = \psi_\Gamma$ as the unique solution to the nonlinear PB equation, which in turn determines the electrostatic free energy $G[\Gamma]$. Appealing the notion and method of shape derivatives, we give a precise definition of the dielectric boundary force and derive formulas for such a force. We notice that in [57] the dielectric boundary force with the Coulomb-field approximation of electrostatic free energy is derived and implemented in the level-set VISM for charged molecules. In this approach, there is no need to estimate the GB radii as usually done in a GB model which is also based on the Coulomb-field approximation in a simple setting. In [11], the Yukawa-field approximation of the electrostatic free energy is proposed and the formula of the corresponding dielectric boundary force is derived. These approaches require no solutions to any partial differential equations. In comparison, our current approach is more accurate analytically but can be less efficient computationally. Related work on electrostatic forces in molecular systems can be found in [29, 32, 38].

We assume that the entire solvation system occupies a bounded region $\Omega \subset \mathbb{R}^3$. It is divided into three disjoint parts: the region of solute Ω_- (e.g., charged biomolecules such as proteins), the region of solvent Ω_+ (e.g., salted water), and the solute-solvent interface or the dielectric boundary Γ that separates Ω_- and Ω_+ . See Figure 1, where n denotes the unit normal to the boundary Γ pointing from Ω_- to Ω_+ and also the exterior unit normal to $\partial\Omega$, the boundary of Ω . The solute region Ω_- is completely contained in the entire system region Ω , i.e., $\overline{\Omega_-} \subset \Omega$, where an over-line denotes the closure. The solute region Ω_- contains all the solute atoms located at X_1, \dots, X_N , carrying charges Q_1, \dots, Q_N , respectively.




Figure 1: The geometry of a solvation system with an implicit solvent. Dots represent solute atoms at X_i carrying charge Q_i ($i = 1, \dots, N$). The solute-solvent interface Γ separates the solute region Ω_- and the solvent region Ω_+ . The corresponding dielectric coefficients are denoted by ε_- and ε_+ , respectively. The unit normal at the interface Γ pointing from Ω_- to Ω_+ and the exterior unit normal at the boundary of the entire solvation region Ω are both denoted by n .

In the PB theory, the electrostatic part of the solvation free energy—the electrostatic free energy—is given by

$$G[\Gamma] = \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma}}{2} |\nabla\psi|^2 + f\psi - \chi_+ \beta^{-1} \sum_{j=1}^M c_j^{\infty} (e^{-\beta q_j \psi} - 1) \right] dX \quad (1.1)$$

through the electrostatic potential $\psi : \Omega \rightarrow \mathbb{R}$ [8, 20, 29, 37, 49]. Here, $\varepsilon_{\Gamma} : \Omega \rightarrow \mathbb{R}$ is the dielectric coefficient defined by

$$\varepsilon_{\Gamma}(X) = \begin{cases} \varepsilon_- & \text{if } X \in \Omega_-, \\ \varepsilon_+ & \text{if } X \in \Omega_+, \end{cases} \quad (1.2)$$

where ε_- and ε_+ are two positive constants. We have $\varepsilon_+ = \varepsilon'_+ \varepsilon_0$ and $\varepsilon_- = \varepsilon'_- \varepsilon_0$, respectively, where ε_0 is the vacuum permittivity, and ε'_+ and ε'_- are the temperature-dependent relative permittivities. Typically, the value of ε'_- is in between 1 and 10 for proteins and that of ε'_+ is close to 80 for water at normal conditions. The function $f : \Omega \rightarrow \mathbb{R}$ is the fixed charge density of charged solute molecules. It is usually the sum of the point charges Q_i located at X_i ($i = 1, \dots, N$). Here we assume that f is an integrable function that approximates these point charges. The function χ_+ is the characteristic function of Ω_+ defined by $\chi_+(X) = 1$ if $X \in \Omega_+$ and $\chi_+(X) = 0$ otherwise. The parameter $\beta > 0$ is the inverse thermal energy, $M \geq 2$ is the number of ionic species in the solvent, and $q_j \in \mathbb{R}$ and $c_j^\infty > 0$ are the charge and bulk concentration, respectively, of the j th ionic species with $j = 1, \dots, M$. Note that all $\varepsilon_+, \varepsilon_-, f, \beta$, and q_j and c_j^∞ ($1 \leq j \leq M$) are input data. We use the SI units of electrostatics. The electrostatic potential ψ is the unique solution of a boundary-value problem of the nonlinear PB equation [1, 20, 25, 37, 48]

$$\nabla \cdot \varepsilon_\Gamma \nabla \psi + \chi_+ \sum_{j=1}^M q_j c_j^\infty e^{-\beta q_j \psi} = -f \quad \text{in } \Omega. \quad (1.3)$$

See also [5, 34, 36, 37, 54, 55, 58] for generalized PB equations to include ionic excluded-volume effects. Eq. (1.3) is the Euler–Lagrange equation of the right-hand side of (1.1) viewed as a functional of ψ . Note that this functional is *concave* in ψ .

Let $V : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a smooth map vanishing outside a small neighborhood of the dielectric boundary Γ . Let $x = x(t, X)$ be the solution map of the dynamical system [6, 21, 33, 51]

$$\frac{dx(t, X)}{dt} = V(x(t, X)) \quad \forall t > 0 \text{ small} \quad \text{and} \quad x(0, X) = X \quad \forall X \in \mathbb{R}^3.$$

The shape derivative of the electrostatic free energy $G[\Gamma]$ in the direction of $V : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is defined to be

$$\delta_{\Gamma, V} G[\Gamma] = \left. \frac{d}{dt} G[\Gamma_t(V)] \right|_{t=0},$$

where $\Gamma_t(V) = \{x(t, X) : X \in \Gamma\}$ and $G[\Gamma_t(V)]$ is defined similarly using $\Gamma_t(V)$ instead of Γ . It will be shown that $\delta_{\Gamma, V} G[\Gamma]$ is an integral of the product of $V \cdot n$ and some function on Γ that is independent of V , where n is the unit normal along Γ (cf. Theorem 4.1). We identify that function on Γ as the shape derivative of $G[\Gamma]$ and denote it by $\delta_\Gamma G[\Gamma]$. We define the dielectric boundary force, or more precisely the normal component of the dielectric boundary force, to be $-\delta_\Gamma G[\Gamma]$ and denote it by F_n . Notice that it is only the normal component, not the tangential components of the boundary force, that determines the motion of the boundary.

Our main result is the following formula of the shape derivative $\delta_\Gamma G[\Gamma] : \Gamma \rightarrow \mathbb{R}$ of the electrostatic free energy $G[\Gamma]$ with respect to the dielectric boundary Γ :

$$\delta_\Gamma G[\Gamma] = \frac{\varepsilon_+}{2} |\nabla \psi^+|^2 - \frac{\varepsilon_-}{2} |\nabla \psi^-|^2 - \varepsilon_+ |\nabla \psi^+ \cdot n|^2 + \varepsilon_- |\nabla \psi^- \cdot n|^2 + B(\psi), \quad (1.4)$$

where ψ is the electrostatic potential, a superscript $+$ or $-$ denotes the restriction onto Ω_+ or Ω_- , respectively, and

$$B(s) = \beta^{-1} \sum_{j=1}^M c_j^\infty (e^{-\beta q_j s} - 1) \quad \forall s \in \mathbb{R}. \quad (1.5)$$

Note that $\varepsilon_+ \nabla \psi^+ \cdot n = \varepsilon_- \nabla \psi^- \cdot n$ on Γ and this common value is denoted by $\varepsilon_\Gamma \nabla \psi \cdot n$. The dependence on the direction of n is in $V \cdot n$ in the integral over Γ . See Theorem 4.1 for the details. A different but useful form of the shape derivative $\delta_\Gamma G[\Gamma] : \Gamma \rightarrow \mathbb{R}$ is

$$\delta_\Gamma G[\Gamma] = \frac{1}{2} \left(\frac{1}{\varepsilon_-} - \frac{1}{\varepsilon_+} \right) |\varepsilon_\Gamma \nabla \psi \cdot n|^2 + \frac{1}{2} (\varepsilon_+ - \varepsilon_-) |(I - n \otimes n) \nabla \psi|^2 + B(\psi), \quad (1.6)$$

where I is the identity matrix. See Corollary 4.1. The vector $(I - n \otimes n) \nabla \psi$ is the tangential component of $\nabla \psi$. It is continuous across the boundary Γ . The corresponding term, the middle term, in the boundary force (1.6), may not be small compared with the first term in (1.6), since a solute-solvent interface can be rough.

We remark that our results hold true for the function $B : \mathbb{R} \rightarrow \mathbb{R}$ that is more general than that defined in (1.5). Our formula (1.4) corrects that in [8] (cf. (3.13) in [8]). If we define $E = -\nabla \psi$ and

$$T = \varepsilon_\Gamma E \otimes E - \frac{\varepsilon_\Gamma}{2} |E|^2 I - \chi_+ B(\psi) I \quad \text{in } \Omega,$$

then it is easy to verify by the formula (1.4) that

$$F_n = -\delta_\Gamma G[\Gamma] = n \cdot T^+ n - n \cdot T^- n.$$

The quantity T is the Maxwell stress tensor of our underlying charged molecular system [29, 38, 40].

A direct consequence of our result (1.6) is that, under the assumption $\varepsilon_- < \varepsilon_+$ which is true in general, we always have

$$F_n < 0 \quad \text{on } \Gamma. \quad (1.7)$$

See Corollary 4.1. This gives a quantitative interpretation, in the framework of implicit solvent, of the following phenomenon described by Debye in 1960s [16]: “Under the combined influence of electric field generated by solute charges and their polarization in the surrounding medium which is electrostatic neutral, an additional potential energy emerges and drives the surrounding molecules to the solutes.” A significant consequence of our mathematical statement (1.7) is as follows: A neutral cavity close to a charged solute (e.g., a protein) will move away from the solute due to the dielectric boundary force. This important charge effect to the hydrophobic interaction in biomolecules has been observed in the recent level-set variational implicit-solvent modeling of BphC, a two-domain protein [57].

In the proof of our main results, we use some uniform bounds of the potential ψ . Such bounds are obtained in the proof of the existence and uniqueness of the solution to the boundary-value problem of the PB equation. We also use the variational principle that the electrostatic free energy $G[\Gamma]$ *maximizes* a corresponding functional of ψ for a fixed boundary Γ . In deriving the formula of boundary force, we do not use an abstract lemma as usually done [21, 51]. Rather we give a direct and simpler derivation using the definition of shape derivatives. Notice that we assume in this work that the fixed charge density f is a compactly supported smooth function that approximates point charges. Numerically such approximation can be made by that of the Dirac delta function [50]. Our analysis does not directly extend to the case of point charges which can be in general more difficult to treat. In fact, with point charges in the PB model, a rigorous definition of the electrostatic free energy must be given with caution. This will be our future work.

The rest of the paper is organized as follows: In Section 2, we recall the boundary-value problem of the PB equation and the related electrostatic free energy. In Section 3, we define the dielectric boundary force using the notion of shape derivative. Finally, in Section 4, we derive our main formula of the dielectric boundary force and show that the force is always attractive to solutes.

2 The Poisson–Boltzmann Equation and Electrostatic Free Energy

We make the following assumptions throughout the rest of the paper:

- A1. All Ω , Ω_- , and Ω_+ are non-empty, bounded, and open subsets of \mathbb{R}^3 with $\overline{\Omega_-} \subset \Omega$ and $\Omega_+ = \Omega \setminus \overline{\Omega_-}$. The boundary $\Gamma = \partial\Omega_- = \overline{\Omega_-} \cap \overline{\Omega_+}$. Both $\partial\Omega$ and Γ are of C^2 . The unit normal at Γ pointing from Ω_- to Ω_+ is denoted by n . The unit exterior normal at the boundary of Ω is also denoted by n . See Figure 1.
- A2. $M \geq 2$ is an integer. All $\beta > 0$, $q_j \in \mathbb{R}$ and $c_j^\infty > 0$ ($1 \leq j \leq M$), $\varepsilon_- > 0$, and $\varepsilon_+ > 0$ are constants. The function $\varepsilon_\Gamma \in L^\infty(\Omega)$ is defined in (1.2). The parameters q_j and c_j^∞ ($1 \leq j \leq M$) satisfy the condition of charge neutrality: $\sum_{j=1}^M q_j c_j^\infty = 0$.
- A3. The fixed charge density $f : \Omega \rightarrow \mathbb{R}$ satisfies that $f \in L^2(\Omega)$ and $\text{supp}(f) \subset \Omega_-$. The boundary data in the Dirichlet boundary condition that we use is given by a function $g \in W^{2,\infty}(\Omega)$. Moreover, we use the notation

$$H_g^1(\Omega) = \{\phi \in H^1(\Omega) : \phi = g \text{ on } \partial\Omega\}.$$

Here and below we use the standard notation for the Sobolev space $W^{k,p}(\Omega)$ which, for any fixed integer $k \geq 1$ and any extended real number $p : 1 \leq p \leq \infty$, consists of all k -times (weakly) differentiable functions $u : \Omega \rightarrow \mathbb{R}$ with the p th power of u or any of its derivatives of order $\leq k$ integral over Ω if $1 \leq p < \infty$ or with u and all its derivatives of order $\leq k$ essentially bounded in Ω .

- A4. There exists a smooth function $B : \mathbb{R} \rightarrow \mathbb{R}$ that satisfies $B(s) > B(0) = 0$ for all $s \neq 0$, $B'(0) = 0$, $\inf_{\mathbb{R}} B'' > 0$ and hence B is strictly convex, $B(\pm\infty) = \infty$, and

$B'(\infty) = \infty$ and $B'(-\infty) = -\infty$.

In the usual PB theory, the function $B : \mathbb{R} \rightarrow \mathbb{R}$ is given by (1.5). One easily verifies using the charge neutrality and Jensen's inequality that for this specific form of B , all the properties listed in A4 are satisfied. A general function $B : \mathbb{R} \rightarrow \mathbb{R}$ as we assumed in A4 above covers other cases such as those including ionic size effects [5, 34, 36, 37, 54, 55].

We define $G[\Gamma, \cdot] : H^1(\Omega) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ by

$$G[\Gamma, \phi] = \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma}}{2} |\nabla \phi|^2 + f\phi - \chi_+ B(\phi) \right] dX. \quad (2.1)$$

We consider the maximization of the functional $G[\Gamma, \cdot]$ and the boundary-value problem of the Poisson–Boltzmann (PB) equation

$$\nabla \cdot \varepsilon_{\Gamma} \nabla \psi - \chi_+ B'(\psi) = -f \quad \text{in } \Omega, \quad (2.2)$$

$$\psi = g \quad \text{on } \partial\Omega. \quad (2.3)$$

Theorem 2.1. (1) *The functional $G[\Gamma, \cdot] : H_g^1(\Omega) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ has a unique maximizer $\psi_0 \in H_g^1(\Omega)$, defined by*

$$G[\Gamma, \psi_0] = \max_{\phi \in H_g^1(\Omega)} G[\Gamma, \phi].$$

Moreover, this maximum value is finite and

$$\|\psi_0\|_{H^1(\Omega)} + \|\psi_0\|_{L^\infty(\Omega)} \leq C_1$$

for some constant $C_1 > 0$ depending on $\varepsilon_-, \varepsilon_+, f, g, B$ and Ω but not on Γ .

(2) *The maximizer ψ_0 is the unique solution to the boundary-value problem of the PB equation (2.2) and (2.3).*

We define the electrostatic free energy corresponding to the dielectric boundary Γ to be

$$G[\Gamma] = \max_{\phi \in H_g^1(\Omega)} G[\Gamma, \phi] = G[\Gamma, \psi_0]. \quad (2.4)$$

Proof of Theorem 2.1. Let $u_0 \in H_g^1(\Omega)$ be such that

$$\int_{\Omega} \varepsilon_{\Gamma} \nabla u_0 \cdot \nabla v dX = \int_{\Omega} f v dX \quad \forall v \in H_0^1(\Omega). \quad (2.5)$$

Standard regularity theory implies that $u_0 \in L^\infty(\Omega)$ and there exists a constant $\hat{C}_1 > 0$, depending possibly on $\varepsilon_-, \varepsilon_+, f, g$, and Ω but not on Γ , such that

$$\|u_0\|_{H^1(\Omega)} + \|u_0\|_{L^\infty(\Omega)} \leq \hat{C}_1, \quad (2.6)$$

cf. [28] (Chapter 8). Let $\phi \in H_g^1(\Omega)$ and $u = \phi - u_0 \in H_0^1(\Omega)$. By (2.1), we have

$$G[\Gamma, \phi] = -I[\Gamma, u] + \int_{\Omega} \left(f u_0 - \frac{\varepsilon_{\Gamma}}{2} |\nabla u_0|^2 \right) dX,$$

where

$$I[\Gamma, u] = \int_{\Omega} \left[\frac{\varepsilon_{\Gamma}}{2} |\nabla u|^2 + \chi_+ B(u + u_0) \right] dX.$$

Hence the maximization of $G[\Gamma, \phi]$ over all $\phi \in H_g^1(\Omega)$ is equivalent to the minimization of $I[\Gamma, u]$ over all $u \in H_0^1(\Omega)$.

By the Poincaré inequality and the fact that $B : \mathbb{R} \rightarrow \mathbb{R}$ is non-negative, there exists a constant $C > 0$ such that

$$I[\Gamma, u] \geq C \|u\|_{H^1(\Omega)}^2 \quad \forall u \in H_0^1(\Omega).$$

Thus $\alpha := \inf_{u \in H_0^1(\Omega)} I[\Gamma, u]$ is finite. Let $u_k \in H_0^1(\Omega)$ ($k = 1, 2, \dots$) be such that

$$\lim_{k \rightarrow \infty} I[\Gamma, u_k] = \alpha.$$

Then $\{u_k\}$ is bounded in $H^1(\Omega)$ and hence it has a subsequence, not relabeled, that weakly converges to some $u_{\infty} \in H_0^1(\Omega)$. Since the embedding $H^1(\Omega) \hookrightarrow L^2(\Omega)$ is compact, up to a further subsequence, again not relabeled, $u_k \rightarrow u_{\infty}$ a. e. in Ω . Therefore, since $B : \mathbb{R} \rightarrow \mathbb{R}$ is continuous and non-negative, Fatou's lemma implies

$$\liminf_{k \rightarrow \infty} \int_{\Omega_+} B(u_k) dX \geq \int_{\Omega_+} B(u_{\infty}) dX.$$

Since

$$u \mapsto \int_{\Omega} \varepsilon_{\Gamma} |\nabla u|^2 dX$$

is convex and $H^1(\Omega)$ -continuous, it is sequentially weakly lower semi-continuous. Consequently,

$$\liminf_{k \rightarrow \infty} I[\Gamma, u_k] \geq I[\Gamma, u_{\infty}].$$

Thus u_{∞} is a minimizer of $I : H_0^1(\Omega) \rightarrow \mathbb{R} \cup \{\infty\}$ and the minimum value is clearly finite. The uniqueness of such a minimizer follows from the strict convexity of $I[\Gamma, \cdot]$. Therefore $\psi_0 = u_0 + u_{\infty} \in H_g^1(\Omega)$ is the unique maximizer of $G[\Gamma, \cdot]$ over $H_g^1(\Omega)$ and the maximum value is finite.

We show now the boundedness of the minimizer u_{∞} of $I[\Gamma, \cdot]$ over $H_0^1(\Omega)$ and hence that of the maximizer $\psi_0 = u_0 + u_{\infty}$ of $G[\Gamma, \cdot]$ over $H_g^1(\Omega)$. By (2.6) and the assumption that $B'(\infty) = \infty$ and $B'(-\infty) = -\infty$, there exists $\lambda > 0$ depending on \hat{C}_1 and B such that

$$B'(u_0 + \lambda) \geq 1 \quad \text{and} \quad B'(u_0 - \lambda) \leq -1 \quad \text{a.e. } \Omega_+.$$

Define $u_{\lambda} : \Omega \rightarrow \mathbb{R}$ by

$$u_{\lambda}(X) = \begin{cases} -\lambda & \text{if } u_{\infty}(X) < -\lambda, \\ u_{\infty}(X) & \text{if } |u_{\infty}(X)| \leq \lambda, \\ \lambda & \text{if } u_{\infty}(X) > \lambda. \end{cases}$$

Clearly $u_\lambda \in H_0^1(\Omega)$ and hence $I[\Gamma, u_\infty] \leq I[\Gamma, u_\lambda]$. This and the fact that $|\nabla u_\lambda| \leq |\nabla u_\infty|$ a.e. in Ω imply

$$\int_{\Omega_+} B(u_0 + u_\infty) dX \leq \int_{\Omega_+} B(u_0 + u_\lambda) dX.$$

Consequently, we have by the convexity of $B : \mathbb{R} \rightarrow \mathbb{R}$ imply that

$$\begin{aligned} 0 &\geq \int_{\{X \in \Omega_+ : u_\infty(X) > \lambda\}} [B(u_0(X) + u_\infty(X)) - B(u_0(X) + \lambda)] dX \\ &\quad + \int_{\{X \in \Omega_+ : u_\infty(X) < -\lambda\}} [B(u_0(X) + u_\infty(X)) - B(u_0(X) - \lambda)] dX \\ &\geq \int_{\{X \in \Omega_+ : u_\infty(X) > \lambda\}} B'(u_0(X) + \lambda) [u_\infty(X) - \lambda] dX \\ &\quad + \int_{\{X \in \Omega_+ : u_\infty(X) < -\lambda\}} B'(u_0(X) - \lambda) [u_\infty(X) + \lambda] dX \\ &\geq \int_{\{X \in \Omega_+ : u_\infty(X) > \lambda\}} [u_\infty(X) - \lambda] dX - \int_{\{X \in \Omega_+ : u_\infty(X) < -\lambda\}} [u_\infty(X) + \lambda] dX \\ &= \int_{\{X \in \Omega_+ : |u_\infty(X)| > \lambda\}} [|u_\infty(X)| - \lambda] dX \\ &\geq 0. \end{aligned}$$

Hence the last integral is 0. This implies that the Lebesgue measure of the set $\{X \in \Omega_+ : |u_\infty(X)| > \lambda\}$ is 0. Therefore $|u_\infty| \leq \lambda$ a.e. Ω_+ .

Since $u_\infty \in H_0^1(\Omega)$ is a minimizer of $I : H_0^1(\Omega) \rightarrow \mathbb{R} \cup \{\infty\}$, it is a weak solution of the corresponding Euler–Lagrange equation, which can be rewritten as

$$\nabla \cdot \varepsilon_\Gamma \nabla u_\infty = \chi_+ B'(u_\infty + u_0) \quad \text{in } \Omega.$$

The right-hand side of this equation is bounded in Ω by a constant depending only on \hat{C}_1 and B but not on Γ . Thus, by the regularity theory, both the $H^1(\Omega)$ and $L^\infty(\Omega)$ norms of u_∞ and hence those of $\psi_0 = u_\infty + u_0$ are bounded by a constant that only depends on \hat{C}_1 and B but not on Γ . This proves the desired boundedness of ψ_0 .

Now routine calculations lead to

$$\left. \frac{d}{dt} \right|_{t=0} G[\Gamma, \psi_0 + t\phi] = \int_{\Omega} [-\varepsilon_\Gamma \nabla \psi_0 \cdot \nabla \phi + f\phi - \chi_+ B'(\psi_0) \phi] dX \quad \forall \phi \in H_0^1(\Omega).$$

Hence ψ_0 is a weak solution to the boundary-value problem of PB equation (2.2) and (2.3).

If $\phi_0 \in H_g^1(\Omega)$ is another weak solution to the boundary-value problem of PB equation (2.2) and (2.3), then

$$\int_{\Omega} \{\varepsilon_\Gamma \nabla(\psi_0 - \phi_0) \cdot \nabla \eta + \chi_+ [B'(\psi_0) - B'(\phi_0)] \eta\} dX = 0$$

for any $\eta \in H_0^1(\Omega)$. Choosing $\eta = \psi_0 - \phi_0 \in H_0^1(\Omega)$ and using the convexity of $B : \mathbb{R} \rightarrow \mathbb{R}$, we obtain $\psi_0 = \phi_0$, proving the uniqueness. **Q.E.D.**

We remark that the boundary-value problem of the PB equation (2.2) and (2.3) is equivalent to the elliptic interface problem

$$\begin{cases} \varepsilon_- \Delta \psi = -f & \text{in } \Omega_-, \\ \varepsilon_+ \Delta \psi - B'(\psi) = 0 & \text{in } \Omega_+, \\ \llbracket \psi \rrbracket = \llbracket \varepsilon_\Gamma \nabla \psi \cdot n \rrbracket = 0 & \text{on } \Gamma, \\ \psi = g & \text{on } \partial\Omega, \end{cases} \quad (2.7)$$

where $\llbracket u \rrbracket = u|_{\Omega_+} - u|_{\Omega_-}$ denotes the jump across Γ of a function u from Ω_+ to Ω_- . See [37] for a proof. In particular, this equivalence implies the continuity of $\varepsilon_\Gamma \partial\psi/\partial n$ across the boundary Γ .

3 Dielectric boundary forces as shape derivatives

Let $V \in C^\infty(\mathbb{R}^3, \mathbb{R}^3)$. Assume that $V(x) = 0$ if $\text{dist}(x, \Gamma) > d$ for some $d > 0$ such that

$$d < \frac{1}{2} \min(\text{dist}(\Gamma, \partial\Omega), \text{dist}(\text{supp}(f), \Gamma)). \quad (3.1)$$

Let $X \in \mathbb{R}^3$ and consider the dynamical system for $x = x(t)$:

$$\begin{cases} \frac{dx}{dt} = V(x) & \forall t > 0, \\ x(0) = X. \end{cases} \quad (3.2)$$

The solution of this dynamical system defines a smooth map from \mathbb{R}^3 to \mathbb{R}^3 at each $t \geq 0$. We shall denote this map by $x = x(t, X) = T_t(X)$ for all $t \geq 0$ and $X \in \mathbb{R}^3$. Each $T_t : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a homeomorphism with both T_t and T_t^{-1} being smooth. Clearly T_0 is the identity map.

Notice that for $t > 0$ small we have by Taylor's expansion that

$$\begin{aligned} T_t(X) &= x(t, X) \\ &= x(0, X) + t \partial_t x(0, X) + O(t^2) \\ &= X + tV(x(0, X)) + O(t^2) \\ &= X + tV(X) + O(t^2). \end{aligned}$$

This means that the perturbation of identity defined by $X \mapsto X + tV(X)$ and the map $T_t(X) = x(t, X)$ agree with each other up to the leading order term in t . Notice also that we only consider $V = V(x)$ instead of a more general $V = V(t, x)$, since the shape derivative defined via $V = V(t, x)$ only depends on $V(0, \cdot)$ by the Structure Theorem [21, 51].

For each $t \geq 0$, we denote

$$\Omega_t = T_t(\Omega), \quad \Omega_{t-} = T_t(\Omega_-), \quad \Omega_{t+} = T_t(\Omega_+), \quad \Gamma_t = T_t(\Gamma).$$

Clearly, all Ω_t , Ω_{t-} , and Ω_{t+} are open sets, and $\Gamma_t = \partial\Omega_{t-} = \overline{\Omega_{t+}} \cap \overline{\Omega_{t-}}$. To indicate the dependence on V , we also write $\Gamma_t = \Gamma_t(V)$. Note that each T_t perturbs Γ locally: $T_t(X) = X$ if $\text{dist}(X, \Gamma) > d$. Since $d > 0$ satisfies (3.1), we have $T_t(X) = X$ if $f(X) \neq 0$ or $X \in \partial\Omega$. In particular, $\Omega_t = T_t(\Omega) = \Omega$ and $T_t(\partial\Omega) = \partial T_t(\Omega) = \partial\Omega$.

We recall that the electrostatic free energy $G[\Gamma]$ is given by (2.4), where the functional $G[\Gamma, \cdot]$ is given in (2.1) and ψ_0 is the weak solution to (2.2) and (2.3). For $t > 0$, we define

$$G[\Gamma_t, \phi] = \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma_t}}{2} |\nabla\phi|^2 + f\phi - \chi_{t+} B(\phi) \right] dx \quad \forall \phi \in H^1(\Omega_t) = H^1(\Omega), \quad (3.3)$$

where $\varepsilon_{\Gamma_t} : \Omega \rightarrow \mathbb{R}$ is defined by

$$\varepsilon_{\Gamma_t}(x) = \begin{cases} \varepsilon_- & \text{if } x \in \Omega_{t-}, \\ \varepsilon_+ & \text{if } x \in \Omega_{t+}, \end{cases} \quad (3.4)$$

and χ_{t+} is the characteristic function of Ω_{t+} . By Theorem 2.1 there exists a unique $\psi_t \in H_g^1(\Omega) \cap L^\infty(\Omega)$ that maximizes $G[\Gamma_t, \cdot]$ over $H_g^1(\Omega)$. The maximum is finite and is given by

$$G[\Gamma_t] = \max_{\phi \in H_g^1(\Omega)} G[\Gamma_t, \phi] = G[\Gamma_t, \psi_t]. \quad (3.5)$$

This is the electrostatic free energy corresponding to Γ_t . In addition, ψ_t is the unique weak solution to the corresponding boundary-value problem of the Poisson–Boltzmann equation

$$\nabla \cdot \varepsilon_{\Gamma_t} \nabla \psi_t - \chi_{t+} B'(\psi_t) = -f \quad \text{in } \Omega, \quad (3.6)$$

$$\psi_t = g \quad \text{on } \partial\Omega. \quad (3.7)$$

Finally,

$$\|\psi_t\|_{H^1(\Omega)} + \|\psi_t\|_{L^\infty(\Omega)} \leq C_1, \quad (3.8)$$

where C_1 is the same constant in Theorem 4.1. In particular, C_1 does not depend on Γ_t .

Definition 3.1. *Let V and $\Gamma_t(V)$ ($t \geq 0$) be given as above. The shape derivative of $G[\Gamma]$ in the direction of V is*

$$\delta_{\Gamma, V} G[\Gamma] = \left. \frac{d}{dt} G[\Gamma_t(V)] \right|_{t=0} = \lim_{t \rightarrow 0^+} \frac{G[\Gamma_t(V)] - G[\Gamma]}{t},$$

if the limit exists.

In general, there exists $w : \Gamma \rightarrow \mathbb{R}$ such that

$$\delta_{\Gamma, V} G[\Gamma] = \int_{\Gamma} w(V \cdot n) dS,$$

where n is the unit normal to Γ . We shall define

$$\delta_\Gamma G[\Gamma] = w$$

and call it the *shape derivative* of $G[\Gamma]$.

We recall some properties of the transformation $T_t(X)$ defined by $V = V(X)$. These properties can be proved by direct calculations, cf. [21] (Chapter 8 and Chapter 9).

- (1) Let $X \in \mathbb{R}^3$ and $t \geq 0$. Let $\nabla T_t(X)$ be the Jacobian matrix of T_t at X defined by $(\nabla T_t(X))_{ij} = \partial_j T_t^i(X)$, where T_t^i is the i th component of T_t ($i = 1, 2, 3$). Let

$$J_t(X) = \det \nabla T_t(X).$$

Then for each X the function $t \rightarrow J_t(X)$ is in C^∞ and at X

$$\frac{dJ_t}{dt} = J_t(\nabla \cdot V) \circ T_t, \quad (3.9)$$

where \circ denotes the composition of functions or maps. Clearly, ∇T_0 is the identity matrix and $J_0 = 1$. The continuity of J_t at $t = 0$ then implies that $J_t > 0$ for $t > 0$ small enough.

- (2) Define $A(t) : \Omega \rightarrow \mathbb{R}$ for $t \geq 0$ small enough by

$$A(t) = J_t (\nabla T_t)^{-1} (\nabla T_t)^{-T}, \quad (3.10)$$

where a superscript T denotes the matrix transpose. We have at each point in Ω that

$$\begin{aligned} A'(t) = & [((\nabla \cdot V) \circ T_t) - (\nabla T_t)^{-1}((\nabla V) \circ T_t)\nabla T_t \\ & - (\nabla T_t)^{-1}((\nabla V) \circ T_t)^T(\nabla T_t)] A(t). \end{aligned} \quad (3.11)$$

- (3) For any $u \in L^2(\Omega)$ and $t \geq 0$, $u \circ T_t \in L^2(\Omega)$ and $u \circ T_t^{-1} \in L^2(\Omega)$. Moreover,

$$\lim_{t \rightarrow 0} u \circ T_t = u \quad \text{and} \quad \lim_{t \rightarrow 0} u \circ T_t^{-1} = u \quad \text{in } L^2(\Omega). \quad (3.12)$$

- (4) Let $t \geq 0$ and $u \in H^1(\Omega)$. Then both $u \mapsto u \circ T_t$ and $u \mapsto u \circ T_t^{-1}$ are one-to-one and onto maps from $H^1(\Omega)$ (resp. $H_g^1(\Omega)$) to $H^1(\Omega_t) = H^1(\Omega)$ (resp. $H_g^1(\Omega)$). Moreover, for any $u \in H^1(\Omega)$,

$$\nabla(u \circ T_t^{-1}) = (\nabla T_t^{-1})^T (\nabla u \circ T_t^{-1}) \quad \text{and} \quad \nabla(u \circ T_t) = (\nabla T_t)^T (\nabla u \circ T_t). \quad (3.13)$$

- (5) For any $u \in H^1(\Omega)$ and $t \geq 0$,

$$\frac{d}{dt}(u \circ T_t) = (\nabla u \cdot V) \circ T_t. \quad (3.14)$$

4 Formulas of the dielectric boundary force

Let $V \in C^\infty(\mathbb{R}^3, \mathbb{R}^3)$ be such that $V(X) = 0$ if $\text{dist}(X, \Gamma) > d$ for some $d > 0$ that satisfies (3.1). Let the transformations T_t ($t \geq 0$) be defined by (3.2). For $t > 0$, the electrostatic free energy $G[\Gamma_t]$ is given by (3.5), where the functional $G[\Gamma_t, \cdot]$ is given in (3.3) and ψ_t is the weak solution to (3.6) and (3.7), respectively. For $t = 0$, the electrostatic free energy $G[\Gamma]$ is given by (2.4), where the functional $G[\Gamma, \cdot]$ is given in (2.1) and ψ_0 is the weak solution to (2.2) and (2.3).

Theorem 4.1. *Assume $f \in H^1(\Omega)$. Then the shape derivative of the electrostatic free energy $G[\Gamma]$ in the direction of V is given by*

$$\begin{aligned} \delta_{\Gamma, V} G[\Gamma] = \int_{\Gamma} \left[\frac{\varepsilon_+}{2} |\nabla \psi_0^+|^2 - \frac{\varepsilon_-}{2} |\nabla \psi_0^-|^2 \right. \\ \left. - \varepsilon_+ |\nabla \psi_0^+ \cdot n|^2 + \varepsilon_- |\nabla \psi_0^- \cdot n|^2 + B(\psi_0) \right] (V \cdot n) dS. \end{aligned} \quad (4.1)$$

Proof. We divide our proof into four steps.

Step 1. Let $t \geq 0$. Since each $\phi \in H_g^1(\Omega_t) = H_g^1(\Omega)$ corresponds uniquely to $\phi \circ T_t^{-1} \in H_g^1(\Omega)$, we have by (3.5) that

$$G[\Gamma_t] = \max_{\phi \in H_g^1(\Omega)} G[\Gamma_t, \phi \circ T_t^{-1}].$$

Let $\phi \in H^1(\Omega) \cap L^\infty(\Omega)$ and $t \geq 0$, and denote

$$z(t, \phi) = G[\Gamma_t, \phi \circ T_t^{-1}]. \quad (4.2)$$

We prove that $\partial_t z(t, \phi)$ exists for $t \geq 0$ small and derive a formula for this derivative.

Recall that $J_t(X) = \det \nabla T_t(X)$. By the continuity of $t \mapsto J_t(X)$ and the fact that $J_0(X) = 1$ at each $X \in \Omega$, there exists $\tau > 0$ such that $J_t(X) > 0$ for all $t \in [0, \tau]$ and all $X \in \Omega$. Now let $t \in [0, \tau]$ and $\phi \in H^1(\Omega) \cap L^\infty(\Omega)$. By the definition of $G[\Gamma_t, \phi]$ (cf. (3.3)), the change of variable $x = T_t(X)$, and (3.13), we obtain

$$\begin{aligned} z(t, \phi) &= G[\Gamma_t, \phi \circ T_t^{-1}] \\ &= \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma_t}}{2} |\nabla(\phi \circ T_t^{-1})|^2 + f(\phi \circ T_t^{-1}) - \chi_{t+}(B(\phi) \circ T_t^{-1}) \right] dx \\ &= \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma}}{2} A(t) \nabla \phi \cdot \nabla \phi + (f \circ T_t) \phi J_t - \chi_+ B(\phi) J_t \right] dX, \end{aligned} \quad (4.3)$$

where $A(t)$ is given in (3.10). By the properties of the transformations T_t ($t \geq 0$), cf. (3.9), (3.11), and (3.14), each term in the above integral is differentiable with respect to t , and

$$\partial_t z(t, \phi) = \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma}}{2} A'(t) \nabla \phi \cdot \nabla \phi + (\partial_t(f \circ T_t)) \phi J_t + (f \circ T_t) \phi \partial_t J_t - \chi_+ B(\phi) \partial_t J_t \right] dX$$

$$\begin{aligned}
&= \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma}}{2} A'(t) \nabla \phi \cdot \nabla \phi + ((\nabla f \cdot V) \circ T_t) \phi J_t + (f \circ T_t) \phi ((\nabla \cdot V) \circ T_t) J_t \right. \\
&\quad \left. - \chi_+ B(\phi) ((\nabla \cdot V) \circ T_t) J_t \right] dX \\
&= \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma}}{2} A'(t) \nabla \phi \cdot \nabla \phi + ((\nabla \cdot (fV)) \circ T_t) \phi J_t \right. \\
&\quad \left. - \chi_+ B(\phi) ((\nabla \cdot V) \circ T_t) J_t \right] dX. \tag{4.4}
\end{aligned}$$

Step 2. Let $t \in (0, \tau]$. Since $\psi_t \in H_g^1(\Omega) \cap L^\infty(\Omega)$ and $\psi_0 \in H_g^1(\Omega) \cap L^\infty(\Omega)$ maximize $G[\Gamma_t, \cdot]$ and $G[\Gamma, \cdot]$, respectively, over $H_g^1(\Omega)$ (cf. (3.5) and (2.4)), we have

$$\begin{aligned}
G[\Gamma_t, \psi_0 \circ T_t^{-1}] &\leq G[\Gamma_t, \psi_t] = G[\Gamma_t], \\
G[\Gamma, \psi_t] &\leq G[\Gamma, \psi_0] = G[\Gamma], \\
G[\Gamma, \psi_t \circ T_t] &\leq G[\Gamma, \psi_0] = G[\Gamma].
\end{aligned}$$

Hence

$$\frac{G[\Gamma_t, \psi_0 \circ T_t^{-1}] - G[\Gamma, \psi_0]}{t} \leq \frac{G[\Gamma_t] - G[\Gamma]}{t} \leq \frac{G[\Gamma_t, \psi_t] - G[\Gamma, \psi_t \circ T_t]}{t}.$$

Consequently, we obtain by (4.2) that

$$\frac{z(t, \psi_0) - z(0, \psi_0)}{t} \leq \frac{G[\Gamma_t] - G[\Gamma]}{t} \leq \frac{z(t, \psi_t \circ T_t) - z(0, \psi_t \circ T_t)}{t}.$$

From Step 1, there exist $\xi(t), \eta(t) \in [0, t]$ for each $t \in (0, \tau]$ such that

$$\partial_t z(\xi(t), \psi_0) \leq \frac{G[\Gamma_t] - G[\Gamma]}{t} \leq \partial_t z(\eta(t), \psi_t \circ T_t) \quad \forall t \in (0, \tau]. \tag{4.5}$$

Step 3. We prove

$$\lim_{t \rightarrow 0} \partial_t z(\xi(t), \psi_0) = \partial_t z(0, \psi_0), \tag{4.6}$$

$$\lim_{t \rightarrow 0} \partial_t z(\eta(t), \psi_t \circ T_t) = \partial_t z(0, \psi_0). \tag{4.7}$$

We only prove (4.7). The proof of (4.6) is similar and in fact simpler.

Since $V \in C^\infty(\mathbb{R}^3, \mathbb{R}^3)$ is compactly supported in Ω , it is easy to see that

$$\begin{aligned}
A'(\eta(t)) &\rightarrow A'(0), \\
J_{\eta(t)} &\rightarrow J_0 = 1,
\end{aligned}$$

uniformly on Ω as $t \rightarrow 0$ which implies $\eta(t) \rightarrow 0$ as $t \rightarrow 0$. By (3.12), we also have as $t \rightarrow 0$ that

$$(\nabla \cdot (fV)) \circ T_{\eta(t)} \rightarrow \nabla \cdot (fV) \quad \text{in } L^2(\Omega),$$

$$(\nabla \cdot V) \circ T_{\eta(t)} \rightarrow \nabla \cdot V \quad \text{in } L^2(\Omega).$$

We shall prove at the end of this step

$$\lim_{t \rightarrow 0} \|\psi_t \circ T_t - \psi_0\|_{H^1(\Omega)} = 0. \quad (4.8)$$

Notice by Theorem 2.1 that ψ_t is uniformly bounded in $L^\infty(\Omega)$ with respect to $t \in [0, \tau]$, cf. (3.8). Thus

$$B(\psi_t \circ T_t) - B(\psi_0) = B'(\lambda(t)) (\psi_t \circ T_t - \psi_0) \rightarrow 0 \quad \text{in } H^1(\Omega),$$

as $t \rightarrow 0$, where $\lambda(t) : \Omega \rightarrow \mathbb{R}$ is in between $\psi_t \circ T_t$ and ψ_0 at each point in Ω and hence $\lambda(t)$ is uniformly bounded in $L^\infty(\Omega)$ with respect to $t \in [0, \tau]$. Applying all these convergence results, we obtain by (4.4) that

$$\begin{aligned} & \partial_t z(\eta(t), \psi_t \circ T_t) \\ &= \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma}}{2} A'(\eta(t)) \nabla(\psi_t \circ T_t) \cdot \nabla(\psi_t \circ T_t) + ((\nabla \cdot (fV)) \circ T_{\eta(t)}) (\psi_t \circ T_t) J_{\eta(t)} \right. \\ & \quad \left. - \chi_+ B((\psi_t \circ T_t)) ((\nabla \cdot V) \circ T_{\eta(t)}) J_{\eta(t)} \right] dX \\ & \rightarrow - \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma}}{2} A'(0) \nabla \psi_0 \cdot \nabla \psi_0 + (\nabla \cdot (fV)) \psi_0 - \chi_+ B(\psi_0) (\nabla \cdot V) \right] dX \\ &= \partial_t z(0, \psi_0) \quad \text{as } t \rightarrow 0, \end{aligned}$$

proving (4.7).

We now prove (4.8). Fix $t > 0$. Since $\psi_t \in H_g^1(\Omega)$ maximizes $G[\Gamma_t, \cdot]$ over $H_g^1(\Omega)$, $\psi_t \circ T_t$ maximizes $z(t, \phi) = G[\Gamma_t, \phi \circ T_t^{-1}]$ over all $\phi \in H_g^1(\Omega)$. Since ψ_t is bounded in Ω by (3.8), we obtain from (4.3) and

$$\left. \frac{d}{d\sigma} \right|_{\sigma=0} z(t, (\psi_t + \sigma \eta \circ T_t^{-1}) \circ T_t) = \left. \frac{d}{d\sigma} \right|_{\sigma=0} z(t, \psi_t \circ T_t + \sigma \eta) = 0 \quad \forall \eta \in H_0^1(\Omega)$$

that

$$\int_{\Omega} [-\varepsilon_{\Gamma} A(t) \nabla(\psi_t \circ T_t) \cdot \nabla \eta + (f \circ T_t) J_t \eta - \chi_+ B'(\psi_t \circ T_t) J_t \eta] dX = 0 \quad \forall \eta \in H_0^1(\Omega).$$

Since $\psi_0 \in H_g^1(\Omega)$ maximizes $G[\Gamma, \cdot]$ over $H_g^1(\Omega)$,

$$\int_{\Omega} [-\varepsilon_{\Gamma} \nabla \psi_0 \cdot \nabla \eta + f \eta - \chi_+ B'(\psi_0) \eta] dX = 0 \quad \forall \eta \in H_0^1(\Omega).$$

Subtracting one of these two equations from the other and choosing $\eta = \psi_t \circ T_t - \psi_0 \in H_0^1(\Omega)$, we deduce by rearranging the terms that

$$\int_{\Omega} \varepsilon_{\Gamma} |\nabla(\psi_t \circ T_t) - \nabla \psi_0|^2 dX + \int_{\Omega_+} [B'(\psi_t \circ T_t) - B'(\psi_0)] (\psi_t \circ T_t - \psi_0) dX$$

$$\begin{aligned}
&= - \int_{\Omega} \varepsilon_{\Gamma} [(A(t) - I)\nabla(\psi_t \circ T_t)] \cdot [\nabla(\psi_t \circ T_t) - \nabla\psi_0] dX \\
&\quad + \int_{\Omega} [(f \circ T_t)J_t - f](\psi_t \circ T_t - \psi_0) dX - \int_{\Omega_+} B'(\psi_t \circ T_t)(J_t - 1)(\psi_t \circ T_t - \psi_0) dX.
\end{aligned}$$

Notice by the convexity of B that

$$[B'(\psi_t \circ T_t) - B'(\psi_0)](\psi_t \circ T_t - \psi_0) \geq 0 \quad \text{in } \Omega_+.$$

Therefore, applying the Poincaré inequality to $\psi_t \circ T_t - \psi_0 \in H_0^1(\Omega)$ and using the Cauchy-Schwarz inequality, we obtain that

$$\begin{aligned}
\|\psi_t \circ T_t - \psi_0\|_{H^1(\Omega)}^2 &\leq C \int_{\Omega} \varepsilon_{\Gamma} |\nabla(\psi_t \circ T_t) - \nabla\psi_0|^2 dX \\
&\quad + C \int_{\Omega_+} [B'(\psi_t \circ T_t) - B'(\psi_0)](\psi_t \circ T_t - \psi_0) dX \\
&\leq C \left[\|A(t) - I\|_{L^\infty(\Omega)} \|\nabla(\psi_t \circ T_t)\|_{L^2(\Omega)} + \|(f \circ T_t)J_t - f\|_{L^2(\Omega)} \right. \\
&\quad \left. + \|B'(\psi_t \circ T_t)(J_t - 1)\|_{L^\infty(\Omega)} \right] \|\psi_t \circ T_t - \psi_0\|_{H^1(\Omega)}.
\end{aligned}$$

Here and below we denote by $C > 0$ a generic constant that can depend on $\varepsilon_-, \varepsilon_+, \Omega, B$ but not on t or ψ_0 . Note that

$$(f \circ T_t)J_t - f = (f \circ T_t - f)J_t + f(J_t - 1).$$

Consequently, the uniform boundedness of ψ_t in $L^\infty(\Omega)$ and in $H^1(\Omega)$ for small $t > 0$ (cf. (3.8)), the uniform convergence $A(t) \rightarrow A(0) = I$ and $J_t \rightarrow 1$ as $t \rightarrow 0$, and the convergence $f \circ T_t \rightarrow f$ in $L^2(\Omega)$ (cf. (3.12)) imply that

$$\begin{aligned}
\|\psi_t \circ T_t - \psi_0\|_{H^1(\Omega)} &\leq C \left[\|A(t) - I\|_{L^\infty(\Omega)} \|\nabla(\psi_t \circ T_t)\|_{L^2(\Omega)} + \|f \circ T_t - f\|_{L^2(\Omega)} \|J_t\|_{L^\infty(\Omega)} \right. \\
&\quad \left. + \|f\|_{L^2(\Omega)} \|J_t - 1\|_{L^\infty(\Omega)} + \|B'(\psi_t \circ T_t)\|_{L^\infty(\Omega)} \|J_t - 1\|_{L^\infty(\Omega)} \right] \\
&\rightarrow 0 \quad \text{as } t \rightarrow 0.
\end{aligned}$$

This is (4.8).

Step 4. It follows from (4.5)–(4.7) that

$$\delta_{\Gamma, V} G[\Gamma] = \frac{d}{dt} \Big|_{t=0} G[\Gamma_t(V)] = \partial_t z(0, \psi_0).$$

We now show that $\partial_t z(0, \psi_0)$ is the right-hand side of (4.1).

It follows from (4.4) with $t = 0$ and $\phi = \psi_0$, and (3.11) with $t = 0$ that

$$\begin{aligned}\partial_t z(0, \psi_0) &= \int_{\Omega} \left[-\frac{\varepsilon_{\Gamma}}{2} A'(0) \nabla \psi_0 \cdot \nabla \psi_0 + (\nabla \cdot (fV)) \psi_0 - \chi_+ B(\psi_0) (\nabla \cdot V) \right] dX \\ &= - \int_{\Omega} \frac{\varepsilon_{\Gamma}}{2} (\nabla \cdot V) \nabla \psi_0 \cdot \nabla \psi_0 dX + \int_{\Omega} \varepsilon_{\Gamma} (\nabla V) \nabla \psi_0 \cdot \nabla \psi_0 dX \\ &\quad + \int_{\Omega} (\nabla \cdot (fV)) \psi_0 dX - \int_{\Omega_+} B(\psi_0) (\nabla \cdot V) dX.\end{aligned}$$

Denote by V^i and n_i the i th components of V and n , respectively. Recall that the unit normal n to Γ points from Ω_- to Ω_+ . By integration by parts and the fact that V vanishes in a neighborhood of $\partial\Omega$, by (2.7) with $\psi = \psi_0$, and using the summation convention, we continue to have

$$\begin{aligned}\partial_t z(0, \psi_0) &= - \int_{\Omega_+} \frac{\varepsilon_+}{2} \partial_{X_i} V^i \partial_{X_j} \psi_0 \partial_{X_j} \psi_0 dX - \int_{\Omega_-} \frac{\varepsilon_-}{2} \partial_{X_i} V^i \partial_{X_j} \psi_0 \partial_{X_j} \psi_0 dX \\ &\quad + \int_{\Omega_+} \varepsilon_+ \partial_{X_j} V^i \partial_{X_j} \psi_0 \partial_{X_i} \psi_0 dX + \int_{\Omega_-} \varepsilon_- \partial_{X_j} V^i \partial_{X_j} \psi_0 \partial_{X_i} \psi_0 dX \\ &\quad - \int_{\Omega} f(V \cdot \nabla \psi_0) dX + \int_{\Omega_+} B'(\psi_0) (\nabla \psi_0 \cdot V) dX + \int_{\Gamma} B(\psi_0) (V \cdot n) dX \\ &= \int_{\Omega_+} \varepsilon_+ V^i \partial_{X_j} \psi_0 \partial_{X_j X_i}^2 \psi_0 dX + \int_{\Gamma} \frac{\varepsilon_+}{2} |\nabla \psi_0^+|^2 (V \cdot n) dS \\ &\quad + \int_{\Omega_-} \varepsilon_- V^i \partial_{X_j} \psi_0 \partial_{X_j X_i}^2 \psi_0 dX - \int_{\Gamma} \frac{\varepsilon_-}{2} |\nabla \psi_0^-|^2 (V \cdot n) dS \\ &\quad - \int_{\Omega_+} \varepsilon_+ V^i \left(\partial_{X_j X_j}^2 \psi_0 \partial_{X_i} \psi_0 + \partial_{X_j} \psi_0 \partial_{X_i X_j}^2 \psi_0 \right) dX \\ &\quad - \int_{\Gamma} \varepsilon_+ V^i n_j \partial_{X_j} \psi_0^+ \partial_{X_i} \psi_0^+ dS \\ &\quad - \int_{\Omega_-} \varepsilon_- V^i \left(\partial_{X_j X_j}^2 \psi_0 \partial_{X_i} \psi_0 + \partial_{X_j} \psi_0 \partial_{X_i X_j}^2 \psi_0 \right) dX \\ &\quad + \int_{\Gamma} \varepsilon_- V^i n_j \partial_{X_j} \psi_0^- \partial_{X_i} \psi_0^- dS \\ &\quad - \int_{\Omega} f(V \cdot \nabla \psi_0) dX + \int_{\Omega_+} B'(\psi_0) (\nabla \psi_0 \cdot V) dX + \int_{\Gamma} B(\psi_0) (V \cdot n) dS \\ &= \int_{\Gamma} \frac{\varepsilon_+}{2} |\nabla \psi_0^+|^2 (V \cdot n) dS - \int_{\Gamma} \frac{\varepsilon_-}{2} |\nabla \psi_0^-|^2 (V \cdot n) dS \\ &\quad - \int_{\Gamma} \varepsilon_+ (\nabla \psi_0^+ \cdot n) (V \cdot \nabla \psi_0^+) dS + \int_{\Gamma} \varepsilon_- (\nabla \psi_0^- \cdot n) (V \cdot \nabla \psi_0^-) dS \\ &\quad - \int_{\Omega_+} [\varepsilon_+ \Delta \psi_0 + f - B'(\psi_0)] (\nabla \psi_0 \cdot V) dX\end{aligned}$$

$$\begin{aligned}
& - \int_{\Omega_-} (\varepsilon_- \Delta \psi_0 + f) (\nabla \psi_0 \cdot V) dX + \int_{\Gamma} B(\psi_0) (V \cdot n) dS \\
= & \int_{\Gamma} \frac{\varepsilon_+}{2} |\nabla \psi_0^+|^2 (V \cdot n) dS - \int_{\Gamma} \frac{\varepsilon_-}{2} |\nabla \psi_0^-|^2 (V \cdot n) dS \\
& - \int_{\Gamma} \varepsilon_+ (\nabla \psi_0^+ \cdot n) (V \cdot \nabla \psi_0^+) dS + \int_{\Gamma} \varepsilon_- (\nabla \psi_0^- \cdot n) (V \cdot \nabla \psi_0^-) dS \\
& + \int_{\Gamma} B(\psi_0) (V \cdot n) dS. \tag{4.9}
\end{aligned}$$

Since $\psi_0^+ = \psi_0^-$ along Γ ,

$$\nabla (\psi_0^+ - \psi_0^-) = (\nabla \psi_0^+ \cdot n - \nabla \psi_0^- \cdot n) n \quad \text{on } \Gamma.$$

By (2.7),

$$\varepsilon_+ \nabla \psi_0^+ \cdot n = \varepsilon_- \nabla \psi_0^- \cdot n = \varepsilon_{\Gamma} \nabla \psi_0 \cdot n \quad \text{on } \Gamma. \tag{4.10}$$

We thus have

$$\begin{aligned}
& \int_{\Gamma} \varepsilon_+ (\nabla \psi_0^+ \cdot n) (V \cdot \nabla \psi_0^+) dS - \int_{\Gamma} \varepsilon_- (\nabla \psi_0^- \cdot n) (V \cdot \nabla \psi_0^-) dS \\
& = \int_{\Gamma} \varepsilon_{\Gamma} (\nabla \psi_0 \cdot n) V \cdot \nabla (\psi_0^+ - \psi_0^-) dS \\
& = \int_{\Gamma} \varepsilon_{\Gamma} (\nabla \psi_0 \cdot n) V \cdot (\nabla \psi_0^+ \cdot n - \nabla \psi_0^- \cdot n) n \\
& = \int_{\Gamma} \left[\varepsilon_+ |\nabla \psi_0^+ \cdot n|^2 - \varepsilon_- |\nabla \psi_0^- \cdot n|^2 \right] (V \cdot n) dS.
\end{aligned}$$

This and (4.9) imply that $\partial_t z(0, \psi_0)$ is the right-hand side of (4.1). **Q.E.D.**

Corollary 4.1. *Under the assumption of Theorem 4.1 we have*

$$\delta_{\Gamma} G[\Gamma] = \frac{1}{2} \left(\frac{1}{\varepsilon_-} - \frac{1}{\varepsilon_+} \right) |\varepsilon_{\Gamma} \nabla \psi_0 \cdot n|^2 + \frac{1}{2} (\varepsilon_+ - \varepsilon_-) |(I - n \otimes n) \nabla \psi_0|^2 + B(\psi_0). \tag{4.11}$$

In particular, if $\varepsilon_- < \varepsilon_+$ then the dielectric boundary force in the normal direction from Ω_- to Ω_+ is always negative: $F_n = -\delta_{\Gamma} G[\Gamma] < 0$ on Γ .

Proof. We have the orthogonal decomposition

$$\nabla \psi_0^{\pm} = (\nabla \psi_0^{\pm} \cdot n) n + (I - n \otimes n) \nabla \psi_0^{\pm} \quad \text{on } \Gamma.$$

Moreover, since ψ_0 is continuous, its tangential derivatives along Γ are continuous. Hence

$$(I - n \otimes n) \nabla \psi_0^+ = (I - n \otimes n) \nabla \psi_0^- \quad \text{on } \Gamma.$$

We denote this common value by $(I - n \otimes n)\nabla\psi_0$. Notice that

$$n \cdot (I - n \otimes n)\nabla\psi_0 = 0.$$

We thus have by (4.10) that

$$\begin{aligned} & \frac{\varepsilon_+}{2}|\nabla\psi_0^+|^2 - \frac{\varepsilon_-}{2}|\nabla\psi_0^-|^2 - \varepsilon_+|\nabla\psi_0^+ \cdot n|^2 + \varepsilon_-|\nabla\psi_0^- \cdot n|^2 \\ &= -\frac{\varepsilon_+}{2}|\nabla\psi_0^+ \cdot n|^2 + \frac{\varepsilon_-}{2}|\nabla\psi_0^- \cdot n|^2 + \frac{\varepsilon_+}{2}|(I - n \otimes n)\nabla\psi_0^+|^2 - \frac{\varepsilon_-}{2}|(I - n \otimes n)\nabla\psi_0^-|^2 \\ &= \frac{1}{2}\left(\frac{1}{\varepsilon_-} - \frac{1}{\varepsilon_+}\right)|\varepsilon_{\Gamma}\nabla\psi_0 \cdot n|^2 + \frac{1}{2}(\varepsilon_+ - \varepsilon_-)|(I - n \otimes n)\nabla\psi_0|^2. \end{aligned}$$

This and (4.1) imply (4.11). If $\varepsilon_- < \varepsilon_+$ then (4.11) implies $F_n = -\delta_{\Gamma}G[\Gamma] < 0$. **Q.E.D.**

Acknowledgment. B. L. was supported by the US National Science Foundation (NSF) through the grant DMS-0811259, by the NSF Center for Theoretical Biological Physics (CTBP) through the NSF grant PHY-0822283, and by the National Institutes of Health through the grant R01GM096188. Z. Z. was supported by a fellowship from the Lu Graduate Education International Exchange Fund, Zhejiang University, China. Part of this work was completed during Li's visit to the Department of Mathematics, Zhejiang University, in the summer of 2010, and during Zhang's visit to the Department of Mathematics, UC San Diego, in Fall 2010 and Winter 2011. The hospitality of these departments is greatly appreciated. The authors thank Dr. Ray Luo for helpful discussions and for sharing with them his unpublished notes on the Maxwell stress tensor.

References

- [1] D. Andelman. Electrostatic properties of membranes: The Poisson–Boltzmann theory. In R. Lipowsky and E. Sackmann, editors, *Handbook of Biological Physics*, volume 1, pages 603–642. Elsevier, 1995.
- [2] N. A. Baker. Improving implicit solvent simulations: a Poisson-centric view. *Curr. Opin. Struct. Biol.*, 15:137–143, 2005.
- [3] D. Bashford and D. A. Case. Generalized Born models of macromolecular solvation effects. *Ann. Rev. Phys. Chem.*, 51:129–152, 2000.
- [4] P. W. Bates, Z. Chen, Y. H. Sun, G. W. Wei, and S. Zhao. Geometric and potential driving formation and evolution of biomolecular surfaces. *J. Math. Biology*, 59:193–231, 2009.
- [5] I. Borukhov, D. Andelman, and H. Orland. Steric effects in electrolytes: a modified Poisson–Boltzmann equation. *Phys. Rev. Lett.*, 79:435–438, 1997.

- [6] D. Bucur and G. Buttazzo. *Variational Methods in Shape Optimization Problems*. Progress in Nonlinear Differential Equations and Their Applications. Birkhäuser, Boston, 2005.
- [7] D. L. Chapman. A contribution to the theory of electrocapillarity. *Phil. Mag.*, 25:475–481, 1913.
- [8] J. Che, J. Dzubiella, B. Li, and J. A. McCammon. Electrostatic free energy and its variations in implicit solvent models. *J. Phys. Chem. B*, 112:3058–3069, 2008.
- [9] J. Chen, C. L. Brooks III, and J. Khandogin. Recent advances in implicit solvent based methods for biomolecular simulations. *Curr. Opin. Struct. Biol.*, 18:140–148, 2008.
- [10] Z. Chen, N. A. Baker, and G. W. Wei. Differential geometry based solvation model I: Eulerian formulation. *J. Comput. Phys.*, 229:8231–8258, 2010.
- [11] H. B. Cheng, L.-T. Cheng, and B. Li. Yukawa-field approximation of electrostatic free energy and dielectric boundary force. *Nonlinearity*, 24:3215–3236, 2011.
- [12] L.-T. Cheng, J. Dzubiella, J. A. McCammon, and B. Li. Application of the level-set method to the implicit solvation of nonpolar molecules. *J. Chem. Phys.*, 127:084503, 2007.
- [13] L.-T. Cheng, B. Li, and Z. Wang. Level-set minimization of potential controlled Hadwiger valuations for molecular solvation. *J. Comput. Phys.*, 229:8497–8510, 2010.
- [14] L.-T. Cheng, Z. Wang, P. Setny, J. Dzubiella, B. Li, and J. A. McCammon. Interfaces and hydrophobic interactions in receptor-ligand systems: A level-set variational implicit solvent approach. *J. Chem. Phys.*, 131:144102, 2009.
- [15] L.-T. Cheng, Y. Xie, J. Dzubiella, J. A. McCammon, J. Che, and B. Li. Coupling the level-set method with molecular mechanics for variational implicit solvation of nonpolar molecules. *J. Chem. Theory Comput.*, 5:257–266, 2009.
- [16] B. Chu. *Molecular Forces, Based on the Lecture of Peter J. W. Debye*. Interscience, John Wiley & Sons, 1967.
- [17] M. L. Connolly. Analytical molecular surface calculation. *J. Appl. Cryst.*, 16:548–558, 1983.
- [18] M. L. Connolly. The molecular surface package. *J. Mol. Graphics*, 11:139–141, 1992.
- [19] C. J. Cramer and D. G. Truhlar. Implicit solvation models: Equilibria, structure, spectra, and dynamics. *Chem. Rev.*, 99:2161–2200, 1999.

- [20] M. E. Davis and J. A. McCammon. Electrostatics in biomolecular structure and dynamics. *Chem. Rev.*, 90:509–521, 1990.
- [21] M. C. Delfour and J.-P. Zolésio. *Shapes and Geometries: Analysis, Differential Calculus, and Optimization*. SIAM, 1987.
- [22] J. Dzubiella, J. M. J. Swanson, and J. A. McCammon. Coupling hydrophobicity, dispersion, and electrostatics in continuum solvent models. *Phys. Rev. Lett.*, 96:087802, 2006.
- [23] J. Dzubiella, J. M. J. Swanson, and J. A. McCammon. Coupling nonpolar and polar solvation free energies in implicit solvent models. *J. Chem. Phys.*, 124:084905, 2006.
- [24] M. Feig and C. L. Brooks III. Recent advances in the development and applications of implicit solvent models in biomolecule simulations. *Curr. Opin. Struct. Biol.*, 14:217–224, 2004.
- [25] F. Fixman. The Poisson–Boltzmann equation and its application to polyelectrolytes. *J. Chem. Phys.*, 70:4995–5005, 1979.
- [26] F. Fogolari and J. M. Briggs. On the variational approach to Poisson–Boltzmann free energies. *Chem. Phys. Lett.*, 281:135–139, 1997.
- [27] F. Fogolari, A. Brigo, and H. Molinari. The Poisson–Boltzmann equation for biomolecular electrostatics: a tool for structural biology. *J. Mol. Recognit.*, 15:377–392, 2002.
- [28] D. Gilbarg and N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Springer-Verlag, 2nd edition, 1998.
- [29] M. K. Gilson, M. E. Davis, B. A. Luty, and J. A. McCammon. Computation of electrostatic forces on solvated molecules using the Poisson–Boltzmann equation. *J. Phys. Chem*, 97:3591–3600, 1993.
- [30] M. Gouy. Sur la constitution de la charge électrique a la surface d’un électrolyte. *J. de Phys.*, 9:457–468, 1910.
- [31] P. Grochowski and J. Trylska. Continuum molecular electrostatics, salt effects and counterion binding—A review of the Poisson–Boltzmann model and its modifications. *Biopolymers*, 89:93–113, 2008.
- [32] W. Im, D. Beglov, and B. Roux. Continuum Solvation Model: computation of electrostatic forces from numerical solutions to the Poisson–Boltzmann equation. *Comput. Phys. Commun.*, 111:59–75, 1998.
- [33] B. Kawohl, O. Pironneau, L. Tartar, and J.-P. Zolésio. *Optimal Shape Design*, volume 1740 of *Lecture Notes in Mathematics*. Springer, 2000.

- [34] V. Kralj-Iglic and A. Iglic. A simple statistical mechanical approach to the free energy of the electric double layer including the excluded volume effect. *J. Phys. II (France)*, 6:477–491, 1996.
- [35] B. Lee and F. M. Richards. The interpretation of protein structures: Estimation of static accessibility. *J. Mol. Biol.*, 55:379–400, 1971.
- [36] B. Li. Continuum electrostatics for ionic solutions with nonuniform ionic sizes. *Non-linearity*, 22:811–833, 2009.
- [37] B. Li. Minimization of electrostatic free energy and the Poisson–Boltzmann equation for molecular solvation with implicit solvent. *SIAM J. Math. Anal.*, 40:2536–2566, 2009.
- [38] B. Lu, X. Cheng, T. Hou, and J. A. McCammon. Calculation of the Maxwell stress tensor and the Poisson–Boltzmann force on a solvated molecular surface using hyper-singular boundary integrals. *J. Chem. Phys.*, 123:084904, 2005.
- [39] B. Z. Lu, Y. C. Zhou, M. J. Holst, and J. A. McCammon. Recent progress in numerical methods for the Poisson–Boltzmann equation in biophysical applications. *Commun. Comput. Phys.*, 3:973–1009, 2008.
- [40] R. Luo. Personal communications. 2010.
- [41] S. Osher and R. Fedkiw. *Level Set Methods and Dynamic Implicit Surfaces*. Springer, New York, 2002.
- [42] S. Osher and J. A. Sethian. Fronts propagating with curvature-dependent speed: Algorithms based on Hamilton-Jacobi formulations. *J. Comput. Phys.*, 79:12–49, 1988.
- [43] F. M. Richards. Areas, volumes, packing, and protein structure. *Annu. Rev. Biophys. Bioeng.*, 6:151–176, 1977.
- [44] T. J. Richmond. Solvent accessible surface area and excluded volume in proteins. Analytical equations for overlapping spheres and implications for the hydrophobic effect. *J. Mol. Biol.*, 178:63–89, 1984.
- [45] B. Roux and T. Simonson. Implicit solvent models. *Biophys. Chem.*, 78:1–20, 1999.
- [46] J. A. Sethian. *Level Set Methods and Fast Marching Methods: Evolving Interfaces in Geometry, Fluid Mechanics, Computer Vision, and Materials Science*. Cambridge University Press, 2nd edition, 1999.
- [47] P. Setny, Z. Wang, L.-T. Cheng, B. Li, J. A. McCammon, and J. Dzubiella. Dewetting-controlled binding of ligands to hydrophobic pockets. *Phys. Rev. Lett.*, 103:187801, 2009.

- [48] K. A. Sharp and B. Honig. Calculating total electrostatic energies with the nonlinear Poisson–Boltzmann equation. *J. Phys. Chem.*, 94:7684–7692, 1990.
- [49] K. A. Sharp and B. Honig. Electrostatic interactions in macromolecules: Theory and applications. *Annu. Rev. Biophys. Biophys. Chem.*, 19:301–332, 1990.
- [50] P. Smereka. The numerical approximation of a delta function with application to level set methods. *J. Comput. Phys.*, 211:77–90, 2006.
- [51] J. Sokolowski and J.-P. Zolésio. *Introduction to Shape Optimization: Shape Sensitivity Analysis*. Springer Series in Computational Mathematics. Springer, 1992.
- [52] W. C. Still, A. Tempczyk, R. C. Hawley, and T. Hendrickson. Semianalytical treatment of solvation for molecular mechanics and dynamics. *J. Amer. Chem. Soc.*, 112:6127–6129, 1990.
- [53] J. Tomasi and M. Persico. Molecular interactions in solution: An overview of methods based on continuous distributions of the solvent. *Chem. Rev.*, 94:2027–2094, 1994.
- [54] G. Treset. Generalized Poisson–Fermi formalism for investigating size correlation effects with multiple ions. *Phys. Rev. E*, 78:061506, 2008.
- [55] E. Trizac and J.-L. Raimbault. Long-range electrostatic interactions between like-charged colloids: Steric and confinement effects. *Phys. Rev. E*, 60:6530–6533, 1999.
- [56] J. A. Wagoner and N. A. Baker. Assessing implicit models for nonpolar mean solvation forces: The importance of dispersion and volume changes. *Proc. Natl. Acad. Sci., USA*, 103:8331–8336, 2006.
- [57] Z. Wang, J. Che, L.-T. Cheng, J. Dzubiella, B. Li, and J. A. McCammon. Level-set variational implicit solvation with the Coulomb-field approximation. *J. Chem. Theory Comput.*, 2011 (submitted).
- [58] S. Zhou, Z. Wang, and B. Li. Mean-field description of ionic size effects with non-uniform ionic sizes: A numerical approach. *Phys. Rev. E*, 84:021901, 2011.