

# IEDB-3D: structural data within the immune epitope database

Julia Ponomarenko<sup>1,2,\*</sup>, Nikitas Papangelopoulos<sup>1</sup>, Dirk M. Zajonc<sup>3</sup>, Bjoern Peters<sup>3</sup>,  
Alessandro Sette<sup>3</sup> and Philip E. Bourne<sup>1,2</sup>

<sup>1</sup>San Diego Supercomputer Center, <sup>2</sup>Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California San Diego, CA 92093 and <sup>3</sup>La Jolla Institute for Allergy and Immunology, La Jolla, CA 92037, USA

Received August 14, 2010; Revised September 16, 2010; Accepted September 19, 2010

## ABSTRACT

**IEDB-3D is the 3D structural component of the Immune Epitope Database (IEDB) available via the 'Browse by 3D Structure' page at <http://www.iedb.org>. IEDB-3D catalogs B- and T-cell epitopes and Major Histocompatibility Complex (MHC) ligands for which 3D structures of complexes with antibodies, T-cell receptors or MHC molecules are available in the Protein Data Bank (PDB). Journal articles that are primary citations of PDB structures and that define immune epitopes are curated within IEDB as any other reference along with accompanying functional assays and immunologically relevant information. For each curated structure, IEDB-3D provides calculated data on intermolecular contacts and interface areas and includes an application, EpitopeViewer, to visualize the structures. IEDB-3D is fully embedded within IEDB, thus allowing structural data, both curated and calculated, and all accompanying information to be queried using multiple search interfaces. These include queries for epitopes recognized in different pathogens, eliciting different functional immune responses, and recognized by different components of the immune system. The query results can be downloaded in Microsoft Excel format, or the entire database, together with structural data both curated and calculated, can be downloaded in either XML or MySQL formats.**

## INTRODUCTION

The Immune Epitope Database (IEDB) (1) catalogs experimentally identified B- and T-cell epitopes and MHC ligands through manual curation of the scientific literature. By the end of 2011, all published epitopes from infectious agents [except HIV, which are maintained

in (2)], allergens and autoimmune diseases should be included in the IEDB. In addition, all epitopes characterized by 3D structures of immune receptors complexed with antigens found in the Protein Data Bank (PDB) (3) will be included, independent of the disease association of the antigen. This expansion in scope is made as to understand the general structural principles of epitope recognition a large dataset is desirable, but detailed 3D structural information on epitope complexes is rare.

The 3D structural component of IEDB, called here IEDB-3D, is embedded seamlessly within the general IEDB, thus ensuring that each reference describing 3D structure is curated within IEDB as any other journal article describing immune epitopes. The focus on curation of immunological and epitope relevant information is the major difference between IEDB and IMGT/3Dstructure-DB (4). Similar to IMGT/3Dstructure-DB (4) and other related databases, such as MPID-T (5), Epitome (6), BEID (7) and CED (8), IEDB provides calculated data on intermolecular contacts and interface areas and includes an application to visualize the structure, EpitopeViewer (9), which is a high-quality graphic and rendering tool. Among the aforementioned databases, only CED and IEDB curate epitope residues from the literature. In IEDB, antibody, MHC and T-cell receptor (TCR) residues interacting with the epitope are also curated if they are provided in the reference. Thus, within IEDB, curated and derived data on epitope and antigen–receptor interactions can be seen and compared side-by-side on the IEDB web page and also through launching EpitopeViewer.

## IEDB-3D OVERVIEW

IEDB-3D is fully integrated within IEDB, with its data structure, ontology and query interface indistinguishable from those of the IEDB database. IEDB-3D is the 3D structural component of the Immune Epitope Database (IEDB) available via the 'Browse by 3D Structure' page

\*To whom correspondence should be addressed. Tel: +1 858 822 5415; Fax: +1 858 822 0873; Email: [jpon@sdsc.edu](mailto:jpon@sdsc.edu)

**IMMUNE EPITOPE DATABASE AND ANALYSIS RESOURCE**

Home | **Browse** | **Search** | Tools | Support | More IEDB

Browse By MHC Allele  
Browse By Source Organism  
**Browse By 3D Structure**

Any  
Linear Peptide: [Enter Search] [Exact Match]  
Discontinuous Peptide  
Non-Peptidic: [Enter Search] [Molecule Finder]

Epitope Source  
Source Organism: [Enter Search] [Organism Finder]  
Source Antigen: [Enter Search] [Molecule Finder]

Immune Recognition Context  
 B Cell Response  
 T Cell Response  
 MHC Binding  
 MHC Ligand Elution  
 Host Organism: [Enter Search] [Organism Finder]  
 MHC Restriction: [Enter Search] [Allele Finder]  
 MHC Class: [Dropdown]  
 [Search] [Clear]

Summary Metric

Summary Metric	Count
Peptidic Epitopes	76590
Non-Peptidic Epitopes	619
T Cell Assays	159192
B Cell Assays	135299
MHC Ligand Elution Assays	2007
MHC Binding Assays	199220
Epitope Source Organisms	2529
Restricting MHC Alleles	559
References	10755

Figure 1. The screen shot of IEDB web page.

at <http://www.iedb.org> (Figure 1). Similar to IEDB, the types of curated data include experiments describing recognition of epitopes or antigens (peptidic or non-peptidic) by TCRs (T-cell assays describing 3D structures of antigens/epitopes in complexes with MHC and TCR), immunoglobulins or antibodies (B-cell assays describing structures of antibody-antigen complexes), and MHC molecules (MHC binding assays describing structures of antigens/epitopes in complex with MHC) (Figure 2).

Curation of 3D complex data is handled like the remaining data in the IEDB, except that all epitopes are considered in scope, including those from HIV or cancer. IEDB does not handle HIV data, if the paper provides only information on immunological assay involving HIV antigen explicitly, since these data are curated in HIV database (2). However, if 3D structural data are available for HIV epitopes, they are curated, but with a lower priority compared to other 3D structural data. In IEDB-3D, in addition to the immunological context curated for all epitopes in IEDB, information on 3D structure is provided in the section '3D Structure of Complex' (Figure 3).

An overview of IEDB-3D content is given in Table 1. Current statistics on the number of distinct epitope structures curated by assay type and source organism can be accessed from the 'Browse by 3D Structure' page (Figure 1, red box).

### CURATION OF 3D STRUCTURAL DATA

The process of curating epitopes from 3D structures is shown in Figure 2. To be curated, the paper describing




Figure 2. Objects, processes and roles as represented in IEDB-3D.

the structure (primary citation in PDB) should be published; likewise the paper describing the structure cannot be curated if the structure has not been deposited in PDB, unless this article describes the epitope in the

Epitope Information		Epitope
Epitope	Epitope ID:	77491
Chemical Type:	Discontinuous protein residues	
Discontinuous Residues:	T85, A86, L87, E89, P91, T92, Y95, N98, Q100, S109, C110, T111, S112, K113, A114, V115, I162, K163, G164	
Source Name:	Major surface antigen p30 precursor	
Source Accession:	129348	
Source Organism ID:	5811	
Source Organism:	Toxoplasma gondii	
<b>Epitope Reference Details</b>		
Epitope Structure Defines:	Epitope containing region/antigenic site	
Epitope Evidence Code:	Exact match to reference information	
Epitope Name:	4F1E12 Fab epitope of SAG1	
Reference Region:	T2037, A2038, L2039, E2041, P2043, T2044, Y2047, N2050, Q2052, S2061, C2062, T2063, S2064, K2065, A2066, V2067, I2114, K2115, G2116	
B Cell Assay Information		Immunization
Host Organism ID:	10090	
Host Organism:	Mus musculus	
<b>1st In Vivo Process</b>		
In Vivo Process Type:	Administration in vivo	
<b>1st In Vivo Process Administration Details</b>		
Route:	Intraperitoneal (i.p.)	
<b>1st Immunogen</b>		
Epitope Relation:	Source Organism	
Object Type:	Organism	
Organism ID:	5811	
Organism:	Toxoplasma gondii	
<b>Immunogen Details</b>		
Immunogen Evidence Code:	Not determined	
Immunogen Reference Name:	Toxoplasma gondii	
<b>Immunization Comments</b>		
Immunization Comments:	mAb 4F1E12 was generated by intraperitoneal infection of mice with cysts from the 76K T. gondii strain, boosted one month later intravenously with formalin-treated, fixed and purified tachyzoites.	
B Cell Assay		Assay
Qualitative Measurement:	Positive	
Assay Type:	X-Ray Crystallography	
Assay Type Group:	Characterization of Ab binding	
Assay Type Units:	Angstroms	
<b>Measurement Details</b>		
Measurement Inequality:	=	
Quantitative measurement:	3.1	
Assayed Antibody		Antibody
Assayed Antibody Source Material:	Purified Immunoglobulin	
Assayed Antibody Immunoglobulin Domain:	Fab	
Assayed Antibody Purification Status:	Monoclonal	
Assayed Antibody Name:	4F1E12	
Assayed Antibody Heavy Chain Type:	IgG1	
Assayed Antibody Light Chain Type:	Kappa	
Antigen		Antigen
Epitope Relation:	Source Antigen	
Chemical Type:	Protein	
Molecule Name:	Major surface antigen p30 precursor	
Molecule Accession:	129348	
Organism ID:	5811	
Organism:	Toxoplasma gondii	
<b>Antigen Details</b>		
Antigen Evidence Code:	Exact match to reference information	
Antigen Conformation:	Native	
3D Structure of Complex		3D Structure of Complex
Complex PDB ID:	1YNT	
Antibody Chain 1 PDB Chain:	B	
Antibody Chain 2 PDB Chain:	A	
Antigen PDB Chain:	F	
Comments:	There is no drastic conformational change in the D1 or D2 domain of SAG1 observed comparing the unbound homodimeric form [PDB: 1K2Q] and Fab-bound monomeric structure.	
<b>Curated Contacts</b>		
Epitope Residues:	F: T2037, A2038, L2039, E2041, P2043, T2044, Y2047, N2050, Q2052, S2061, C2062, T2063, S2064, K2065, A2066, V2067, I2114, K2115, G2116	
Antibody Residues Interacting with Antigen:	A: Y32:L1, Y50:L2, R53:L2, N92:L3, T93:L3, L94:L3; B: T530:H1, D531:H1, G533:H1, I550:H2, S552:H2, T553:H2, Y554:H2, S555:H2, G556:H2, D557:H2, A558:H2, S559:H2, S599:H3, S600:H3, T601:H3, W602:H3, Y603:H3;	
Contact Area for Antibody:	1330	
View 3D Structure:	<a href="#">[View 3D Structure]</a> <a href="#">[View Curated Contacts XML file]</a>	
<b>Calculated Contacts</b>		
Epitope Residues:	F: T2037, A2038, L2039, E2041, P2043, T2044, Y2047, N2050, Q2052, C2062, T2063, S2064, K2065, A2066, V2067, T2068, I2114, K2115, G2116	
Antibody Residues Interacting with Antigen:	A: 530, Y32, Y50, N92, T93, L94; B: T530, D531, Y532, G533, S552, T553, Y554, S555, D557, A558, S559, Y560, S599, S600, T601, W602, Y603	
Contact Area for Antigen:	907.3	
Contact Area for Antibody:	842.1	
View 3D Structure:	<a href="#">[View 3D Structure]</a> <a href="#">[View Calculated Contacts XML file]</a>	
<b>Assay Reference Details</b>		
Assay Comments:	The structure contains two identical complexes in the asymmetric unit; therefore, only one (chains A,B,F) was curated. SAG1 monomer observed in the Fab-SAG1 complex crystal structure is independent of the binding of Fab. Epitope is located at one extremity of the SAG1 monomer N-terminal D1 domain that is not involved in dimer formation. Chain F in the structure is the rest part of the source antigen. Protein L (chain E) was used for crystallization purposes.	

**Figure 3.** Example of the B-cell response page. Detailed information is provided for the epitope (blue box) and B-cell response, including data on the immunization (purple box), assay (teal box), antibody (yellow box), antigen used in the X-ray crystallography experiment (green box) and the 3D structure of antigen-antibody complex (red box). The antigen-antibody interactions were curated from (12) and calculated from the coordinates provided in the PDB file [PDB: 1YNT]. The results of clicking on the links ‘View 3D Structure’ and ‘View Contacts XML file’ (red arrows) are shown in Figure 4.

**Table 1.** IEDB-3D content

	Epitope type	Total number of references <sup>a</sup>	Curated references <sup>b</sup>
<b>IEDB category</b>			
Infectious diseases	Peptidic	131	131
	Non-peptidic	11	4
Allergy	Peptidic	8	8
	Non-peptidic	5	3
Autoimmunity	Peptidic	39	30
	Non-peptidic	11	11
Cancer	Peptidic	40	16
	Non-peptidic	8	5
Transplant	Peptidic	22	18
	Non-peptidic	2	1
Other	Peptidic	241	130
	Non-peptidic	82	2
HIV		55	4
Total		655	365
<b>Structural category</b>			
B-cell epitopes	Peptidic	311	168
	Non-peptidic	108	20
T-cell epitopes	Peptidic	217	177
	Non-peptidic	19	0
Total		655	365

<sup>a</sup>PDB and PubMed search as of January 7, 2010.

<sup>b</sup>Curation status as of January 9, 2010.

context of immunological assays. Upon curation, both a PDB structure and journal article, which is a primary reference of the structure are considered as sources of information for curation.

Similar to other assays, for 3D structures, data on epitope molecular structure, epitope source, immunization that led to recognition of the epitope, assay, antigen, antibody, TCR or MHC are curated according to IEDB general procedure (1). An example of a B-cell response is given in Figure 3. The difference of IEDB-3D data compared to regular IEDB curations is the information captured under the ‘3D Structure of Complex’ heading.

In general, studies often do not provide the exact epitope structure recognized by an antibody/B-cell receptor, especially in the case of discontinuous B-cell epitopes. Therefore, discontinuous epitopes are often curated as partial epitopes, with only a few key residues captured. In cases where the epitope is curated from the structure this can be avoided; if the authors of the paper stated that only residues for part of the epitope were provided, we calculate the epitope residues from the PDB structure as the antigen residues separated from the antibody by 4 Å atomic distance (T-cell epitopes are curated as whole peptides or non-peptidic structures). The mapping between the numbers of the epitope residues provided in the paper and the external database is done manually, using the mapping between the PDB and UniProt numberings provided on the PDB website. For example, the epitope shown in Figure 3 had different residue numbers in the paper (‘Reference Region’ field at the top in Figure 3) compared to the numbering that matches the source

antigen sequence stored in the external database ('Discontinuous Residues' field).

Information on the 3D structure is provided in the section '3D Structure of Complex'. There are three types of tables with different fields for B-cell, T-cell and MHC-binding, respectively. The fields curated for B-cell responses, or structures of antibody-antigen complexes, are shown in Figure 3. The following structural information is curated: PDB ID, antibody and antigen chain IDs, epitope and antibody residues in PDB numbering, as well as CDR loops for the antibody residues and contact surface areas for the antibody and antigen if they are provided in the paper.

Interacting residues and contact areas are also calculated. The former are defined based on 4-Å atomic distance, according to the definition from (10), the latter are calculated using the NACCESS program (11). Pairwise atomic contacts are provided in the XML files available via hyperlinks (Figure 3) 'View Curated Contacts XML file' and 'View Calculated Contacts XML file'. The first link is provided only if the contacts were specified in the journal article and curated, the latter link allows the user to view and download the file providing calculated antibody-antigen (antigen-MHC, or antigen-TCR and MHC-TCR) inter-molecular contacts (Figure 4C). The following types of contacts are calculated: hydrogen bonds, salt bridges, van der Waals, hydrophobic and 4-Å interactions (interactions are defined in the EpitopeViewer tutorial at [http://spdc.sdsc.edu/iedb/epitopeViewer/EpitopeViewerTutorial\\_v2.0.htm](http://spdc.sdsc.edu/iedb/epitopeViewer/EpitopeViewerTutorial_v2.0.htm); this link is also available inside the EpitopeViewer application). The calculation is done during curation, using a php program that takes as an input curated data and PDB file and outputs two XML files with the interacting residues, contact areas and pairwise contacts. These files are also used as input for EpitopeViewer (Figure 4). The curators use the php tool and EpitopeViewer as part of the curation process and to check for errors.

EpitopeViewer allows the user to visualize, render and analyze the structure and save structural and contact views as high-quality pictures for publication (9). Figure 4 shows an example of how the EpitopeViewer can be used to analyze specific inter-molecular contacts. The residue Arg53 of the antibody light chain (Figure 3) was curated as part of the paratope; however, it can be seen that it is located relatively far from the nearest epitope atom (5.4 Å) (Figure 4A). At the same time, another light chain residue, Ser30, which was not curated as part of the paratope, contacts the epitope residues through van der Waals and 4-Å interactions (Figure 4B).

Additional detailed information on the 3D structure is captured in free text form in the 'Assay Comments' field (bottom of Figure 3). This can, for example, provide information on antigen and receptor conformational changes and comparison with other relevant structures if this information is mentioned in the paper. Also, the user can get further details on the PDB website via the PDB ID hyperlink in the 'Complex PDB ID' field.

## QUERYING IEDB-3D

Since IEDB-3D is fully integrated within IEDB, structural data, both curated and calculated, and all accompanying information can be queried using the multiple search capabilities implemented as part of the IEDB web site and described in this section.

On the IEDB home page (Figure 1), epitopes can be searched by epitope sequence, epitope source organism, source antigen and the immune recognition context, including the type of response (B-cell, T-cell or MHC-binding), host organism and MHC allele. If epitopes curated from 3D structures satisfy the search criteria, they will be retrieved together with all other epitopes curated in IEDB.

Using the keyword search option on the IEDB home page (the box is located in the right top corner of the page, Figure 1, red arrow), data are retrieved by keywords, PDB ID or SwissProt/GenBank ID; the keyword search runs throughout all fields in the database. In addition, the user can explicitly qualify what identifiers to search for when using the option 'Search by Identifier' from the 'Search' dropdown menu on the home page (Figure 1, yellow box). This type of search can be done, using IEDB internal identifiers, PDB ID, PubMed ID and CHEBI ID.

The 'Search' dropdown menu on the home page allows advanced search of any specified field in the database, including the fields related to 3D structure. For example, on the 'B Cell Search' webpage, the search fields related to 3D structure are made visible by clicking the '+' sign next to '3D Structure of Complex' within the 'B Cell Assay' subsection. Figure 5A shows how to search on the 'B Cell Search' page for the structures of antigens in complex with antibodies obtained via *in vivo* administration/immunization with *Toxoplasma gondii* and containing tryptophan (W) in the paratope (residues in the antibody interacting with the antigen). The result of this query is a single epitope that was curated from the structure with the PDB ID 1YNT (Figure 3).

Alternatively, epitopes for which 3D structural information is available can also be searched for by specifying a particular type of assay. For example, the epitope shown in Figure 3 can be found by specifying the assay type 'X-ray crystallography' using the 'Assay Finder' on the 'B Cell Search' form. Additionally, the resolution for the 3D structure can be specified.

In addition to the keyword, simple and advanced searches, epitopes curated from 3D structures can be accessed via the 'Browse by 3D Structure' page provided in the 'Browse' dropdown menu on the IEDB home page (Figure 1, red box). By expanding the 'B Cell', 'T Cell' and 'MHC Binding' trees the epitopes can be browsed by the organism that is the source of the antibody, T cell, and MHC molecule, respectively. This page also provides an up to date overview on the number of distinct epitope structures curated in IEDB-3D.

When clicking on an epitope ID (Figure 5C), all assays in which that epitope is characterized are returned.



**B Cell Search**

**Reference**

**Epitope** A

**Immunization**

Host Organism

**Host Details**

**1st In Vivo Process**  may or may not  MUST  MUST NOT be present in search results.

**1st Immunogen**  may or may not  MUST  MUST NOT be present in search results.

Epitope Relation

Type

Organism

**Immunogen Details**

**2nd In Vivo Process**  may or may not  MUST  MUST NOT be present in search results.

**2nd Immunogen**  may or may not  MUST  MUST NOT be present in search results.

**Immunization Comments**

**Adoptive Transfer**  may or may not  MUST  MUST NOT be present in search results.

**B Cell Assay**

Qualitative Measurement

Assay Type

**Measurement Details**

**Assayed Antibody**

**Antigen**

**3D Structure of Complex**

Complex PDB ID

Antibody Chain 1 PDB Chain

Antibody Chain 2 PDB Chain

Antigen PDB Chain

Viewer Flag  Any  Yes  No

Comments

**Curated Contacts**

**Calculated Contacts**

Epitope Residues

Antibody Residues Interacting with Antigen

Search Parameters:

- 1st In Vivo Process is present
- 1st Immunogen is present
- In Vivo1 Immunogen Epitope Relation equals 'Source Organism'
- In Vivo1 Immunogen Object Subtype equals 'Organism'
- In Vivo1 Immunogen Object Organism Name contains 'Toxoplasma gondii'
- Complex Calculated Antibody Antigen Residues contains 'W'

Epitopes	Positive*	Negative**	All
Peptidic	1	0	1
Non-Peptidic	0	0	0

Assays	Positive	Negative	All
T Cell Response	0	0	0
B Cell Response	1	0	1
MHC Ligand Elution	0	0	0
MHC Binding	0	0	0

B

**IMMUNE EPITOPE DATABASE AND ANALYSIS RESOURCE**

Home Browse Search Tools Support

Epitope

1 item(s) found, displaying 1 to 1 (Click the column headers to adjust the sorting)

Export all results:  (full)

Epitope ID	Structure	Source Antigen	Source Organism
77491	T85, A86, L87, E89, P91, T92, Y95, N98, Q100, S109...	Major surface antigen p30 precursor	Toxoplasma gondii

1 item(s) found, displaying 1 to 1

Export all results:  (full)

C

**IMMUNE EPITOPE DATABASE AND ANALYSIS RESOURCE**

Home Browse Search Tools Support More IEDB

Epitope Information

Distinct Epitope

Epitope ID:	77491
Discontinuous Residues:	T85, A86, L87, E89, P91, T92, Y95, N98, Q100, S109, C110, T111, S112, K113, A114, V115, I162, K163, G164
Source Antigen:	Major surface antigen p30 precursor
Source Organism:	Toxoplasma gondii

D

Source (1) Reference (1) B Cell Assay (2)

2 item(s) found, displaying 1 to 2 (Click the column headers to adjust the sorting)

Export all results:  (compact) |  (full)

B Cell ID	Reference	Epitope	Host	Immunization	Assay Antigen	Antigen Epitope Relation	Assay Description
1245875	Marc Gralle; J Mol Biol 2005	T85, A86, L87, E89, P91, T92, Y95, N98, Q100, S109... Major surface antigen p30 precursor Toxoplasma gondii	Mus musculus	Administration in vivo with Toxoplasma gondii (Source Organism)	Toxoplasma gondii Toxoplasma gondii	Source Organism	Ab Competition of Ab Binding Detection of Ab/Ag binding <b>Positive</b>
1246572	Marc Gralle; J Mol Biol 2005	T85, A86, L87, E89, P91, T92, Y95, N98, Q100, S109... Major surface antigen p30 precursor Toxoplasma gondii	Mus musculus	Administration in vivo with Toxoplasma gondii (Source Organism)	Major surface antigen p30 precursor Major surface antigen p30 precursor Toxoplasma gondii	Source Antigen	X-Ray Crystallography Characterization of Ab binding <b>Positive</b>

2 item(s) found, displaying 1 to 2

Export all results:  (compact) |  (full)

View 3D Structure

**Figure 5.** Example of the IEDB-3D query. (A) The advanced B-cell search page showing the query for 3D structures of antigens in complex with antibodies obtained via *in vivo* administration/immunization with *Toxoplasma gondii* and containing tryptophan (W) in the paratope calculated as the antibody residues interacting with the antigen at <4Å atomic distance. The result of clicking on the button 'Search' (red box) is shown in B. (B) The result of the search shown in A. The result of clicking on the number of epitopes (red box) is shown in C. (C) The list of epitopes as the result of the search shown in A. The result of clicking on the epitope ID (red box) is shown in D. (D) The list of assays for the epitope found using the query shown in A. The button 'View 3D Structure' (red arrow) launches the EpitopeViewer for curated data (Figure 4A). The result of clicking on the assay ID (red box) is shown in Figure 3.

## FUNDING

Funding for open access charge: National Institute of Health/National Institute of Allergy and Infectious Diseases (contract number HHSN2662004000 0 6C).

*Conflict of interest statement.* None declared.

## REFERENCES

1. Vita,R., Zarebski,L., Greenbaum,J.A., Emami,H., Hoof,I., Salimi,N., Damle,R., Sette,A. and Peters,B. (2010) The immune epitope database 2.0. *Nucleic Acids Res.*, **38**, D854–D862.
2. Korber,B.T.M., Brander,C., Haynes,B.F., Koup,R., Moore,J.P., Walker,B.D. and Watkins,D.I. (2007) HIV Molecular Immunology. *Los Alamos National Laboratory, Theoretical Biology and Biophysics, Los Alamos, New Mexico*, LA-UR 07-4752.
3. Berman,H.M., Westbrook,J., Feng,Z., Gilliland,G., Bhat,T.N., Weissig,H., Shindyalov,I.N. and Bourne,P.E. (2000) The Protein Data Bank. *Nucleic Acids Res.*, **28**, 235–242.
4. Ehrenmann,F., Kaas,Q. and Lefranc,M.P. (2010) IMGT/3Dstructure-DB and IMGT/DomainGapAlign: a database and a tool for immunoglobulins or antibodies, T cell receptors, MHC, IgSF and MhcSF. *Nucleic Acids Res.*, **38**, D301–D307.
5. Tong,J.C., Kong,L., Tan,T.W. and Ranganathan,S. (2006) MPID-T: database for sequence-structure-function information on T-cell receptor/peptide/MHC interactions. *Appl. Bioinformatics*, **5**, 111–114.
6. Schlessinger,A., Ofra,Y., Yachdav,G. and Rost,B. (2006) Epitome: database of structure-inferred antigenic epitopes. *Nucleic Acids Res.*, **34**, D777–D780.
7. Tong,J.C., Song,C.M., Tan,P.T., Ren,E.C. and Sinha,A.A. (2008) BEID: Database for sequence-structure-function information on antigen-antibody interactions. *Bioinformatics*, **3**, 58–60.
8. Huang,J. and Honda,W. (2006) CED: a conformational epitope database. *BMC Immunol.*, **7**, 7.
9. Beaver,J.E., Bourne,P.E. and Ponomarenko,J.V. (2007) EpitopeViewer: a Java application for the visualization and analysis of immune epitopes in the Immune Epitope Database and Analysis Resource (IEDB). *Immunome Res.*, **3**, 3.
10. Greenbaum,J.A., Andersen,P.H., Blythe,M., Bui,H.H., Cachau,R.E., Crowe,J., Davies,M., Kolaskar,A.S., Lund,O., Morrison,S. *et al.* (2007) Towards a consensus on datasets and evaluation metrics for developing B-cell epitope prediction tools. *J. Mol. Recognit.*, **20**, 75–82.
11. Hubbard,S.J. and Thornton,J.M. (1993) NACCESS. *Department of Biochemistry and Molecular Biology, University College, London*.
12. Graille,M., Stura,E.A., Bossus,M., Muller,B.H., Letourneur,O., Battail-Poirot,N., Sibai,G., Gauthier,M., Rolland,D., Le Du,M.H. *et al.* (2005) Crystal structure of the complex between the monomeric form of *Toxoplasma gondii* surface antigen 1 (SAG1) and a monoclonal antibody that mimics the human immune response. *J. Mol. Biol.*, **354**, 447–458.