

MINIREVIEW

Seed biopriming with plant growth promoting rhizobacteria: a review

Ahmad Mahmood^{1,2,*}, Oğuz Can Turgay¹, Muhammad Farooq³ and Rifat Hayat⁴

¹Department of Soil Science and Plant Nutrition, Faculty of Agriculture, Ankara University, 06110 Ankara, Turkey, ²Institute of Biochemical Plant Pathology, German Research Center for Environmental Health, Ingolstaedter Landstrasse 1, D-85764 Neuherberg, Munich, Germany, ³Department of Agronomy, University of Agriculture, Faisalabad 38040, Pakistan and ⁴Department of Soil Science and Soil Water Conservation, PMAS-Arid Agriculture University, Rawalpindi 46300, Pakistan

*Corresponding author: Department of Soil Science and Plant Nutrition, Faculty of Agriculture, Ankara University, 06110 Ankara, Turkey.

Tel: +00905070332247; E-mail: ahmadmahmood91@gmail.com

One sentence summary: The article reviews the potential of seed priming with plant growth promoting bacteria over conventional methods of bacterial application to the soil in improving plant productivity.

Editor: Gerard Muyzer

ABSTRACT

Beneficial microbes are applied to the soil and plant tissues directly or through seed inoculation, whereas soil application is preferred when there is risk of inhibitors or antagonistic microbes on the plant tissues. Insufficient survival of the microorganisms, hindrance in application of fungicides to the seeds and exposure to heat and sunlight in subsequent seed storage in conventional inoculation methods force to explore appropriate and efficient bacterial application method. Seed priming, where seeds are hydrated to activate metabolism without actual germination followed by drying, increases the germination, stand establishment and stress tolerance in different crops. Seed priming with living bacterial inoculum is termed as biopriming that involves the application of plant growth promoting rhizobacteria. It increases speed and uniformity of germination; also ensures rapid, uniform and high establishment of crops; and hence improves harvest quality and yield. Seed biopriming allows the bacteria to enter/adhere the seeds and also acclimatization of bacteria in the prevalent conditions. This review focuses on methods used for biopriming, and also the role in improving crop productivity and stress tolerance along with prospects of this technology. The comparison of methods being followed is also reviewed proposing biopriming as a promising technique for application of beneficial microbes to the seeds.

Keywords: biopriming; plant growth promoting rhizobacteria; inoculation

INTRODUCTION

Soil microbes since their discovery in late 18th century have been used extensively in crop production. Advent of technology allowed the researchers to study more about the microbial populations, and Kloepper and Schroth (1978) first time used the term plant growth promoting rhizobacteria (PGPR) explain-

ing them as bacteria which are closely related to rhizosphere. Kloepper, Lifshitz and Zablutowicz (1989) also used the term rhizobacteria. Functions and mechanisms of growth promotion by these microbes have been discussed, and microorganisms have been categorized in different classes (Hayat et al. 2010). Microbes actively involved in crop production are generally termed as

plant growth promoting bacteria (PGPB), whereas the bacteria isolated from the root zone are termed as plant growth promoting rhizobacteria (Kloepper and Schroth 1978). Major functions of these beneficial microbes are supply of nutrients to crops, stimulation of plant growth namely producing phytohormones, biocontrol of phytopathogens, improving soil structure, bioaccumulation of inorganic compounds and bioremediation of metal contaminated soils (Brierley 1985; Davison 1988; Ehrlich 1990; Middeldorp, Briglia and Salkinoja-Salonen 1990; Wilson and Lindow 1993; Burd, Dixon and Glick 2000; Zaidi et al. 2006).

Interaction between beneficial soil microbes and plants determines the plant health and soil fertility (Jeffries et al. 2003). The concept of sustainable agriculture has given much importance to the use of rhizospheric bacteria to help plants easy nutrient uptake and solubilization of fixed nutrients such as phosphorus (Hayat et al. 2010). It is the need of the time to reduce the agricultural inputs through combining beneficial microorganisms for better and sustainable agriculture. Several symbiotic such as *Rhizobium* spp. and *Frankia* spp. and asymbiotic bacteria such as *Azotobacter*, *Azospirillum*, *Bacillus* and *Klebsiella* spp. are used throughout the world to increase the crop growth and yield (Staley and Drahos 1994). Bacteria inhabiting plant rhizosphere are called PGPR which can promote the growth and productivity of plants through various mechanisms (Kloepper, Lifshitz and Zablutowicz 1989; Cleyet-Marcel et al. 2001). These rhizosphere inhabiting bacteria have also been categorized as nodule promoting rhizobacteria which is an important interaction of microbes with plants and plant health promoting rhizobacteria (Burr and Caesar 1984). PGPRs can also be categorized based on their relationships i.e. symbiotic and free-living soil inhabiting bacteria (Khan 2005). Gray and Smith (2005) also classified intercellular PGPR (iPGPR) called symbiotic bacteria and extracellular PGPR which are free-living bacteria. Rhizobia are famous iPGPR as they produce nodules in leguminous plants (Sriprang et al. 2003).

Podile and Kishore (2006) conclude several plant growth promoting (PGP) mechanisms of PGPR such as modification and increased branches in root hair, improvement in germination of seeds, enhanced and faster nodule performance, increase in leaf area per plant, release of certain phytohormones, augmented nutrients and water uptake by plants, increased biomass of the plants with more vigor growth and better carbohydrate accumulation which increases the growth of plant species. On the other hand, Glick (2003) categorizes the bacterial assisted plant growth in three different ways, including plant hormone production (Dobbelaere, Vanderleyden and Okon 2003), bacterial assisted better nutrient uptake by plants (Çakmakçi et al. 2006) and avoiding the diseases in plants through biological control (Saravanakumar et al. 2008). Dey et al. (2004) suggest the need of exploring other mechanisms of plant growth promotion by PGPR apart from the list already studied. Listing all the explored and investigated mechanisms of PGPR, following can be included: (a) solubilization and mineralization of nutrients notably phosphorus (Richardson 2001; Banerjee and Yesmin 2002); (b) nitrogen fixation through symbiosis and asymbiosis (Kennedy, Choudhury and Kecskés 2004); (c) release of certain plant hormones such as gibberellic acid and cytokinins (Dey et al. 2004), indole acetic acid (Patten and Glick 2002) and abscisic acid (Dobbelaere, Vanderleyden and Okon 2003); (d) production of 1-aminocyclopropane-1-carboxylate (ACC)-deaminase helping to lower ethylene level in roots this increasing length and vigor of roots (Li et al. 2000; Penrose and Glick 2001); (e) antagonism toward plant pathogens by producing substances such as cyanides and antibiotics (Glick and Pasternak 2003); (f) in-

creasing the availability of nutrients specifically of iron through chelating by producing siderophores (Glick and Pasternak 2003); (g) tolerance against several abiotic stresses such as oxidative (Stajner et al. 1995, 1997) and drought stress (Alvarez, Sueldo and Barassi 1996); (h) water soluble vitamin production including biotin, niacin, thiamine and riboflavin (Revillas et al. 2000); (i) detoxification of heavy metals (Ma et al. 2011); (j) tolerance of salinity (Tank and Saraf 2010); and (k) biological control of pests and insects (Russo et al. 2008).

Several studies have documented beneficial effects of certain rhizobial strains in improving growth of legumes as well as non-legumes. Second, inoculation of rhizobium in consortium with free-living rhizospheric bacteria has also given excellent results in improving crop growth and productivity (Kishore, Pande and Podile 2005; Tilak, Ranganayaki and Manoharachi 2006; Wani, Khan and Zaidi 2007). These PGPRs can be used effectively to meet the nutrient-deficient conditions and their use can be favorable to reduce the uses of chemical fertilizers and support of environment friendly crop productivity (Herrera, Salamanka and Barea 1993; Requena et al. 1997). The beneficial and plant growth enhancing effects of PGPR are well reported and explained. PGPR inoculation has increased different crop yields in normal and stress conditions. From the recent literature, PGPR inoculation increased the stress resistance and production of the crops, including tomato (Almaghrabi, Massoud and Abdelmoneim 2013), lettuce (Kohler et al. 2009), wheat (Jaderlund et al. 2008; Chakraborty et al. 2013; Nadeem et al. 2013; Islam et al. 2014; Kumar, Maurya and Raghuwanshi 2014), rice (Bal et al. 2013; Jha, Saxena and Sharma 2013; Lavakush et al. 2014), soybean (Masciarelli, Llanes and Luna 2014), groundnut (Paulucci et al. 2015), broad bean (Younesi and Moradi 2014), maize (Rojas-Tapias et al. 2012) and chickpea (Patel et al. 2012). The increase in yields and other yield parameters can be different in different crops and environments and normally range from 25% to 65%. Local reviews also indicate the growth promotion of crops by application of PGPR including wheat and barley (Ozturk, Caglar and Sahin 2003; Salantur et al. 2005; Turan, Çakmak and Şahin 2013), sugar beet (Sahin, Çakmakçi and Kantar 2004), strawberry (Esitken et al. 2010), apple (Aslantas, Çakmakçi and Sahin 2007), grapes (Köse, Güleriyüz and Demirtaş 2005) and raspberry (Orhan et al. 2006).

Bacterial inoculation to enhance the productivity of different crops is being practiced since the discovery of beneficial effects of these bacteria. The methods used for augmentation of the beneficial bacteria include seed coating, pelleting, foliar application and direct soil application where most commonly used is inoculation. Every method has been used with modifications according to the requirements. However, inoculation is most commonly used because it is easy to use and is practiced since the advent of this technique. Availability of sticking agent although is a limitation in this method but is still the most trusted method throughout the world. PGPR application through seed priming, soaking the seeds for premeasured time in liquid bacterial suspension, starts the physiological processes inside the seed while radicle and plumule emergence is prevented (Anitha et al. 2013) until the seed is sown. The start of physiological process inside the seed enhances the abundance of PGPR in the spermosphere (Taylor and Harman 1990). This proliferation of antagonist PGPR inside the seeds is 10-fold than attacking pathogens which enables the plant to survive those pathogens (Callan, Mathre and Miller 1990) increasing the use of biopriming for biocontrol too.

Method of application contributes mainly to the survival efficiency of the bacteria in the soil and on the seeds. Most common methods developed and explored include seed treatment, soil amendment and roots dipping in the bacterial suspensions

before transplanting particularly in rice. Other uncommon methods include foliar spray or application of bacteria through drip irrigation (Podile and Kishore 2006). PGPR are applied to the soil or seeds and/or to the plant parts when there is risk of inhibitors or antagonistic microbes on the plant tissues (Gindrat 1979). Diverse carrier materials have been tried and are being used depending on their quality to keep the bacteria viable for longer times as well as to reduce the desiccation chances along with the adhesive ability to the plant parts (Chao and Alexander 1984; Elegba and Rennie 1984). Proper inoculation procedures are followed as survival of bacterial cells depends mainly on the environmental conditions. Several carriers such as broth cultures, agar cultures and powder carriers have been used (Strijdom and Deschodt 1976; Thompson 1980), yet peat-based inoculants have shown good results and have been used widely but there are issues with peat-based inoculants such as exposure of peat based inoculants to high temperatures or water scarcity, presence of antagonist microorganisms and quality of peat strictly affects the bacterial viability (Chao and Alexander 1984). Exposure of the inoculated seed to sun causes the death of the bacterial cells as well as its exposure to environment can lead to contamination. Therefore, peat-involving inoculants are not yet considered as the best option as there can be pathogenic microbes causing plant diseases.

Use of proper carrier strongly influences the survival and colonization ability of the bacteria in the soil as well as in the roots. Peat soil is most preferred carrier material being used for inoculation of bacteria but its availability is a major limitation (Boonkerd and Singleton 2002). Similarly, rice husk is also being used in Asian countries. Trevors et al. (1992) found that mixing of bentonite clay in the carrier increased the survival of bacteria in fine textured soils. A similar study also suggested that mixing of 1% bentonite clay in fresh grown or freeze-dried *Rhizobium leguminosarum* suspension enhanced the bacterial survival markedly when compared to no amendment (Heijnen, Hok-A-Hin and Van Veen 1992). Heijnen, Hok-A-Hin and Van Elsas (1993) also reported that fresh cells showed less survival ability and colonization as compared to starved bacterial cells. Different soil amendments and chemical polymers have also been tried to entrap the bacteria in the carrier material, but a more promising report is use of barley straw which increased the survival of bacteria and also improved the root colonizing ability of the strains (Stephens 1994). Inoculation techniques are yet to be explored as there is scarce information available regarding the delivery and application of bacteria to the soil or the seeds. However, it is quite clear that population of the bacteria in soil is mainly dependent on initial stack of inoculums on the seed (Milus and Rothrock 1993). Hebbar et al. (1992) stated that application of more inoculums per seed can increase the efficiency but results are not always steady. Bacteria need to compete with other microbes to colonize so it can be concluded that introduced bacteria should be competitive enough to efficiently compete and colonize the roots.

Variety of methods have been used and studied by researchers for producing better inoculums which can survive better in the soil. The simplest strategy as explained by Paau (1989), local strains should be selected which are competent, adapted and dominant in a particular geographical area, and then mutant from the parent strain having more nitrogen fixing and competitive ability should be used in that particular area. This method or strategy is also used in preparation of microbial pesticides (Watrud et al. 1985) where mutation has been used. Effectiveness of the inoculum in the soil depends on the conditions after the release in the soil and if the conditions are optimum,

inoculum will survive better. Several laboratory microcosm and field studies have been conducted on bacterial survival potential in soil (van Elsas and Heijnen 1990). Presence of microniches in the soil enables the bacteria to survive after their application to soil otherwise reduction in bacterial number has been observed (van Elsas et al. 1986; Heijnen et al. 1988; Postma, Scheffers and van Dijken 1988; van Elsas and Heijnen 1990). Other factors affecting the bacterial survival in the soil include certain abiotic factors including soil temperature and moisture, nutrient presence and pH of the soil (García et al. 2010). Several studies have documented the effects of biotic and abiotic factors on survival of bacteria in soil (Bashan et al. 1995; Bashan and Vazquez 2000; Oliveira et al. 2004). Environmental factors affect the survival of bacteria in the soil, as an example fluorescent pseudomonad strain survived 10-fold better in sandy loam soil as compared to clay loam (Bahme and Schroth 1987; Pathma, Kennedy and Sakthivel 2011). Amending the soil with bentonite mineral increased the bacterial survival in loamy sand soil (Heijnen et al. 1988) through protection against protozoa (van Elsas and Heijnen 1990). It can be concluded that both biotic and abiotic factors affect bacterial survival and root colonization by bacteria in the soil (Campbell and Ephgrave 1983; Postma, Hok-A-Hin and Van Veen 1990). Most of studies indicate bottlenecks in various techniques of bacterial application either to the soil or to the plant tissues. Among different methods being used for introducing beneficial bacteria include seed coating and covering, root dipping, foliar application, direct soil application and seed inoculation which have various merits and demerits reviewed in Table 1.

Seed coating and covering is a general term where liquids or suspending solids are applied to the seed coat, prospectively to cover it homogeneously. This method requires use of adhesives to ensure proper coating of the seed which hinder the further application of pesticides to the seeds (Bardin and Huang 2003). Bacterial survival and nitrogen fixation was reduced when pelleting of the molybdenum was carried out along with bacterial inoculation and 99% bacteria were dead after 4 days (Burton and Curley 1966). Campo, Araujo and Hungria (2009) has also reported the drawback of applying micronutrient to seeds and inoculants together. Seed coating also hinders the gaseous exchange to the leguminous seeds which causes reduction in nitrogen fixation (Duarte et al. 2004), along with problems such as reducing the number of bacteria on the seeds due to desiccation. This technique is usually used for application of biocontrol agents (Paulitz, Zhou and Rankin 1992).

Dipping the roots in bacterial suspension has been used for biocontrol, and very few evidences can be found. Srinivasan et al. (2009) applied this technique and found that it is possible option in controlling *Fusarium wilt* of tomato. Munif, Hallmann and Sikora (2013) studied the effect of endophytic bacteria against *Meloidogyne incognita* using root dipping technique and found less number of galls on treated plants. Another report is from Esitken et al. (2010), who used root dipping along with foliar application and have reported significant increases in yield parameters of strawberry. Root dipping however needs prepared plant nursery which is not very economical in most of the crops.

Foliar application is not widely practice by the researchers or the farmers for the augmentation of these significant bacteria to the plants. However, in some cases such as biocontrol of fungus, this application has been used (Obradovic et al. 2004). Another research group has applied the PGPR through both root dipping and foliar application and have concluded that it increased the yield and yield parameters of fruits such as strawberry (Esitken et al. 2010), apricot (Esitken et al. 2002), sweet cherry (Esitken et al.

Table 1. Advantages and disadvantages of different application methods.

Method	Advantages	Disadvantages	Reference
Career-based inoculation	Easy availability Easy to prepare Lower cost	Contamination of the inoculants by unwanted microbes from career such as peat No uniformity on the career Short-term storage ability	Brockwell (1977); Brockwell, Gault and Chase (1977); Bezdicek et al. (1978); Gault, Chase and Brockwell (1982); Bashan, Levanony and Ziv-Vecht (1987); Brockwell, Holliday and Pilka (1988); Rice and Olsen (1988); Kosanke et al. (1992); Smith (1992); Bashan (1998)
Seed coating and covering	Easier to apply No specific machinery needed Practiced by farmers in case of pesticide application to seeds	Application of pesticides to the seeds Sticking agents harmful to bacteria Flexibility in seeding is less	Brockwell (1977); Brockwell, Holliday and Pilka (1988); Bashan and Levanony (1990); Bashan and Carrillo (1996); Bashan and Holguin (1997); Bashan (1998)
Pelleting	Easy to apply Favored by farmers Flexibility in seeding and application Lime pellets can be used for acid soils	Survival of bacteria is hindered due to lower moisture levels Special machinery needed to prepare thus increases cost	Brockwell (1977); Bezdicek et al. (1978); Bordeleau and Prevost (1981); Bashan and Levanony (1990); Bashan (1998)
Direct soil application	Injection in the root zone is possible Easy and simple	Exposure to the sun Desiccation problems Needs more volume	Brockwell (1977); Bordeleau and Prevost (1981); Bashan and Levanony (1990); Bashan (1998)
Root dipping	Nursery required Simple and easy	Large amount of liquid media and bacterial cells needed Contamination from environment quite normal	Brockwell (1977); Bordeleau and Prevost (1981); Bashan and Levanony (1990); Bashan (1998)

2006) and apple (Pirlak et al. 2007). Sudhakar et al. (2000) also investigated the effect of foliar application of *Azotobacter*, *Azospirillum* and *Beijerinckia* on mulberry and have reported positive effects.

Application of the inoculum directly to the soil is favored when there is threat of presence of antagonistic microbes or pesticides on the plant tissues (Gindrat 1979). Presence of inhibitory compounds on the plant tissues also inhibits plant part inoculation. Soil application needs large amount of inoculants which contradicts with the economics of the farming. Solid inoculants are easy but if there are liquid inoculants, it needs special care from the transportation and after the application to the field.

Application of beneficial bacteria to the seeds is generally called as inoculation. It is the most common method been used since the beneficial bacteria have been studied and discovered. Seed inoculation involves use of carrier material for better transportation and application, use of adhesives to ensure the sticking of bacteria to the seeds and sometimes other materials avoiding desiccation of the inoculum (Elegba and Renne 1984). Peat-based inoculants are most common and extensively used since the discovery of rhizobium for leguminous crops. Peat being easily available and a cheap source is sterilized and milled so used as carrier for most of the inoculation material (Walker, Rossall and Asher 2004). Most favored and commonly used method of inoculation includes application of adhesive agents on the seeds followed by inoculum spreading under shade (Vincent, Thompson and Donovan 1962). Among the adhesive agents, most commonly used are Arabic gum, sugar solution, methylcellulose, polyvinylpyrrolidone, caseinate salts and polyvinylacetate (Deaker, Roughly and Kennedy 2004). Inoculation usually produces favorable results with rhizobia; however, their development limit has been reached (Burton 1976; Thomp-

son 1980). As discussed above, extreme environmental factors such as high temperatures decrease the viable cell count in the inoculum (Chao and Alexander 1984). Apart from this, it has several drawbacks depending on the nature and type of peat and issues of peat availability in different countries (Bashan et al. 2002). Polymer-based inoculants can be used over peat-based inoculants but they are expensive and need more biotechnical handling (Fages 1992). As far as polymer-based inoculants are concerned, they are also being opposed as they are hazardous to environment (Cassidy, Lee and Trevors 1996). Merits and demerits of different application methods are described in Table 1.

MECHANISMS INVOLVED IN SEED COLONIZATION

Efficient colonization supports better functioning of plant beneficial bacteria (Compant, Clément and Sessitsch 2010). Diverse endophytic bacteria which spend part of their life inside the plant tissue without causing any disease (Döbereiner 1992); colonize different parts of plants without causing any damage (Bacon and Hinton 2006; Ali et al. 2014) and similar to phytopathogens, they enter the plants through various mechanisms. Entry through wounded plant parts (Agarwal and Shende 1987), stomatal openings (Roos and Hattingh 1983), lenticels (Scot et al. 1996), germinating radicles (Gagné et al. 1987) and root cracks (Sørensen and Sessitsch 2006) includes different colonization processes where root cracks entry helps root inoculation by bacteria (Ali et al. 2014).

Bacteria after soil application tend to colonize rhizosphere (Gamalero et al. 2003) followed by adherence to root surfaces and finally to the rhizodermis making a string of bacteria (Hansen

et al. 1997). Bacteria form biofilms or microcolonies on the rhizodermal cells where the colonization occurs (Benizri, Baudoin and Guckert 2001). Rhizosphere colonization is linked to photosynthates translocation to roots and exudation (Lugtenberg and Dekkers 1999; Bais et al. 2006) along with root mucilage (Knee et al. 2001). These exudates include diverse kind of organic acids, amino acids and carbohydrates which serve as food for most of bacteria inhabiting rhizosphere (Walker et al. 2003). This chemotaxis helps bacteria in multiplication along with colonization (Lugtenberg and Kamilova 2009) but when limited results in reduced root colonization (de Weert et al. 2002). Concentration and composition of root exudates also influences colonization where colonization occurs on different levels proportional to concentration of exudates (Gamalero et al. 2004). Soil characteristics and nutrient availability have also been reported as factors influencing colonization (Krafczyk, Trolldenier and Beringer 1984; Paterson and Sim 2000). Plant pathogenic infection also affects the root colonization processes, e.g. plant released malic acid to attract bacteria against the infection of pathogen where the bacteria protected the roots of the plants by creating biofilm (Rudrappa et al. 2008).

Plant beneficial bacteria also have to compete with the local bacteria and other soil organisms in the root zone for colonization (Walker et al. 2003) and under severe competitive conditions, PGPB also secrete siderophores and lytic enzymes to limit growth of plant pathogens (Compant, Clément and Sessitsch 2010), metabolites (Haas and Défago 2005), and they also release certain antibiotic compounds for better colonization (van Loon and Bakker 2005). Production of several other compounds such as amino acids, vitamins, enzymes and polysaccharides has also been reported enhancing root colonization (Vesper 1987; de Weger et al. 1989; Simons et al. 1997; Dekkers et al. 1998; Camacho et al. 2002). Physically, flagella of the bacteria help them making contact with exudates (Turbull et al. 2001) but are not always important for root colonization (Scher et al. 1988). Quorum sensing based on cell density is also involved in colonization of rhizosphere and rhizoplane (Soto, Sanjuán and Olivares 2006), which might be linked to enhancing competitive ability of PGPB (Compant, Clément and Sessitsch 2010).

Legumes show symbiosis with members of Rhizobiaceae family and this symbiosis needs exchange of resources (Giordano and Hirsch 2004; Ahemad and Kibret 2014). Endophytic colonization needs penetration of bacteria inside the plant tissues which then show the PGP traits (Hallmann and Berg 2006). Nodulating bacteria have evolved certain processes of entry like introduction through cortex and lateral root fissures and intercellular cracks forming specialized organs called nodules by penetrating in the roots through utilizing flavonoids and nod genes from such microbes (Garg and Geetanjali 2007; Compant, Clément and Sessitsch 2010). This type of colonization involves physical (Böhm, Hurek and Reinhold-Hurek 2007) and further chemical mechanisms including cell-wall-degrading enzyme production (Lodewyckx et al. 2002). Most of the *Rhizobium* species have been found to produce indole acetic acid (Ahemad and Khan 2012), which is essential for process of nodule formation through cell division and differentiation along with vascular tissue formation (Ahemad and Kibret 2014). Thus, higher auxin levels in legume plants are responsible for nodule formation (Spaepen, Vanderleyden and Remans 2007; Glick 2012) and symbiotic relationships. The PGPR showing non-symbiotic interaction with plants often contribute very small amount of nitrogen (Glick 2012). Diazotrophs being free-living nitrogen-fixing soil bacteria show non-obligate relationship with the non-leguminous plants (Glick et al. 1999).

HOST SPECIFICITY IN PGPR APPLICATIONS

Host specificity depends on particular bacterial strains to non-specific traits of host plant or non-specific bacterial strains to particular traits of the host plants, but evolution has played its role in preferential interaction between host and bacterial strains (Drogue et al. 2013). Bacterial association with particular hosts involves interaction and recognition process (Benizri, Baudoin and Guckert 2001). The recognition process involves root exudates concentration and composition where composition of root exudates depends on the cultivars, stress condition and plant growth stage (Haichar et al. 2008). In other studies, claiming no particular host specificity found in *Azospirillum* indicated that chemotaxis was however strain specific, and the bacteria showed preference toward exudates of their isolated host (Bacilio-Jiménez et al. 2003; Pedraza et al. 2010). Chemical signals from the plants as root exudates serve as attractants to microbes (Doty 2011). Host specificity can be influenced by chemotaxis and metabolic activities can be its determinant and provision of nutrients by the host plant also plays important role in specifying bacteria to the plants (Reinhold, Hurek and Fendrik 1985; Buyer, Roberts and Russek-Cohen 2002; Reis, dos Santos Teixeira and Pedraza 2011). Plant genetic makeup is important in determining microbiome associations with roots and rhizosphere (Bulgarelli et al. 2015). In detailed study regarding genome wide study in *Arabidopsis-Pseudomonas*, it was observed and concluded that plants genetics is the core element of benefiting from PGPR and identification of genes responsible for host specificity is needed (Wintermans, Bakker and Pieterse 2016). Several genes responsible for chemotaxis, flagella formation, transportation and metabolic pathways are involved in root colonization which complete recognition and chemotaxis (Compant, Clément and Sessitsch 2010). Studies regarding host specificity and microbial presence in the plant roots have been made easy through use of next-generation sequencing techniques in recent years (Bai et al. 2015), but yet extensive research in this area is lacking. Nitrogen-fixing bacteria can be applied to unrelated plants (Doty 2011), and application of such bacteria in non-leguminous plants enhanced growth and productivity (Bhattacharjee, Singh and Mukhopadhyay 2008). Different PGPR can promote the growth and productivity of diverse crops depending on genetics of the host and exudates released, and also ability of beneficial bacteria to compete and colonize rhizosphere and roots (Vessey 2003).

Keeping in mind the prospects of biopriming, this review focuses on (i) the comparison of past bacterial application methods with their drawbacks, (ii) suggesting the technique biopriming as a promising method for bacterial application in increasing stress resistance and crop productivity.

BIOPRIMING

Since the advent of seed priming, a lot of work has been done on this aspect of seed treatment and is now common in most of the areas for delayed sowing and to obtain vigorous plant growth. As defined by McDonald (1999), seed priming is soaking the seeds in any solution containing our required priming agent followed by redrying the seeds which result into start of germination process except the radicle emergence. Among different priming techniques, hydration using any biological compound is termed as biopriming (Ashraf and Foolad 2005). Seed priming creates ideal conditions for the bacterial inoculation and colonization in the seed (McQuilken, Rhodes and Halmer 1998). Soaking the seeds in the bacterial suspension for precalculated period of time to

Table 2. Role of biopriming in different PGP activities.

Strains under study	Crop	PGP activities	Reference
<i>Azotobacter chroococcum</i> , <i>A. lipoferum</i>	Barley	Increase in 1000-grain weight, dry matter accumulation, grain yield, biological yield and harvest index	Mirshekari et al. (2012)
<i>Pseudomonas</i> spp.	Safflower	Increased number of branches, heads per plant, diameter of head, grain number per head, grains per plant, 1000 grain weight, oil content and grain yield	Sharifi (2012)
<i>Azotobacter chroococcum</i> <i>Azospirillum lipoferum</i> , <i>A. chroococcum</i> <i>A. lipoferum</i>	Maize	Grain yield, crop growth rate and dry matter accumulation	Sharifi (2011)
<i>Azotobacter</i> and <i>Azospirillum</i> spp.	Maize	Increase in grain yield, plant height, number of kernels per ear and number of grains per ear row	Sharifi and Khavazi (2011)
<i>Pseudomonas fluorescens</i>	Sunflower	Shoot height, root length and seedling weight	Moeinzadeh et al. (2010)
<i>Clonostachys rosea</i> , <i>P. chlororaphis</i> , <i>P. fluorescens</i> , <i>T. harzianum</i> , <i>T. viride</i>	Carrot and onion	Increase in emergence and yield	Bennett, Mead and Whipps (2009)

allow the bacterial imbibition into the seed is known as biopriming (Abuamsha, Salman and Ehlers 2011). Reddy (2013) explained biopriming more in biocontrol aspect as application of beneficial bacterial inoculum to the seeds and their hydration to protect seeds against disease control. This soaking of seeds in bacterial suspension initiates the physiological processes in the seed where plumule and radicle emergence is prevented (Anitha et al. 2013), until the seeds have temperature and oxygen after being sown. PGPR keep on multiplying in the seed and proliferate in the spermosphere (Taylor and Harman 1990) even before sowing. Seed biopriming is being focused as it ensures the entrance of endophytic bacteria into the sides along with avoiding the effect of high temperature. Biopriming treatment is potentially able to promote quick and even germination as well as better plant growth (Moeinzadeh et al. 2010). Biopriming with rhizospheric bacteria has been reported in crops such as carrot (Jensen et al. 2002), sweet corn (Callan, Mathre and Miller 1990, 1991) and tomato (Harman and Taylor 1988; Legro and Satter 1995; Warren and Bennett 1999). In case of efficacy and survival of biological agents, priming has been reported beneficial and been reported to enhance the plant growth and yield (Harman, Taylor and Stasz 1989; Callan, Mathre and Miller 1990, 1991; Warren and Bennett 1999). Germination and enhanced seedling establishment is obtained through seed priming with PGPR (Anitha et al. 2013). Bio-osmopriming can significantly enhance the uniformity of the germination and plant growth traits when associated with bacterial coating (Bennett 1998). Uniformity in germination and better stand establishment options when considered, biopriming is favored method. Biopriming has been practiced and explained by different researchers (Callan, Mathre and Miller 1991; Bennett, Mead and Whipps 2009; Moeinzadeh et al. 2010; Chakraborty et al. 2011; Sharifi 2011, 2012; Sharifi and Khavazi 2011; Gururani et al. 2012; Mirshekari et al. 2012) in several ways, but still is an ambiguous approach which needs to be explored and discussed.

There are different methods used explaining biopriming varying in the temperature and time duration of soaking the seeds (Miché and Balandreau 2001; Gholami, Shahsavani and Nezarat 2009; Abuamsha, Salman and Ehlers 2011; Sharifi and Khavazi 2011; Sharifi, Khavazi and Gholipouri 2011; Carrozzi et al. 2012; Firuzsalari, Mirshekari and Khochebagh 2012; Saber et al.

2012; Kasim et al. 2013; Reddy 2013). Some of the researchers have also surface disinfected the seeds before soaking into the bacterial suspension (Sharifi, Khavazi and Gholipouri 2011; Firuzsalari, Mirshekari and Khochebagh 2012; Saber et al. 2012; Reddy 2013).

BIOPRIMING AND CROP PRODUCTIVITY

Biopriming contributes to various PGP activities which has been studied by researchers as reviewed in Table 2. Saber et al. (2012) used the technique biopriming with a commercial biofertilizer having different bacterial species including *Bacillus lentus*, *B. subtilis*, *Pseudomonas fluorescens*, *P. putida* and *Azospirillum* spp. They observed increase in several agromorphological traits of wheat plants. In addition, they also postulate that requirement of nitrogen and phosphorus was decreased in bioprimed plants as compared to control plants. Stem and total seedling fresh weight was increased with the priming of PGPR in maize seedlings in a laboratory experiment (Gholami, Shahsavani and Nezarat 2009). Barley seed priming with a consortium of *Azotobacter chroococcum* and *Azospirillum lipoferum* in combination with 80 kg ha⁻¹ urea and 60 kg ha⁻¹ P₂O₅ significantly increased the yield attributes such as thousand grain weight, dry matter accumulation, biological yield, grain yield and harvest index (Mirshekari et al. 2012). In maize, different *Azotobacter* and *Azospirillum* strains were used for biopriming of the seeds, and the results showed that biopriming significantly increased the crop growth rate, dry matter accumulation and grain yield (Sharifi 2011). Different bacterial strains were also investigated for biopriming in safflower, and it was observed that seed priming with *Pseudomonas* strain 186 with coapplication of 180 kg ha⁻¹ increased the number of branches, heads per plant, diameter of head, grain number per head, grains per plant, 1000 grain weight and grain yield of the plants (Sharifi 2012).

ROLE OF BIOPRIMING IN RESISTANCE AGAINST ABIOTIC STRESSES

Kasim et al. (2013) used the technique biopriming to document its effects against drought stress. They used two strains including *A. brasilense* and *B. amyloliquefaciens* and observed that

Table 3. Role of biopriming in abiotic stress tolerance.

Strains under study	Mechanism of action	Crop	Role in stress tolerance	PGP activities	Reference
<i>Bacillus pumilus</i> , <i>B. firmus</i>	ACC-deaminase activity, IAA production, phosphate solubilization, phytate mineralization, siderophore production	Potato	Salinity, drought, heavy metal stress tolerance	Increase in plant height, No. of leaves plant ⁻¹ , No. of tubers plant ⁻¹ , tuber yield plant ⁻¹	Gururani et al. (2012)
<i>Bacillus cereus</i>	Phosphate solubilization, IAA, catalase, protease, chitinase and siderophore production, nitrate reduction, starch hydrolysis	Rice, mungbean, chickpea	Salinity tolerance	Increase in seedling height, number and length of leaves, root and shoot biomass	Chakraborty et al. (2011)
<i>Agrobacterium rubi</i> , <i>Burkholderia gladii</i> , <i>P. putida</i> , <i>B. subtilis</i> , <i>B. megaterium</i>	–	Radish	Improved seed germination under saline conditions	Increase in seed germination	Kaymak et al. (2009)

Table 4. Role of biopriming in biotic stress tolerance.

Strains under study	Crop	Role in stress tolerance	Reference
<i>Trichoderma harzianum</i>	Maize	<i>Fusarium verticillioides</i> and fumonisins tolerance	Nayaka et al. (2010)
<i>Pseudomonas fluorescens</i>	Sunflower	<i>Alternaria blight</i> tolerance	Rao et al. (2009)
<i>Clonostachys rosea</i>	Carrot	<i>Alternaria dauci</i> and <i>Al. radicina</i> tolerance	Jensen et al. (2004)
<i>Pseudomonas fluorescens</i>	Pearl millet	Downy mildew tolerance	Raj, Shetty and Shetty (2004)
<i>Pseudomonas aureofaciens</i>	Sweet corn	<i>Pythium ultimum</i> tolerance	
<i>Pseudomonas fluorescens</i>	Sweet corn	Damping-off tolerance	Callan, Mathre and Miller (1991)

biopriming with these strains increased drought tolerance in wheat plants through upregulation of genes related to stress. Role of biopriming has been studied in various crops using different PGPR as compiled in Table 3. Role of biopriming in salinity stress tolerance is widely studied and promising results have been recorded. Most notable genus used in abiotic stress tolerance is *Bacillus* which is used in potato (Gururani et al. 2012), radish (Kaymak et al. 2009) rice, mungbean and chickpea (Chakraborty et al. 2011).

ROLE OF BIOPRIMING IN RESISTANCE AGAINST BIOTIC STRESSES

Biopriming has been applied in various crops for the biocontrol of several diseases (Table 4). Abuamsha, Salman and Ehlers (2011) applied *Serratia plymuthica* and *P. chlororaphis* to the different oilseed rape cultivars for the control of a pathogen *Leptosphaeria maculans* causing blackleg disease, and it was observed that disease extent was reduced up to 71.6% by *S. plymuthica* and 54% by *P. chlororaphis*. Seed biopriming gave the highest control over *Verticillium longisporum* as compared to coating the bacteria on the seeds (Müller and Berg 2008). Biopriming has been reported to control damping-off disease in various crops such as cucumber (Pill et al. 2009), maize (Callan, Mathre and Miller 1990), pea (Taylor, Harman and Nielsen 1994) and soybeans. Rao et al. (2009) applied a biocontrol agent *P. fluorescens* through seed biopriming and observed that incidence of *Alternaria blight* was reduced and biopriming helped the plants to tolerate the disease incidence efficiently. In maize, biocontrol agent *Trichoderma harzianum* was applied which resulted in better control of *F. verticillioides* and *fumonisins* (Nayaka et al. 2010). Similarly, different biocontrol agents were applied to the seeds through bioprim-

ing, and better biocontrol was observed in radish (Kaymak et al. 2009), carrot (Jensen et al. 2004), sweet corn (Bennett 1997) and pearl millet (Niranjan, Shetty and Shetty 2004).

Microbes capable of colonizing the rhizosphere and plant roots can protect the plants to pathogens through antagonistic interaction (Buchenauer 1998, Berg et al. 2001, Whipps 2001). They can also induce systemic resistance to the plants which can reduce the fungal infection (Compant et al. 2005). During the seed germination, successful antagonizing microbe colonization helps in reducing the pathogenic attack on the plant (Weller 1983). They can also induce systemic resistance to the plants which can reduce the fungal infection (Compant et al. 2005). Jensen et al. (2004) reported that death of carrot plants due to seedborne pathogens such as *Al. radicina* and *Al. dauci* was significantly reduced with biopriming of the seeds with *Clonostachys rosea* and was as effective as use of fungicide iprodione. Root rot caused by different pathogens such as *Macrophomina phaseolina*, *F. solani* and *Rhizoctonia solani* was reduced in cowpea through biopriming of the seeds with *T. harzianum* by 56.3%–64% at the pre-emergence and 57.1%–64% at the post-emergence stage (El-Mohamedy, Abd-Alla and Badiaa 2006). In faba bean, biopriming with different bacterial strains was tested to reduce the incidence of root rot, and it was observed that use of the biopriming technique can be used as economical, safe and easy to apply biocontrol method (El-Mougy and Abdel-Kader 2008). *Trichoderma harzianum* is the main focus of the researchers in terms of biopriming and has been used widely in different crops. Another evidence of *T. harzianum* with coapplication of *P. fluorescens* and *B. subtilis* as biopriming significantly reduced the incidence of root rot pathogenic disease caused by *F. solani* and *R. solani* in pea under greenhouse and field conditions (El-Mohamedy and Abd-El-Baky 2008).

ECONOMICS OF BIOPRIMING

Several workers have encouraged this technique being a cost-effective approach for the biocontrol of different pathogenic microbes (Rao *et al.* 2009) and application of beneficial bacteria to the soil. Along with the crop productivity, biopriming can also be favored as the potential technique for biocontrol of several plant pathogens. Control of these plant pathogens is usually carried out by using costly pesticides where we can promote this technique as dual purpose technology enhancing the plant productivity and stress resistance side by side.

CONCLUSION AND FUTURE PROSPECTS

Regarding the application of the bacteria, it has been explained by the scientists that biopriming can be used effectively in application of the bacteria as it gives enough number of bacteria in the seeds. Competition of the our desired inoculants with local bacteria is also a problem which can be addressed by biopriming as our desired bacteria will already be inside the seeds reducing the chance of desiccation as well as harmful effects of any pesticides applied to the field. On the other basis, it can also be an alternative approach for the application of bacteria to small seeded crops which can imbibe the bacterial suspension resulting in entrance of bacteria inside the seed. Biopriming gives equal or better control against several root rot diseases so can be used commercially as an alternative to fungicides successfully. In the application, there is need to search for the more better media for application due to cost hurdles which can definitely be reduced by further research. Second, this method can be implied to other crops yet not experimented which will give better picture of potential of this technology.

Conflict of interest. None declared.

REFERENCES

- Abuamsha R, Salman M, Ehlers R. Effect of seed priming with *Serratia plymuthica* and *Pseudomonas chlororaphis* to control *Leptosphaeria maculans* in different oilseed rape cultivars. *Eur J Plant Pathol* 2011;**130**:287–95.
- Agarwhal S, Shende ST. Tetrazolium reducing microorganisms inside the root of Brassica species. *Curr Sci* 1987;**56**:187–8.
- Ahemad M, Khan MS. Productivity of greengram in tebuconazole-stressed soil, by using a tolerant and plant growth-promoting Bradyrhizobium sp. MRM6 strain. *Acta Physiol Plant* 2012;**34**:245–54.
- Ahemad M, Kibret M. Mechanisms and applications of plant growth promoting rhizobacteria: Current perspective. *J King Saud Univ-Sci* 2014;**26**:1–20.
- Ali S, Duan J, Charles TC *et al.* A bioinformatics approach to the determination of genes involved in endophytic behavior in Burkholderia spp. *J Theor Biol* 2014;**343**:193–8.
- Almagrabi OA, Massoud SI, Abdelmoneim TS. Influence of inoculation with plant growth promoting rhizobacteria (PGPR) on tomato plant growth and nematode reproduction under greenhouse conditions. *Saudi J Biol Sci* 2013;**20**:57–61.
- Alvarez MI, Sueldo RJ, Barassi CA. Effect of *Azospirillum* on coleoptile growth in wheat seedlings under water stress. *Cereal Res Commun* 1996;**24**:101–8.
- Anitha D, Vijaya T, Reddy NV *et al.* Microbial endophytes and their potential for improved bioremediation and biotransformation: a review. *Indo Am J Pharmaceutical Res* 2013;**3**:6408–17.
- Ashraf M, Foolad MR. Pre-sowing seed treatment-a shotgun approach to improve germination growth and crop yield under saline and none-saline conditions. *Adv Agron* 2005;**88**:223–71.
- Aslantas R, Cakmakci R, Sahin F. Effect of plant growth promoting rhizobacteria on young apples trees growth and fruit yield under orchard conditions. *Sci Hortic* 2007;**111**:371–7.
- Bacilio-Jiménez M, Aguilar-Flores S, Ventura-Zapata E *et al.* Chemical characterization of root exudates from rice (*Oryza sativa*) and their effects on the chemotactic response of endophytic bacteria. *Plant Soil* 2003;**249**:271–7.
- Bacon CW, Hinton DM. Bacterial endophytes: The endophytic niche, its occupants, and its utility. In: Gnanamanickam SS (ed). *Plant-Associated Bacteria*. Dordrecht, The Netherlands: Springer, 2006, 155–94.
- Bahme JB, Schroth MN. Spatial-temporal colonization patterns of a rhizobacterium on underground organs of potato. *Phytopathology* 1987;**77**:1093–100.
- Bai Y, Müller DB, Srinivas G *et al.* Functional overlap of the Arabidopsis leaf and root microbiota. *Nature* 2015;**528**:364–9.
- Bais HP, Weir TL, Perry LG *et al.* The role of root exudates in rhizosphere interactions with plants and other organisms. *Ann Rev Plant Biol* 2006;**57**:233–66.
- Bal HB, Nayak L, Das S *et al.* Isolation of ACC deaminase PGPR from rice rhizosphere and evaluating their plant growth promoting activity under salt stress. *Plant Soil*, 2013;**366**:93–105.
- Banerjee MR, Yesmin L. Sulfur oxidizing rhizobacteria: an innovative environment friendly soil biotechnological tool for better canola production. *Proc Agroenviron*, Cairo, Egypt, 2002,1–7.
- Bardin SD, Huang HC. Efficacy of stickers for seeds treatment with organic matter or microbial agents for the control of damping-off of sugar beet. *Plant Path Bul* 2003;**12**:19–26.
- Bashan Y. Inoculants for plant growth-promoting bacteria in agriculture. *Biotechnol Adv* 1998;**16**:729–70.
- Bashan Y, Carrillo A. Bacterial inoculants for sustainable agriculture. In: Pérez-Moreno J, Ferrera-Cerrato R (eds). *New Horizons in Agriculture: Agroecology and Sustainable Development*. Proceedings of the 2nd international symposium on agroecology, sustainable agriculture and education. San Luis Potosi, Mexico, 16-18.11.1994. Montecillo, Mexico: Colegio de Postgraduados en ciencias agrícolas, 1996, 125–55.
- Bashan Y, Holguin G. Short- and medium- term avenues for *Azospirillum* inoculation. In: Ogoshi A, Kobayashi K, Homma Y *et al.* (eds). *Plant Growth-Promoting Rhizobacteria -Present Status and Future Prospects*. Sapporo, Japan: Faculty of Agriculture, Hokkaido University, 1997, 130–49.
- Bashan Y, Levanony H. Current status of *Azospirillum* inoculation technology: *Azospirillum* as a challenge for agriculture. *Can J Microbiol* 1990;**36**:591–608.
- Bashan Y, Levanony H, Ziv-Vecht O. The fate of field-inoculated *Azospirillum brasilense* Cd in wheat rhizosphere during the growing season. *Can J Microbiol* 1987;**33**:1074–9.
- Bashan Y, Li CY, Lebsky VK *et al.* Primary colonization of volcanic rocks by plants in arid Baja California, Mexico. *Plant Biol* 2002;**4**:392–402.
- Bashan Y, Puente ME, Rodriguez-Mendoza MN *et al.* Survival of *Azospirillum brasilense* in the bulk soil and rhizosphere of 23 soil types. *Appl Environ Microb* 1995;**61**:1938–45.
- Bashan Y, Vazquez P. Effect of calcium carbonate, sand, and organic matter levels on mortality of five species of *Azospirillum* in natural and artificial bulk soils. *Biol Fert Soils* 2000;**30**:450–9.
- Benizri E, Baudoin E, Guckert A. Root colonization by inoculated plant growth rhizobacteria. *Biocontrol Sci Techn* 2001;**11**:557–74.

- Bennett AJ, Mead A, Whipps JM. Performance of carrot and onion seed primed with beneficial microorganisms in glasshouse and field trials. *Biol Control* 2009;51:417–26.
- Bennett MA. Biological seed treatments for improved corn germination and emergence. In: *Proceeding Of the 51st Annual ASTA Corn and Sorghum Research Conference* 51. Washington, DC: American Seed Trade Association, Inc., 1997, 200–11.
- Bennett MA. The use of biologicals to enhance vegetable seed quality. *Seed Technol* 1998;20:198–208.
- Berg G, Fritze A, Roskot N et al. Evaluation of potential biocontrol rhizobacteria from different host plants of *Verticillium dahliae* Kleb. *J Appl Microbiol* 2001;156:75–82.
- Bezdicke DF, Evans DW, Abeda B et al. Evaluation of peat and granular inoculum for soybean yield and N₂ fixation under irrigation. *Agron J* 1978;70:865–8.
- Bhattacharjee RB, Singh A, Mukhopadhyay SN. Use of nitrogen-fixing bacteria as biofertilizer for non-legumes: prospects and challenges. *Appl Microbiol Biot* 2008;80:199–209.
- Böhm M, Hurek T, Reinhold-Hurek B. Twitching motility is essential for endophytic rice colonization by the N₂-fixing endophyte *Azoarcus* sp. strain BH72. *Mol Plant-Microbe In* 2007;20:526–33.
- Boonkerd N, Singleton P. Production of rhizobium biofertilizer. In: Kannaiyan S (ed). *Biotechnology of Biofertilizers*. New Delhi, India: Narosa Publishing House, 2002, 122–8.
- Bordeleau LM, Prevost D. Quality of commercial legume inoculants in Canada. In: Clarke KW, Stephens JHG (eds). *Proceedings of the 8th North American Rhizobium Conference*. Winnipeg: University of Manitoba, 1981, 562–5.
- Brierley JA. Use of microorganisms for mining metals. In: Halvorson HO, Pramer D, Rogul M (eds). *Engineered Organisms in The Environment: Scientific Issues*. Washington DC: ASM Press, 1985, 141–6.
- Brockwell J. Application of legume seed inoculants. In: Hardy RWF, Gibson AH (eds). *A Treatise on Dinitrogen Fixation. IV. Agronomy and Ecology*. New York: John Wiley & Sons, 1977, 277–309.
- Brockwell J, Gault RR, Chase DL. Inoculating soybeans with root-nodule bacteria. *Farmers' Newsl*. In: *Irrigation Research and Extension Committee*, Vol. 104, NSW, Australia: Griffith University, 1977, 7–12.
- Brockwell J, Holliday RA, Pilka A. Evaluation of the symbiotic fixing potential of soils by direct microbiological means. *Plant Soil* 1988;108:163–70.
- Buchenaer H. Biological control of soil-borne diseases by rhizobacteria. *J Plant Dis Protect* 1998;105:329–48.
- Bulgarelli D, Garrido-Oter R, Münch PC et al. Structure and function of the bacterial root microbiota in wild and domesticated barley. *Cell Host Microb* 2015;17:392–403.
- Burd GI, Dixon DG, Glick BR. Plant growth-promoting bacteria that decrease heavy metal toxicity in plants. *Can J Microbiol* 2000;46:237–45.
- Burr TJ, Caesar A. Beneficial plant bacteria. *Crit Rev Plant Sci* 1984;2:1–20.
- Burton JC. Methods of inoculating seeds and their effect on survival of rhizobia. In: Nutman PS (ed). *Symbiotic Nitrogen Fixation in Plants*. London: Cambridge University Press, 1976, 175–85.
- Burton JC, Curley RL. Compatibility of *Rhizobium japonicum* and sodium molybdate when combined in a peat carrier medium. *Agron J* 1966;58:327–30.
- Buyer JS, Roberts DP, Russek-Cohen E. Soil and plant effects on microbial community structure. *Can J Microbiol* 2002;48:955–64.
- Çakmakçı R, Dönmez F, Aydın A et al. Growth promotion of plants by plant growth-promoting rhizobacteria under greenhouse and two different field soil conditions. *Soil Biol Biochem* 2006;38:1482–7.
- Callan NW, Mathre DE, Miller JB. Bio-priming seed treatment for control of *Pythium ultimum* pre emergence damping-off in sh-2 sweet corn. *Plant Dis* 1990;74:368–72.
- Callan NW, Mathre DE, Miller JB. Field performance of sweet corn seed bio-primed and coated with *Pseudomonas fluorescens* AB254. *Hortscience* 1991;26:1163–5.
- Camacho MMC, Wijffes AHM, Mulders IHM et al. Characterization of NADH dehydrogenases of *Pseudomonas fluorescens* WCS365 and their role in competitive root colonization. *Mol Plant-Microbe In* 2002;15:662–71.
- Campbell R, Ephgrave YM. Effect of bentonite clay on the growth of *Gaeumannomyces graminis* var. tritici and its interactions with antagonistic bacteria. *J Gen Microbiol* 1983;129:771–7.
- Campo RJ, Araujo RS, Hungria M. Nitrogen fixation with the soybean crop in Brazil: compatibility between seed treatment with fungicides and Bradyrhizobial inoculants. *Symbiosis* 2009;48:154–63.
- Carrozzi LE, Creus CM, Barassi CA et al. Reparation of aged lettuce (*Lactuca sativa*) seeds by osmotic priming and *Azospirillum brasilense* inoculation. *Botany* 2012;90:1093–102.
- Cassidy MB, Lee H, Trevors JT. Environmental applications of immobilized microbial cells: a review. *J Ind Microbiol* 1996;16:79–101.
- Chakraborty AP, Dey P, Chakraborty B et al. Plant growth promotion and amelioration of salinity stress in crop plants by a salt-tolerant bacterium. *Rec Res Sci Technol* 2011;3:61–70.
- Chakraborty U, Chakraborty BN, Chakraborty AP et al. Water stress amelioration and plant growth promotion in wheat plants by osmotic stress tolerant bacteria. *World J Micro Biot* 2013;29:789–803.
- Chao WL, Alexander M. Mineral soils as carriers for *Rhizobium* inoculants. *Appl Environ Microb* 1984;47:94–7.
- Cleyet-Marcel JC, Larcher M, Bertrand H et al. Plant growth enhancement by rhizobacteria. In: Morot Gaudry JF (ed). *Nitrogen Assimilation by Plants: Physiological, Biochemical And Molecular Aspects*. Plymouth, UK: Science Publishers Inc., 2001, 185–97.
- Compant S, Clément C, Sessitsch A. Plant growth-promoting bacteria in the rhizo- and endosphere of plants: their role, colonization, mechanisms involved and prospects for utilization. *Soil Biol Biochem* 2010;42:669–78.
- Compant S, Duffy B, Nowak J et al. Use of plant growth-promoting bacteria for biocontrol of Plant Diseases: principle, mechanisms of action, and future prospects. *Appl Environ Microb* 2005;71:4951–9.
- Davison J. Plant beneficial bacteria. *Nat Biotechnol* 1988;6:282–6.
- de Weert S, Vermeiren H, Mulders IHM et al. Flagella-driven chemotaxis towards exudate components is an important trait for tomato root colonization by *Pseudomonas fluorescens*. *Mol Plant-Microbe In* 2002;15:1173–80.
- de Weger LA, Bakker PAHM, Schippers B et al. *Pseudomonas* spp. with mutational changes in the O-antigenic side chain of their lipopolysaccharide are affected in their ability to colonize potato roots. In: Lugtenberg BJJ (ed). *Signal Molecules in Plants and Plant-Microbe Interactions*. Berlin: Springer, 1989, 197–202.
- Deaker R, Roughly RJ, Kennedy IR. Legume seed inoculation technology—a review. *Soil Biol Biochem* 2004;36:1275–88.
- Dekkers LC, van der Bij AJ, Mulders IHM et al. Role of the O antigen of lipopolysaccharide, and possible roles of growth rate

- and of NADH: ubiquinone oxidoreductase (nuo) in competitive tomato root-tip colonization by *Pseudomonas fluorescens* WCS365. *Mol Plant-Microbe In* 1998;11:763–71.
- Dey R, Pal KK, Bhatt DM et al. Growth promotion and yield enhancement of peanut (*Arachis hypogaea* L.) by application of plant growth-promoting rhizobacteria. *Microbiol Res* 2004;159:371–94.
- Dobbelaere S, Vanderleyden J, Okon Y. Plant growth-promoting effects of diazotrophs in the rhizosphere. *Crit Rev Plant Sci* 2003;22:107–49.
- Döbereiner J. Recent changes in concepts of plant bacteria interactions: endophytic N₂ fixing bacteria. *Ciência Cultura* 1992;44:310–3.
- Doty SL. Nitrogen-fixing endophytic bacteria for improved plant growth. In: *Bacteria in Agrobiological Plant Growth Responses*. Springer, 2011, 183–99.
- Drogue B, Sanguin H, Borland S et al. Host specificity of the plant growth-promoting cooperation between *Azospirillum* and rice. In: *Symposium on Bacterial Genetics and Ecology*. Slovénie: Ljubljana, 2013, 1.
- Duarte CR, Neto JLV, Lisboa MH et al. Experimental study and simulation of mass distribution of the covering layer of soybean seeds coated in a spouted bed. *Braz J Chem Eng* 2004;21:59–67.
- Ehrlich HL. *Geomicrobiology*, 2nd edn. New York: Marcel Dekker, 1990, 646.
- Elegba MS, Rennie RJ. Effect of different inoculant adhesive agents on rhizobial survival, nodulation, and nitrogenase (acetylene-reducing) activity of soybeans (*Glycine max* (L.) Merrill). *Can J Soil Sci* 1984;64:631–6.
- El-Mohamedy RSR, Abd-Alla MA, Badiia RI. Soil amendment and bio-priming treatments as alternative fungicides for controlling root rot diseases on cowpea plants in Nobria province. *Res J Agr Biol Sci* 2006;2:391–8.
- El-Mohamedy RSR, Abd-El-Baky MMH. Effect of seed treatment on control of root rot disease and improvement growth and yield of pea plants. *Middle Eastern Russian J Plant Sci Biotech* 2008;2:84–90.
- El-Mougy NS, Abdel-Kader MM. Long-term activity of bio priming seed treatment for biological control of faba bean root rot pathogens. *Australas Plant Path* 2008;37:464–71.
- Esitken A, Karlidag H, Ercisli S et al. Effects of foliar application of *Bacillus subtilis* OSU-142 on the yield, growth and control of shot-hole disease (*Coryneum* blight) of apricot. *Gartenbauwissenschaft* 2002;67:139–42.
- Esitken A, Pirlak L, Turan M et al. Effects of floral and foliar application of plant growth promoting rhizobacteria (PGPR) on yield, growth and nutrition of sweet cherry. *Sci Hortic* 2006;110:324–7.
- Esitken A, Yildiz HE, Ercisli S et al. Effects of plant growth promoting bacteria (PGPB) on yield, growth and nutrient contents of organically grown strawberry. *Sci Hortic* 2010;124:62–6.
- Fages J. An industrial view of *Azospirillum* inoculants: formulation and application technology. *Symbiosis* 1992;13:15–26.
- Firuzsalari SM, Mirshekari B, Khochebagh SB. Effect of seed inoculation with bio-fertilizer on germination and early growth of corn. *Int Res J App Basic Sci* 2012;3:1097–102.
- Gagné S, Richard C, Rouseau H et al. Xylem-residing bacteria in alfalfa roots. *Can J Microbiol* 1987;33:996–1000.
- Gamalero E, Lingua G, Berta G et al. Methods for studying root colonization by introduced beneficial bacteria. *Agronomie* 2003;23:407–18.
- Gamalero E, Lingua G, FG Capri et al. Colonization pattern of primary tomato roots by *Pseudomonas fluorescens* A6RI characterized by dilution plating, flow cytometry, fluorescence, confocal and scanning electron microscopy. *FEMS Microbiol Ecol* 2004;48:79–87.
- García R, Bælum J, Fredslund L et al. Influence of temperature and predation on survival of *Salmonella enterica* Serovar Typhimurium and expression of *invA* in soil and manure-amended soil. *Appl Environ Microb* 2010;76:5025–31.
- Garg N, Geetanjali. Symbiotic nitrogen fixation in legume nodules: process and signaling. A review. *Agron Sustain Dev* 2007;27:59–68.
- Gault RR, Chase DL, Brockwell J. Effects of spray inoculation equipment on the viability of *Rhizobium* spp. in liquid inoculants for legumes. *Aust J Exp Agr Anim Hus* 1982;22:299–309.
- Gholami A, Shahsavani S, Nezarat S. The effect of plant growth promoting rhizobacteria (PGPR) on germination, seedling growth and yield of maize. *Proc World Acad Sci Eng Technol* 2009;49:19–24.
- Gindrat D. *Alternaria radicina*, an important parasite of market garden Umbelliferae. *Revue Suisse de Viticulture, d'Arboriculture et d'Horticulture* 1979;11:257–67.
- Giordano W, Hirsch AM. The expression of MaEXP1, a *Melilotus alba* expansin gene, is upregulated during the sweetclover-Sinorhizobium meliloti interaction. *Mol Plant Microbe* 2004;17:613–22.
- Glick BR. Phytoremediation: synergistic use of plants and bacteria to cleanup the environment. *Biotechnol Adv* 2003;21:383–93.
- Glick BR. *Plant Growth-Promoting Bacteria: Mechanism and Applications*, Vol. 2012. Scientifica: Hindawi Publishing Corporation, Article ID 963401, 2012.
- Glick BR, Patten CL, Holguin G et al. *Biochemical and genetic mechanisms used by plant growth promoting bacteria*, World Scientific, 1999.
- Glick BR, Pasternak JJ. Plant growth promoting bacteria. In: Glick BR, Pasternak JJ (eds). *Molecular Biotechnology—Principles and Applications of Recombinant DNA*, 3rd edn. Washington, DC: ASM Press, 2003, 436–54.
- Glick BR, Patten CL, Holguin G, Penrose D. *Biochemical and genetic mechanisms used by plant growth promoting bacteria*. World Scientific 1999.
- Gray EJ, Smith DL. Intracellular and extracellular PGPR: commonalities and distinctions in the plant-bacterium signaling processes. *Soil Biol Biochem* 2005;37:395–412.
- Gururani MA, Upadhyaya CP, Baskar V et al. Plant growth-promoting rhizobacteria enhance abiotic stress tolerance in *Solanum tuberosum* through inducing changes in the expression of ROS-scavenging enzymes and improved photosynthetic performance. *J Plant Growth Regul* 2012;32:245–58.
- Haas D, Défago G. Biological control of soil-borne pathogens by fluorescent *Pseudomonads*. *Nat Rev Microbiol* 2005;3:307–19.
- Haichar FZ, Marol C, Berge O et al. Plant host habitat and root exudates shape soil bacterial community structure. *ISME J* 2008;2:1221–30.
- Hallmann J, Berg G. Spectrum and population dynamics of bacterial root endophytes. In: Schulz B, Boyle C, Sieber N (eds). *Soil Biology*, Vol. 9. Berlin, Germany: Springer, 2006, 15–31.
- Hansen ML, Kregelund L, Nybroe O et al. Early colonization of barley roots by *Pseudomonas fluorescens* studied by immunofluorescence technique and confocal laser scanning microscopy. *FEMS Microbiol Ecol* 1997;23:353–60.
- Harman GE, Taylor AG. Improved seedling performance by integration of biological control agents at favorable pH levels with solid matrix priming. *Phytopathology* 1988;78:520–5.

- Harman GE, Taylor AG, Stasz TE. Combining effective strains of *Trichoderma harzianum* and solid matrix priming to improve biological seed treatments. *Plant Dis* 1989;73:631–7.
- Hayat R, Ali S, Amara U et al. Soil beneficial bacteria and their role in plant growth promotion: a review. *Ann Microbiol* 2010;60:579–98.
- Hebbar P, Davey AG, Merrin J et al. *Pseudomonas cepacia*, a potential suppressor of maize soil borne diseases: seed inoculation and maize root colonization. *Soil Biol Biochem* 1992;24:999–1007.
- Heijnen CE, Hok-A-Hin CH, Van Elsas JD. Root colonization by *Pseudomonas fluorescens* introduced into soil amended with bentonite. *Soil Biol Biochem* 1993;25:239–46.
- Heijnen CE, Hok-A-Hin CH, Van Veen JA. Improvements to the use of bentonite clay as a protective agent increasing survival levels of bacteria introduced into soil. *Soil Biol Biochem* 1992;24:533–8.
- Heijnen CE, Van Elsas JJ, Kuikman PJ et al. Dynamics of *Rhizobium leguminosarum* biovar trifolii introduced to soil; the effect of bentonite clay on predation by Protozoa. *Soil Biol Biochem* 1988;20:483–8.
- Herrera MA, Salamanka CP, Barea JM. Inoculation of woody legumes with selected arbuscular mycorrhizal fungi and rhizobia to recover desertified Mediterranean ecosystems. *Appl Environ Microb* 1993;59:129–33.
- Islam F, Yasmeen T, Ali Q et al. Influence of *Pseudomonas aeruginosa* as PGPR on oxidative stress tolerance in wheat under Zn stress. *Ecotox Environ Safe* 2014;104:285–93.
- Jaderlund L, Arthurson V, Granhall U et al. Specific interactions between arbuscular mycorrhizal fungi and plant growth-promoting bacteria: as revealed by different combinations. *FEMS Microbiol Lett* 2008;287:174–80.
- Jeffries P, Gianinazzi S, Perotto S et al. The contribution of arbuscular mycorrhizal fungi in sustainable maintenance of plant health and soil fertility. *Biol Fert Soils* 2003;37:1–16.
- Jensen B, Knudsen IMB, Madsen M et al. Biopriming of infected carrot seed with antagonist, *Clonostachys rosea*, selected for control of seedborne *Alternaria* spp. *Phytopathology* 2004;94:551–60.
- Jensen B, Poulsen FV, Knudsen IMB et al. Combining microbial seed treatment with priming of carrot seeds for control of seed borne *Alternaria* spp. In: Elad Y, Freeman S, Monte E (eds). *Biocontrol Agents: Mode of Action and Interaction with Other Means of Control*. Cited in IOBC WPRS Bulletin 24, Dijon: INRA, 2002, 197–201.
- Jha A, Saxena J, Sharma V. An investigation on phosphate solubilization potential of agricultural soil bacteria as affected by different phosphorus sources, temperature, salt and pH. *Commun Soil Sci Plan* 2013;44:2443–58.
- Kasim WA, Osman ME, Omar MN et al. Control of drought stress in wheat using plant growth-promoting bacteria. *J Plant Growth Regul* 2013;32:122–30.
- Kaymak HC, Güvenç I, Yaralı F et al. The effects of biopriming with PGPR on germination of radish (*Raphanus sativus* L.) seeds under saline conditions. *Turkish J Agr Forest* 2009;33:173–9.
- Kennedy IR, Choudhury A, Kecskés ML. Non-symbiotic bacterial diazotrophs in crop-farming systems: can their potential for plant growth promotion be better exploited? *Soil Biol Biochem* 2004;36:1229–44.
- Khan AG. Role of soil microbes in the rhizospheres of plants growing on trace metal contaminated soils in phytoremediation. *J Trace Elem Med Bio* 2005;18:355–64.
- Kishore GK, Pande S, Podile AR. Phylloplane bacteria increase seedling emergence, growth and yield of field-grown groundnut (*Arachis hypogaea* L.). *Lett Appl Microbiol* 2005;40:260–8.
- Kloepper JW, Lifshitz R, Zablutowicz RM. Free-living bacterial inocula for enhancing crop productivity. *Trends Biotechnol* 1989;7:39–43.
- Kloepper JW, Schroth MN. Plant growth promoting rhizobacteria on radishes. In: *Proceedings of the Fourth International Conference on Plant Pathogen Bacteria*, Vol. 2. INRA, Gilbert-Clarey, Tours, France, 1978, 879–82.
- Knee EM, Gong FC, Gao M et al. Root mucilage from pea and its utilization by rhizosphere bacteria as a sole carbon source. *Mol Plant-Microbe In* 2001;14:775–84.
- Kohler J, Hernández JA, Caravaca F, Roldán A. Induction of antioxidant enzymes is involved in the greater effectiveness of a PGPR versus AM fungi with respect to increasing the tolerance of lettuce to severe salt stress. *Environ Exp Bot* 2009;65:245–52.
- Kosanke J, Osborn R, Shuppe G et al. Slow rehydration improves the recovery of dried bacterial populations. *Can J Microbiol* 1992;38:520–5.
- Köse C, Güleriyüz MŞF, Demirtaş I. Effects of some plant growth promoting rhizobacteria (PGPR) on graft union of grapevine. *J Sustain Agr* 2005;26:139–47.
- Krafczyk I, Trolldenier G, Beringer H. Soluble exudates of maize: influence of potassium supply and rhizosphere microorganisms. *Soil Biol Biochem* 1984;16:315–22.
- Kumar A, Maurya BR, Raghuvanshi R. Isolation and characterization of PGPR and their effect on growth, yield and nutrient content in wheat (*Triticum aestivum* L.). *Biocatalysis Agric Biotechnol* 2014;3:121–8.
- Lavakush YJ, Verma JP, Jaiswal DK et al. Evaluation of PGPR and different concentration of phosphorus level on plant growth, yield and nutrient content of rice (*Oryza sativa*). *Ecol Eng* 2014;62:123–8.
- Legro B, Satter H. Biological control of *Pythium* through seed coating and seed priming with *Trichoderma*. In: Bradford K, Hartz T (eds). *Monterey Proceedings of the 4th National Symposium on Stand Establishment of Horticultural Crops*, Monterey, California, 1995, 235–7.
- Li J, Ovakim DH, Charles TC et al. An ACC-Deaminase minus mutant of *Enterobacter cloacae* UW4 no longer promotes root elongation. *Curr Microbiol* 2000;41:101–5.
- Lodewyckx C, Vangronsveld J, Porteous F et al. Endophytic bacteria and their potential applications. *Crit Rev Plant Sci* 2002;21:583–606.
- Lugtenberg B, Kamilova F. Plant-growth-promoting rhizobacteria. *Ann Rev Microbiol* 2009;63:541–56.
- Lugtenberg BJJ, Dekkers LC. What makes *Pseudomonas* bacteria rhizosphere competent? *Environ Microbiol* 1999;1:9–13.
- Ma Y, Prasad MNV, Rajkumar M et al. Plant growth promoting rhizobacteria and endophytes accelerate phytoremediation of metalliferous soils. *Biotechnol Adv* 2011;29:248–58.
- McDonald MB. Seed deterioration: physiology, repair and assessment. *Seed Sci Technol* 1999;27:177–237.
- McQuilken MP, Rhodes DJ, Halmer P. Application of microorganisms to seeds. In: Burges HD (ed). *Formulation of Microbial Biopesticides, Beneficial Microorganisms, Nematodes and Seed Treatments*. Dordrecht: Kluwer Academic Publishers, 1998, 255–85.
- Masciarelli O, Llanes A, Luna V. A new PGPR co-inoculated with *Bradyrhizobium japonicum* enhances soybean nodulation. *Microbiol Res* 2014;169:609–15.

- Miché L, Balandreau J. Effects of rice seed surface sterilization with hypochlorite on inoculated *Burkholderia vietnamiensis*. *Appl Environ Microb* 2001;**67**:3046–52.
- Middeldorp PJM, Briglia M, Salkinoja-Salonen MS. Biodegradation of pentachlorophenol in natural polluted soil by inoculated *Rhodococcus chlorophenolica*. *Microb Ecol* 1990;**20**:123–39.
- Milus EA, Rothrock CS. Rhizosphere colonization of wheat by selected soil bacteria over diverse environments. *Can J Microbiol* 1993;**39**:335–41.
- Mirshakari B, Hokmalipour S, Sharifi RS et al. Effect of seed bio-priming with plant growth promoting rhizobacteria (PGPR) on yield and dry matter accumulation of spring barley (*Hordeum vulgare* L.) at various levels of nitrogen and phosphorus fertilizers. *J Food Agric Environ* 2012;**10**:314–20.
- Moeinzadeh A, Sharif-Zadeh F, Ahmadzadeh M et al. Biopriming of sunflower (*Helianthus annuus* L.) seed with *Pseudomonas fluorescens* for improvement of seed invigoration and seedling growth. *Aust J Crop Sci* 2010;**4**:564.
- Müller H, Berg G. Impact of formulation procedures on the effect of the biocontrol agent *Serratia plymuthica* HRO-C48 on *Verticillium* wilt in oilseed rape. *Biol Control* 2008;**53**:905–16.
- Munif A, Hallmann J, Sikora RA. The influence of endophytic bacteria on *Meloidogyne incognita* infection and tomato plant growth. *J ISSAAS* 2013;**19**:68–74.
- Nadeem SM, Zaheer ZA, Naveed M et al. Mitigation of salinity-induced negative impact on the growth and yield of wheat by plant growth-promoting rhizobacteria in naturally saline conditions. *Ann Microbiol* 2013;**63**:225–32.
- Nayaka SC, Niranjana SR, Shankar ACU et al. Seed biopriming with novel strain of *Trichoderma harzianum* for the control of toxigenic *Fusarium verticillioides* and *fumonisin* in maize. *Arch Phytopathol PFL* 2010;**43**:264–82.
- Niranjana SR, Shetty NP, Shetty HS. Seed bio-priming with *Pseudomonas fluorescens* isolates enhances growth of pearl millet plants and induces resistance against downy mildew. *Int J Pest Manage* 2004;**50**:41–8.
- Obradovic A, Jones JB, Momol MT et al. Management of tomato bacterial spot in the field by foliar applications of bacteriophages and SAR inducers. *Plant Dis* 2004;**88**:736–40.
- Oliveira ALM, Canuto EL, Silva EE et al. Survival of endophytic diazotrophic bacteria in soil under different moisture levels. *Braz J Microbiol* 2004;**35**:295–9.
- Orhan E, Esitken A, Ercisli S et al. Effects of plant growth promoting rhizobacteria (PGPR) on yield, growth and nutrient contents in organically growing raspberry. *Sci Hortic* 2006;**111**:38–43.
- Ozturk A, Caglar O, Sahin F. Yield response of wheat and Barley to inoculation of plant growth promoting rhizobacteria at various levels of nitrogen fertilizers. *J Plant Nutr Soil Sc* 2003;**166**:262–6.
- Paau AS. Improvement of rhizobium inoculants. *Appl Environ Microb* 1989;**55**:862–5.
- Patel HA, Patel RK, Khristi SM et al. Isolation and characterization of bacterial endophytes from *Lycopersicon esculentum* plant and their plant growth promoting characteristics. *Nepal J Biotechnol* 2012;**2**:37–52.
- Paterson E, Sim A. Effect of nitrogen supply and defoliation on loss of organic compounds from roots of *Festuca rubra*. *J Exp Bot* 2000;**51**:1449–57.
- Pathma J, Kennedy RK, Sakthivel N. Mechanisms of fluorescent pseudomonads that mediate biological control of phytopathogens and plant growth promotion of crop plants. In: Maheshwari DK (ed). *Bacteria in Agrobiology: Plant Growth Responses*. Berlin: Springer, 2011, 86.
- Patten CL, Glick BR. Role of *Pseudomonas putida* indole acetic acid in development of host plant root system. *Appl Environ Microb* 2002;**48**:3795–801.
- Paulitz TC, Zhou T, Rankin L. Selection of rhizosphere bacteria for biological control of *Pythium aphanidermatum* on hydroponically-grown cucumber. *Biol Control* 1992;**2**:226–37.
- Paulucci NS, Gallarato LA, Reguera YB et al. *Arachis hypogaea* PGPR isolated from Argentine soil modifies its lipids components in response to temperature and salinity. *Microbiol Res* 2015;**173**:1–9.
- Pedraza OR, Mentel IM, Ragout LA et al. Plant growth-promoting bacteria: the role of chemotaxis in the association *Azospirillum brasilense*-plant. In: Frank C (ed). *Chemotaxis: Types, Clinical Significance, and Mathematical Models*, Chapter 2. New York, USA: Nova Science Publishers, Inc., 2010, 1–31.
- Penrose DM, Glick BR. Levels of 1-aminocyclopropane-1-carboxylic acid (ACC) in exudates and extracts of canola seeds treated with plant growth-promoting bacteria. *Can J Microbiol* 2001;**47**:368–72.
- Pill WG, Collins CM, Goldberger B et al. Responses of non-primed or primed seeds of “Marketmore 76” cucumber (*Cucumis sativus* L.) slurry coated with *Trichoderma* species to planting in growth media infested with *Pythium aphanidermatum*. *Sci Hortic* 2009;**121**:54–62.
- Pirlak L, Turan M, Sahin F et al. Floral and foliar application of plant growth promoting rhizobacteria (PGPR) to apples increases yield, growth, and nutrition element contents of leaves. *J Sustain Agr* 2007;**30**:145–55.
- Podile AR, Kishore GK. Plant growth promoting rhizobacteria. In: Gnanamanickam SS (ed.). *Plant Associated Bacteria*. Dordrecht, The Netherlands: Springer, 2006, 195–230.
- Postma E, Scheffers WA, van Dijken JP. Adaptation of the kinetics of glucose transport to environmental conditions in the yeast *Candida utilis* CBS 621: a continuous culture study. *J Gen Microbiol* 1988;**134**:1109–16.
- Postma J, Hok-A-Hin CH, Van Veen JA. Role of microniches in protecting introduced *Rhizobium leguminosarum* biovar trifolii against competition and predation in soil. *Appl Environ Microb* 1990;**56**:495–502.
- Raj SN, Shetty NP, Shetty HS. Note: Proline—an inducer of resistance against pearl millet downy mildew disease caused by *Sclerospora graminicola*. *Phytoparasitica* 2004;**32**:523–7.
- Rao MS, Ramachandran N, Sowmya DS et al. Biological control of nematode induced disease complex in certain vegetable crops. *Abstracts of International Conference on Horticulture*. Bangalore, India, 2009, 213.
- Reddy PP. *Recent Advances in Crop Protection*. India: Springer, 2013, 83.
- Reinhold B, Hurek T, Fendrik I. Strain-Specific chemotaxis of *Azospirillum* spp. *J Bacteriol* 1985;**162**:190–5.
- Reis VM, dos Santos Teixeira KR, Pedraza RO. What is expected from the genus *Azospirillum* as a plant growth promoting bacteria? In: Maheshwari DK (ed). *Bacteria in Agrobiology: Plant Growth Responses*. Berlin: Springer, 2011, 123–38.
- Requena BN, Jimenez I, Toro M et al. Interaction between plant growth promoting rhizobacteria (PGPR) arbuscular fungi and *Rhizobium* spp. in the rhizosphere of *Anthyllis cytisioides*, a model legume for regeneration in Mediterranean semi-arid ecosystem. *New Phytol* 1997;**136**:667–77.
- Revillas JJ, Rodelas B, Pozo C et al. Production of B-group vitamins by two *Azotobacter* strains with phenolic compounds as sole carbon source under diazotrophic and adiazotrophic conditions. *J Appl Microbiol* 2000;**89**:486–93.

- Rice WA, Olsen PE. Soil inoculants for alfalfa grown on moderately acid soil. *Commun Soil Sci Plan* 1988;19:947–56.
- Richardson AE. Prospects for using soil microorganisms to improve the acquisition of phosphorus by plants. *Aust J Plant Physiol* 2001;28:897–906.
- Rojas-Tapias D, Moreno-Galván A, Pardo-Díaz S et al. Effect of inoculation with plant growth-promoting bacteria (PGPB) on amelioration of saline stress in maize (*Zea mays*). *Appl Soil Ecol* 2012;61:264–72.
- Roos IMM, Hattingh MJ. Scanning electron microscopy of *Pseudomonas syringae* pv: morspennorum on sweet cherry leaves. *J Phytopathol* 1983;108:18–25.
- Rudrappa T, Czymmek KJ, PW Paré et al. Root-secreted malic acid recruits beneficial soil bacteria. *Plant Physiol* 2008;148:1547–56.
- Russo A, Vettori L, Felici S et al. Enhanced micropropagation response and biocontrol effect of *Azospirillum brasilense* SP245 on *Prunus cerasifera* L. clone Mr.S 2/5 plants. *J Biotechnol* 2008;134:312–9.
- Saber Z, Pirdashti H, Esmaili M et al. Response of wheat growth parameters to co-inoculation of plant growth promoting rhizobacteria (PGPR) and different levels of inorganic nitrogen and phosphorus. *World Appl Sci J* 2012;16:213–9.
- Sahin F, Çakmakçı R, Kantar F. Sugar beet and barley yields in relation to inoculation with N₂-fixing and phosphate solubilizing bacteria. *Plant Soil* 2004;265:123–9.
- Salantur A, Ozturk A, Akten S et al. Effect of inoculation with non-indigenous and indigenous rhizobacteria of Erzurum (Turkey) origin on growth and yield of spring barley. *Plant Soil* 2005;275:147–56.
- Saravanakumar D, Lavanya N, Muthumeena B et al. *Pseudomonas fluorescens* enhances resistance and natural enemy population in rice plants against leaf folder pest. *J Appl Entomol* 2008;132:469–79.
- Scher FM, Klopper JW, Singleton C et al. Colonization of soybean roots by *Pseudomonas* and *Serratia* species: relationship to bacterial motility, chemotaxis and generation time. *Phytopathology* 1988;78:1055–9.
- Scot RI, Chard JM, Hocart MJ et al. Penetration of potato tuber lenticels by bacteria in relation to biological control of blackleg disease. *Potato Res* 1996;39:333–44.
- Sharifi RS. Study of grain yield and some of physiological growth indices in maize (*Zea mays* L.) hybrids under seed biopriming with plant growth promoting rhizobacteria (PGPR). *J Food Agric Environ* 2011;189:3–4.
- Sharifi RS. Study of nitrogen rates effects and seed biopriming with PGPR on quantitative and qualitative yield of Safflower (*Carthamus tinctorius* L.). *Tech J Eng Appl Sci* 2012;2:162–6.
- Sharifi RS, Khavazi K. Effects of seed priming with plant growth promoting rhizobacteria (PGRP) on yield and yield attribute of maize (*Zea mays* L.) hybrids. *J Food Agric Environ* 2011;9:496–500.
- Sharifi RS, Khavazi K, Gholipouri A. Effect of seed priming with plant growth promoting Rhizobacteria (PGPR) on dry matter accumulation and yield of maize (*Zea mays* L.) hybrids. *Int Res J Biochem Bioinf* 2011;1:76–83.
- Simons M, Permentier HP, de Weger LA et al. Amino acid synthesis is necessary for tomato root colonization by *Pseudomonas fluorescens* strain WCS365. *Mol Plant-Microbe In* 1997;10:102–6.
- Smith RS. Legume inoculant formulation and application. *Can J Microbiol* 1992;38:485–92.
- Sørensen J, Sessitsch A. Plant-associated bacteria lifestyle and molecular interactions. In: van Elsas JD, Jansson JK, Trevors JT (ed.). *Modern Soil Microbiology*. 2nd edition. CRC Press, 2006, 211–36.
- Soto MJ, Sanjuán J, Olivares J. Rhizobia and plant-pathogenic bacteria: common infection weapons. *Microbiology* 2006;152:3167–74.
- Spaepen S, Vanderleyden J, Remans R. Indole-3-acetic acid in microbial and microorganism-plant signaling. *FEMS Microbiol Rev* 2007;31:425–48.
- Srinivasan K, Gilardi G, Garibaldi A et al. Bacterial antagonists from used rockwool soilless substrates suppress *Fusarium* wilt of tomato. *J Plant Pathol* 2009;91:147–54.
- Sriprang R, Hayashi M, Ono H et al. Enhanced accumulation of Cd²⁺ by a *Mesorhizobium* sp. transformed with a gene from *Arabidopsis thaliana* coding for phytochelatase. *Appl Environ Microb* 2003;69:1791–6.
- Stajner D, Gasaić O, Matković B et al. Metolachlor effect on antioxidant enzyme activities and pigments content in seeds and young leaves of wheat (*Triticum aestivum* L.). *Agric Med* 1995;125:267–73.
- Stajner D, Kevrean S, Gasaić O et al. Nitrogen and *Azotobacter chroococcum* enhance oxidative stress tolerance in sugar beet. *Biol Plantarum* 1997;39:441–5.
- Staley TE, Drahos DJ. Marking soil bacteria with lacZY. In: Weaver RW, Angel JS, Bottomley PJ (eds). *Methods of Soil Analysis. Part 2. Microbiological and Biochemical Properties*. Soil Science Society of America, Madison, WI, 1994, 689–706.
- Stephens PM. Recent methods to improve root colonization by PGPR strains in soil. In: Ryder MH, Stephens PM, Bowen GD (eds). *Improving Plant Productivity with Rhizosphere Bacteria, Commonwealth Scientific and Industrial Research Organization*. Australia: Adelaide, 1994.
- Strijdom BW, Deschodt CC. Carriers of rhizobia and the effects of prior treatment on the survival of rhizobia. In: Nutman PS (ed.). *Symbiotic Nitrogen Fixation in Plants. International Biological Programme No. 7*. Cambridge, UK: Cambridge University Press, 1976, 151–68.
- Sudhakar P, Chattopadhyay GN, Gangwar SK et al. Effect of foliar application of *Azotobacter*, *Azospirillum* and *Beijerinckia* on leaf yield and quality of mulberry (*Morus alba*). *J Agric Sci* 2000;134:227–34.
- Tank N, Saraf M. Salinity-resistant plant growth promoting rhizobacteria ameliorates sodium chloride stress on tomato plants. *J Plant Int* 2010;5:51–58.
- Taylor AG, Harman GE. Concept and technologies of selected seed treatments. *Ann Rev Phytopathol* 1990;28:321–39.
- Taylor AG, Harman GE, Nielsen PA. Biological seed treatments using *Trichoderma harzianum* for horticultural crops. *Hort Technol* 1994;4:105–9.
- Thompson JA. Production and quality control of legume inoculants. In: Begerson FJ (ed). *Methods for Evaluating Biological Nitrogen Fixation*. New York: John Wiley & Sons, 1980, 489–533.
- Tilak KVBR, Ranganayaki N, Manoharachari C. Synergistic effect of plant growth promoting rhizobacteria and *Rhizobium* on nodulation and nitrogen fixation by pigeonpea (*Cajanus cajan*). *Eur J Soil Sci* 2006;57:67–71.
- Trevors JT, van Elsas JD, Lee H et al. Use of alginate and other carriers for encapsulation of microbial cells for use in soil. *Microbial Releases* 1992;1:61–9.
- Turan MMG, Çakmak R, Şahin F. Effect of plant growth promoting rhizobacteria strain on freezing injury and antioxidant enzyme activity of wheat and barley. *J Plant Nutr* 2013;36:731–48.
- Turnbull GA, Morgan JAW, Whipps JM et al. The role of bacterial motility in the survival and spread of *Pseudomonas*

- fluorescens* in soil and in the attachment and colonization of wheat roots. *FEMS Microbiol Ecol* 2001;**36**:21–31.
- van Elsas JD, Heijnen CE. Methods for the introduction of bacteria into soil: a review. *Biol Fert Soils* 1990;**10**:127–33.
- van Elsas JD, Kijstra AR, Govaert JM et al. Survival of *Pseudomonas fluorescens* and *Bacillus subtilis* introduced into two soils of different texture in field microplots. *FEMS Microbiol Ecol* 1986;**38**:151–60.
- van Loon LC, Bakker PAHM. Induced systemic resistance as a mechanism of disease suppression by rhizobacteria. In: Siddiqui ZA (ed). *PGPR: Biocontrol and Biofertilization*. Dordrecht, The Netherlands: Springer, 2005, 39–66.
- Vesper SJ. Production of pili (fimbriae) by *Pseudomonas fluorescens* and a correlation with attachment to corn roots. *Appl Environ Microb* 1987;**53**:1397–405.
- Vessey JK. Plant growth promoting rhizobacteria as biofertilizers. *Plant Soil* 2003;**255**:571–86.
- Vincent JM, Thompson JA, Donovan KO. Death of root nodule bacteria on drying. *Aust J Agr Res* 1962;**13**:258–70.
- Walker R, Rossall S, Asher MJC. Comparison of application methods to prolong the survival of potential biocontrol bacteria on stored sugar-beet seed. *J Appl Microbiol* 2004;**97**:293–305.
- Walker TS, Bais HP, Grotewold E et al. Root exudation and rhizosphere biology. *Plant Physiol* 2003;**132**:44–51.
- Wani PA, Khan MS, Zaidi A. Effect of metal tolerant plant growth promoting *Bradyrhizobium* sp. (vigna) on growth, symbiosis, seed yield and metal uptake by green gram plants. *Chemosphere* 2007;**70**:36–45.
- Warren JE, Bennett MA. Bio-osmopriming tomato (*Lycopersicon esculentum* Mill.) seeds for improved stand establishment. *Seed Sci Technol* 1999;**27**:489–99.
- Watrud LS, Perlak FJ, Tran MT et al. Cloning of the *Bacillus thuringiensis* subsp. *kurstaki* delta-endotoxin gene into *Pseudomonas fluorescens*: molecular biology and ecology of an engineered microbial pesticide. In: Halvorson HO, Pramer D, Rogul M (eds). *Engineered Organisms in the Environment*. Washington, DC: American Society for Microbiology, 1985, 40–46.
- Weller DM. Colonization of wheat roots by fluorescent *Pseudomonas* suppressive to take-all. *Phytopathology* 1983;**73**:1548.
- Whipps J. Microbial interactions and biocontrol in the rhizosphere. *J Exp Bot* 2001;**52**:487–511.
- Wilson M, Lindow SE. Effect of phenotypic plasticity on epiphytic survival and colonization by *Pseudomonas syringae*. *Appl Environ Microb* 1993;**59**:410–6.
- Wintermans PCA, Bakker P, Pieterse CMJ. Natural genetic variation in *Arabidopsis* for responsiveness to plant growth-promoting rhizobacteria. *Plant Mol Biol* 2016;**90**:623–34.
- Younesi O, Moradi A. Effects of plant growth-promoting rhizobacterium (PGPR) and arbuscular mycorrhizal fungus (AMF) on antioxidant enzyme activities in salt-stressed bean (*Phaseolus vulgaris* L.). *Agriculture (Poľnohospodárstvo)* 2014;**60**:10–21.
- Zaidi S, Usmani S, Singh BR et al. Significance of *Bacillus subtilis* strain SJ-101 as a bioinoculant for concurrent plant growth promotion and nickel accumulation in *Brassica juncea*. *Chemosphere* 2006;**64**:991–7.