

Meiotic study of *Zea mays* ssp. *mays* ($2n = 40$) x *Tripsacum dactyloides* ($2n = 72$) hybrid and its progeny

María del Carmen Molina*

Instituto Fitotécnico de Santa Catalina
Facultad de Ciencias Agrarias y Forestales
Universidad Nacional de la Plata
CIGen (UNLP-CONICET-CIC)
Garibaldi 3400, CC4, 1836 Llavallol
Buenos Aires, Argentina
Tel/fax: 54 11 4282 0233
E-mail: mcmgen@yahoo.com

María Dina García

Facultad de Ciencias Agrarias
Universidad Nacional de Lomas de Zamora
Ruta 4 Km. 2, CC 95, 1832 Lomas de Zamora
Buenos Aires, Argentina
Tel/fax: 54 11 4282 6263/7860/7896
E-mail: mdinaga@hotmail.com

Silvia Elena Chorzempa

Facultad de Ciencias Agrarias
Universidad Nacional de Lomas De Zamora
Ruta 4 Km. 2, CC 95, 1832 Lomas de Zamora
Buenos Aires, Argentina
Tel/fax: 54 11 4282 6263/7860/7896
E-mail: chorzempa2000@yahoo.com.ar

Website: <http://www.agro.unlp.edu.ar>

Website: <http://www.unlz.edu.ar>

Financial support: FCAyF-UNLP (grant A129), FCA-UNLZ and CONICET (grant 4650).

Keywords: chromosome number, corn, eastern gamagrass, remote hybridisation.

Abbreviations: MPC: mother pollen cells
pZmT: progeny of ZmT plants
ZmT: *Z. mays* ssp. *mays* x *Tripsacum dactyloides* F1 hybrid

Maize ($2n = 40$) x *Tripsacum dactyloides* ($2n = 72$) F1 hybrid plants ($2n = 56$) were obtained by embryo rescue and induction of somatic embryogenesis/organogenesis. Hybrid plants showed *Tripsacum*-like phenotypes, tolerance to stresses such as NaCl salinity and low temperatures. The more frequent meiotic configurations were 28 II (24%), 24 II + 2 IV (19%) and 26 II + 1 IV (12%), with an average per cell of 0,55 I + 25,18 II + 1,19 IV. Significant differences between plants were not observed. Pollen fertility ranged from 0% to 50%. After pollination with maize or *Tripsacum*, 20% of F1 plants have developed viable seeds, which originated the progeny. Thirty five percent of the progeny showed $2n = 56$ chromosomes and F1 like-phenotypes, which suggests they have apomictic origin. The remaining

plants were fertile and they showed maize-like phenotypes and different chromosome numbers ($2n = 22, 24, 26, 28$ and 30), because they kept the complete maize chromosome complement and some of the *Tripsacum* chromosomes. Meiotic cells showed pairing between chromosomes from both parental species, which suggests the possibility of genetic recombination between them.

Interspecific hybridisation of remotely related plant species offers a great potential to improve efficiency in crop breeding and production. Introduction of novel genetic variability and apomictic seed development into crops have been recognizable as two important achievements of the

* Corresponding author




Figure 1. Meiotic configurations of maize x *Tripsacum* F1 hybrid plants.

(a) Diakinesis with $2n = 56$.

(b) Chromosomes of maize (external chromosomes in diakinesis) and *Tripsacum* (internal chromosomes in pachytene) disposed in two groups.

(c) Anaphase I showing 7 and 8 univalent chromosomes in each pole.

(d) Anaphase I showing 2 univalent chromosomes (arrowed) and bridges.

manipulation of reproductive systems in plants (Matzk et al. 1997).

Corn (*Zea mays ssp. mays*) is an annual crop that shows susceptibility to adverse environmental conditions such as low temperatures (Hodges et al. 1995) and NaCl salinity (Pasternak et al. 1995). *Tripsacum dactyloides* is a perennial forage which presents high palatability and productivity (Faix et al. 1980), tolerance to different environmental stresses (Foy, 1997; Clark et al. 1998; Ray et al. 1998) and apomictic reproduction (Burson et al. 1990; Leblanc et al. 1995a; Kindiger et al. 1996a; Grimanelli et al. 1998). Such characteristics make this species an interesting source of genetic variability for corn improvement.

Different authors have investigated the possibility to introgress some characteristics from *Tripsacum* to maize, for example, disease tolerance (Bergquist, 1981), constitutive aerenchyma (Ray et al. 1999), and gametophytic apomixis (Leblanc et al. 1995b; Kindiger et

al. 1996b; Lukina and Chistyakova, 1999).

Hybrids between maize and *Tripsacum* can be obtained with more or less success, according to the maize genotype used as female parent, through embryo rescue and induction of organogenesis or somatic embryogenesis (Furini and Jewell, 1995; García and Molina, 1997). Maize x *T. dactyloides* F1 hybrid plants are sexually sterile, but low percentages of viable seeds have been obtained mainly by apomixis (Leblanc et al. 1995a; Sokolov et al. 2000). Regenerated hybrid plants showed salinity tolerance (Pesqueira et al. 2003). Further, F1 hybrid plants grown under field conditions or culture chambers showed high tolerance to low temperatures (-2°C and 5°C , respectively), although maize and *Tripsacum* showed great susceptibility under the same conditions (Jatimlansky et al. 2004).

The objectives of this work were to evaluate pollen fertility and production of viable seeds of *Z. mays ssp. mays* ($2n = 40$) x *T. dactyloides* ($2n = 72$) regenerated hybrid plants, and to analyse meiosis of fertile F1 hybrid plants and the

progeny of these plants.

MATERIALS AND METHODS

Inbred plants of *Z. mays* ssp. *mays* N107B ($2n = 40$), supplied by Maize Genetics Coop. Stock Center, Urbana, Illinois) were pollinated with *Tripsacum dactyloides* ($2n = 72$). Immature embryos originated embryogenic and organogenic callus cultures on the basic medium (García et al. 1992) supplemented with $4 \mu\text{molL}^{-1}$ 2,4 dichlorophenoxyacetic acid. Regenerated shoots were rooted on the basic medium free of plant growth regulators. The 130 F1 plants analysed in this work arose from one embryo. The progeny was obtained from free pollination of F1 plants with maize or *Tripsacum*.

Immature tassels were fixed in Farmer solution (3:1 absolute ethanol-acetic acid) and coloured with a 2% solution of acetic haematoxylin. Pairing configurations were observed at diakinesis-metaphase I. Pollen fertility was determined using Lugol solution. Pollen was counted as fertile if at least 75% of the grain was coloured.

RESULTS AND DISCUSSION

Z. mays ssp. *mays* ($2n = 40$) x *T. dactyloides* ($2n = 72$) F1 hybrid plants showed a chromosome number $2n = 56$, *Tripsacum*-like phenotypes, an average height of 2.10 m, 3 to 20 tillers per plant and 4 or 5 ears per stem. Their flowering period lasted from November to June or July. Half of the plants showed an annual cycle and the rest were biannual or perennial ones. Further, tolerance to low temperatures and NaCl salinity was observed in these plants (Pesqueira et al. 2003; Jatimliansky et al. 2004). Pollen fertility ranged from 0% (most of the plants) to 50%. Eighty percent of the plants did not produce viable seeds;

the other 20% developed 2 to 10 viable seeds per ear. Normal microspore development but complete male sterility was observed in 56-chromosome maize x *Tripsacum* backcross hybrids (Kindiger, 1993).

Over 40 plants were evaluated from the progeny, 35% showed $2n = 56$ chromosomes (Figure 1a) and F1-like phenotypes. Pollen fertility ranged from 0% to 50%. The other 65% of the plants resembled maize more than *Tripsacum* phenotype, except that they showed up to 3 tillers per plant and continuous flowering on very prolific stems during the growing season. Pollen fertility ranged from 40 to 96%.

Tetraploid ($2n = 4x = 72$) and triploid ($2n = 3x = 54$) eastern gamagrass reproduced by diplosporous pseudogamy, *Antennaria*-type megagametogenesis and diploid parthenogenesis (Burson et al. 1990). Chromosome number of $2n = 56$ observed in 35% of the maize/*Tripsacum* F1 progeny suggests that these plants were originated by apomixis as described in polyploid *Tripsacum* species (Burson et al. 1990). However, reduction of *Tripsacum* chromosome complement was observed in 65% of maize/*Tripsacum* F1 progeny plants. A previous report (Kindiger et al. 1996b) also showed partial loss of *Tripsacum* chromosomes in backcrosses of an apomictic maize/*Tripsacum* F1 hybrid ($2n = 56$).

The most frequent meiotic configurations in diakinesis - metaphase I (% of analysed cells), average number of uni-, bi-, tri-, and tetraivalents and chiasmata cell⁻¹ (Table 1) were the following:

Zea mays ssp. *mays* ($2n = 40$) showed 10IV (30%) and 9IV + 2II (24%), an average number of 8.15IV + 3.27II and

Table 1. Cytogenetic study of *Zea mays*, *Tripsacum dactyloides*, *Z. mays* x *T. dactyloides* F1 hybrid plants (ZmT) and their progeny (pZmT): chromosome number, averages of uni-, bi-, tri and tetraivalents per cell (PMC) and chiasmata cell⁻¹.

Genotype	2n	Meiotic configurations				Chiasmata cell ⁻¹	N° of PMC*
		I	II	III	IV		
<i>Zea mays</i>	40	0.04	3.15	0.04	8.27	33.75	143
<i>T. dactyloides</i>	72	0.01	20.65	0.01	6.80	58.50	153
ZmT	56	0.27	25.28	0	1.11	40.63	129
ZmT	56	0.62	25.16	0	1.30	39.07	111
ZmT	56	0.63	23.70	0	1.94	44.50	108
ZmT	56	0.70	26.58	0	0.41	42.80	105
pZmT	22	0.04	10.70	0	0.14	30.52	96
pZmT	22	0.17	10.63	0	0.03	20.13	105
pZmT	22	0.07	10.60	0	0.18	20.09	118
pZmT	24	0.06	11.35	0	0.31	22.70	146
pZmT	26	0.04	10.20	0	1.20	24.00	84
pZmT	26	1.00	10.00	0	0.92	21.00	89
pZmT	26	0.08	12.35	0	0.31	24.26	105
pZmT	26	0.06	12.06	0	0.45	22.15	121
pZmT	26	0.16	12.43	0	0.24	22.78	92
pZmT	28	0.05	10.89	0	1.54	26.19	104
pZmT	28	0.03	11.20	0	1.39	25.32	78
pZmT	30	0.04	13.76	0	0.58	24.22	100

33.75 chiasmata cell⁻¹. Cells showed normal anaphase I, i.e., a regular migration of 20 chromosomes to each pole.

T. dactyloides (2n = 72) showed 5IV + 26II (21%) and 10IV + 16II (19%), an average number of 6.80IV + 20.65II and 58 chiasmata cell⁻¹.

F1 hybrid (2n = 56) showed 28II (24%), 24II + 2IV (19%) and 26II + 1IV (12%), an average number of 0.55I + 25.18II + 1.19IV and 42 chiasmata cell⁻¹. No significant differences were observed among analysed plants. Cells in anaphase I showed multipolar spindle, lagging chromosomes (Figure 1c) and one or two bridges (Figure 1d). Other meiotic abnormalities were also observed in F1 plants, such as 2 or more nuclei in different stages in the same cytoplasm, spatial separation and asynchronous stages of *Tripsacum* and maize chromosomes (Figure 1b). Some of the hybrid plants showed one or two tetravalents, one of them constituted by two chromosomes from maize and two from *Tripsacum* and the other (if there were two) by homoeologous *Tripsacum* chromosomes.

F1 hybrid progeny plants with 2n = 56 and F1-like phenotypes showed cytogenetic characteristics similar to that described in F1 plants. Instead, the F1 progeny maize-like plants showed different chromosome numbers (2n = 22, 24, 26, 28 and 30) and produced viable seeds. The different chromosome numbers were the result of 26 to 34 *Tripsacum* chromosome elimination in metaphase I. Meiosis was normal in cells with 22 or 24 chromosomes. On the contrary, cells with 26 or more chromosomes showed lagging chromosomes, spatial separation of chromosomes in two groups by species and subsequent loss of *Tripsacum* chromosomes. The number of *Tripsacum* chromosomes represented as univalents was small in all MI cells of reduced maize-*Tripsacum* hybrids with 29 chromosomes (4.63 ± 0.17 to 6.72 ± 0.3 univalents per cell) (Lukina and Chistyakova, 1999) but higher than the number of univalents per cell observed in the present work for maize x *Tripsacum* F1 hybrid (2n = 56) plants and their progeny (0.01 to 1.0 univalents per cell, Table 1).

In conclusion, the study of maize x *T. dactyloides* F1 hybrid plants and their progeny showed that: i) Hybrids between different genera may express phenotypic characteristics absent in their progenitor species; ii) Fertility can be restored after most of the *Tripsacum* chromosomes elimination, iii) Pairing between parental species suggest the possibility of genetic recombination between them.

REFERENCES

BERGQUIST, R.R. Transfer from *Tripsacum dactyloides* to corn of a major gene locus conditioning resistance to *Puccinia sorghi*. *Phytopathology*, 1981, vol. 71, no. 5, p. 518-520.

BURSON, B.; VOIGTH, P.W.; SHERMAN, R.A. and DEWALD, C.L. Apomixis and sexuality in Eastern

Gamagrass. *Crop Science*, 1990, vol. 30, no. 1, p. 86-89.

CLARK, R.B.; ALBERTS, E.E.; ZOBEL, R.W.; SINCLAIR, T.R.; MILLER, M.S.; KEMPER, W.D. and FOY, C.D. Eastern gamagrass (*Tripsacum dactyloides*) root penetration into and chemical properties of claypan soils. *Plant and Soil*, 1998, vol. 200, no. 1, p. 33-45.

FAIX, J.J.; KAISEER, C.J. and HINDS, F.C. Quality, yield, and survival of Asiatic bluestems and an eastern gamagrass in Southern Illinois. *Journal of Range Management*, 1980, vol. 33, no. 5, p. 338-390.

FOY, C.D. Tolerance of eastern gamagrass to excess Aluminium in acid soil and nutrient solution. *Journal of Plant Nutrition*, 1997, vol. 20, no. 9, p. 1119-1136.

FURINI, A. and JEWELL, C. Somatic embryogenesis and plant regeneration of maize/*Tripsacum* hybrids. *Maydica*, 1995, vol. 40, no. 2, p. 205-210.

GARCÍA, María Dina; MOLINA, María del Carmen and CASO, Osvaldo Héctor. El cultivo de callos organogénicos como fuente de variabilidad genética para el mejoramiento del maíz. In: *Proceedings of the V Congreso Nacional del Maíz- II Reunión Suramericana de Maiceros* (11th - 13th November, 1992, Pergamino, Buenos Aires, Argentina) vol. 1, p. 61-69.

GARCÍA, M.D. and MOLINA, M.C. Embriogénesis somática y regeneración de plantas a partir de embriones híbridos de *Zea mays* ssp. *mays* (2n = 20 y 40) and *Tripsacum dactyloides*. In: *Proceedings of the VI Congreso Nacional de Maíz* (12th - 14th November 1997, Pergamino, Buenos Aires, Argentina) vol. 1, p. 79-86.

GRIMANELLI, D.; LEBLANC, O.; ESPINOSA, E.; PEROTTI, E.; GONZÁLEZ DE LEÓN, D. and SAVIDAN, I. Mapping diplosporous apomixis in tetraploid *Tripsacum*: one gene or several genes? *Heredity*, 1998, vol. 80, no. 1, p. 33-39.

HODGES, D.M.; HAMILTON, R.I. and CHAREST, C. A chilling response test for early growth phase maize. *Agronomy Journal*, 1995, vol. 87, no. 5, p. 970-974.

JATIMLIANSKY, J.R.; GARCÍA, M.D. and MOLINA, M.C. Response to chilling of *Zea mays*, *Tripsacum dactyloides* and their hybrid. *Biologia Plantarum*, 2004, vol. 48, no. 4, p. 561-567.

KINDIGER, B. Aberrant microspore development in hybrids of maize x *Tripsacum dactyloides*. *Genome*, 1993, vol. 36, no. 5, p. 987-997.

KINDIGER, B.; BAI, D. and SOKOLOV, V. Assignment of a gene(s) conferring apomixis in *Tripsacum* to a chromosome arm: cytological and molecular evidence. *Genome*, 1996a, vol. 39, no. 6, p. 1133-1141.

KINDIGER, B.; SOKOLOV, V. and KHATYPOVA, J.V. Evaluation of apomictic reproduction in a set of 39 chromosome maize-*Tripsacum* backcross hybrids. *Crop Science*, 1996b, vol. 36, no. 5, p. 1108-1113.

LEBLANC, O.; PEEL, M.D.; CARMAN, J.G. and SAVIDAN, Y. Megasporogenesis and megagametogenesis in several *Tripsacum* species (Poaceae). *American Journal of Botany*, 1995a, vol. 82, no. 1, p. 57-63.

LEBLANC, O.; GRIMANELLI, D.; GONZÁLEZ-DE-LEÓN, D. and SAVIDAN, Y. Detection of the apomictic mode of reproduction in maize-*Tripsacum* hybrids using maize RFLP markers. *Theoretical and Applied Genetics*, 1995b, vol. 90, no. 7-8, p. 1198-1203.

LUKINA, L.A. and CHISTYAKOVA, A.K. Abnormal behavior of chromosomes in meiosis of reduced maize-*Tripsacum* hybrids. *Apomixis Newsletter* [online]. 1999, vol. 11 [cited June 1999]. Available from Internet: <http://www.cimmyt.org/abc/researchprojects/apomixis/apomixisnews11/htm/APOMIXISNews11-4.htm>.

MATZK, F.; OERTEL, C.; ALTENHOFER, P. and SCHUBERT, I. Manipulation of reproductive systems in Poaceae to increase the efficiency in crop breeding and production. *Trends in Agronomy*, 1997, vol. 1, no. 1, p. 19-34.

PASTERNAK, D.; SAGIH, N.; DEMALACH, Y.; KEREN, Y. and SHAFFER, A. Irrigation with brackish water under desert conditions XI. Salt tolerance in sweet-corn cultivars. *Agricultural water management*, 1995, vol. 28, no. 4, p. 325-334.

PESQUEIRA, J.; GARCÍA, M.D. and MOLINA, M.C. NaCl tolerance in maize (*Zea mays* ssp. *mays* L.) x *Tripsacum dactyloides* L. hybrid calli and regenerated plants. *Spanish Journal of Agricultural Research*, 2003, vol. 1, no. 2, p. 59-63.

RAY, J.D.; KINDIGER, B.; DEWALD, C.L. and SINCLAIR, T.R. Preliminary survey of root aerenchyma in *Tripsacum*. *Maydica*, 1998, vol. 43, no. 1, p. 49-53.

RAY, J.D.; KINDIGER, B. and SINCLAIR, T.R. Introgressing root aerenchyma into maize. *Maydica*, 1999, vol. 44, no. 2, p. 113-117.

SOKOLOV, V.A.; DEWALD, C.L. and KHATYPOVA, I.V. The genetic programs of nonreduction and parthenogenesis in corn-gamagrass hybrids are inherited and expressed in an independent manner. *Maize Newsletter*, 2000, vol. 74, no. 1, p. 55-57.