



VOJNOSANITETSKI PREGLED

VOJNOMEDICINSKA AKADEMIJA

Crnotravska 17, 11 000 **Beograd, Srbija**

Tel/faks: +381 11 2669689

vsp@vma.mod.gov.rs

ACCEPTED MANUSCRIPT

Accepted manuscripts are the articles in press that have been peer reviewed and accepted for publication by the Editorial Board of the *Vojnosanitetski Pregled*. They have not yet been copy edited and/or formatted in the publication house style, and the text could still be changed before final publication.

Although accepted manuscripts do not yet have all bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: article title, the author(s), publication (year), the DOI.

Please cite this article: **ASTHMA AND PERIODONTAL HEALTH IN CHILDREN
ASTMA I PARODONTALNO ZDRAVLJE KOD DJECE**

Authors: **Bojana Davidović***, **Mirjana Ivanović[†]**, **Dejan Bokonjić[‡]**, **Svjetlana Janković***, **Jelena Erić***, **Jelena Lečić***; *Vojnosanitetski pregled* (2017); Online First April, 2017.

UDC:

DOI: <https://doi.org/10.2298/VSP170221063D>

When the final article is assigned to volumes/issues of the Journal, the Article in Press version will be removed and the final version appear in the associated published volumes/issues of the Journal. The date the article was made available online first will be carried over.

ASTHMA AND PERIODONTAL HEALTH IN CHILDREN

Astma i parodontalno zdravlje kod djece

Bojana Davidović*, Mirjana Ivanović†, Dejan Bokonjić‡, Svjetlana Janković*, Jelena Erić*, Jelena Lečić*.

* University of East Sarajevo, Faculty of Medicine, Dentistry Program, Foca, Bosnia and Herzegovina;

† University of Belgrade, Faculty of Dental Medicine, Department of Pediatric and Preventive Dentistry, Belgrade, Serbia;

‡ University Hospital in Foca, Pediatric Clinic, Foca, Bosnia and Herzegovina;

Correspondence to:

Bojana DAVIDOVIC, Faculty of Medicine, University of East Sarajevo, Studentska 5, 73300 Foca, Bosnia and Herzegovina. Tel/Fax: +387 58 211 906, E-mail: boki1310@yahoo.com

Asthma and periodontal health in children

Abstract

Introduction/Objective: Oral health is an important part of overall health. Good oral health is important for oral diseases prevention and health maintenance of respiratory system. The aim of the study was to evaluate oral hygiene and periodontal health parameters of asthmatic children and to compare them with children without asthma, as well as to evaluate those parameters according to type of used medications and time of taking medications in children with asthma.

Material and methods: This epidemiological study included 68 children with asthma and 68 children without asthma or any other chronic disease aged from 6 to 16 years. Parameters used in this study were Greene-Vermillion index, Löe-Silness gingival index and Community Periodontal Index (CPI).

Results: Good oral hygiene (31.1%) was more present in children without asthma whereas poor hygiene (20.0%) was more frequent in children with asthma ($p < 0.001$). Healthy gingiva was more frequent in children without asthma (25%) while mild (58.8%) and moderate gingival inflammation (5.9%) were more frequent in the group of children with asthma ($p < 0.01$). Mean CPI values were higher in children with asthma ($p < 0.001$). Mean values of Plaque Index, Gingival Index and CPI did not show statistically significant difference in relation to type of administered medication. However, taking of medications in the afternoon was related to higher mean values of Plaque Index and Gingival Index ($p < 0.05$) within the group of children with asthma.

Conclusion: Children with asthma had poorer oral hygiene and were diagnosed with greater values of oral hygiene and periodontal indices compared with the group of children without asthma. For this reason, it is necessary to promote oral health and establish good oral hygiene habits in asthmatic children.

Keywords: children; asthma; oral hygiene; gingivitis; periodontal parameters

Kratak sadržaj:

Uvod/Cilj: Oralno zdravlje je važan dio opšteg zdravlja. Dobro oralno zdravlje je važno kako za prevenciju oralnih oboljenja tako i za održavanje zdravlja respiratornog sistema. Cilj istraživanja je bio da se procjeni oralna higijena, stanje zdravlja parodontalnih tkiva djece oboljele od astme, te da se te vrijednosti uporede sa pronađenim vrijednostima kod djece bez astme, kao i da se dobijene vrijednosti procjene i u odnosu na vrstu i vrijeme upotrebljenog lijeka u grupi djece sa astmom.

Materijal i metode: U ovu epidemiološku studiju uključeno je 68 djece sa astmom i 68 djece bez astme ili bilo koje druge hronične bolesti, uzrasta od 6 do 16 godina. Parametri korišteni u ovoj studiji su plak indeks po Greene-Vermillionu, Löe-Silnessov gingivalni indeks i Indeks stanja parodonticijuma u zajednici (Community Periodontal Index (CPI)).

Rezultati: Dobra oralna higijena je prisutnija kod djece bez astme (31,1%), dok je loša češća kod djece sa astmom (20,0%) ($p < 0,001$). Zdrava gingiva je češća kod djece bez astme (25%), dok je blaga (58,8%) i umjerena (5,9%) upala desni prisutnija u grupi djece sa astmom ($p < 0,01$). Prosječne vrijednosti CPI indeksa bile su veće kod djece sa astmom ($p < 0,001$). Prosječne vrednosti plak indeksa, gingivalnog indeksa i CPI nisu ukazali na značajnu razliku u odnosu na vrstu primenjenog leka. Međutim, uzimanje lijekova u popodnevnim satima je povezan sa višim vrednostima plak i gingivalnog indeksa ($p < 0,05$) u grupi dece sa astmom.

Zaključak: Djeca sa astmom su imala lošiju oralnu higijenu, i kod njih su utvrđene veće vrijednost plak i parodontalnih indeksa u poređenju sa grupom djece bez astme. Iz tog razloga neophodno je promovisanje oralnog zdravlja kao i uspostavljanje dobrih oralno higijenskih navika kod djece sa astmom.

Ključne riječi: djeca, astma, oralna higijena, gingivitis, parodontalni parametri

Introduction

Asthma is the most frequent chronic disease and represents not only a health but a social problem, as well. It appears relatively early, in childhood, and its symptoms aren't often recognized and/or aren't treated in a proper way what might present aggravating circumstance for these patients.

Asthma is the leading cause of hospitalizations in children. Children who are hospitalized for an illness may have lethargy or malaise, so that their oral hygiene may be neglected. Having recognized this problem, Blevins proposed oral health protection program for these children based on the fact that pediatric nurses can as well encourage children with asthma to regular oral hygiene and promote the importance of good oral health [1].

Parents are poorly aware of potential oral diseases in asthmatic children and preventive actions what results in increased incidence of the oral diseases. It is proved that if parents are less educated, their children are more exposed to oral diseases [2].

The dental plaque control is the basis for good oral hygiene. Good oral health is important not only to prevent oral disease but also to maintain healthy respiratory system. The evidence also suggests that complete removal of the dental plaque may lead to the improvement of the respiratory function in children with asthma [3].

Children with asthma usually prefer to consume sweet soft drinks and food, while neglecting adequate oral hygiene, the use of fluorides and regular dental checks. Also, some of the drugs used for treatment of asthma usually contain artificial sweeteners [1] that may affect teeth, other oral tissues and promote presence of oral diseases. Poor oral hygiene followed by soft and hard tooth debris accumulation leads to gingivitis that not necessarily has to turn into periodontal diseases. However, it has been reported that it may progress to severe periodontal disease and suggested that high-risk group identification should be given greater attention [4].

Scannapieco and Ho [5] emphasized the connection between chronic respiratory disease and participants with poor oral hygiene. The oral cavity bacteria might have an important part in exacerbations of chronic obstructive pulmonary diseases, as the dental plaque might serve as storage for respiratory pathogenic micro-organisms [5].

Recent studies very seriously considered relationship between oral pathogens, which cause gingivitis and periodontal diseases and asthma [6, 7]. At the other hand, chronic

disease (like asthma) may be considered as a risk factor for the periodontal diseases [8] as well.

The aim of the study was to evaluate oral hygiene and periodontal health parameters of asthmatic children and to compare it with children without asthma, as well as to evaluate oral hygiene and periodontal health parameters according to type of used medications and time of taking medications in children with asthma.

Material and methods

Study population

This one-year long study was conducted according to Declaration of Helsinki of 1975, as revised in 2000. The study was approved by the Ethical Committee of our Institution (No. 01-8/37). Before the examination, the parents and children involved in the study were informed about the aim of the study and applied methodology. The parents were asked to sign written approval for the participation of their children in this study, and children were permitted to leave process in any time during study.

The study included two groups of children, aged from 6 to 16 years. The first group consisted of children with asthma (AG), without any other systemic disease, who had symptoms 2 of asthma and were treated or examined in University Hospital Foca or Primary care facility in Foca. On the basis of present symptoms of asthma AG group was split into two subgroups [9]. The division was conducted by specialist in pediatric pulmonology. First subgroup consisted of children with good controlled asthma (GCA), while second subgroup consisted of children with partly controlled asthma (PCA). Asthma is well controlled if: children have symptoms no more than 2 days a week, these symptoms don't wake them from sleep more than 1 or 2 nights a month; they can do all normal activities; they take quick-relief medicines no more than 2 days a week; they do not have more than one asthma attack a year that requires them to take systematic steroids and their peak flow doesn't drop below 80 percent of their personal best value. Asthma is partly controlled if: children have symptoms more than 2 days a week; they can't perform normal physical activities and they have nocturnal symptoms; they have more asthma attack a year that requires them to take asthma drugs and their peak flow drop below 80 percent of their personal best values.

The second group consisted of children without asthma (NAG) or any other chronic disease and was matched by gender and age to the children with asthma.

Examination

Prior to the examination general data along with medical history data were recorded. The planned epidemiologic examination was performed by recommendation by the World Health Organization – for epidemiologic explorations of National Oral Health Survey [10]. All participants were examined in Department of Pediatric and Preventive Dentistry, Faculty of Medicine Foca. The dental examination was performed by the use of dental or periodontal probe, mirror and artificial light. Data were recorded according to plan prepared for this study.

Clinical measurements

Oral hygiene was evaluated by Greene-Vermillion index for soft deposits or “plaque index” (PI) [11]. The presence of the plaque was noticed on the certain surfaces of completely erupted representative teeth.

The Löe-Silness gingival index (GI) was used for the evaluation of the gingival status [12]. The clinical examination of the gingiva according to mentioned index included the evaluation of gingival status by inspection and probing at vestibular, mesial, oral and distal side of each present tooth.

Community Periodontal Index (CPI) [10] was used to evaluate the periodontal state of health, following WHO recommendation for participants under 15 years.

Statistical analysis

The study analysis was carried out with Statistical Package for Social Sciences (SPSS version 19.0 for Windows, SPSS Inc., Chicago, IL, USA) using chi-square test, Independent-Samples T Test, Mann-Whitney Test, One-Way Anova, in accordance with particular parameters. The values $p < 0.05$ were considered statistically significant.

Results

The research included 136 children, aged between 6 and 16 (10.5 ± 3.3), (AG group ($n=68$; age 10.5 ± 3.3) and NAG ($n=68$; age 10.5 ± 3.3)). In this study, there was statistically significant ($p < 0.05$) higher percentage of male patients (77.9%) than female (22.1%).

AG group consisted of GCA subgroup which was 64.7% of the AG group (mean age 10.3 ± 3.5) and PCA subgroup 35.3% of the AG group (mean age 10.9 ± 2.8). There was no statistical significance related to age between two subgroups ($p > 0.05$) within the AG group.

The results revealed that 63.0% participants were using inhalator steroids, 13.2% bronchodilators and 23.5% combination of inhaler steroids and bronchodilators.

The average value of the PI was 1.2 in AG group (Table 1). Excellent hygiene (19.7%) was more present in NAG group whereas poor hygiene (47.5%) was more frequent

in AG group (Table 2). The obtained values showed no statistical difference in relation to the applied medicine (Table 3). However, PI values showed statistically significant difference related to the time of the day of asthma drugs administration (Table 4).

Healthy gingiva was more present in NAG group (25%), while the presence of mild (58.8%) and moderate (5.9%) inflammation was more frequent in the AG group (Table 5). Statistical significance in GI values was found between AG and (0.5) and NAG groups (0.3) (Table 1). GI values were higher in AG group who were using anti-asthmatic therapy in the afternoon hours (Table 4). PCA children showed higher values of GI when treated with bronchodilators (Table 3).

As far as periodontal health of the experimental group concerns, the average CPI value was 0.8 (Table 1). The healthy periodontal tissue was found in 78.7 % of NAG group (Table 6). The presence of hard debris on teeth, prominent filling edges or the presence of periodontal pockets with the depth of 4-5 mm, were more frequent in AG group. The values of the average CPI values were higher in PCA children, who were using the therapy in the afternoon (Table 4).

Discussion

This study included elementary school age children. This various group might indicate whether asthma or the drugs used in asthma treatment can affect health of periodontal tissues. In addition, evaluation of PI and CPI could have been recorded against completely erupted permanent teeth and GI against all present teeth, what could present limiting factor of this study.

Some of authors support finding that children with asthma have higher values of the plaque index and poorer oral hygiene [13-17] what is in accordance with the results of this study that speak in favor of the close connection between poorer hygiene and the presence of asthma. In present study, three quarters of the AG group participants had poor and very poor oral hygiene. That was even more frequent in children who were using anti-asthmatic drugs in the afternoon as well as in PCA group. Anandhan et al. reported that children with advanced forms of asthma had poorer oral hygiene and that girls had better oral hygiene than boys [18].

However, Ehsani et al. didn't notice any difference in the amount of the plaque between children with asthma and children without asthma [19].

Apart from the statements mentioned above, there is a group of authors who point that a lower PI was registered among the asthmatic children [20, 21], thanks to good oral

hygiene, frequent parental surveillance and regular diet [21]. Having in mind that the diet has an important role in the dental plaque formation, Mazzoleni et al. reported that children with asthma showed to have similar dietary habits but better oral hygiene related habits in comparison to children without asthma of the same age [20].

Saliva has an important role in the limitation of periodontal diseases. However, drugs prescribed for asthma treatment that can cause reduced salivary secretion, greatly affect the severity of periodontal diseases of these group of patients [22]. Higher average values of the GI in participants with asthma compared with the healthy group were found in Attavar, India [13] and Jonkoping, Sweden [23, 24] what is in accordance with the results of our study.

The protective mechanisms within saliva balance the interactions between bacterial and immunological factors and help maintain periodontal health. Mouth-breathing, frequently observed in asthmatic patients due to the obstruction of respiratory system [25], along with a decrease in salivary flow caused by long term use of asthma medication, reduce saliva's protective qualities and therefore place the patient at greater risk of developing periodontal disease [26].

Asthma is a disease, which is very prevalent among young people, whereas periodontal diseases are more frequent in older people. Some authors consider that as a reason why periodontal diseases are rarer in young people with asthma, while other studies showed that periodontal diseases were more frequent in people with asthma than in healthy people [8].

This study revealed poorer state of health of the periodontal tissues in children with asthma and 18.3% of them were diagnosed with the presence of incorrect dental works, hard tooth debris, while the presence of shallow periodontal pockets were recorded in 1.7% of the participants. Considering that the most of children with asthma suffered from gingival bleeding after probing, which is a clear sign of gingivitis, a necessary treatment which consists of instructions on proper and regular oral hygiene and motivation / re-motivation of the patients is required. Higher values of observed indices were recorded with children who used drugs in the afternoon hours what might be explained by inadequate oral hygiene, larger sweet drinks consumption and reduced salivation as well.

Positive correlation between asthma and the presence of the periodontal diseases in children and adolescents (age of 13–17) were not found in some studies [27, 28]. Shulman et al. [27] think that these results may be false due to hormonal status of the observed

groups, the fact that the depth of pockets isn't measured on children younger than the age of 13 as well as the fact that the majority of the studied group was consisted of adolescents with mild and moderate types of asthma.

Belgian scientists confirmed that there was no difference in the gingival health status in children with or without asthma, regardless duration of asthma, or type of used drugs [28]. The absence of difference in gingival or periodontal status of the participants with or without asthma was confirmed in studies in Iran [19] and Italy [29].

Conclusion

In conclusion, results of this study demonstrated that children with asthma had poorer oral hygiene and poorer state of health of the periodontal tissues compared with children without asthma. And that PI and GI values obtained within the group of children with asthma were higher in patients who were taking their asthma drugs in the afternoon hours. However, results of previously mentioned studies are somewhat conflicting, dental clinical protocols for prevention of periodontal tissue deterioration in children with asthma do not exist and the literature does not provide sufficient data on preventive programs achievements. Due to complexity and importance of this matter, further studies that might provide sufficient data, better understanding and would help in creating most efficient protocol for prevention and dental treatment of asthmatic child patients in future are needed.

REFERENCES

1. *Blevins JY*. Oral Health Care For Hospitalized Children. *Pediatr Nurs* 2011;37:229-235.
2. *Miller E, Lee JY, DeWalt DA, Vann WF Jr*. Impact of caregiver literacy on children's oral health outcomes. *Pediatrics* 2010;126:107-114.
3. *Pambudi W, Fabiola I, Indrawati R, Utomo H, Endaryanto A, Harsono A*. Changes in bacterial profiles after periodontal treatment associated with respiratory quality of asthmatic children. *Paediatr Indones* 2008;48:327-337.
4. *FDI World Dental Federation*. The Challenge of Oral Disease – A call for global action. The Oral Health Atlas. 2nd ed. Geneva: FDI World Dental Federation; 2015. Available from: [http://www.fdiworldental.org/publications/oral-health-atlas/oral-health-atlas-\(2015\)](http://www.fdiworldental.org/publications/oral-health-atlas/oral-health-atlas-(2015)).

5. *Scannapieco FA, Ho AW*. Potential associations between chronic respiratory disease and periodontal disease: analysis of National Health and Nutrition Examination Survey III. *J Periodontol* 2001;72:50-56.
6. *Arbes SJJr, Matsui EC*. Can oral pathogens influence allergic disease? *J Allergy Clin Immunol* 2011;127:1119-1127.
7. *Singh Uppal R, Brar R, Goel A*. Association between asthma and chronic periodontitis: a clinical study. *Pakistan oral & dental journal* 2015; 35(3):448-451.
8. *Georgiou TO, Marshall RI, Bartold PM*. Prevalence of systemic diseases in Brisbane general and periodontal practice patients. *Aust Dent J* 2004;49:177-184.
9. *Global initiative for asthma*. Pocket guide for asthma management and prevention (for adults and children older than 5 years). A pocket guide for physicians and nurses: Global initiative for asthma; 2012. Available from:http://www.farm.ucl.ac.be/Benin/2014/pharmacologie-speciale/6-systeme-respiratoire/GINA_Pocket2013_May15.pdf
10. *World Health Organization*. Oral health surveys: Basic methods. 4th ed. Geneva: World Health Organization; 1997.
11. *Greene JC, Vermillion JR*. The Simplified Oral Hygiene Index. *J Am Dent Assoc* 1964;68:7-13.
12. *Löe H*. The gingival index, the plaque index and the retention index systems. *J Periodontol* 1967;38:Suppl:610-616.
13. *Mehta A, Sequeira PS, Sahoo RC, Kaur G*. Is bronchial asthma a risk factor for gingival diseases? A control study. *N Y State Dent J* 2009;75:44-46.
14. *Botelho MPJ, Maciel SM, Cerci Neto A, Dezan CC, Fernandes KBP, de Andrade FB*. Cariogenic microorganisms and oral conditions in asthmatic children. *Caries Res* 2011;45:386-392.
15. *Santos NC, Jamelli S, Costa L, Baracho Filho C, Medeiros D, Rizzo J.A, Sarinho E*. Assessing caries, dental plaque and salivary flow in asthmatic adolescents using inhaled corticosteroids. *Allergol Immunopathol (Madr)* 2012;40:220-224.
16. *Marković D, Perić T, Sovtić A, Minić P, Petrović V*. Stanje oralnog zdravlja kod dece obolele od astme. *Srp Arh Celok Lek* 2015;143:539-544.(Serbian)
17. *Chakiri H, Bahije L, Fawzi R*. The effects of the asthma and its treatments on oral health of children: A case control study. *Pediatr Dent Care* 2016;1(4):120.

18. *Anandhan V, Bharathan R, Venkataraghavan K, Reddy NV.* The prevalence and severity of dental caries and oral hygiene status of asthmatic children between the age group of 6 and 12 years: A cross-sectional study. *World J Dent* 2012;3:250–254.
19. *Ehsani S, Moin M, Meighani G, Pourhashemi S.J, Khayatpisheh H, Yarahmadi N.* Oral health status in preschool asthmatic children in Iran. *Iran J Allergy Asthma Immunol* 2013;12:254–261.
20. *Mazzoleni S, Stellini E, Cavaleri E, Volponi AA, Ferro R, Colombani SF.* Dental caries in children with asthma undergoing treatment with short-acting β_2 -agonists. *Eur J Paediatric Dent* 2008;9:132-138.
21. *Lindemeyer RG, Satpute NS, Katz SH.* Evaluation of bronchial asthma as risk factor for early childhood caries. *N Y State Dent J* 2011;77:18-21.
22. *Godara N, Godara R, Khullar M.* Impact of inhalation therapy on oral health. *Lung India* 2011;28:272–275.
23. *Stensson M, Wendt LK, Koch G, Nilsson M, Oldaeus G, Birkhed D.* Oral health in pre-school children with asthma-followed from 3 to 6 years. *Int J Paediatr Dent* 2010;20:165-172.
24. *Stensson M, Wendt LK, Koch G, Oldaeus G, Lingström P, Birkhed D.* Caries prevalence, caries-related factors and plaque pH in adolescents with long-term asthma. *Caries Res* 2010;44:540-546.
25. *Steinbacher DM, Glick M.* The dental patient with asthma. An update and oral health considerations. *J Am Dent Assoc* 2001;132:1229-1239.
26. *Harrington N, Prado N, Barry S.* Dental treatment in children with asthma - a review. *Br Dent J* 2016;220(6):299-302.
27. *Shulman JD, Nunn ME, Taylor SE, Rivera-Hidalgo F.* The prevalence of periodontal-related changes in adolescents with asthma: results of the Third Annual National Health and Nutrition Examination Survey. *Pediatr Dent* 2003;25:279-284.
28. *Eloot AK, Vanobbergen JN, De Baets F, Martens LC.* Oral health and habits in children with asthma related to severity and duration of condition. *Eur J Paediatric Dent* 2004;5:210–215.
29. *Ferrazzano GF, Sangianantoni G, Cantile T, Amato I, Ingenito A, Noschese P.* Dental health in asthmatic children: a South Italy study. *J Dent Child (Chic)* 2012;79:170-175.

Table 1

Mean (SD) values of plaque index, gingival index and Community Periodontal Index in observed groups

	AP ($\bar{x}\pm SD$)	AG ($\bar{x}\pm SD$)	NAG ($\bar{x}\pm SD$)	GCA ($\bar{x}\pm SD$)	PCA ($\bar{x}\pm SD$)	p
PI	0.9±0.6	1.2±0.7	0.7±0.5	1.1±0.6	1.2±0.8	<0.001* NS†
GI	0.4±0.4	0.5±0.4	0.3±0.3	0.5±0.4	0.5±0.5	<0.001* NS†
CPI	0.5±0.7	0.8±0.7	0.3±0.5	0.8±0.8	0.8±0.7	<0.001* NS†

* statistical significance between AG and NAG; † statistical significance between GCA and PCA

PI-Plaque index; GI-Gingival index; CPI – Community Periodontal Index; AP – All Participants; AG – Children with asthma; NAG – Children without asthma; GCA – Good Controlled Asthma; PCA – Partly Controlled Asthma; p – level of statistical significance (Independent-Samples T-Test), NS – non significant.

Table 2

Comparison of difference in oral hygiene according by Plaque Index

Plaque Index (%)	AP (n=121)	AG (n=60)	NAG (n=61)	GCA(n=37)	PCA (n=23)
Excellent hygiene	14.9	10.0	19.7	5.4	17.4
Good hygiene	23.1	15.0	31.1	16.2	13.0
Poor hygiene	51.2	55.0	47.5	62.2	43.5
Very poor hygiene	10.7	20.0	1.6	16.2	26.1
p			<0.001*		NS†

* statistical significance between AG and NAG; † statistical significance between GCA and PCA

AP – All Participants; AG – Children with asthma; NAG – Children without asthma; GCA – Good Controlled Asthma; PCA – Partly Controlled Asthma; n – number of subjects; % – percentage of subjects; p – level of statistical significance (Mann-Whitney Test), NS – non significant.

Table 3

Mean (SD) of Plaque Index, Gingival Index and Community Periodontal Index in relation of type administered medication

Type of applied medications	Plaque Index ($\bar{x}\pm SD$)			Gingival Index ($\bar{x}\pm SD$)			CPI ($\bar{x}\pm SD$)		
	AG (n=60)	GCA (n=37)	PCA (n=23)	AG (n=68)	GCA (n=24)	PCA (n=24)	AG (n=60)	GCA (n=37)	PCA (n=23)
Inhalator steroids	1.2±0.6	1.1±0.6	1.2±0.7	0.5±0.4	0.5±0.4	0.5±0.4	0.9±0.7	0.9±0.8	0.8±0.7
Bronchodila- tors	1.2±0.5	0.7±0.4	1.6±0.3	0.7±0.5	0.4±0.4	1.2±0.3	1.1±0.8	1.0±1.2	1.3±0.5
Combination of drugs	1.2±0.7	1.3±0.7	0.9±1.2	0.4±0.4	0.4±0.3	0.5±0.7	0.5±0.6	0.5±0.7	0.4±0.5
P	NS	NS	NS	NS	NS	<0.05	NS	NS	NS

AG – Children with asthma; GCA – Good Controlled Asthma; PCA – Partly Controlled Asthma; CPI – Community Periodontal Index; n – number of subjects; p – level of statistical significance (One-Way ANOVA), NS – non significant.

Table 4

Mean (SD) of Plaque Index, Gingival Index and Community Periodontal Index in relation of time of medication administration

Time of taking medication	Plaque Index ($\bar{x}\pm SD$)			Gingival Index ($\bar{x}\pm SD$)			CPI ($\bar{x}\pm SD$)		
	AG (n=60)	GCA (n=37)	PCA (n=23)	AG (n=68)	GCA (n=44)	PCA (n=24)	AG (n=60)	GCA (n=37)	PCA (n=23)
Only in the morning	1.0±0.6	0.9±0.6	1.5±0.1	0.5±0.4	0.4±0.4	0.6±0.2	0.8±0.9	0.8±1.0	1.0±0.0
Only in the afternoon	2.3±0.5	0.0±0.0	2.3±0.5	1.1±0.1	0.0±0.0	1.1±0.1	2.0±0.0	0.0±0.0	2.0±0.0
In the morning and evening	1.1±0.7	1.2±0.6	0.9±0.8	0.4±0.4	0.5±0.4	0.4±0.5	0.6±0.7	0.7±0.7	0.5±0.6
Before sleeping	1.4±0.4	1.3±0.4	1.5±0.3	0.7±0.5	0.6±0.4	1.0±0.5	1.1±0.7	1.2±0.8	1.0±0.7
P	<0.05	NS	<0.05	<0.05	NS	NS	NS	NS	<0.05

AG – Children with asthma; GCA- Good Controlled Asthma; PCA – Partly Controlled Asthma; CPI – Community Periodontal Index; n – number of subjects; p – level of statistical significance (One-Way ANOVA), NS – non significant.

Table 5

Comparison of difference in gingival health status according to Gingival Index

Gingival Index(%)	AP (n=136)	AG (n=68)	NAG (n=68)	GCA(n=44)	PCA(n=24)
Healthy gingiva	21.3	17.6	25.0	13.6	25.0
Mild gingivitis	64.0	58.8	69.1	68.2	41.7
Moderate gingivitis	14.7	23.5	5.9	18.2	33.3
p		<0.01*		NS†	

* statistical significance between AG and NAG; † statistical significance between GCA and PCA

AP – All Participants; AG – Children with asthma; NAG – Children without asthma; GCA – Good Controlled Asthma; PCA – Partly Controlled Asthma; n – number of subjects; % – percentage of subjects; p – level of statistical significance (Mann-Whitney Test), NS – non significant.

Table 6

Comparison of difference in periodontal status according to Community Periodontal Index

CPI (%)	AP (n=121)	AG (n=60)	NAG (n=61)	GCA (n=37)	PCA (n=23)
0	58.7	38.3	78.7	40.5	34.8
1	29.8	41.7	18.0	37.8	47.8
2	10.7	18.3	3.3	18.9	17.4
3	0.8	1.7	0	2.7	0
P			<0.001*		NS†

* statistical significance between AG and NAG; † statistical significance between GCA and PCA

CPI – Community Periodontal Index ; AP – All Participants; AG – Children with asthma; NAG – Children without asthma; GCA – Good Controlled Asthma; PCA – Partly Controlled Asthma; n – number of subjects; % – percentage of subjects; p – level of statistical significance (Mann-Whitney Test), NS – non significant.

Received on February 21, 2017.

Revised on April 03, 2017.

Accepted on April 13, 2017.

Online First April, 2017.