

Endoscopic findings around the gastroesophageal junction: an experience from a tertiary hospital in Korea

Ji Hyun Kim, M.D.², Jin Ki Hwang, M.D.¹, Juhjung Kim, M.D.¹,
Sehe Dong Lee, M.D.¹, Beom Jae Lee, M.D.¹,
Jae Seon Kim, M.D.¹ and Young-Tae Bak, M.D.¹

*Department of Gastroenterology, Korea University Guro Hospital, Seoul, Korea¹;
Department of Internal Medicine, Inje University Busan Paik Hospital, Busan, Korea²*

Background/Aims: Important lesions related to gastroesophageal reflux disease (GERD) are located around the gastroesophageal junction (GEJ). This study examined the distribution of endoscopic findings around the GEJ and elucidated their relationship to each other and esophageal manometric features.

Methods: Endoscopic data were collected prospectively from 2,450 consecutive diagnostic upper gastrointestinal endoscopies. The presence and degree of hiatal hernia (HH), columnar-lined esophagus (CLE), and reflux esophagitis (RE) were recorded. Esophageal manometric data were collected from 181 patients.

Results: The prevalence of HH, CLE, and RE was 9.8, 18.8, and 9.9%, respectively. Of all HH and CLE cases, 62.8 and 98.9%, respectively, were of the short-segment variety. Of all RE cases, 95.0% were mild. Younger age, male gender, the presence of HH, and a higher gastroesophageal flap valve (GEFV) grades were associated with the presence of RE. Increased ZAP grades were associated with increased prevalence and grades of HH, CLE, and RE. Higher GEFV grades were associated with increased prevalence and grades of HH, CLE, and RE. Lower esophageal sphincter pressure (LESF) decreased in patients with HH or RE compared to those without HH or RE.

Conclusions: Endoscopic findings around the GEJ revealed that a substantial proportion of our patients showed features potentially related to GERD. In combination with other recent reports, our study implies that Korea is no longer a very-low-prevalence area of GERD, although it may predominate in silent or milder forms.

Key Words: Columnar-lined esophagus; gastroesophageal flap valve; hiatal hernia; reflux esophagitis; Z-line appearance

INTRODUCTION

The gastroesophageal junction (GEJ) plays a crucial role as a gatekeeper against gastroesophageal reflux, and most clinically important lesions related to gastroesophageal reflux disease (GERD) are located in this area. Therefore, the importance of careful examination of the GEJ during routine diagnostic upper gastrointestinal (UGI) endoscopy cannot be overemphasized.

Several frequently used classification systems describe aspects of endoscopic findings around the GEJ, including the Los Angeles (LA) classification for reflux esophagitis (RE)¹ and grading systems for the Z-line appearance (ZAP)² and gastroesophageal flap valve (GEFV) morphology³. This study explored the distribution of endoscopic findings around the GEJ and elucidated their relationship to each other and esophageal manometric findings.

• Received: February 9, 2008

• Accepted: June 16, 2008

• Correspondence to: Young-Tae Bak, M.D., Department of Gastroenterology, Korea University Guro Hospital, 97 Gurodong-gil, Guro-gu, Seoul 152-703, Korea Tel: 82-2-2626-1778, Fax: 82-505-115-1778, E-mail: drbakyt@korea.ac.kr

MATERIALS AND METHODS

Study population

Data were collected prospectively from 2,450 consecutive patients visiting the gastroenterology clinic of one author (YTB) and undergoing their first routine diagnostic UGI endoscopy for various reasons at Korea University Guro Hospital, Seoul, Korea from March 2004 to April 2006. Those who underwent urgent or therapeutic endoscopies or who had previously undergone UGI endoscopies at Korea University Guro Hospital were excluded, as were patients with gastric outlet obstruction.

Endoscopic assessment

All endoscopies were performed by one experienced endoscopist (YTB). Endoscopic findings around the GEJ were described separately, including the presence or absence and degree of hiatal hernia (HH), columnar-lined esophagus (CLE), and RE.

HH and CLE were assessed during insertion of the endoscope with minimum air inflation. The GEJ was defined endoscopically as the distal end of longitudinally arrayed palisade capillaries⁴⁾, or, if it was unclear, the proximal end of the gastric type mucosal folds⁵⁾. The patient was diagnosed with HH if the GEJ was located at least 0.5 cm proximal to the level of the diaphragmatic pinchcock action. HH was divided into two grades: short-segment HH if the length was shorter than 2 cm, and long-segment HH if otherwise. The patient was diagnosed with CLE if columnar-looking mucosa was observed proximal to the GEJ⁴⁾ and was at least 0.5 cm in length. CLE was divided into two grades, short-segment CLE if the length was shorter than 3 cm, and long-segment CLE if otherwise.

Reflux esophagitis and ZAP were observed during both insertion and withdrawal of the endoscope, with maximal air inflation of the lower esophagus in full inspiration if possible. Reflux esophagitis was diagnosed and graded according to the LA classification¹⁾. The Z-line was observed carefully for any tongue-like projections or islands of columnar epithelium proximal to the Z-line. ZAP was defined and graded according to Wallner et al²⁾. GEFV with a mucosal fold or ridge surrounding an endoscopic shaft was observed in the retroflexed view and was graded according to Hill et al³⁾.

Esophageal manometry

Routine esophageal manometry was performed in patients showing symptoms suggestive of esophageal motility disorders, such as dysphagia or chest pain. The manometric data were transferred to a computer via a polygraph (Polygraf ID; Medtronic Functional Diagnostics, Skovlunde, Denmark) and pressure transducers (DPT-6000; Smiths Medical Deutschland, Kirshseeon, Germany), which were connected to an

eight-channel water perfusion catheter (Medtronic Functional Diagnostics) and a low-compliance pneumohydraulic capillary infusion system (PIP 3-8; Sandhill Scientific, Colorado, USA). The manometric data were reviewed using analysis software (Polygram NET version 4.1; Medtronic Functional Diagnostics), and the lower esophageal sphincter pressure (LESP) was measured using the station pull-through technique. Esophageal peristalsis was evaluated after each of the last 10 of 15 swallows of 5 mL of water, and the distal esophageal peristaltic amplitude (DEA) was measured at 3 and 8 cm proximal to the lower esophageal sphincter.

Statistical analysis

Statistical analysis was performed using SPSS (version 13.0 for Windows, Chicago, IL, USA). Data were expressed as the mean±standard deviation (SD). One-way analysis of variance (ANOVA) was used to assess statistical significance for age, body mass index (BMI), and esophageal manometry parameters according to HH, CLE, ZAP, and GEFV status. Age, BMI, and esophageal manometry parameters were compared among patients with or without HH, CLE, and RE, and among groups with different ZAP and GEFV grades, using unpaired *t*-tests. Differences in HH, CLE, and RE according to ZAP and GEFV grades and the relationship between ZAP and GEFV grades were assessed using the chi-square test for trend. Spearman's rank correlation coefficient was calculated to examine the correlation between GEFV and RE grades, ZAP and RE grades, and HH and RE grades. To evaluate the ability of the variables of interest to predict CLE and RE, a logistic regression analysis model was used. The LESP and DEA values were compared using one-way ANOVA or unpaired *t*-tests using log-transformed data. $p < 0.05$ was considered statistically significant.

RESULTS

Endoscopic data were collected from 2,450 patients (1,152 men and 1,298 women, age 51.4±13.7 years). The prevalence of endoscopic HH, CLE, and RE was 9.8, 18.8, and 9.9%, respectively. Short-segment HH made up 62.8% of all HH cases, whereas short-segment CLE made up 98.9% of all CLE cases. Of all RE cases, 95.0% showed grades LA-A or B. ZAP grade I and GEFV grade II were the most common types, occurring in 51.1 and 47.6% of patients, respectively (Table 1).

The age and BMI of patients with endoscopic CLE were significantly higher than in those without endoscopic CLE, although the differences were not large. Male gender, the presence of HH or RE, and higher ZAP (grade II or III) or GEFV (grade III or IV) grades were all significantly associated

Table 1. Endoscopic findings around the gastroesophageal junction (n=2,450).

Endoscopic findings	n (%)
Hiatal hernia, n=239 (9.8%)	
Short segment (< 2 cm)	150 (6.1%)
Long segment (≥ 2 cm)	89 (3.6%)
Columnar-lined esophagus, n=460 (18.8%)	
Short-segment (< 3 cm)	455 (18.6%)
Long-segment (≥ 3 cm)	5 (0.2%)
Reflux esophagitis, ^{1) n=242 (9.9%)}	
LA-A	145 (5.9%)
LA-B	85 (3.5%)
LA-C	11 (0.4%)
LA-D	1 (0.0%)
Z-line appearance ²⁾	
0	873 (35.6%)
I	1253 (51.1%)
II	289 (11.8%)
III	5 (0.2%)
Unknown	30 (1.2%)
Gastroesophageal flap valve ³⁾	
I	635 (25.9%)
II	1166 (47.6%)
III	347 (14.2%)
IV	234 (9.6%)
Unknown	68 (2.8%)

with the presence of endoscopic CLE in the univariate and multivariate analyses (Table 2).

Patients with RE were significantly younger than those without RE, although the difference was not large. No significant difference in BMI was observed between those with or without RE. Male gender, the presence of HH or CLE, and high GEFV grades (III or IV) were significantly associated with the presence of RE in the univariate and multivariate analyses. Higher ZAP grade (II or III) was significantly associated with the presence of RE in the univariate analysis, but not in the multivariate analysis (Table 3).

Patients with ZAP grades of I or II were significantly older

than those with ZAP grades of 0, and males were more likely to have high ZAP grades. Patients with ZAP grade III had a higher BMI than those with ZAP grades of 0. An increase in ZAP grade was also significantly associated with the increased prevalence of HH, CLE, and RE (Table 4). Increased HH, CLE, and RE grades were all correlated with increased ZAP grades (Spearman's rank correlation, $r=0.075$ and $p<0.001$, $r=0.548$ and $p<0.001$, and $r=0.141$ and $p<0.001$, respectively).

No difference was observed in GEFV grade according to age, although increased GEFV grade was significantly associated with male gender. BMI was also significantly higher in patients with GEFV grades of IV compared to those with lower GEFV grades, and increased GEFV grades were significantly associated with the increased prevalence of HH, CLE, and RE (Table 5). Higher HH, CLE, and RE grades were all significantly correlated with higher GEFV grades (Spearman's rank correlation, $r=0.325$ and $p<0.001$, $r=0.198$ and $p<0.001$, and $r=0.231$ and $p<0.001$, respectively). Higher GEFV grades were correlated with higher ZAP grades (Spearman's rank correlation, $r=0.271$, $p<0.001$; Table 6).

Esophageal manometric data from 181 patients (68 men and 113 women) that underwent both upper endoscopy and esophageal manometry were analyzed. LESP was significantly lower in patients with HH or RE than in patients without HH or RE. Patients with high GEFV grades (III or IV) had significantly lower LESP and DEA values. LESP and DEA did not differ between patients with or without endoscopic CLE or between patients with lower versus higher ZAP grades (Table 7).

In patients with LA-B grade RE, LESP was significantly lower than in patients without RE, and there was a negative correlation between LESP and GEFV grade (Table 8). No difference in LESP was observed according to the grade of HH, CLE, or ZAP.

Table 2. Clinical and endoscopic features of patients with or without endoscopic columnar-lined esophagus (n=2,450).

	Columnar-lined esophagus		<i>p</i> value, univariate	<i>p</i> value, multivariate	Multivariate adjusted OR (95% CI)
	Present (n=460)	Absent (n=1,990)			
Age, years	52.9±13.0	51.0±13.8	0.008	0.018	1.01 (1.00–1.03)
Male gender, n (%)	310 (67.4)	842 (42.3)	< 0.001	< 0.001	2.38 (1.71–3.31)
BMI, kg/m ²	24.0±3.0	23.2±3.5	< 0.001	0.001	1.06 (1.03–1.11)
HH, n (%)	88 (19.1)	151 (7.6)	< 0.001	0.001	3.58 (2.38–5.37)
RE, n (%)	77 (16.7)	165 (8.3)	< 0.001	0.012	1.60 (1.04–2.46)
ZAP II or III, n (%)	269 (58.5)	25 (1.3)	< 0.001	< 0.001	126.21 (78.25–203.56)
GEFV III or IV, n (%)	186 (40.4)	395 (19.8)	< 0.001	< 0.001	1.56 (1.10–2.22)

OR, odds ratio; CI, confidence interval; BMI, body mass index; HH, hiatal hernia; RE, reflux esophagitis; ZAP, Z-line appearance²⁾; GEFV, gastroesophageal flap valve³⁾.

Table 3. Clinical and endoscopic features of patients with or without reflux esophagitis (n=2,450)

	Reflux esophagitis		<i>p</i> value, univariate	<i>p</i> value, multivariate	Multivariate adjusted OR (95% CI)
	Present (n=242)	Absent (n=2208)			
Age, years	48.3±14.4	51.7±13.5	< 0.001	< 0.001	0.98 (0.97–0.99)
Male gender, n (%)	165 (68.2)	987 (44.7)	< 0.001	0.013	1.72 (1.25–2.38)
BMI, kg/m ²	23.3±3.4	23.6±3.5	0.144	0.327	1.02 (0.98–1.07)
HH, n (%)	71 (29.3)	168 (7.6)	< 0.001	< 0.001	2.84 (1.95–4.12)
CLE, n (%)	77 (31.8)	383 (17.3)	< 0.001	0.032	1.59 (1.04–2.44)
ZAP II, or III, n (%)	42 (17.4)	252 (11.4)	< 0.001	0.931	0.98 (0.59–1.63)
GEFV III or IV, n (%)	123 (50.8)	458 (20.7)	< 0.001	< 0.001	2.42 (1.75–3.35)

OR, odds ratio; CI, confidence interval; BMI, body mass index; HH, hiatal hernia; CLE, columnar-lined esophagus; ZAP, Z-line appearance²; GEFV, gastroesophageal flap valve³.

Table 4. Case profiles and Z-line appearance (n=2,420)

	Z-line appearance				<i>p</i> value
	0 (n=873)	I (n=1253)	II (n=289)	III (n=5)	
Age, years	49.8±13.9 ^{a,b}	52.2±13.4 ^a	52.4±12.7 ^b	52.8±26.1	0.001
Male gender, n (%)	268 (30.7)	678 (54.1)	186 (64.4)	4 (80.0)	< 0.001
BMI, kg/m ²	22.7±3.7 ^c	23.6±3.3	23.7±2.9	25.4±5.5 ^c	< 0.001
HH, n (%)	54 (6.2)	152 (12.2)	28 (9.7)	2 (40.0)	< 0.001
< 2.0 cm	31	96	20	2	
≥ 2.0 cm	23	56	8	0	
CLE, n (%)	13 (1.5)	177 (14.1)	264 (91.3)	5 (100.0)	< 0.001
< 3.0 cm	13	177	264	0	
≥ 3.0 cm	0	0	0	5	
RE, n (%)	36 (4.1)	158 (12.7)	40 (13.8)	2 (40.0)	< 0.001
LA-A	27	99	18	1	
LA-B	8	51	21	1	
LA-C	1	8	0	0	
LA-D	0	0	1	0	

a, b, and c, *p*<0.05 versus the same letter by Tukey's multiple comparison; BMI, body mass index; RE, reflux esophagitis according to the Los Angeles (LA) classification; CLE, columnar-lined esophagus; HH, hiatal hernia.

DISCUSSION

Recent epidemiologic studies indicate that RE is on the rise in Asia, with a reported prevalence of 3~16%⁶⁻⁸. However, most cases (80~95%) present in a relatively mild form (LA-A or B)^{6,9}. In our study, the prevalence of RE was 9.9%, and 95.0% of these cases were mild. Although our data may not represent the general Korean population, recent reports from other Korean cities obtained similar data for the prevalence of erosive RE^{10, 11}.

Younger age, male sex, the presence of HH or CLE, and higher GEFV grade were significantly associated with the presence of RE in our study. There are several reported pathogenic factors for RE, including low LESP, short LES length (especially the intra-abdominal portion), disturbed esophageal acid clearance, and transient LES relaxation¹²⁻¹⁶. Our data suggest that patients with RE were younger than those without RE; although the explanation for this is unclear, it may be due to an alpha error resulting from a very large sample size. In

addition, although the difference was statistically significant, it may not be large enough to be clinically relevant. Several studies have reported that LESP and esophageal motor activity become progressively impaired with increasing severity of RE^{17, 18}. We found a significantly lower LESP in patients with RE compared to those without RE. In addition, our study demonstrated a progressive decrease in LESP with increasing RE grade, suggesting a probable correlation between lower LESP and RE.

Hiatal hernia is thought to predispose patients to GERD via various mechanisms, such as low LESP and impaired acid clearance from the esophagus^{19, 20}. Our results are consistent with previous reports and demonstrate correlations between the presence of HH and RE and lower LESP.

The definition of endoscopic CLE is controversial as it depends on the definition of the precise point at which the esophagus ends and the stomach begins. Generally, the endoscopic GEJ is defined as the proximal end of the gastric

Table 5. Case profiles and the gastroesophageal flap valve (n=2,382)

	Gastroesophageal flap valve				p value
	I (n=635)	II (n=1166)	III (n=347)	IV (n=234)	
Age, years	52.4±13.8	51.1±12.8	51.81±4.2	49.8±15.6	0.056
Male gender, n (%)	142 (22.4)	553 (47.4)	241 (69.5)	179 (76.5)	< 0.001
BMI, kg/m ²	22.7±3.5 ^{a,b,c}	23.5±3.3 ^{a,d}	23.4±3.0 ^{b,e}	24.1±4.0 ^{c,d,e}	< 0.001
HH, n (%)	6 (0.9)	73 (6.3)	58 (16.7)	96 (41.6)	< 0.001
< 2.0 cm	4	54	41	46	
≥ 2.0 cm	2	19	17	50	
CLE, n (%)	66 (10.4)	202 (17.3)	102 (29.4)	84 (36.1)	< 0.001
< 3.0 cm	66	198	102	83	
≥ 3.0 cm	0	4	0	1	
RE, n (%)	16 (2.5)	97 (8.3)	60 (17.5)	63 (27.0)	< 0.001
LA-A	13	63	35	31	
LA-B	3	30	21	29	
LA-C	0	4	4	2	
LA-D	0	0	0	1	

a, b, c, d, and e, *p*<0.05 versus the same letter by Duncan's multiple comparison; BMI, body mass index; RE, reflux esophagitis according to the Los Angeles classification; CLE, columnar-lined esophagus; HH, hiatal hernia.

Table 6. Z-line appearance (ZAP) and the gastroesophageal flap valve (GEFV) (n=2,371)[†]

	GEFV I (n=635)	GEFV II (n=1161)	GEFV III (n=345)	GEFV IV (n=230)
ZAP 0 (n=855)	356 (56.1%)	382 (32.9%)	85 (24.6%)	32 (13.9%)
ZAP I (n=1,226)	230 (36.2%)	647 (55.7%)	204 (59.1%)	145 (63.0%)
ZAP II (n=285)	49 (7.7%)	128 (11.0%)	56 (16.2%)	52 (22.6%)
ZAP III (n=5)	0 (0.0%)	4 (0.3%)	0 (0.0%)	1 (0.4%)

[†]*p*<0.001, Spearman's rank correlation.

Table 7. Lower esophageal sphincter pressure (LESP) and distal esophageal contraction amplitude (DEA) according to endoscopic findings

	LESP (mmHg)	p value [†]	DEA (mmHg)	p value [†]
Hiatal hernia (n=181)		0.023		0.759
Present (n=27)	27.4±8.6		79.4±36.6	
Absent (n=154)	34.3±15.4		81.8±40.1	
Columnar-lined esophagus (n=179)		0.249		0.132
Present (n=34)	30.1±10.8		70.6±28.4	
Absent (n=145)	33.9±15.4		84.0±41.3	
Reflux esophagitis (n=181)		0.017		0.066
Present (n=31)	28.4±13.9		67.7±29.8	
Absent (n=150)	34.0±14.8		84.1±40.6	
Z-line appearance (n=177)		0.173		0.492
0 or I (n=155)	33.7±15.0		82.8±40.9	
II or III (n=22)	29.4±12.3		73.4±26.8	
Gastroesophageal flap valve (n=172)		< 0.001		0.034
I or II (n=134)	35.2±14.9		84.2±40.4	
III or IV (n=38)	26.7±13.2		70.9±37.4	

[†]Independent t-test on log-transformed data.

folds in addition to the distal end of the fine longitudinal vessels^{4, 5, 21}. We diagnosed endoscopic CLE if the length of columnar-lined mucosa proximal to the distal margin of the

palisade zone on the lower esophagus was 0.5 cm or longer; if it was unclear, the proximal end of the gastric type mucosal folds was used as an indicator of the GEJ instead. We adopted

Table 8. Lower esophageal sphincter pressure (LESP) according to grade of reflux esophagitis and gastroesophageal flap valve

	LESP (mmHg)	p value [†]
Reflux esophagitis (n=181)		0.032
Negative (n=150)	34.0±14.8 ^a	
LA-A (n= 20)	31.7±16.5	
LA-B (n=8)	21.4±4.1 ^a	
LA-C (n=3)	27.0±8.0	
Gastroesophageal flap valve (n=172)		< 0.001
I (n=57)	38.1±18.2 ^{d,e}	
II (n=77)	33.1±11.5 ^f	
III (n=26)	30.1±14.4 ^{d,g}	
IV (n=12)	19.5±5.9 ^{e,f,g}	

a, b, c, d, e, f, and g, $p < 0.05$ between the same characters.

[†]One-way ANOVA using log-transformed data.

an arbitrary threshold of 0.5 cm to avoid overdiagnosis, considering a potential gap of several millimeters between the "true" GEJ and the distal margin of the palisade zone⁵). The prevalence of endoscopic CLE in our study was 18.8%, and almost all patients showed the short-segment variety. Based on our previous experience, about one-third of endoscopic CLE cases would be diagnosed as Barrett's esophagus after histological examination with non-targeted biopsy specimens²²).

In this study, older age, male gender, higher BMI, the presence of HH, the presence of RE, and higher GEFV grades were significantly associated with the presence of endoscopic CLE. These findings have implications for the association between gastroesophageal reflux and the presence of endoscopic CLE. Again, the discrepancy in age between RE and CLE cannot be explained clearly. However, as noted in patients with RE, the difference in age according to the presence or absence of CLE may not be large enough to be clinically significant. In addition, we did not detect a correlation between the presence of endoscopic CLE and manometric parameters, such as LESP and DEA.

Wallner et al.¹⁾ first proposed the ZAP classification, a 4-grade system for the appearance of the endoscopic Z-line. This system is correlated with the prevalence of intestinal metaplasia in the GEJ and GERD symptom frequency²³⁻²⁵). In our study, the presence of HH, CLE, higher RE grades, and higher GEFV grades were all significantly correlated with ZAP grades. These results suggest that the ZAP classification is an indicator of pathological changes in the GEJ and gastroesophageal reflux.

The GEFV, the sling or oblique fibers of the stomach located below the LES, contribute to the barrier function via a flap valve mechanism. Since Hill et al.²⁾ developed an endoscopic grading system to assess the GEFV, several reports have noted a correlation between increased GEFV grades and the increased

prevalence of GERD and Barrett's esophagus^{10, 26, 27}). We found that male gender was associated with higher GEFV grades and our data confirm the association between an increased GEFV grade and the increased prevalence and grade of HH, CLE, and RE. These results suggest that GEFV grading may permit the endoscopic evaluation of gastroesophageal reflux status. We also found a negative correlation between GEFV grade and LESP level, implying that increasing GEFV is associated with the deterioration of LES function.

In conclusion, endoscopic findings around the GEJ revealed that a substantial proportion of our patients showed features potentially related to GERD. Combined with other reports^{10, 11}), these results suggest that Korea is no longer a very-low-prevalence region for GERD, although the disease may predominate in silent or milder forms.

REFERENCES

- 1) Armstrong D, Bennett JR, Blum AL, Dent J, De Dombal FT, Galmiche JP, Lundell L, Margulies M, Richter JE, Spechler SJ, Tytgat GN, Wallin L. *The endoscopic assessment of esophagitis: a progress report on observer agreement. Gastroenterology* 111:85-92, 1996
- 2) Wallner B, Sylvan A, Stenling R, Janunger KG. *The esophageal Z-line appearance correlates to the prevalence of intestinal metaplasia. Scand J Gastroenterol* 35:17-22, 2000
- 3) Hill LD, Kozarek RA, Kraemer SJ, Aye RW, Mercer CD, Low DE, Pope CE 2nd. *The gastroesophageal flap valve: in vitro and in vivo observations. Gastrointest Endosc* 44:541-547, 1996
- 4) Choi DW, Oh SN, Baek SJ, Ahn SH, Chang YJ, Jeong WS, Kim HJ, Yeon JE, Park JJ, Kim JS, Byun KS, Bak YT, Lee CH. *Endoscopically observed lower esophageal capillary patterns. Korean J Intern Med* 17:245-248, 2002
- 5) Boyce HW. *Endoscopic definitions of esophagogastric junction regional anatomy. Gastrointest Endosc* 51:586-592, 2000
- 6) Goh KL. *Changing epidemiology of gastroesophageal reflux disease in the Asian-Pacific region: an overview. J Gastroenterol Hepatol* 19(Suppl 3):S22-S25, 2004
- 7) Ho KY, Chan YH, Kang JY. *Increasing trend of reflux esophagitis and decreasing trend of Helicobacter pylori infection in patients from a multiethnic Asian country. Am J Gastroenterol* 100:1923-1928, 2005
- 8) Wong BC, Kinoshita Y. *Systemic review on epidemiology of gastroesophageal reflux disease in Asia. Clin Gastroenterol Hepatol* 4:398-407, 2006
- 9) Okamoto K, Iwakiri R, Mori M, Hara M, Oda K, Kanjo A, Ootani A, Sakata H, Fujimoto K. *Clinical symptoms in endoscopic reflux esophagitis: evaluation in 8031 adult subjects. Dig Dis Sci* 48:2237-2241, 2003
- 10) Kim GH, Kang DH, Song GA, Kim TO, Heo J, Cho M, Kim JS, Lee BJ, Wang SG. *Gastroesophageal flap valve is associated with gastroesophageal and gastropharyngeal reflux. J Gastroenterol* 41:654-661, 2006
- 11) Lee JY, Hong WK, Sohn KM, Kim DY, Kim YM, Choi HH, Choi CS,

- Kim SJ, Baek KH, Kim JB, Kim DJ. *Prevalence and clinical characteristics of gastroesophageal reflux disease and reflux esophagitis in Chuncheon city: health care examination study. Korean J Med* 74:385-390, 2008
- 12) Mittal RK, Balaban DH. *Esophagogastric junction. N Engl J Med* 336:924-932, 1997
 - 13) Meining A, Fackler A, Tzavella K, Storr M, Allescher HD, Klauser A, Heldwein W. *Lower esophageal sphincter pressure in patients with gastroesophageal reflux disease and posture and time patterns. Dis Esophagus* 17:155-158, 2004
 - 14) Behar J, Biancani P, Sheahan DG. *Evaluation of esophageal tests in the diagnosis of reflux esophagitis. Gastroenterology* 71:9-15, 1976
 - 15) Katzka DA. *Motility abnormalities in gastroesophageal reflux disease. Gastroenterol Clin North Am* 28:905-915, 1999
 - 16) Mittal RK, Holloway RH, Penagini R, Blackshaw LA, Dent J. *Transient lower esophageal sphincter relaxation. Gastroenterology* 109:601-610, 1995
 - 17) Csendes A, Maluenda F, Burdiles P, Henriquez A, Quesada MS, Csendes P. *Prospective study of esophageal motor abnormalities in patients with gastroesophageal disease reflux according to severity of endoscopic esophagitis. Hepatogastroenterology* 43:394-399, 1996
 - 18) Zaninotto G, DeMeester TR, Bremner CG, Smyrk TC, Cheng SC. *Esophageal function in patients with reflux-induced strictures and its relevance to surgical treatment. Ann Thorac Surg* 47:362-370, 1989
 - 19) Sloan S, Rademaker AW, Kahrilas PJ. *Determinants of gastroesophageal junction incompetence: hiatal hernia, lower esophageal sphincter, or both? Ann Intern Med* 117:977-982, 1992
 - 20) Mittal RK, Lange RC, McCallum RW. *Identification and mechanism of delayed esophageal acid clearance in subjects with hiatal hernia. Gastroenterology* 92:130-135, 1987
 - 21) Spechler SJ. *Columnar-lined esophagus: definitions. Chest Surg Clin N Am* 12:1-13, 2002
 - 22) Kim JY, Kim YS, Jung MK, Park JJ, Kang DH, Kim JS, Song CW, Lee SW, Bak YT. *Prevalence of Barrett's esophagus in Korea. J Gastroenterol Hepatol* 20:633-636, 2005
 - 23) Wallner B, Sylvan A, Stenling R, Janunger KG. *The esophageal Z-line appearance correlates to the prevalence of intestinal metaplasia. Scand J Gastroenterol* 35:17-22, 2000
 - 24) Wallner B, Sylvan A, Stenling R, Janunger KG. *The Z-line appearance and prevalence of intestinal metaplasia among patients without symptoms or endoscopic signs indicating gastroesophageal reflux. Surg Endosc* 15:886-889, 2001
 - 25) Wallner B, Sylvan A, Janunger KG, Bozoky B, Stenling R. *Immuno-histochemical markers for Barrett's esophagus and associations to esophageal Z-line appearance. Scand J Gastroenterol* 36:910-915, 2001
 - 26) Fujiwara Y, Higuchi K, Shiba M, Watanabe T, Tominaga K, Oshitani N, Matsumoto T, Arakawa T. *Association between gastroesophageal flap valve, reflux esophagitis, Barrett's epithelium, and atrophic gastritis assessed by endoscopy in Japanese patients. J Gastroenterol* 38:533-539, 2003
 - 27) Lin BR, Wong JM, Chang MC, Liao CC, Yang PM, Jeng JH, Liao WR, Wang TH. *Abnormal gastroesophageal flap valve is highly associated with gastroesophageal reflux disease among subjects undergoing routine endoscopy in Taiwan. J Gastroenterol Hepatol* 21:556-562, 2006