

## Review

# No more recalcitrant: Chickpea regeneration and genetic transformation

Firoz Anwar<sup>2\*</sup>, P. Sharmila<sup>1</sup> and P. Pardha Saradhi<sup>1</sup>

<sup>1</sup>Department of Environmental Biology, North Campus, University of Delhi, Delhi, India.

<sup>2</sup>Plant Production Department, Biotechnology Division, King Saud University, Riyadh Kingdom of Saudi Arabia.

Accepted 28 December, 2009

Chickpea production is limited worldwide because of abiotic and biotic stresses. Efforts to overcome these production constraints through traditional breeding are difficult due to limited genetic variation. Novel regeneration is pre-requisite for genetic transformation offers the opportunity to overcome hybridization barriers and introduce novel genes for resistance. Although direct gene transfer via direct DNA transfer has been reported, *Agrobacterium* mediated transformation is the preferred method and standard protocols have been established for the production of transgenic plantlets derived from co-cultivation of embryonic axes. This was soon adopted due to difficulties associated with regeneration of whole plants from callus. Only few reports have been reported using genetic transformation/transgene(s) against abiotic stress tolerance transgenic chickpea plants. Transgenic chickpea using bacterial *codA* gene tolerance against abiotic stresses have been developed. Chickpea improvement and application of genomics tools to the study of the chickpea genome will be enhanced through the use of genetic transformation.

**Key words:** *Cicer arietinum* L., TDZ, Indole butyric acid, (IBA), efficient rooting, aeration, transplantation, transformation.

## INTRODUCTION

Chickpea (*Cicer arietinum* L.) is an important food crop in much of the developing world and ranks third in production among food legumes. India accounts for 75% of the world's production and is grown in rotation with cereal crops to break disease cycles, allow grassy weed control and improve soil nutrient status through the return of atmospheric nitrogen fixed through symbiosis. Chickpea is good as a source of carbohydrate (48.2 - 67.6%), protein (12.4 - 31.5%), fat (6%) and nutritionally important minerals. Among the legumes, chickpea is the best hypocholesteremic agent, followed by black gram and green gram. Direct shoot organogenesis and establishment of plantlets from different explants of chickpea was reported earlier (Polisetty et al., 1996, 1997; Paul et al., 2000; Rizvi and Singh, 2000; Chauhan et al., 2003; Jayanand et al., 2003; Chakraborti et al., 2006; Table 1). Plantlets were developed through direct somatic embryogenesis and through callus from different explants of chickpea

{Barna and Wakulu, 1993; Sagare et al., 1993; Suhasini et al., 1994; Kumar et al. 1994, 1995; Rizvi and Singh, 2000; Chauhan et al., 2002; Kar et al., 1996, 1997; Kiran et al., 2005 (Table 2)}.

In spite of several reports of successful regeneration, chickpea is widely considered to be highly recalcitrant (Shri and Davis, 1992; Vani and Reddy, 1996; Rizvi and Singh, 2000; Polowick et al., 2004). Surprisingly, majority of published/reported chickpea regeneration protocols often are either not repeatable or work only in certain research laboratories, making researchers to believe that chickpea regeneration is highly recalcitrant. Two major hurdles that limit *in vitro* regeneration of chickpea are (i) induction and development of strong root system and (ii) establishment of *in vitro* raised plantlets in pots. In order to escape from these hurdles, researchers have preferred to go for grafting (Krishnamurthy et al., 2000; Sarmah et al., 2004; Senthil et al., 2004; Tewari-Singh et al., 2004; Sanyal et al., 2005). In general, grafting is tedious and time consuming requiring special skills. Moreover, grafting besides requiring additional seed lot, also might promote emergence of shoots/branches from axillary

\*Corresponding author. E-mail: [anfiroz@gmail.com](mailto:anfiroz@gmail.com).

**Table 1.** Regeneration in various genotypes of chickpea.

Cultivar	Medium	Growth regulators	Explants	Rooting	Responses	References
B115, C-235,	MMS	BAP, NAA, IAA	Single cotyledon with 1/2 embryo	Grafting	Establishment of plants	Chakraborti et al., 2006
C-235	MS	TDZ, 2-iP, Kinetin, GA <sub>3</sub> , IBA	Embryonic axis	MS + IBA 5 μM/ 10 mM Pulse	Establishment of plants	Jayanand et al., 2003
C-235, k-850, BG-256, PGD- 84-10	MS MS salt+B <sub>5</sub>	IAA, NAA, IBA Kinetin, BAP	a. Mature embryo b. Immature embryo	MS + 10.7 μM NAA + 9.8 μM IBA	Establishment of plants	Chauhan et al., 2003
P-256	MS	TDZ, Proline	Whole Seedling (cotyledonary notch region)	MS + 2.5 μM NAA	Establishment of plants	Rizvi et al., 2002
P-256	MS	TDZ, BAP, IBA, Spermidine, Putrescine, Potassium phosphate	Immature leaflet	MS + 2.5 μM IBA	Establishment of plants	Rizvi and Singh, 2000
C-235, BG-329	MS salts + B <sub>5</sub> vitamins	BAP	Excised root ip from two day old seedlings	Modified MS	Establishment of plants	Paul et al., 2000
BG-362, BG-329, BG-267, BG-256, C-235	MS Salts + B <sub>5</sub> Vitamins	Different concentrations of BA	a) Embryo axis with both cotyledons b) as in (a) except the tip of radicle c) As in (a) except one of the cotyledons d) as in (c) except the tip of radicle	1/4 <sup>th</sup> MS	Multiple shoots and thin plants were developed	Polisetty et al., 1997
BG-362, BG-329 BG-267, BG-256 C-235	MS	NAA, IBA	Shoot tips	1/4 <sup>th</sup> MS	Establishment of plants	Polisetty et al., 1996
ICCV-1, ICCV-6 and Desi Variety	MS Macro + 4xMS Micro + B <sub>5</sub> Vitamins	BAP, NAA, IBA	Embryonic axis without apical root and shoot part	MS + 0.01 mg l <sup>-1</sup> IBA	Formation of plants	Kar et al., 1996
JG-62 C-235, PGC1	B <sub>5</sub>	BA, Kinetin, IAA	Epicotyl	B <sub>5</sub> + 4 mg l <sup>-1</sup> / IAA + 0.5 mg l <sup>-1</sup> KN	Multiple Shoots and shoots were rooted	Vani and Reddy, 1996
C-235	MS	2.4D, BAP, NAA, IBA, Picloram	Leaflet	MS + 1 μM IBA	Formation of plants	Barna and Wakhlu, 1994
Annigeri, ICCV6	B <sub>5</sub>	BAP, IBA, Kin,	a. Meristem tips (apical meristem and leaf primordia), b. Cotyledonary-nodes	MS + 2.5 μM IBA	Multiple shoots and shoots were rooted	Brandt and Hess, 1993

buds that are often retained in the axils of cotyledons and the later might dominate over the grafted shoots.

The following section is a brief overview on chickpea regeneration and genetic transformation and confirmed

Table 1. Contd.

Cultivar	Medium	Growth regulators	Explants	Rooting	Responses	References
Pusa-256	MS	NAA , IBA	Shoot tip	MS + 2 mg l <sup>-1</sup> NAA + 2 mg l <sup>-1</sup> IBA	Multiple shoots and individual shoots were rooted	Chandra et al., 1993
Local Ecotype	MS	Kinetin, BAP, NAA, IAA	Shoot tips	MS + 0.5 mg l <sup>-1</sup> IAA + 0.05 mg l <sup>-1</sup> Kin	Multiple shoots and individual shoots were rooted	Fontana et al., 1993
C-235	MS	2.4D, BAP, NAA	Leaflet	-	Multiple shoot	Barna and Wakhlu 1993
C-235, H-208	B <sub>5</sub>	BAP, NAA	a) White Seed b) Shoot tip c) Cotyledon	-	Multiple shoots	Prakash et al., 1992
Kabuli	MS salts + B <sub>5</sub> Vitamins	TDZ, NAA	Mature Seeds	-	Spontaneous shoot differentiation on prolonged culture of seedlings	Malik and Saxena. 1992.
ICC 640	B <sub>5</sub>	Zeatin, IAA, BA	Immature Cotyledons with excised embryonal axis	-	White cotyledon like structures (CLS) formed at the proximal end of the cotyledons; multiple shoots initiation near the base of CLS.	Shri and Davis 1992
ICCC-4 Annigeri, H-08	B <sub>5</sub>	BAP, Kinetin, IAA	a) Hypocotyl b) Shoot tip	-	Multiple shoots	Neelam et al., 1986
G543, G130, L550	MS	2, 4 -D, IAA, NAA, Kinetin, BA	Shoot tip meristem	0.1 mg l <sup>-1</sup> NAA + 0.02 mg l <sup>-1</sup> BA	Shoots with root	Bajaj and Dhanju, 1979.

that chickpea is no more recalcitrant.

## REGENERATION OF CHICKPEA

### Direct shoot organogenesis from different explants of chickpea


A procedure to initiate shoot regeneration from various explants derived from mature seeds germinated on a medium containing cytokinins or cytokinins like substances and/ or auxin has been developed using *in vitro* culture of seeds in the presence of TDZ. Among the different cytokinins used in this study, a higher frequency of adventitious shoot buds/multiple shoots formation was observed in TDZ containing medium (Anwar et al., 2008). TDZ induces high frequency of somatic embryogenesis/ adventitious bud formation in some plant species, either alone or in combination with other growth regulator/s (Murthy et al., 1998, 1995; Chen and Chang, 2006; Delvin, 1989; Ganeshan et al. 2003; Huettelman and Preece, 1993; Onamu, 2003; Mroginski, 2004; Shan, 2000; Sharma, 2005; Tang and Newton, 2005, 2006 and reference cited in). A substituted phenyl urea derivative was found to be very effective for induction of multiple

shoots from nodal and basal regions of the primary epicotyl (Malik and Saxena, 1992). MS salts and B<sub>5</sub> vitamins supplemented with 5 µM benzyl aminopurine (BAP) has been found to be a highly effective medium for multiple shoot formation from intact seedlings (Polisetty et al., 1997). This procedure excludes the selection and preparation of explants and complex manipulation of cultural condition. Multiple shoots have also been regenerated directly from immature cotyledons on B<sub>5</sub> basal medium containing zeatin (13.7 µM) and indole acetic acid (IAA) (0.2 µM) (Shri and Davis, 1992).

Regeneration of multiple shoots has been achieved from seedling. Shoot tips excised from field grown plantlets gave rise to multiple shoots when cultured on MS medium supplemented with IAA (2 mg l<sup>-1</sup>) and Kinetin (0.5 mg l<sup>-1</sup>) (Bajaj and Dhanju, 1979). Multiple shoots have been induced from shoots tips cultured on MS medium supplemented with naphthalene acetic acid (NAA) (2 mg l<sup>-1</sup>) and indole butyric acid (IBA) (2 mg l<sup>-1</sup>) (Chandra et al., 1993; Polisetty et al., 1996). Epicotyl explants gave rise to multiple shoots when cultured on B<sub>5</sub> medium containing BA (1 mg l<sup>-1</sup>) and kinetin (1 mg l<sup>-1</sup>) + IAA (0.5 mg l<sup>-1</sup>) (Vani and Reddy, 1996). Various workers have achieved multiple shoots from embryo axis apical meristem cultured on MS containing kinetin (1 mg l<sup>-1</sup>)

**Table 2.** Somatic embryogenesis and plant development in various genotypes of chickpea.

Cultivar	Medium	Growth regulators	Explants	Responses	References
<b>Direct Somatic Embryogenesis</b>					
1. ICCV-10, Annigeri	MS	Picloram, 2, 4 -D, 2, 4, 5 -T, NAA, BAP, Kinetin	Hypocotyls	Plants developed from somatic embryos	Kiran et al., 2005
2. PG12, C-235	MS	2, 4, 5 -T, 2, 4 -D, BAP, Kinetin	a. Immature (IM)-cotyledon, b. IM embryo, c. Mature embryo, d. Leaf	Somatic embryo	Sagare et al., 1999
3. PG-5 PG-12 C-235	MS	2, 4, 5 -T, 2, 4 -D	Mature embryo	Plants developed from somatic embryo	Sagare et al., 1993
4. PG12, C-235	MS	2, 4, 5 -T, 2, 4 -D, BAP	Immature-cotyledon	Plants developed from somatic embryos	Sagare et al., 1993
5. PG12	MS	2, 4, 5 -T	Mature embryo axis	Plants developed from somatic embryos	Suhasini et al., 1994
6. Kabuli	MS salts + B <sub>5</sub> Vitamins	TDZ , L-Proline	Mature seeds	Somatic embryos	Murthy et al., 1996
<b>Somatic Embryogenesis via Callus</b>					
C235, K-850, BG-256	MS MS + B <sub>5</sub>	2, 4 -D, NAA, IBA, BAP, Kinetin,	Embryonal axis	Plants developed from somatic embryos	Chauhan et al., 2002
P-256	MS	TDZ, Proline	Whole Seedling (cotyledonary notch region)	Plants developed from somatic embryos	Rizvi et al., 2002
6153, CM72	MS	NAA, BAP, 2, 4 -D, Kinetin, IAA, IBA	Hypocotyls	Somatic embryo	Hussain et al., 2000
PG12	MS	2, 4, 5 -T	Mature embryo	Somatic embryo	Sagare et al., 1999
ICCV-4918	B <sub>5</sub>	2, 4, 5 -T, BAP	Immature cotyledons	Somatic embryos	Ramana et al., 1996
BG256	MS, B <sub>5</sub>	2, 4 -D, BAP, Picloram, IAA	Leaf	Plant from somatic embryo	kumar et al., 1995
C-235, JG-262 P-144, P-209	MS	2, 4 -D, Kinetin	Leaf	Plants developed from somatic embryos	Kumar et al., 1994
C-235	MS, B <sub>5</sub> , MS salt+B <sub>5</sub> vitamins	2, 4 -D, Kinetin, BAP, IBA	Leaf	Plants developed from somatic embryos	Kumar et al., 1994
Phule-G5	MS	2, 4 -D, Picloram, Kinetin	a. Root, b. cotyledonary nodes, c. epicotyl, d. nodes, internodes and leaf	Somatic embryo	Shanker and Mohanram, 1993
JG-62, Gaurav, Annger, BG-267 C-235	MS	2, 4 -D, BAP, Kinetin, TDZ	Immature leaflet	Plants developed from somatic embryos	Barna and Wakhlu, 1993
BG- 256	MS	2, 4 -D, BAP	Leaflet	Multiple shots/somatic embryos	Rao and Chopra VL, 1989



**Figure 1.** A: Seeds de-coated after 14 h imbibition, on shoot induction medium (SIM) (MS medium supplemented with 4  $\mu\text{M}$ , 10  $\mu\text{M}$  2-iP and 2  $\mu\text{M}$  Kin); B: Chickpea seedlings (6 d after incubation of de-coated seeds on SIM) used for obtaining explants for regeneration and transformation; D & E: Swollen embryonal axis attached to cotyledon showing the initiation of shoot induction {6 - 8 days after incubating the explants on SIM}; D: Two weeks old culture of embryonal axis with single cotyledon, showing the emergence of large number of adventitious buds {15 days after incubating 6 - 8 days old cultures from SIM on MS medium}.


(Fontana et al., 1993) or on medium containing MS macro 4 X MS micro, B<sub>5</sub> vitamins, BAP (3 mg l<sup>-1</sup>) and NAA (0.004 mg l<sup>-1</sup>) (Kar et al., 1996, 1997). C-235 and BG-329 seedling explant were used in MS salt + B<sub>5</sub> vitamins + 5  $\mu\text{M}$  BAP and checked for the effect of BAP induction period on shoot differentiation. With increase in the duration of culture, there was increase in the number of shoot buds with a simultaneous decrease in further elongation in the elongation medium (Paul et al., 2000). Direct regeneration from mature and immature embryo as explants led to maximum regeneration frequency as observed from mature embryo in MS salts + B<sub>5</sub> vitamins + 4% sucrose which was further elongated on MS + 0.6  $\mu\text{M}$  IBA + 8.8  $\mu\text{M}$  BAP + 4% sucrose (95% regeneration) followed by 50% on MS + 0.6  $\mu\text{M}$  IBA + 4.4  $\mu\text{M}$  BAP + 4% sucrose. Immature embryo regenerated better in liquid medium using filter paper bridge when compared to solid medium. Regeneration frequency of 52% was observed in liquid medium (MS + 2.8  $\mu\text{M}$  IAA + 2.3  $\mu\text{M}$ ) (Chauhan et al., 2003). Over 100 multiple shoots were observed in embryonal axis explants from seedling (C-235) in induction medium (MS + 4  $\mu\text{M}$  TDZ + 10  $\mu\text{M}$  2-iP + 2  $\mu\text{M}$  kinetin, pH 5 - 5.5) for about 6 - 7 weeks, multiple shoots were elongated on MS containing 5  $\mu\text{M}$  2-iP + 2  $\mu\text{M}$  kinetin, pH 5 - 5.5 (SEM1), non elongated shoots on SEM1 were transferred on SEM2 (MS + 2  $\mu\text{M}$  GA<sub>3</sub>, pH 5-5.5) (Jayanand et al., 2003; Sharma et al., 2007). Maximum 26 shoots were induced from single cotyledon with half embryonal axis in MS medium supplemented with 1.6 mg l<sup>-1</sup> BA, 0.04 mg l<sup>-1</sup> NAA and shoots elongate in the

presence of 0.2 mg l<sup>-1</sup> IAA (Chakrabarti et al., 2006); > 100 shoots per explant (single cotyledon with half embryonal axis) in MS medium supplemented with 4  $\mu\text{M}$ , 10  $\mu\text{M}$  2-iP and 2  $\mu\text{M}$  Kin (Anwar, 2007; Anwar et al., 2008, 2009) (Figure 1C).

Explants consisting of single cotyledon with half embryonal axis obtained from 6 day old seedlings (Figure 1B) on MS medium supplemented with 4  $\mu\text{M}$  TDZ, 2  $\mu\text{M}$  kinetin and 10  $\mu\text{M}$  2-iP was found to be good for producing optimal shoot/adventitious buds from the explants. Embryonal axis attached to cotyledon showed significant swelling and exhibited initiation of shoot induction within a duration of 6 - 8 days (Figures 1C - E). Use of TDZ with purine ring containing cytokinins such as kinetin, BAP and N6-[2-Isopentyl]adenine (2-iP) has been shown to promote the formation of a large number of healthy shoots (Eisinger, 1983; Mroginski and Kartha, 1984; Radhika et al., 2006). In order to avoid negative impact of TDZ in the formation of shoots, it was found wise to transfer the explants immediately after shoot induction event to a medium devoid of TDZ. Cytokinin such as 2-iP and kinetin are well known to promote rapid shoot multiplication (Jayanand et al., 2003; Kiran et al., 2005). Even in this investigation, TDZ at 4  $\mu\text{M}$  in combination with 2-iP (10  $\mu\text{M}$ ) and kinetin (2  $\mu\text{M}$ ) was found to be optimal for the expansion of meristematic zone followed by shoot induction. Subsequent to the transfer of the cultures onto MS basal medium that is growth regulator free medium, emergence of a large number of adventitious shoots/ buds from all over the surface of the swollen embryonal axis was recorded within 7 - 15 days. In general, 50 - 100 shoots arose from each explant within a time period of 15 days (Figures 1F - G). The induced multiple shoot/adventitious buds were excised from the bunch, without any callus or globular structures and cultured on the shoot elongation medium (SEM) consisting of MS medium supplemented with 5  $\mu\text{M}$  2-iP and 2  $\mu\text{M}$  kinetin for 10 days. They were then routinely sub-cultured at an interval of 10 - 15 days on SEM (Figure 2).

Gibberellins promote elongation of shoots (Jayanand et al., 2003; Sharma et al., 2007), but during the present investigation, it was realized that the shoots transferred to plant growth regulator free medium have potential to synthesize and maintain desired endogenous levels of gibberellins and other auxins. Although, elongation of shoots was promoted on MS medium supplemented with various cytokinins in combination with GA<sub>3</sub>, stronger/ healthy elongated shoots were obtained on MS medium supplemented with kinetin and 2-iP in the absence of GA<sub>3</sub>. GA<sub>3</sub> promoted elongation of the shoots but the shoots were weak with more inter-nodal elongation. Therefore, for subsequent experiments, shoot elongation was achieved on MS medium supplemented with kinetin in combination with 2-iP.

In general, it is widely accepted that apical meristems are strong zones for synthesis of auxins (IAA). Therefore, exogenous application is often found to be deleterious.



**Figure 2.** Shoots on shoot elongation medium (MS medium with 5  $\mu\text{M}$  2-iP and 2  $\mu\text{M}$  kinetin) (10-14 d of different stage after transfer of shoots from SIM).

IAA is photosensitive and gets degraded under light. Interestingly, Chakraborti et al. (2006) have reported that IAA at low concentration can bring about shoot elongation. Based on the findings of the present investigation it can be authentically stated that exogenous application of neither auxins nor gibberellic acid is necessary and should not be encouraged. Therefore, the use of IAA in the medium for the elongation of shoot is not advisable.

#### **Indirect shoot organogenesis via callus from different explants of chickpea**

Regeneration of shoots via callus phase has been achieved from whole cotyledon and shoot tip explants by culturing on  $B_5$  supplemented with BAP (1  $\text{mg l}^{-1}$ ) and NAA (1  $\text{mg l}^{-1}$ ) (Prakash et al., 1992). Multiple shoots were induced from calli derived from hypocotyls cultured on  $B_5$  + 2% sucrose + BAP (1  $\text{mg l}^{-1}$ ) + Kinetin (1  $\text{mg l}^{-1}$ ) + IAA (0.5  $\text{mg l}^{-1}$ ) and subsequently transferred to MS + 2% sucrose + kinetin (2  $\text{mg l}^{-1}$ ) + NAA (2  $\text{mg l}^{-1}$ ) (Neelam et al., 1986b, 1986c). Induction of multiple shoots from cotyledonary node explants cultured on  $B_5$  medium containing BAP (0.5 to 1  $\text{mg l}^{-1}$ ) or NAA (0.5  $\text{mg l}^{-1}$ ) has been reported (Khan and Ghosh, 1984). Callus derived from immature leaflets cultured on MS + 2, 4 -D (25  $\mu\text{M}$ ) or MS + 2, 4 -D (2, 4 -Dichlorophenoxyacetic acid) (5  $\mu\text{M}$ ) gave rise to multiple shoots on subsequent transfer to MS + BAP (10  $\mu\text{M}$ ) (Barna and Wakhlu, 1994). Immature leaflet explants of chickpea (P-256) developed callus on MS medium supplemented with 1.25  $\mu\text{M}$  2, 4 -D and 10  $\mu\text{M}$  BAP and calli were regenerated on MS + 5  $\mu\text{M}$  BAP + 10  $\mu\text{M}$  TDZ + 0.1  $\mu\text{M}$  IBA. The addition of spermidine enhanced the number of shoots by 73.8% (Rizvi and Singh, 2000). Up to date review of direct and indirect shoot organogenesis from different cultivar and explants of chickpea are listed in Table 1.

#### **Somatic embryogenesis**

Somatic embryogenesis involves *in vitro* formation of embryos from somatic tissues cultured on media supplemented with various hormones. These embryos develop from single cells and pass through globular, heart shaped, torpedo shaped, nodulated and cotyledonary embryo stages to give rise to complete plantlets. The whole process may proceed on a single medium or may require medium alterations specific for a specific developmental stage. Direct callus somatic embryogenesis and plant development in various genotypes of chickpea are listed in Table 2.

Somatic embryogenesis in chickpea has been involved directly or through an intervening callus phase in different explants that were recalcitrant. Some report on somatic embryogenesis has been reported. Direct somatic embryogenesis was achieved by culturing, mature seeds on MS salts +  $B_5$  vitamins + TDZ (10  $\mu\text{M}$ ) + proline (1  $\text{mM}$ ) (Murthy et al., 1996). Immature cotyledons gave rise to somatic embryos when cultured on  $B_5$  medium supplemented with 2, 4, 5 -T (2, 4, 5 -Trichlorophenoxyacetic acid) (7.8  $\mu\text{M}$ ) + BAP (4.4  $\mu\text{M}$ ) (Ramana et al., 1996). Somatic embryos obtained by culturing immature cotyledons and embryo axes on MS + 2, 4, 5 -T (3.0  $\text{mg l}^{-1}$ ), gave rise to complete plantlets on subsequent transfer to half strength MS medium supplemented with zeatin (1  $\text{mg l}^{-1}$ ) (Sagare et al., 1993). Mature embryo axes cultured on solid or liquid MS medium supplemented with 2, 4, 5 -T (3  $\text{mg l}^{-1}$ ), directly induced somatic embryos which were transferred to half strength MS medium containing 0.1  $\text{mg l}^{-1}$  abscisic acid from maturation and subsequently sub cultured on half strength MS medium containing zeatin (1  $\text{mg l}^{-1}$ ) (Suhasini et al., 1994). Mature embryo axis of chickpea cultivar PG-12 was cultured on MS supplemented with 3  $\text{mg l}^{-1}$  2, 4, 5 -T (Sagare et al., 1999).

Complete plantlets have been regenerated through somatic embryogenesis via callus by culturing mature leaflets on MS + 2, 4 -D (0.5  $\text{mg l}^{-1}$ ) + BAP (0.5  $\text{mg l}^{-1}$ ) in dark and subsequent transfer to MS + IAA (0.1  $\text{mg l}^{-1}$ ) + BAP (3  $\text{mg l}^{-1}$ ) (Rao and Chopra, 1989) or immature leaflets on MS + 2, 4 -D (25  $\mu\text{M}$ ) during induction. MS + ABA during maturation and MS +  $\text{GA}_3$  (15  $\mu\text{M}$ ) + IBA (1  $\mu\text{M}$ ) during conversion (Barna and Wakhlu, 1993), gave a maximum number (17) of somatic embryos from callus with 15 *de novo* shoots from whole seedling (hypocotyledonary notch region) on MS + 10  $\mu\text{M}$  TDZ + 5  $\text{mM}$  proline (Rizvi et al., 2002). Creamy callus was observed on MS medium supplemented with 1.25  $\text{mg l}^{-1}$  2, 4 -D and 0.25  $\text{mg l}^{-1}$  kinetin and whitish callus was on MS + 0.5  $\text{mg l}^{-1}$  2, 4 -D and 1.4  $\text{mg l}^{-1}$  2, 4 -D + 0.25  $\text{mg l}^{-1}$  kinetin. Medium with 1.25  $\text{mg l}^{-1}$  2, 4 -D and 0.25  $\text{mg l}^{-1}$  kinetin was found to be best for embryo induction and somatic embryos were best matured on MS salt +  $B_5$  vitamins + 2  $\text{mg l}^{-1}$  BAP + 1.25  $\text{mg l}^{-1}$  IBA for 4 weeks and elongated on MS. C-235 genotypes showed better maturation of somatic embryo followed by BG-256 and K-850.

Complete plants regeneration via somatic embryogenesis was carried out by culturing leaves on MS + 2, 4-D ( $1.25 \text{ mg l}^{-1}$ ) + Kinetin ( $0.25 \text{ mg l}^{-1}$ ) in the dark for embryogenesis and subsequently transferred to light on medium containing MS salts + B<sub>5</sub> vitamins + IBA ( $0.125 \text{ mg l}^{-1}$ ) + BAP ( $2 \text{ mg l}^{-1}$ ) for maturation, and then to B<sub>5</sub> + BAP ( $0.25 \text{ mg l}^{-1}$ ) during conversion and B<sub>5</sub> basal for plantlet formation (Kumar et al., 1994). The same protocol was repeated for 3 more cultivars (Kumar et al., 1995).

### Anther culture

Anthers or pollen grains on culture produce haploid plants, that is, plants with gametic chromosome number. These were particularly useful in plant breeding both for rapid production of homozygous lines following chromosome doubling to original ploidy level and for the detection and selection of recessive mutants. Another culture has also been reported (Gosal and Bajaj, 1979; Khan and Ghosh, 1983; Bajaj and Gosal, 1987; Gosal and Bajaj, 1988; Huda et al., 2001; Kennedy et al., 2002).

### Protoplast culture

Plant protoplasts (single cell system) culture has been exciting possibilities in the field of somatic cell genetics and crop improvement. Isolated protoplasts serve as the field of somatic cell cloning and development of mutant lines (Bhojawani and Razdan, 1996). Isolation and regeneration of chickpea protoplasts from hypocotyl derived protoplasts cultured on V47 medium supplemented with NAA ( $1.5 \text{ mg l}^{-1}$ ) and BAP ( $0.5 \text{ mg l}^{-1}$ ) were able to produce microcalli, but microcalli failed to undergo differentiation and organogenesis to produce plantlets (Sagare and Krishnamurthy, 1991).

### *In vitro* rooting and grafting of shoots

Several protocols of regeneration of chickpea have been suggested using shoot meristems through organogenesis and somatic embryogenesis but rooting in regenerated shoots was a problem. The shoots transferred to a medium containing auxin have been used for rooting. In a protocol provided by Prakash et al. (1992), shoots were transferred firstly to B<sub>5</sub> medium + gelling agent ( $8 \text{ g l}^{-1}$ ) + NAA ( $0.1 \text{ mg l}^{-1}$ ) and then to B<sub>5</sub> + gelling agent ( $8 \text{ g l}^{-1}$ ) + NAA ( $1 \text{ mg l}^{-1}$ ) and then B<sub>5</sub> + gelling agent ( $12 \text{ g l}^{-1}$ ) + NAA ( $2 \text{ mg l}^{-1}$ ). An increase in concentration of agar-agar (1.2%) and inclusion of activated charcoal (0.2%) in the second medium increased percentage of root differentiation. In another method suggested by Malik and Saxena (1992), shoots were cultured for 1 - 3 weeks on basal culture medium containing MS macro- and micro-

nutrients, B<sub>5</sub> vitamins, 3% sucrose and 0.25% gelrite, then transferred to basal medium containing  $2.5 \mu\text{M}$  NAA. *In vitro* regenerated shoots cultured on B<sub>5</sub> +  $0.5 \text{ mg l}^{-1}$  NAA also resulted in profuse rooting (Khan and Ghosh, 1984). Auxins, generally IBA were used in media for rooting. Shoots were cultured in a root induction medium having MS salts and vitamins and  $0.05 \text{ mg l}^{-1}$  IBA (Kar et al., 1996). B<sub>5</sub> medium containing  $1 \text{ mg l}^{-1}$  IAA in combination with  $0.05 \text{ mg l}^{-1}$  kinetin has also been used for root induction (Neelam et al., 1986).

The rooting can be facilitated by another method which gave good results with reduction in amount of salts used in the medium. Polisetty et al. (1996) used shoots (2 - 3 cm) which were initially cultured with  $\frac{1}{4}$ <sup>th</sup> strength MS + 0.75% sucrose + 0.8% agar without hormonal supplement for 30 days and rooting was achieved in 90 - 100% shoots.

Rooting and successful transplantation were major hurdle for chickpea plantlet formation. Therefore, drafting was carried according to the method described by Pickardt et al. (1995) for the establishment of putative transgenics (Krishnamurthy et al., 2000; Senthil et al., 2004; Sarmah et al., 2004; Singh et al., 2004; Sanyal et al., 2005) with minor modification, hence it could not be referred to as a true transgenic.

Regenerated shoots were rooted in MS solid medium containing  $2.5 \mu\text{M}$  IBA (Rizvi and Singh, 2000) and  $2.5 \mu\text{M}$  NAA (Rizvi et al., 2002). 93.3 and 80% rooting were recorded in  $\frac{1}{4}$ <sup>th</sup> MS +  $10.4 \mu\text{M}$  NAA + 2% sucrose and  $\frac{1}{4}$ <sup>th</sup> MS +  $10.7 \mu\text{M}$  NAA + 1% sucrose (Chauhan et al., 2003). Shoots were rooted on liquid medium (MS + 9.4 mM KNO<sub>3</sub> + 2% sucrose +  $5 \mu\text{M}$  IBA) for 2 weeks on filter paper bridge (60 - 70% rooted), un-rooted shoots were pulse treated with  $100 \mu\text{M}$  IBA and culture on filter paper bridge in liquid MS for 2 weeks (10 - 20% rooted) and the remaining were transferred to hydroponic system containing  $\frac{1}{4}$ <sup>th</sup> Arnon solution +  $3 \mu\text{M}$  IBA for 2 - 3 weeks (10-15% rooted). This is more time consuming, for growth chamber is required for rooting and maintenance (Jayanand et al., 2003).

Elongated putative transformed shoots were rooted and grafted; shoots were rooted according to Murfet (1971), Morton et al. (2001), for 1 week or shoots were dip in  $1 \text{ mg l}^{-1}$  IBA and cultured on MS medium supplemented with  $1 \text{ mg l}^{-1}$  IAA + 10 mM MES for 2 weeks (Sarmah et al., 2004). Elongated putative transformed shoots were rooted on B<sub>5</sub> +  $1 \mu\text{M}$  NAA + 150 mg/l Kan for 1 - 3 weeks (Polowick et al., 2004). Elongated putative transformed shoots were rooted on MB (MS salt + B<sub>5</sub> vitamins) +  $2.5 \mu\text{M}$  NAA for 3 - 4 weeks. Rooted shoot were transferred to  $\frac{1}{2}$  MB. Grafting also followed because of the inconsistent performance of *in vitro* rooted plants in the glass house and extended culture period. 65 - 75% of grafted shoots were established in glass house (Senthil et al., 2004). Selected shoots were rooted on MB +  $4.9 \mu\text{M}$  IBA +  $100 \text{ mg l}^{-1}$  kan +  $5 \text{ mg l}^{-1}$  PPT or 2 mM of each of lycine and threonine. Shoots (3 - 5 cm) were also grafted onto



**Figure 3.** Shoots exposed to 10 sec 100 mM IBA pulse treatment, on liquid MS medium A: 14 d; B: 24 d after transfer of the elongated shoots (please note strong root system).

10 - 12 days old seedlings (Singh et al., 2004).

Putative shoots were rooted on  $\frac{1}{2}$  MS + 0.5 mg l<sup>-1</sup> IBA + 0.02 g l<sup>-1</sup> AgNO<sub>3</sub> + 1% sucrose in 0.5% (w/v) agar. Micro-grafting of individual shoots were performed on 7 - 8 days old rootstock of same genotype prepared by transversely cutting the epicotyl region of the germinated seedling (Sanyal et al., 2005).

Thus, increasing the auxin concentration could induce rooting. Normal rooting could be induced only when the shoots were hard and not fragile and succulent or else, they tended to form callus. Roots developing from the shoots should arise directly from the base of the shoot and intermediary callus should be avoided as this does not have well defined vasculature (vascular connection between shoot and root). Tap root is preferred over other kinds of roots.

One of the major hurdles that limit *in vitro* regeneration of chickpea is the induction and development of strong root system. This compelled several researchers to adopt to grafting (Krishnamurthy et al., 2000; Sarmah et al., 2004; Senthil et al., 2004; Sanyal et al., 2005; Chakraborti et al., 2006). However, grafting is time consuming, requires special skills and the success rates vary significantly. Roots induced from cut ends of shoots of chickpea were shorter in length in semisolid medium as compared to liquid medium, similar to earlier reports (Jackson et al., 1991; Cournac et al., 1991; Ebrahim and Ibrahim 2000; Hazarika, 2006). The retardation in root length in the semisolid medium may be attributed to the relatively lower aeration in agar-gelled medium as indicated by earlier researchers (Hazarika, 2006; Pati et al., 2006; Rout et al., 2006).

Kat et al. (1996, 1997) rooted chickpea shoots in rooting medium consisting of MS salts, MS vitamins and 0.05 mg/l IBA and subsequently transferred the plantlets to half-strength MS medium containing 20 g/l sucrose.

Polowick et al. (2004) used rooting medium consisting of B5 basal salts and vitamins supplemented with 1  $\mu$ M NAA for rooting chickpea shoots and subsequently shoots with short roots which were transferred to Magenta vessels containing B5 salts and vitamins and 0.7% agar until root system was well established in 1 - 3 weeks while those with roots longer than 3 cm were transferred directly to soil. However, frequency of rooting was only 10 - 60% (Polowick et al., 2004). Although, Jayanand et al. (2003) reported high frequency rooting and transplantation success, the protocol adopted by them was time consuming involving three specific phases viz. i) rooting of shoots on liquid MS medium with 9.4 mM KNO<sub>3</sub>, 2% sucrose and 5  $\mu$ M IBA for 2 weeks on filter paper bridge, ii) pulse treatment of un-rooted shoots with 100  $\mu$ M IBA and culturing on filter paper bridges in liquid MS for 2 weeks and iii) transfer to hypotonic system containing  $\frac{1}{4}$ <sup>th</sup> Arnon solution with 3  $\mu$ M IBA for 2 - 3 weeks. Even during the present investigations, shoots could be rooted in MS medium with 5  $\mu$ M IBA, but the roots developed were relatively weaker and required 15 - 20 days. However, 5 - 10 s pulse treatment of cut ends of shoots of chickpea with 100 mM (that is, 100  $\mu$ mol/ml) IBA led to root induction within 4 days and subsequently resulted in the development of strong root system within 10 - 12 day in liquid MS basal medium (Figure 3) (Anwar et al., 2008, 2009). This much high concentration of IBA is 100% effective and first time reported in any crop (Anwar et al., 2008, 2009). This idea can be applicable in case of other crop from which *in vitro* rooting is difficult. *In vitro* chickpea rooting and grafting of different cultivar are listed in Table 3.

### Transplantation / establishment of plantlet

After *in vitro* rooting of shoots, the plantlets have to be established in soil and for this, various methods have been employed. Prior to transfer to soil, hardening of the plantlets was done. In case of plants regenerated from leaflet callus via somatic embryogenesis, complete plantlets in tubes were subjected to the following hardening procedure.

Cotton plugs were loosened and kept for 4 - 5 days prior to transferring to  $\frac{1}{4}$ <sup>th</sup> strength liquid MS medium without sucrose for 24 h and they were transferred to sterile soilrite (Kumar et al., 1994) or sterile vermiculite for one week (Barna and Wakhlu, 1993) and then transferred to pots in the ratio of 3:1:1; sterile soil: farmyard manure: sand (Kumar et al., 1994) or to pots containing garden soil (Barna and Wakhlu, 1993).

Plantlets regenerated via direct somatic embryogenesis induced from immature cotyledons and immature embryo axes were established in the soil of green house containing a mixture in the ratio of 1:1:1; peat: vermiculite: soil (Sagare et al., 1993).

Plantlets regenerated from callus cultures of hypocotyls were established in plastic pots containing vermiculite

**Table 3.** Current status of chickpea (*Cicer arietinum* L.) transformation.

Cultivar	Explant	<i>Agrobacterium</i> strain/gene/vector	Mode of transformation /selection	Expression of transgene/s	Rooting/Grafting	Phenotype obtained	References
ICC10943 ICC10386	Embryo axis	LBA4404/ pCAMBIA1305.2/p3 5SGUS	<i>Agrobacterium</i> / <i>Hyg</i>	PCR ( <i>gus</i> ), southern ( <i>gus</i> )	MSB5 + 0.9 $\mu$ M NAA + 1.2 $\mu$ M IAA + 0.9 $\mu$ M Kin	Fertile transgenic plants	Pathak and Hamzah, 2008
Gokce, Er, Akcin, Uzunlu, Kusmen	Seeds	C58C1/EHA101/pTJ K136/pTHW136/uid A/nptII	<i>Agrobacterium</i> / <i>Kan</i>	PCR ( <i>nptII</i> ), <i>gus</i> assay	-	-T <sub>0</sub> Direct <i>gus</i> assay and <i>nptII</i> PCR	Akbulut et al., 2008
ICCC37, PR-12	Epicotyls	<i>pHS 102/gus</i>	Particle bombardment/ <i>Kan</i>	PCR, Southern, <i>gus</i> assay of T <sub>0</sub> plants	-	To plants	Shivani et al., 2007
C-235, BG-256, P-362, P-372	Cotyledonary nodes (20d old seedling)	LBA 4404/ <i>cry1Ac</i> / <i>p35SGUS-INT</i>	<i>Agrobacterium</i> / <i>Kan</i>	PCR ( <i>nptII</i> and <i>cry1Ac</i> ), southern ( <i>nptII</i> and <i>cry1Ac</i> ) ELISA ( <i>cry1Ac</i> ), Insect bioassay	Grafting $\frac{1}{2}$ MS + 0.5 mg/l IBA + 0.02 g l <sup>-1</sup> Ag NO <sub>3</sub> + 1% sucrose	Fertile transgenic plants showed affective against Pod Borer, 14.5-23.5 ng/mg <i>cry1Ac</i> protein accumulation. ....	Sanyal et al., 2005
P-362, P-1042 P-1043	Decapitated embryo with one cotyledon (overnight socked seed)	EHA101/AK/ <i>pIBGUS</i> / <i>pBI121-dsAK</i>	<i>Agrobacterium</i> / <i>Kan</i> Particle bombardment/ <i>Kan</i>	<i>Gus</i> , LP assay, PCR ( <i>gus</i> ), southern ( <i>nptII</i> probe), <i>AR</i> activity	Grafted MS+B5 org. + 4.9 $\mu$ M IBA + 1.5% sucrose	Fertile transgenic plants; 1 <sup>st</sup> chickpea transgenic report of AK/LT selection system, over expression of lysine and threonine...	Singh et al., 2004
ICCV5, H208, ICCL87322, K850 Annigeri	Slices from plumule (2-5h socked seed)	<i>AGL1/uidA, bar</i> / <i>pGIN1</i> or <i>pGIP</i>	<i>Agrobacterium</i> / <i>Kan, PPT</i>	PCR ( <i>gus</i> ), <i>gus</i> and LP assay, southern ( <i>gus</i> )	MS+B5 org. + 2.5 $\mu$ M NAA	Fertile transgenic plants; <i>gus</i> and LP assay confirmed upto T2 generation. ....	Senthil et al., 2004
CDC Yuma- kabuli	Longitudinal slide of embryonal axis	LBG66/ <i>uidA</i> , <i>nptII/pPBI3010</i>	<i>Agrobacterium</i> / <i>Kan</i>	PCR ( <i>gus</i> and <i>nptII</i> ), <i>gus</i> assay, southern ( <i>gus</i> )	B5 + 0.18 mg l <sup>-1</sup> NAA	Fertile transgenic plants; <i>gus</i> expression upto 4 <sup>th</sup> generation....	Polowick et al., 2004
Semsen	Half embryonal axis with one cotyledon	<i>AGL1/<math>\alpha</math>-A1</i> / <i>pRM50</i>	<i>Agrobacterium</i> / <i>Kan</i>	PCR ( <i>nptII</i> ), <i>nptII</i> dot blot, southern ( $\alpha$ -A1 probe), insect bioassay and $\alpha$ -amylase activity	Grafting MS + 1 mg l <sup>-1</sup> IBA	Fertile T1 and T2 transgenic, $\alpha$ -A1 strongly inhibited the development of insect. ...	Sarmah et al., 2004

and soil mixture (Neelam et al., 1986) and plantlets derived from callus culture of immature leaflets were established in pure vermiculite (Barna and Waklu, 1994). Plantlets regenerated

from embryonal axis were transferred to 8 cm (diameter) pots and the plants were covered with polypropylene bags and gradually open after 7-10 days period or the rooted shoots were suspended

in Magenta Jars containing  $\frac{1}{4}$ <sup>th</sup> Arnon solutions and in the second stage, the hardened plants were transferred to 20 cm (diameter) pots containing 3:2; sand: black soil + Cell Rich (5%) +

Table 3. Contd.

Cultivar	Explant	<i>Agrobacterium</i> strain/gene/vector	Mode of transformation /selection	Expression of transgene/s	Rooting/ Grafting	Phenotype obtained	References
Gokce, Akcin91, Izmir92	One week old seedlings	<i>A. rhizogenes</i> pRi15834	<i>Agrobacterium</i>	Wild type	-	Callogenic mass	Khawar and Ozcan, 2004
PG1, PG12 Chafa	Embryo (from overnight soaked seed)	*C58C1/uidA, nptII/p35SGUSINT *GV2260/uidA, nptII/p35SGUSINT *EHA101/uidA, nptII/pBGUS	<i>Agrobacterium</i> / Kan	PCR (nptII), gus assay, southern (nptII probe)	Grafting	Four t1 transgenic plants shown gus only not nptII amplicon. ....	Krishnamurthy et al., 2000
6153, CM72	Hypocotyl	nptII/pBI121/ gus	Particle bombardment/Kan	Gus assay	-	Somatic embryo	Hussai et al., 2000
Red chickpea Canitez 87 MB-10	Shoot primordial of mature embryo (from 24h seedlings)	<i>A. tumefaciens</i> 4404/uidA, nptII/pBI121 <i>A. rhizogenes</i> 9402/uidA, nptII/pBI-121	<i>Agrobacterium</i> / Kan	Southern (gus)	-	Hairy root . ...	Altinkut et al., 1997
ICCV1, ICCV6	Embryonal axes without root and shoot meristem	<i>Cry1Ac</i> /pcrylAc BDC2	Particle bombardment/ Kan	Positive (crylAc), Southern blot (crylAc) and insect bioassay	MS + 0.05 mg l <sup>-1</sup> IBA	Cry1Ac transgenic chickpea. ....	Kar et al., 1997
ICCV-1 ICCV-6 Desi (Local Variety)	Embryonal axes without root and shoot meristem	LBA 4404/uidA, nptII/pBI121	<i>Agrobacterium</i> / Kan	PCR (nptII), gus assay, Southern (nptII)	MS + 0.05 mg l <sup>-1</sup> IBA	Transgenic plantlets transplanted to greenhouse. ....	Kar et al., 1996
ICC-4918	Immature cotyledon	LBA 4404/uidA/pBin19,GUS int)	<i>Agrobacterium</i>	gus assay	1 mg/l Zeatin	Plantlet. . ....	Ramana et al., 1996
Local ecotype (Italy)	Embryonal axis lacking apical meristem	LBA 4404/uid, npt/pBI121	<i>Agrobacterium</i> / Kan	nptII dot blot, Southern (nptII)	0.5 mg l <sup>-1</sup> IAA + 0.05 mg l <sup>-1</sup> Kin	Transgene expression of plantlet	Fontanna et al. 1993
Pusa-256	Leaf and stem explants	R1601/nptII/pTVK291	<i>Agrobacterium</i> / Kan	Southern (nptII probe)	-	Formation of transformed callus	Srinivasan et al., 1991

rice straw compost (5%). Stage 1 and 2 required a day and night temperature of  $20 \pm 2$  and  $15 \pm 1^\circ\text{C}$ , respectively, and 50% humidity during day time and 80% at night and 10,000 - 12,000 lux light intensity during hardening process. This is more time consuming and for maintenance, growth chamber will be required (Jayanand et al., 2003).

Plantlets derived from direct regeneration of shoot tip explants where cultured on 1/4<sup>th</sup> strength MS without agar for elicitation of root growth. Root portion was then washed with 0.1% mercuric chloride and tap water.

These plantlets were transferred to 1/4<sup>th</sup> strength Hoagland solution for 20 days and then established in vermiculite (Polisetty et al., 1996). Rooting was a major hurdle for chickpea plantlet formation, therefore, drafting was carried out according to method described by Pickardt et al. (1995) for the establishment of putative transgenics (Krishnamurthy et al., 2000; Senthil et al., 2004; Sarmah et al., 2004; Sanyal et al., 2005) with minor modification, hence we could not say it was true transgenic.

*In vitro* rooted shoots were transferred to plastic pots



**Figure 4.** Sand and Potting mixture used for planting *in vitro* raised plantlets, respectively. Plantlets transferred to small earthen pots with a mixture of garden soil, sand and bio-manure in equal proportion. Freshly transferred plantlets covered with polythene bags.

containing soil mixture (Sunshine No. 4. Sun Gro Horticulture, Bellevue, Wash.), 20/15°C (day/night) and 16/8 h (light/dark) photo period with 200  $\mu$  mol quanta/m<sup>2</sup>/s, established plants were transferred into bigger pots of same soil mixture and grown under greenhouse, 16/8 h (light/dark) photo period. Young plantlets were covered with transparent beakers to maintain humidity for 3 - 4 days and slowly acclimatized (Polowick et al., 2004). Sarmah et al. (2004) reported that *in vitro* shoot were grafted and grown under glass house condition. 65-75% of grafted shoots were established in glass house (Senthil et al., 2004). *In vitro* rooted and grafted shoots were transferred to pots containing soil: vermiculite; 1:1 and covered with transparent plastic bags. A cut was gradually extended over a 3 week period until the bag was completely open. After 15 - 20 days of transplantation, the plants were transferred to glass house (Singh et al., 2004).

Micrografting of individual shoots were performed on 7 - 8 days old rootstock of same genotype prepared by transversely cutting the epicotyl region of the germinated seedling. The scion of putative shoots was inserted into vertical incision made in the epicotyl region of the stock. The resulting grafts were initially wrapped with micropore tape and cultured in pots containing soilrite that were irrigated with  $\frac{1}{2} \times$  MS for 15 days in culture room and then transferred into the pots containing soilrite: loam: leaf manure; 1:1:1 in green house maintained at 22 - 24°C, 60 - 80% humidity and 14 h photoperiod (Sanyal et al., 2005).

Another major hurdle limiting chickpea regeneration is the establishment of *in vitro* raised plantlets in pots/field. This is another major reason behind preference for graf-

ting. The plantlets with shorter length have higher potential to withstand transplantation shock and establish better in pots. It has been reported earlier that the root to shoot ratio has an important role in the successful establishment of *in vitro* raised plantlets (Subhan et al., 1998). Understandably, shorter shoot area will reduce excessive loss of water leading to rapid loss in turgidity of the plantlets (Munns, 2002). In the present investigations, plantlets with shoot length of 3 to 5 cm survived better than the plantlets with longer shoot length. Many of the plantlets with longer shoot length failed to establish even if the shoots were stouter, as they tend to collapse/ lodge and die within few days after transplantation to pots.

For transplantation of plantlets, various potting mixtures were used. These include autoclaved/sterile (Indurker et al., 2007) and non-autoclaved/non-sterile soil, soilrite, manure, vermiculite, soilrite + garden soil (in the ratio of 1:1), soilrite + garden soil + manure (in the ratio of 1:1:1), commercial soil mixture (Sunshine No. 4, Sun Gro Horticulture, Bellevue) (Polowick et al., 2004). In all these cases, the successful establishment rate was very low. Failure was accounted in these particular cases of excess moisture and high degree of water holding capacity of potting mixture. In the present investigations, potting mixture consisting of garden soil mixed with sand (gravel) and bio-manure (Khadi and Village Commission, Govt. of India) in the ratio of 1:1:1 is most suitable for achieving percent transplantation success. Chickpea is well known to be susceptible to flooding and excess moisture (Yadav et al., 2006). Under high moisture condition, chickpea is prone to fungus and wilt diseases (Yadav et al., 2006). During the present investigations, autoclaved soil mixture was not found in any way to be an essential requirement for the establishment of *in vitro* raised plantlets of chickpea. Instead it was realized that improving aeration/porosity of potting mixture by mixing sand is highly beneficial. Chickpea grows best on fertile sandy, loam soils with good internal drainage (Yadav et al., 2006). Good drainage is necessary because even short period of flooding, water logging soil and moisture reduces growth and increases susceptibility to root and stem rots and heavy rainfall season shows reduced yields due to disease outbreaks and stem lodging problem from excessive vegetative growth (Yadav et al., 2006). Some of the stages involved in transplantation of *in vitro* raised plantlets and their successful establishment in smaller and larger earthen pots is shown in Figures 4 - 8.

Another important factor that often limits regeneration and successful establishment of *in vitro* raised plantlets of chickpea is the season (that is, month of the year) when these are attempted. Best and percentage establishment success followed by good seed set is achieved when the *in vitro* raised plantlets of chickpea are transplanted during the months of October and November (that is, beginning of winter). Over all, vegetative growth including number of branches followed by flowering, pod setting



**Figure 5.** Acclimatized plantlets.



**Figure 6.** Plants of chickpea established in large earthen pots with a mixture of garden soil, sand and bio-manure in equal proportion.



**Figure 7.** Plants of chickpea established in large earthen pots with a mixture of garden soil, sand and bio-manure in equal proportion.



**Figure 8.** Plants of chickpea established in large earthen pots with a mixture of garden soil, sand and bio-manure in equal proportion.



**Figure 9.** Plants showing pods and seed setting.



**Figure 10.** Harvesting stage of seeds after drying of pods and plants.

and seed production (Figures 9 - 10) (Anwar et al., 2008, 2009) was found to be superior in the plants obtained from the plantlets transferred to pots during October and

November.

In summary, a simple and perfect protocol for successful rooting of shoots and establishment of *in vitro* raised plantlets of chickpea has been developed. Successful establishment of *in vitro* raised well rooted plantlets depends upon two basic factors viz., (i) shoot length of the plantlets and root to shoot ratio and (ii) potting-mixture with good aeration and lesser capacity to retain water. During the investigations, the percentage of plantlets with shoot length of 3 to 5 cm potted in garden soil mixed with sand (gravel) and bio manure in equal proportion, established perfectly and set seed (Anwar et al. 2008, 2009) (Figure 9).

## GENETIC TRANSFORMATION OF CHICKPEA

Plant genetic manipulation is the insertion of functional/ desired DNA directly into the genome of plant cell and the regeneration of whole plants from such cells by plant tissue culture techniques, molecular biology and microbiological methods which allows a chosen plant variety to be modified in a small but highly specific manner (Fenning and Gartland, 1995). Up to date genetic transformation of chickpea has been listed in Table 3. Genetic transformation of plants can be carried out using any one of several methods available. Indirect methods involve the use of a vector, which may be a bacterium; *Agrobacterium* or a virus; CaMV or Gemini virus. Direct methods of gene transfer are particle bombardment, electroporation, PEG mediated DNA uptake and microinjection (Potrykus, 1990).

Genetic transformations for incorporation of foreign traits have been achieved in some grain legumes. Success has also been achieved in the transfer of agronomically important traits to improve plants in terms of quantity and quality. But there are very few reports on agronomically important gene transfer in chickpea (Table 3). Both particle inflow gun (PIG) and *Agrobacterium* mediated gene transfer have been employed for engineering chickpea (Srinivasan et al., 1991; Fontanna et al., 1993; Seifkes-Boer et al., 1995; Kar et al., 1996; Kar et al., 1997; Husnain et al., 1997; Krishnamurthy et al., 2000; Sarmah et al., 2004; Singh et al., 2004; Senthil et al., 2004; Polowick et al., 2004; Sanyal et al., 2005; Jiang et al., 2005; Akbulut et al., 2008; Pathak and Hamzah, 2008).

Transgenic callus were raised through inoculating 12-15 days old leaf and stem explants with wild as well as disarmed strains of *Agrobacterium tumefaciens* (Srinivasan et al., 1991). Regeneration of transformed chickpea plants was done using disarmed *A. tumefaciens* strain LBA4404 containing *pBI121* carrying *nptII* and *gus* genes (Fontanna et al., 1993). 24 h old embryonic axis deprived of shoot apex were inoculated with *Agrobacterium* ( $5 \times 10^8$  cells  $\text{ml}^{-1}$ ) for 20 min and cultured on MS + kinetin ( $1 \text{ mg l}^{-1}$ ). Shoots arising from epicotyl region of embryo axes were selected on medium containing Kanamycin ( $50 \text{ mg l}^{-1}$ ) and *gus* expression was observed in leaves and roots of transformed plants. Kar et al. (1996) also used LBA 4404 strain to inoculate embryo axes deprived of root and shoot meristems and obtained multiple shoots by culturing explants on a medium containing MS macro, 4x MS micro, B<sub>5</sub> vitamins, BAP ( $3 \text{ mg l}^{-1}$ ), NAA ( $0.04 \text{ mg l}^{-1}$ ) and 3% (W/V) sucrose. Putative transformed shoots were selected on Kanamycin ( $50 \text{ mg l}^{-1}$ ) containing medium and rooted on MS supplemented with IBA ( $0.05 \text{ mg l}^{-1}$ ). Leaf and roots of transformants were positive for *gus* expression and southern blot for *nptII* gene.

Several groups have reported *Agrobacterium rhizogenes* mediated transformation systems for some

chickpea cultivars (Srinivas et al., 1991; Prakash et al., 1992b; Seifkes-Boer et al., 1995; Altinkut et al., 1997; Khawar and Ozcan, 2004). Hairy roots have been induced in 4 chickpea cultivars by infection with 7 different wild strains of *A. rhizogenes* (Seifkes-Boer et al., 1995). Wild type *A. rhizogenes* by direct inoculation showed a callogenic mass instead of hairy root (Khawar and Ozcan, 2004). Expression of marker genes *nptII* and *gus* was studied employing particle inflow gun (PIG) and *Agrobacterium* to effect gene transfer into zygotic embryos of chickpea (Husnain et al., 1997). Efficiency of transient expression of marker genes by PIG was shown to be dependent on DNA to tungsten ratio, distance between the filter disk and plant tissue and use of a nylon screen. *GUS* under control of Actin promoter gave better results as compared under CaMV and Win promoters. *Agrobacterium* strain A 281 was found to be more virulent than C58. Use of PIG in combination with *Agrobacterium* has been suggested for chickpea transformation.

An agronomically important *cryIAC* gene conferring resistance towards pod borer *Heliothis armigera* and *Helicoverpa armigera* has been reported to be transferred to chickpea by Kar et al. (1997), Sanyal et al. (2005) and Shivani et al. (2007). Embryo or embryonal axes deprived of root and shoot apices were co-transformed with *cryIAC* and *nptII* gene using a biolistic 1000/He particle gun. Explants gave rise to transformed multiple shoots on medium containing MS macro, 4X MS micro, B<sub>5</sub> vitamins,  $3 \text{ mg l}^{-1}$  BAP,  $0.004 \text{ mg l}^{-1}$  NAA, 3% w/v sucrose and  $50 \text{ mg l}^{-1}$  Kanamycin. molecular analyses by Southern and Northern blots revealed the presence of *cryIAC* gene and its expression was confirmed by inhibition or larval development on feeding transgenic shoots. The co-transformation frequency of *nptII* and *cryIAC* was estimated to be 45.8%. Inheritance of transgene/s was shown by PCR of T<sub>1</sub> (first filia) plants. Even the insect bioassay for pest resistance has been performed using stem of the plant instead of pods/seeds (Kar et al., 1997). Embryonal axis explants were used from four accessions of chickpea treated with *A. tumefaciens* strains C58C1/p35SGUSINT/GV2260 and pIBGUS/EHA101. For multiple shoot formation and elongation of shoots, MS medium supplemented with  $0.5 \text{ mg l}^{-1}$  Kanamycin or  $10 \text{ mg l}^{-1}$  phosphinothricin were selected. Putative transformed shoots were grafted according to Pickardt. T<sub>0</sub> (parent) plants showed PCR positive for both *nptII* and *GUS* and southern positive. Four T<sub>1</sub> plants showed PCR positive for *nptII* not for GUS (Krishnamurthy et al., 2000). It could be due to improper/unstable integration of T-DNA into the plant genome.

Longitudinal slices from embryonal axis of imbibed, mature seeds were used for *Agrobacterium* transformation (Schroeder et al. 1995) vector LBG66 (*pPBI3008*) containing binary vector *pPBI3010* (as described by Polowick et al., 2000). B<sub>5</sub> medium was used for co-cultivation and for induction as well as for selection B<sub>5</sub>

medium supplemented with 3 mg l<sup>-1</sup> BAP and 50 mg l<sup>-1</sup> kanamycin. For elongation of shoots, B<sub>5</sub> medium containing 1 mg l<sup>-1</sup> BAP and 50 and 75 mg l<sup>-1</sup> kanamycin for the two subsequent cycle and putative shoots were rooted in B<sub>5</sub> medium containing 0.18 mg l<sup>-1</sup> NAA and 150 mg l<sup>-1</sup> kanamycin. Putative transformed plantlets were established in green house. Molecular analyses of T<sub>0</sub> and T<sub>1</sub> by *GUS* activity, *MUG* assay and Southern blots revealed the presence of *npfl* gene, single insert which showed the 3:1 Mendelian inheritance pattern in T<sub>1</sub> population. Same patten of integration in T<sub>0</sub> and T<sub>1</sub> plants were found that confirmed that T-DNA was stably inherited to plant genome (Polowick et al., 2004). The transformation efficiency was almost 60% higher in the SAAT method than a simple *Agrobacterium* infection without sonication and sonicated wounding of the plant tissue and use of actively growing young decapitated embryo axis as an explant for *Agrobacterium* transformation play an important role in increasing the efficiency of transformation (Pathak and Hamzah, 2008).

In all the protocols, except that reported by Pathak and Hamzah (2008), the transformation frequency was very low. An efficient method to be employed for routine transformation experiments has been developed. Genetic transformations of a plant have to be achieved only if the transgene/s is stably inherited and expressed in subsequent progenies of plant. Only few report has been published on abiotic stress tolerance genetic transformation in chickpea. Hence, a lot still remains to be worked out in chickpea in order to get an economically superior germplasm with respect to the existing cultivated varieties.

## REFERENCES

- Akbulut M, Yucel M, Oktem HA (2008). Analysis and organization of DNA delivery into chickpea (*Cicer arietinum* L.) seedlings by *Agrobacterium tumefaciens*. *Afr. J. Biotech.* 7: 1011-1017.
- Altinkut A, Bajrovic K, Gozukirmizi N (1997a). High percentage of regeneration and transformation in chickpea. *Acta. Hort.* 447: 319-320.
- Altinkut A, Bajrovic K, Gozukirmizi N (1997b). Regeneration and hairy root formation of chickpea using callus derived plantlets and seedlings. *Int. Chickpea Pigeonpea Newslett.* 4: 30-31.
- Anwar F (2007). Genetic transformation of chickpea with bacterial *codA* gene enhancing drought tolerance. Ph.D. Thesis, University of Delhi, Delhi, India.
- Anwar F, Sharmila P, Pardha Saradhi P (2008). An ideal protocol for *in vitro* regeneration, efficient rooting and stable transplantation of chickpea. *Physiol. Mol. Biol. Plants*, 14: 329-335.
- Anwar F, Sharmila P, Pardha Saradhi P (2009). No more hurdle: *In vitro* chickpea rooting and cent-percent transplantation. *Australian journal of Basic and Applied Science*. Accepted for Publication.
- Bajaj YPS, Dhanju MJ (1979). Regeneration of plants from apical meristem tips of some legumes. *Curr. Sci.* 48: 906-907.
- Bajaj YPS, Gosal SS (1987). Pollen embryogenesis and chromosomal variation in cultured anthers of chickpea. *Int. Chickpea Newsl* 17: 12-13.
- Barna KS, Wakhlu AK (1993). Somatic embryogenesis and plant regeneration from callus cultures of chickpea (*Cicer arietinum* L.). *Plant Cell Rep.* 12: 521-524.
- Barna KS, Wakhlu AK (1994). Whole plant regeneration of *Cicer arietinum* from callus cultures via organogenesis. *Plant Cell Rep.* 13: 510-513.
- Bhojwani SS, Razdan MK (1996). Clonal Propagation. In: Bhojwani S.S. and Razdan M.K. (Eds.). *Plant Tissue Culture: Theory and Practice*, A Revised Edition. Elsevier, Amsterdam. pp. 483-536.
- Chakraborti D, Sarkar A, Das S (2006). Efficient and rapid *in vitro* plant regeneration system for Indian cultivars of chickpea (*Cicer arietinum* L.). *Plant cell tissue organ cult.* 86: 117-123.
- Chandra R, Chatrath A, Polisetty R, Khetarpal S (1993). Differentiation of *in vitro* grown explants of chickpea (*Cicer arietinum* L.). *Ind. J. Plant Physiol.* 36: 121-124.
- Chauhan R, Singh NP (2002). Plant regeneration via somatic embryogenesis in chickpea (*Cicer arietinum* L.). *Ind. J. Genet.* 62: 319-321.
- Chauhan R, Tiwari A, Singh NP (2003). Differential requirement of mature and immature embryo of chickpea (*Cicer arietinum* L.) for *in vitro* regeneration. *Ind. J. Plant Physiol.* 8: 28-33.
- Cournac L, Dimon B, Carrier P, Lahou A, Chagvardieff P (1991). Growth and photosynthetic characteristics of *Solanum tuberosum* plantlets cultivated *in vitro* in different conditions of aeration, source supply and CO<sub>2</sub> enrichment. *Plant Physiol.* 97: 112-117.
- Devlin RM, Zbiec II, Nowicka SE (1989). The effect of TDZ on some plant growth systems. *Proc. Plant Growth Regul. Soc.* 16: 99-103.
- Ebrahim MKH, Ibrahim AI (2000). Influence of medium solidification and pH value on *in vitro* propagation of *Maranta leuconeura* cv Kerchoviana. *Sci. Hortic.* 86: 211-221.
- Eisinger W (1983). Regulation of pea internode expansion by ethylene. *Ann. Rev. Plant Physiol.* 34: 225-240.
- Fenning TM, Gartland KMA (1995). Transformation protocols for broadleaved trees. *Meth. Mol. Biol.* 44: 149-166.
- Fontana GS, Santini L, Caretto S, Frugis G, Mariotti (1993). Genetic transformation in the grain legume *Cicer arietinum* L. *Plant Cell Rep.* 12: 194-198.
- Ganeshan S, Baga M, Harvey BL, Rosnagel BG, Scoles GJ, Chibbar RN (2003). Production of multiple shoots from thidiazuron-treated mature embryos and leaf-base/apical meristems of barley (*Hordeum vulgare*). *Plant cell tissue organ cult.* 73: 57-64.
- Gosal SS, Bajaj YPS (1988). Pollen embryogenesis and chromosomal variation in anther culture of three food legumes- *Cicer arietinum*, *Pisum sativum* and *Vigna mungo*. *SABRAO J.* 20: 51-58.
- Gosal SS, Bajaj YPS (1979). Establishment of callus tissue cultures and the induction of organogenesis in some grain legumes. *Crop Improv.* 6:154-160.
- Hazarika BN (2006). Morpho-physiological disorders in *in vitro* culture of plants. *Sci. Hortic.* 108: 105-120.
- Huda S, Islam R, Bari MA, Asaduzzaman M (2001). Anther culture of Chickpea. *Int. Chickpea Pigeonpea Newsl.* 8: 24-26.
- Huetteman CA, Preece JE (1993). Thidiazuron: a potent cytokinin for woody plant tissue culture. *Plant cell tissue organ cult.* 33: 105-119.
- Husnain T, Malik T, Rizauddin S, Gordon MP (1997). Studies on the expression of marker genes in chickpea. *Plant Cell Tissue Organ Cult.* 49: 7-16.
- Hussain T, Fatima T, Islam R and Riazuddin S (2000). Plant regeneration and expression of Beta-glucuronidase gene in hypocotyls tissue of chickpea (*Cicer arietinum* L.). *Pak. J. Biol. Sci.* 3(5): 842-845.
- Indurker S, Misra HS, Eapen S (2007). Genetic transformation of chickpea (*Cicer arietinum* L.) with insecticidal crystal protein gene using particle gun bombardment. *Plant Cell Rep.* 26: 755-763.
- Jackson MB, Abbott AJ, Belcher AR, Hall KC, Butler R, Cameron J (1991). Ventilation in plant tissue culture and effects of poor aeration on ethylene and carbon dioxide accumulation, oxygen depletion and explants development. *Ann. Bot.* 67: 229-237.
- Jayanand B, Sudarsanam G, Sharma KK (2003). An efficient protocol for the regeneration of whole plants of chickpea (*Cicer arietinum* L.) by using axillary meristem explants derived from *in vitro* germinated seedlings. *In Vitro: Cell. Dev. Biol. Plant*, 39: 171-179.
- Jiang B, Yang Y, Guo Y, Guo Z, Chen Y (2005). Thidiazuron-induced *in vitro* shoot organogenesis of the medicinal plant *Arnebia euchroma* (Royle) Johnst. *In Vitro: Cell Dev. Biol. Plant*, 41: 677-681.
- Kar S, Basu D, Das S, Ramakrishnan NA, Mukherjee P, Nayak P, Sen SK (1997). Expression of *CryIA(C)* gene of *Bacillus thuringiensis* in

- transgenic chickpea plants inhibits development of pod borer (*Heliothis armigera*) larvae. *Transgenic Res.* 6: 177-185.
- Kar S, Johnson TM, Nayak P, Sen SK (1996). Efficient transgenic plant regeneration through *Agrobacterium*-mediated transformation of Chickpea (*Cicer arietinum* L.). *Plant Cell Rep.* 16: 32-37.
- Kennedy B, Collard BCY, Brouwer JB, Taylor PWJ, Pang ECK (2002). Doubled haploid production in chickpea via anther culture. Proceedings of the 12th Australasian Plant Breeding Conference, Perth, Australia, 15-20th September 2002. pp. 84-87.
- Khan SK, Ghosh PD (1983). *In vitro* induction of androgenesis and organogenesis in *Cicer arietinum* L. *Curr. Sci.* 52: m891-893.
- Khan SK, Ghosh PD (1984). Plant regeneration from cotyledonary nodes of chickpea. *Int. Chickpea Newsl.* 11: 22-24.
- Khawar KM, Ozcan S (2004). Hairy root transformation in Turkish chickpea (*Cicer arietinum* L.) cultivars. *Biotechnol. Eq.* 18(3): 51-54.
- Kiran G, Kaviraj CP, Jogeswar G, Kishor KVK, Rao S (2005). Direct and high frequency somatic embryogenesis and plant regeneration from hypocotyls of chickpea (*Cicer arietinum* L.), a grain legume. *Curr. Sci.* 89: 1012-1018.
- Krishnamurthy KV, Suhasani K, Sagare AP, Meixner M, Kathen de A, Pickardt T, Schieder O (2000). *Agrobacterium* mediated transformation of chickpea (*Cicer arietinum* L.) embryo axes. *Plant Cell Rep.* 19: 235-240.
- Kumar PA, Bisaria S, Pai RA, Sharma RP (1995). Comparative shoot regeneration in different genotypes of chickpea, *Cicer arietinum* L. *Ind. J. Exp. Biol.* 33: 77-78.
- Kumar VD, Kirti PB, Sachan JKS, Chopra VL (1994). Plant regeneration via somatic embryogenesis in chickpea (*Cicer arietinum* L.). *Plant Cell Rep.* 13: 468-472.
- Malik KA, Saxena PK (1992). Thidiazuron induces high frequency shoots regeneration in intact seedlings of pea (*Pisum sativum*), Chickpea (*Cicer arietinum*) and lentil (*Lens culinaris*). *Aust. J. Plant Physiol.* 19: 731-740.
- Morton RL, Gollasch S, Schroeder HE, Bateman KS, Higgins TJV (2001). Genetic technology in peas for improved field performance and enhanced grain quality. In: *Handbook of Transgenic Food Plants*. Khachatorians GG, McHughen A, Scorza R, Nip WK and Hui YH (eds.), Marcel Dekker, New York, USA. pp. 197-205.
- Mroginiski E, Rey HY, Gonzalez AM, Mroginiski LA (2004). Thidiazuron promotes *in vitro* plant regeneration of *Arachis correntina* (*Leguminosae*) via organogenesis. *Plant Growth Regul.* 23: 129-134.
- Mroginiski LA, Kartha KK (1984). Tissue culture of legumes for crop improvement. *Plant Breed. Rev.* 2: 215-264.
- Munns R (2002). Comparative physiology of salt and water stress. *Plant Cell Environ.* 25: 239-250.
- Murfet IC (1971). Flowering in Pisum: reciprocal grafts between known genotypes. *Aust. J. Biol. Sci.* 24: 1089-1101.
- Murthy BNS, Murch SJ, Saxena PK (1995). TDZ-induced somatic embryogenesis in intact seedlings of peanut (*Arachis hypogaea*): endogenous growth regulator levels and significance of cotyledons. *Physiol. Plant.* 94: 268-276.
- Murthy BNS, Murch SJ, Saxena PK (1998). Thidiazuron: a potent regulator of *in vitro* plant morphogenesis. *In Vitro: Cell Dev. Biol. Plant.* 34: 267-275.
- Murthy BNS, Victor J, Singh RP, Fletcher RA, Saxena PK (1996). *In vitro* regeneration of chickpea (*Cicer arietinum* L.): stimulation of direct organogenesis and somatic embryogenesis by thidiazuron. *Plant Growth Regul.* 19: 233-240.
- Neelam A, Reddy CS, Reddy GM (1986a). Growth and differentiation in tissue cultures of *Cicer arietinum* L. *Int. Chickpea Newsl.* 14: 9-12.
- Neelam A, Reddy CS, Reddy GM (1986b). Plantlet regeneration from callus cultures of *Cicer arietinum* L. *Int. Chickpea Newsl.* 14: 12-13.
- Neelam A, Reddy CS, Reddy GM (1986c). Multiple shoots and plantlet regeneration from shoot apex and hypocotyl explants of *C. arietinum* L. *Int. Chickpea Newsl.* 14: 13-16.
- Onamu R, Obukosia SD, Musembi N and Hutchinson MJ (2003). Efficacy of thidiazuron in *in vitro* propagation of carnation shoot tips: Influence of dose and duration of exposure. *Afr. Crop Sci. J.* 11: 125-132.
- Pathak MR, Hamzah RY (2008). An effective method of sonicated-assisted *Agrobacterium*-mediated transformation of chickpea. *Plant Cell Tissue Organ Cult.* 93: 65-71.
- Pati PK, Rath SP, Sharma M, Sood A, Ahuja PS (2006). *In vitro* propagation of rose-a review. *Biotechnol. Adv.* 24: 94-114.
- Paul V, Chandra R, Khetarpal S and Polisetty R (2000). Effect of BAP induction period on shoot differentiation from seeding explants of chickpea (*Cicer arietinum* L.). *J. Plant Biol.* 27: 235-239.
- Pickardt T, Saalbach I, Waddell D, Meixner M, Müntz K and Schieder O (1995). Seed specific expression of the 2S albumin gene from Brazil nut (*Bertholletia excelsa*) in transgenic *Vicia narbonensis*. *Mol. Breed.* 1: 295-301.
- Polisetty R, Patil P, Deveshwar JJ, Khetarpal S, Chandra R (1996). Rooting and establishment of *in vitro* shoot tip explants of chickpea (*Cicer arietinum* L.). *Ind. J. Exp. Biol.* 34: 806-809.
- Polisetty R, Patil P, Deveshwar JJ, Khetarpal S, Suresh K, Chandra R (1997). Multiple shoot induction by benzyladenine and complete plant regeneration from seed explants of chickpea (*Cicer arietinum* L.). *Plant Cell Rep.* 16: 565-571.
- Polowick PL, Baliski DS, Mahon JD (2004). *Agrobacterium tumefaciens*-mediated transformation of chickpea (*Cicer arietinum* L.): gene integration, expression and inheritance. *Plant Cell Rep.* 23: 485-491.
- Potrykus I (1990). Gene transfer to plants: assessment and perspectives. *Physiol. Plant.* 79: 125-134.
- Prakash S, Chowdhary JB, Jain RK, Chowdhary VK (1992a). Factors affecting plant regeneration in chickpea, *Cicer arietinum* L. *Ind. J. Exp. Biol.* 30: 1149-1153.
- Prakash S, Yadav RC, Chowdhary JB and Sareen PK (1992b). transformation of *Cicer arietinum* L. with *Agrobacterium rhizogenes*. *Int. Chickpea Newsl.* 26: 13-14.
- Radhika K, Sujatha M, Rao TN (2006). Thidiazuron stimulates adventitious shoot regeneration in different safflower explants. *Biol. Plant.* 50: 174-179.
- Ramana RV, Venu CH, Jayasree T, Sadandam A (1996). Direct somatic embryogenesis and transformation in *Cicer arietinum* L. *Ind. J. Exp. Biol.* 34: 716-718.
- Rao BG, Chopra VL (1989). Regeneration in chickpea (*Cicer arietinum* L.) through somatic embryogenesis. *J. Plant Physiol.* 134: 637-638.
- Rizvi SMH, Singh RP (2000). *In vitro* plant regeneration from immature leaflet-derived callus cultures of *Cicer arietinum* L. via organogenesis. *Plant Cell Biotechnol. Mol. Biol.* 1: 109-114.
- Rizvi SMH, Jaiwal PK, Singh RP (2002). A possible involvement of cellular polyamine level in thidiazuron induced somatic embryogenesis in chickpea. In: *Role of Plant Tissue Culture in Biodiversity Conservation Economic Development*. Nandi SK, Palni LMS and Kumar A (eds.), Gyanodaya Prakashan, Nainital, India. pp. 163-175.
- Rout GR, Mohapatra A, Jain SM (2006). Tissue culture of ornamental pot plant: A critical review on present scenario and future prospects. *Biotechnol. Adv.* 24: 531-560.
- Sagare AP, Krishnamurthy KV (1991). Protoplast regeneration in chickpea (*Cicer arietinum* L.). *Indian J. Exp. Biol.* 29: 930-932.
- Sagare AP, Suhasani K, Krishnamurthy KV (1993). Plant regeneration via somatic embryogenesis in chickpea (*Cicer arietinum* L.). *Plant Cell Rep.* 12: 652-655.
- Sagare AP, Suhasani K, Krishnamurthy KV (1999). Comparative study of the development of zygotic and somatic embryos of chickpea (*Cicer arietinum* L.). In: *Plant Tissue Culture and Biotechnology Emerging Trends*. Kavi Kishore PB (ed.). University Press (India) Ltd. Hyd. pp. 56-63.
- Sanyal I, Singh AK, Kaushik M, Amla DV (2005). *Agrobacterium*-mediated transformation of chickpea (*Cicer arietinum* L.) with *Bacillus thuringiensis cry1Ac* gene for resistance against pod borer insect *Helicoverpa armigera*. *Plant Sci.* 168: 1135-1146.
- Sarmah BK, Moore A, Tate W, Molving L, Morton RL, Ress DP, Chiaiese P, Chrispeels MJ, Tabe LM, Higgins TJV (2004). Transgenic chickpea seeds expressing high level of a bean  $\alpha$ -amylase inhibitor. *Mol. Breed.* 14: 73-82.
- Schroeder HE, Gollash S, Moore A (1995). Bean  $\alpha$ -amylase inhibitor confers resistance to the pea weevil (*Bruchus pisorum*) in transgenic peas (*Pisum sativum* L.). *Plant Physiol.* 107: 1233-1239.
- Senthil G, Williamson B, Dinkins RD, Ramsay G (2004). An efficient transformation system for chickpea (*Cicer arietinum* L.). *Plant Cell Rep.* 23: 297-303.
- Shan X, Li D, Qu R (2000). Thidiazuron promotes *in vitro* regeneration

- of wheat and barley. *In Vitro: Cell Dev. Biol. Plant*, 36: 207-210.
- Sharma KK, Mathur PB, Jatanand B (2007). Chickpea (*Cicer arietinum* L.). *In: Methods in Molecular Biology*. Vol. 343: Agrobacterium Protocol, 2/e, vol. 1. Wang K Ed., humana Inc., Tootowa N.J.
- Sharma VK, Hansch R, Mendel RR, Schulze J (2005). Influence of picloram and thidiazuron on high frequency plant regeneration in elite cultivars of wheat with long-term retention of morphogenecity using meristemic shoot segments. *Plant Breed*. 124: 242-246.
- Shivani I, Hari M, Susan E (2007). Genetic transformation of chickpea (*Cicer arietinum* L.) with insecticidal crystal protein gene using particle gun bombardment *Plant Cell Rep*. 26: 755-763.
- Shri PV, Davis TM (1992). Zeatin-induced shoot regeneration from immature chickpea (*Cicer arietinum* L.) cotyledons. *Plant Cell Tissue Organ. Cult*. 28: 45-51.
- Seifkes-Boer HJ, Noonan MJ, Bullock DW, Conner AJ (1995). Hairy root transformation system in large-seeded grain legumes. *Isr. J. Plant Sci*. 43: 1-5.
- Singh NT, Sen J, Kiesecker H, Reddy VS, Jacobsen HJ, Mukherjee SG (2004). Use of herbicide or lysine plus threonine for non-antibiotic selection of transgenic chickpea. *Plant Cell Rep*. 22: 576-583.
- Srinivasan M, Mohaptra ST, Sharma RP (1991). *Agrobacterium* mediated genetic transformation of Chickpea (*Cicer arietinum* L.). *Ind. J. Exp. Biol*. 29: 758-761.
- Subhan S, Sharmila P, Pardha Saradhi P (1998). *Glomus fasciculatum* alleviates transplantation shock of micropropagation *Sesbania*. *Plant Cell Rep*. 17: 268-272.
- Suhasini K, Sagare AP, Krishnamurthy KV (1994). Direct somatic embryogenesis from mature embryo axes in chickpea (*Cicer arietinum* L.). *Plant Sci*. 102: 189-194.
- Tang W, Newton RJ (2005). Peroxidase and catalase activities are involved in direct adventitious shoot formation induced by thidiazuron in eastern white pine (*Pinus strobus* L.) zygotic embryos. *Plant Physiol. Biochem*. 43: 760-769.
- Tang W, Newton RJ, Charles TM (2006). Plant regeneration through multiple adventitious shoot differentiation from callus cultures of slash pine (*Pinus elliottii*). *J. Plant Physiol*. 163: 98-101.
- Tewari-Singh N, Sen J, Kiesecker H, Reddy VS, Jacobsen HJ, Mukherjee SG (2004). Use of herbicide or lysine plus threonine for non-antibiotic selection of transgenic chickpea. *Plant Cell Rep*. 22: 576-583.
- Vani AKS, Reddy VD (1996). Morphogenesis from callus cultures of chickpea, (*Cicer arietinum* L.). *Ind. J. Exp. Biol*. 34: 285-289.
- Yadav SS, Kumar J, Yadav SK, Singh S, Yadav VS, Turner NC and Redden R (2006). Evaluation of *Helicoverpa* and drought resistance in desi and kabuli chickpea. *Plant Genet. Res*. 4: 198-203.