

The long-term oral administration of thyroxine: effects on blood hematological and biochemical features in broiler breeder hens

R. Keshavarz,* A. Akhlaghi,*¹ M. J. Zamiri,* M. R. Jafarzadeh Shirazi,* F. Saemi,* A. A. Akhlaghi,#
M. Zhandi,[§] M. Afrouziyeh,[†] and M. J. Zuidhof[†]

*Department of Animal Science, School of Agriculture, Shiraz University, Shiraz 71441-65186, Fars Province, Iran; #Department of Epidemiology and Reproductive Health, Reproductive Epidemiology Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran 16635-148, Tehran Province, Iran; §Department of Animal Science, University College of Agriculture and Natural Resources, University of Tehran, Karaj 31587-77871, Tehran Province, Iran; and †Department of Agriculture, Food and Nutritional Science, University of Alberta, Edmonton, AB T6G 2P5, Canada

ABSTRACT Published data on the beneficial effect of short-term administration of thyroxine (T_4) in broiler breeder hens to reduce the ascites incidence in their progeny chicks raises the question as to what extent might the long-term maternal administration of T_4 affect the blood hematological and biochemical attributes in breeder hens. A total of 70 broiler breeder hens (47-wk-old) were randomly allotted to control or thyroxine treated (T_4) groups. Pure T_4 (0.3 mg/bird per day) was orally administered to T_4 birds for 14 successive weeks, whereas the control group received the drinking water only. Blood samples were obtained from the brachial vein prior to the initiation of the trial as well as weeks 50, 53, 55, 57, 59, and 61 of age. Body weight was decreased but egg production was not affected by T_4 treatment. Plasma concentration of T_4 , but not triiodothyronine (T_3), was increased in T_4 -treated hens ($P < 0.05$). The total number of leukocytes and erythrocytes were also higher in T_4 birds. A significant effect of time was observed for erythrocyte number

and plasma cholesterol concentration ($P < 0.05$). The long-term administration of T_4 did not affect the concentrations of serum calcium and plasma total protein, albumin, globulin, cholesterol, triglyceride, high density lipoprotein, low density lipoprotein, very low density lipoprotein, alanine amino transferase, and aspartate amino transferase ($P > 0.05$). However, serum concentrations of phosphorus, glucose, and alkaline phosphatase were higher in T_4 hens as compared to their control counterparts. In spite of differences in circulatory concentrations of a number of traits between the experimental groups, the recorded values were within their reference ranges. Therefore, the administration of T_4 for an extended period of time had no apparent adverse effect on the clinical profile in subjected hens, which may practically support the implementation of this preventative treatment as an approach to decrease the ascites incidence; however, a lower incidence rate in the progeny chicks produced from hens receiving T_4 for long-term periods of time remains to be elucidated.

Key words: biochemistry, blood enzyme, hematology, hyperthyroidism, thyroid hormone

2019 Poultry Science 0:1–6
<http://dx.doi.org/10.3382/ps/pez331>

INTRODUCTION

The ascites syndrome, as a consequence of high oxygen demands of the fast growing tissues in meat type chicken (Julian, 1993) is accompanied by a marked increase in broiler mortality rate (Genget al., 2007). Thyroid hormones concentration has been reported to be reduced in ascitic chickens prior to death and thyroxine (T_4) administration was suggested to diminish the mortality rate from the ascites syndrome in broiler chickens

(Luger et al., 2001, 2002). Accordingly, Akhlaghi et al. (2012) found that maternal administration of T_4 for 4 wk was associated with a decreased ascites incidence in their cold-exposed progeny chicks. The same treatment was also reported to be associated with enhanced early immune responses (Akhlaghi et al., 2013a) with no adverse effect on growth performance and intestinal morphology (Akhlaghi et al., 2013b) in their offspring.

Despite the importance of the preventive effect of maternal hyperthyroidism on ascites incidence rate and its enhancing effect on the immune function in the offspring, the effects of long-term administration of T_4 have not been adequately addressed. Most recently, the long-term effects of T_4 -induced hyperthyroidism on the histology of oviduct (Saemi et al., 2018a) and

© 2019 Poultry Science Association Inc.

Received February 3, 2019.

Accepted May 24, 2019.

¹Corresponding author: aakhlaghi@shirazu.ac.ir

reproductive performance (Saemi et al., 2018b) in broiler breeder hens have been reported. To our knowledge, the influence of long-term induced hyperthyroidism on the clinical profile of broiler breeder hens has not been dealt with previously. Practically, plausible adverse effects of long-term maternal administration of T_4 on hematology and biochemistry of blood would limit the use of this treatment in preventing the ascites incidence in the progeny chicks. Therefore, the aim of the present study was to ascertain whether the long-term administration of T_4 for 14 consecutive week might be associated with any changes (possibly the adverse ones) in blood hematological and biochemical attributes in Cobb 500 broiler breeder hens to provide valuable recommendations for T_4 administration in reducing the ascites incidence.

MATERIALS AND METHODS

Birds and Experimental Treatments

All procedures in the current study were approved by the Animal Care and Welfare Committee of the Department of Animal Science, College of Agriculture, Shiraz University (Shiraz, Iran). A total of seventy 47-wk-old Cobb 500 breeder hens (weighing ~3890 g) were randomly allotted to control or extra- T_4 treated (T_4) groups with 5 replicates of 7 birds for each treatment group. The hens in the T_4 treatment group individually received an oral administration of pure T_4 (0.3 mg/bird per day; Iran Hormone Pharmaceutical Company, Tehran, Iran) dissolved in 1 mL of water for 14 successive weeks. A sham operation was conducted for the control counterpart. The birds were reared under the same conditions (a 15.5L:8.5D photoschedule photoperiod and 21°C ambient temperature) and fed a corn-soybean based diet (155 g/d per bird; 2700 kcal metabolizable energy/kg, and 14.0, 2.99, and 0.36% crude protein, calcium, and phosphorus, respectively; Table 1) with a free access to fresh drinking water.

Blood Sampling and Hematology

Body weight and egg production were recorded on a weekly basis. Blood samples (5 mL) were obtained from the brachial vein prior to the experiment (week 47) as well as weeks 50, 53, 55, 57, 59, and 61 of age following weighing the birds. A fraction of each sample was collected in anticoagulant-free tubes to obtain serum samples and the remaining blood was collected in EDTA-coated tube to separate the plasmas. Prior to centrifugation, the whole blood samples were used to quantify the total erythrocyte and leukocyte number by hemocytometer, using the Natt–Herrick's solution (Buitenhuis et al., 2006). To determine the percentages of lymphocytes, monocytes, heterophils, eosinophils, basophils, and heterophil to lymphocyte (**H:L**) ratio, duplicate blood smears were air-dried, stained with

Table 1. Ingredients and chemical composition of the experimental diet fed to breeder hens (DM basis).

| Ingredient | % |
|-----------------------------|-------|
| Corn grain | 36.60 |
| Wheat grain | 25.00 |
| Barley grain | 13.40 |
| Soybean meal (44%) | 15.76 |
| Oyster shell | 7.06 |
| Dicalcium phosphate | 1.48 |
| Sodium chloride | 0.18 |
| Sodium bicarbonate | 0.16 |
| Vitamin premix ¹ | 0.10 |
| Mineral premix ² | 0.10 |
| DL-Methionine | 0.095 |
| L-Lys | 0.040 |
| L-Thr | 0.025 |
| Composition | |
| ME (kcal/kg) | 2700 |
| CP (%) | 14.00 |
| Ca (%) | 2.99 |
| P (%) | 0.36 |

¹Supplied per kg diet: vitamin A, 14,000 IU; vitamin D3, 3000 IU; niacin, 50 mg; vitamin E, 35 mg; calcium pantothenate, 20 mg; vitamin K₃, 4 mg; riboflavin, 7.0 mg; pyridoxine, 5.7 mg; vitamin B₁₂, 25 µg, and biotin, 50 µg.

²Supplied per kg diet: Fe (FeSO₄·H₂O), 85 mg; Mn (MnSO₄·H₂O), 90 mg; Zn (ZnO), 67.3 mg; Cu (CuSO₄·5H₂O), 11.1 mg, and Se (Na₂SeO₃), 0.19 mg.

Wright's-Giemsa (Saikin Kagaku Institute Co. Ltd., Sendai, Japan), and counted to a total of 100 cells/slide, using a Zeiss (Jena, Germany) compound light microscope (×1000 magnification).

Blood Biochemical Indices

Following the sampling for blood cells enumeration, the remaining blood samples were centrifuged (1800 × *g* for 12 min, 12°C), using a rotating centrifuge (International Equipment Co., Needham Heights, MA). Plasma/serum samples were then decanted and divided into 3 aliquots and stored at -20°C until analyzed for T_3 and T_4 , using commercially available kits (Padtan Elm, Tehran, Iran) and an ELISA reader (Anthos2020, Biochrom Co, England). The procedures were validated for parallelism and recovery rate in chicken samples as described previously (Akhlaghi et al., 2012). Briefly, the samples were diluted at a rate of 1 to 5 with the dilution buffer. The concentrations of T_3 and T_4 were calculated from a standard curve, which ranged between 0.5 to 300.0 and 0.03 to 10.0 ng/mL for T_4 and T_3 , respectively ($r > 0.99$). In order to control for linearity, serial dilutions at the ratios of 1:2, 1:4, 1:8, and 1:16 were provided for 4 plasma samples with given T_4 levels, using a calibrator solution ($T_4 = 0.00$ ng/mL). The samples were evaluated in duplicate. The intra- and inter-assay coefficients of variation were 12.6 and 13.2% for T_3 , and 7.6 and 2.2% for T_4 , respectively.

The concentrations of total protein, albumin, globulin, total cholesterol, high-(**HDL**), low-(**LDL**), very low-(**VLDL**) density lipoprotein, triglycerides, aspartate aminotransferase (**AST**), alanine aminotransferase

Table 2. Effects of long-term induced hyperthyroidism on body weight, egg production, and hematological attributes in Cobb 500 broiler breeder hens (least squares means).¹

| Trait | Treatment | | | P-value | | |
|---|-----------|--------------|------|-----------|--------|------------------|
| | Control | Hyperthyroid | SE | Treatment | Time | Treatment × time |
| Body weight (g) | 4369* | 4257 | 38.6 | 0.040 | NS | NS |
| Egg production (%) | 46.01 | 45.84 | 1.09 | NS | 0.0003 | <0.001 |
| Leukocytes number (10 ³ /mL) | 10.7 | 11.5* | 0.14 | <0.001 | NS | NS |
| Erythrocytes number (10 ³ /mL) | 2400 | 2600* | 0.01 | <0.001 | <0.001 | 0.002 |
| Lymphocytes (%) | 49.4 | 50.0 | 0.29 | NS | NS | NS |
| Monocytes (%) | 6.3 | 6.1 | 0.18 | NS | NS | NS |
| Heterophils (%) | 42.5 | 42.1 | 0.27 | NS | NS | NS |
| Eosinophils (%) | 1.8 | 1.6 | 0.15 | NS | NS | NS |
| Heterophils/Lymphocytes | 0.9 | 0.8 | 0.01 | NS | NS | NS |

¹The hens in the hyperthyroid group individually received an oral administration of thyroxine (0.3 mg/bird per day) dissolved in 1 mL of water for 14 successive weeks. A sham operation was conducted for the control group. Blood sampling was done from 47 to 61 wk of age (n = 35 hens/treatment). Blood sampling was done from 47 to 61 wk of age (n = 35 hens/treatment).

*Significant difference ($P \leq 0.05$).

NS: Not significant ($P > 0.05$).

(ALT), and alkaline phosphatase (ALP) were quantified in the plasma samples (Pars Azmoon Co., Tehran, Iran). The concentration of glucose, calcium, phosphorus, and ALP were determined in serum samples. The biochemical assessments were carried out by spectrophotometric methods (Butler and Laqua, 1995). The quantification protocol for each trait was also validated for chicken plasma/serum according to the procedure described for T₃ and T₄.

Statistical Analysis

The experiment was conducted as a completely randomized design. Levene's and Kolmogorov-Smirnov tests were respectively used to test the data for equality of variances and normality. The percentage data were arc-sin transformed and the data subjected to the Proc MIXED (SAS, 2002). Body weight was included in the model as a covariate for analysis of variance. The means were compared by the least squares means and the level of significance was set at $P \leq 0.05$.

RESULTS

The concentration of T₄ was elevated in the T₄ hens (27.08 ng/mL) as compared with that in the control birds (10.24 ng/mL; $P = 0.003$); whereas plasma T₃ concentration was not influenced by the long-term administration of T₄ ($P > 0.05$). No bird died during the study. The effects of long-term induced hyperthyroidism on body weight, egg production, and hematological attributes in broiler breeder hens are presented in Table 2. A decrease in body weight was recorded in T₄ birds (4,369 and 4,257 g for control and T₄ birds, respectively; $P = 0.040$). No differences in egg production were found between the experimental groups ($P > 0.05$), although time ($P = 0.003$) and treatment by time interaction effect ($P < 0.001$; Figure 1) influenced the percentage of egg production.




Figure 1. Thyroxine × time interaction effect on weekly egg production percentage in Cobb 500 broiler breeder hens. The hens in the hyperthyroid group individually received an oral administration of pure thyroxine (0.3 mg/bird per day) dissolved in 1 mL of water for 14 successive weeks. A sham operation was conducted for the control group. ^{a,b}Within each week, means with different letters differ significantly ($P \leq 0.05$). Differences were found in weeks 48, 52, 53, 55, 58, and 59.

Thyroxine administration increased the total number of leukocytes ($11.5 \times 10^3/\text{mL}$) compared to the control ($10.7 \times 10^3/\text{mL}$) birds ($P < 0.001$). Likewise, a higher total erythrocyte number was found in hyperthyroid hens (2600 vs. $2400 \times 10^3/\text{mL}$ for control and T₄ birds, respectively; $P < 0.001$). Treatment × time (wk) interacted to affect the number of erythrocytes, where higher numbers in the T₄ birds were recorded as compared to the control treatment group from 50 to 61 wk of age (Figure 2). Leukocyte differential counts and H:L ratio, however, were not affected by T₄ administration, time, and their interaction effects (Table 2).

The effects of long-term induced hyperthyroidism on blood biochemical attributes in the experimental birds are presented in Table 3. Blood phosphorus level showed an increase in the T₄ hens (5.77 mg/dL) in comparison to that in the control birds (5.17 mg/dL; $P = 0.0121$), although no significant effects were found for time and treatment by time interaction effects ($P > 0.05$). Birds




Figure 2. Thyroxine \times time interaction effect on red blood cells count in Cobb 500 broiler breeder hens. The hens in the hyperthyroid group individually received an oral administration of pure thyroxine (0.3 mg/bird per day) dissolved in 1 mL of water for 14 successive weeks. A sham operation was conducted for the control group. Blood sampling was done from 47 to 61 wk of age ($n = 35$ hens/treatment). ^{a,b}Within each week, means with different letters differ significantly ($P \leq 0.05$). No differences were found in weeks 47 and 61.




Figure 3. Thyroxine \times time interaction effect on blood glucose concentration in Cobb 500 broiler breeder hens. The hens in the hyperthyroid group individually received an oral administration of pure thyroxine (0.3 mg/bird per day) dissolved in 1 mL of water for 14 successive weeks. A sham operation was conducted for the control group. Blood sampling was done from 47 to 61 wk of age ($n = 35$ hens/treatment). ^{a,b}Within each week, means with different letters differ significantly ($P \leq 0.05$). No differences were found in weeks 47 and 61.

subjected to extra T_4 recorded a higher serum glucose level (256.9 mg/dL) when compared to that in their control counterparts (219.3 mg/dL; $P < 0.0001$). The glucose level was also affected by time ($P < 0.0017$) and treatment \times time interactive ($P < 0.0011$) effects, where the hyperthyroid treatment group recorded the higher levels during the trial, except for values recorded for 47 and 61 wk of age (Figure 3). Long-term administration of T_4 was associated with an increased serum level of ALP, where the T_4 birds recorded the value of 20.03 U/L as compared with that in the control birds (18.53 U/L; $P = 0.0051$). Extended period of exposure to T_4 did not influence the serum calcium and plasma concentrations of total protein, albumin, globulin, total cholesterol, triglycerides, HDL, LDL, VLDL, AST, and ALT ($P > 0.05$; Table 3).

DISCUSSION

Considering the role of thyroid hormones in regulating the ascites incidence in broilers (Suvarnaet al., 1993) and the lower cold-induced ascites incidence in progeny chicks produced by hens subjected to extra T_4 for a limited period (Akhlaghi et al., 2012), we hypothesized that the long-term administration of T_4 would affect the blood attributes in broiler breeder hens. Although the percentage of egg production was not affected by T_4 in the present work, comparable treatment in our most recent report was associated with a decrease in albumen height, Haugh unit, and egg weight as well as an increase in albumen pH (Rostami et al., 2019). Although the T_4 concentration was increased in the T_4

Table 3. Effects of long-term induced hyperthyroidism on blood biochemical attributes in Cobb 500 broiler breeder hens (least squares means).¹

| Trait | Treatment | | SE | P-value | | |
|---------------------------|-----------|--------------|------|-----------|---------|-------------------------|
| | Control | Hyperthyroid | | Treatment | Time | Treatment \times time |
| Calcium (mg/dL) | 11.51 | 11.88 | 0.18 | NS | NS | NS |
| Phosphorus (mg/dL) | 5.17 | 5.77* | 0.16 | 0.0121 | NS | NS |
| Glucose (mg/dL) | 219.3 | 256.9* | 3.54 | <0.0001 | <0.0017 | <0.0011 |
| Total protein (mg/dL) | 3.52 | 3.46 | 0.05 | NS | NS | NS |
| Albumin (mg/dL) | 1.81 | 1.86 | 0.02 | NS | NS | NS |
| Globulin (mg/dL) | 1.70 | 1.59 | 0.06 | NS | NS | NS |
| Total cholesterol (mg/dL) | 138.51 | 137.78 | 1.48 | NS | 0.0051 | NS |
| Triglyceride (mg/dL) | 81.58 | 81.64 | 1.69 | NS | NS | NS |
| HDL (mg/dL) | 70.51 | 70.41 | 1.13 | NS | NS | NS |
| LDL (mg/dL) | 51.68 | 51.04 | 1.67 | NS | NS | NS |
| VLDL (mg/dL) | 16.32 | 16.33 | 0.34 | NS | NS | NS |
| ALP (U/L) | 18.53 | 20.03* | 0.37 | 0.0051 | NS | NS |
| ALT (U/L) | 14.62 | 14.69 | 0.29 | NS | NS | NS |
| AST (U/L) | 151.55 | 150.68 | 2.31 | NS | NS | NS |

¹The hens in the hyperthyroid group individually received an oral administration of thyroxine (0.3 mg/bird per day) dissolved in 1 mL of water for 14 successive weeks. A sham operation was conducted for the control group. Blood sampling was done from 47 to 61 wk of age ($n = 35$ hens/treatment).

*Significant difference ($P \leq 0.05$).

NS: Not significant ($P > 0.05$).

HDL: High density lipoprotein; LDL: Low density lipoprotein; VLDL: Very low density lipoprotein; ALP: Alkaline phosphatase; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase.

hen in the current study, there was no effect of T_4 administration on plasma T_3 concentration which can be ascribed to the conversion of T_4 to inactive reversed T_3 (Decuyper et al., 1987).

Red blood cells (Maxon et al., 1975), white blood cells (Fabris, 1973; Bachman and Mashaly, 1987), and blood concentrations of calcium (Jowsey and Detenbeck, 1969), total proteins (Goodman, 2009), albumin (Harrison and Harrison, 1986), cholesterol (Campbell and Coles, 1986), triglycerides (Davey et al., 2001), lipoproteins (Duntas, 2002), and several serum enzymes (Huang and Liaw, 1995) were reported to be influenced by thyroid hormones. Although T_4 treatment increased the total number of leukocytes, we found no effect of long-term administration of T_4 on the relative number of lymphocytes, monocytes, heterophils, and eosinophils. Klecha et al. (2006) showed that thyroid hormones have an important role in regulating lymphocyte reactivity. Fabris (1973) suggested that thyroidectomy caused a decrease in total leukocytes number as a result of decreased lymphocyte number. Consistent to the present findings, Chatterjee and Chandel (1983) showed that hyperthyroidism was associated with increased white blood cells and lymphocytes numbers, and there was a positive association between thyroid hormone levels and lymphocyte numbers in birds. Hyperthyroidism was also shown to be associated with an increased lymphocyte number in human (Ullmann et al., 1963). Increased erythrocyte number in T_4 hens may be ascribed to the erythropoietic effect of thyroid hormones (Golde et al., 1977), although erythrocyte concentration was within reference ranges in the current study.

Bone mobilization in hyperthyroid individuals, as a result of increased osteoclastic activity, is associated with a trend to hypercalcemia and decreased parathyroid hormone secretion (Mosekilde et al., 1990); however, T_4 treatment over a relatively long period did not affect the serum calcium concentration in the present study. Serum phosphorous in the T_4 group was higher than that in the control treatment group. It has been shown that hyperthyroid individuals show hyperphosphatemia as a result of increased bone catabolism and decreased phosphorous clearance (Mosekilde and Christensen, 1977). Therefore, the higher levels of phosphorous in T_4 hens may be due to the direct effect of thyroid hormones on tissue phosphate metabolism and renal phosphate clearance (Dhanwal, 2011).

Alkaline phosphatase, a biochemical marker of bone formation and resorption, was increased in hyperthyroid patients (Pantazi and Papapetrou, 2000). Increased alkaline phosphatase concentration in the T_4 hens in the present work may be attributed to the stimulation of membrane-bound fragment of alkaline phosphatase in osteoblasts by T_4 (Banovac and Koren, 2000).

Birds subjected to extra T_4 recorded a higher serum glucose level in the present study. Thyroid hormones have an important role in regulation of glucose

homeostasis (Brenta, 2010). Hyperthyroidism increases blood glucose levels, and thyrotoxic patients show diabetic glucose tolerance after a reversal to the euthyroid condition (Maxon et al., 1975). Thyroid hormones influence metabolism in the small intestine with T_4 in vitro for 72 h increasing glucose active transport in the duodenum of chick embryos (Black, 1988). Thyroid hormones elevate the expression of the enzymes involved in gluconeogenesis, phosphoenolpyruvatecarboxykinase (Park et al., 1999), and glucose-6-phosphatase (Suh et al., 2013), which are involved in hepatic glucose production. Hyperthyroidism causes activation of hepatic gluconeogenesis by enhancing alanine transport and its transformation to glucose (Park et al., 1999). In hyperthyroid individuals, production of endogenous glucose is elevated and does not respond to the repressive effect of insulin (Holness and Sugden, 1987).

Overall, the long-term administration of T_4 in broiler breeder hens had no adverse effects on blood hematological and biochemical characteristics, and apart from increases in the concentration of some attributes, the recorded values were within their reference ranges. According to unaffected H:L ratio, the birds orally administered by extra T_4 did not experience stress during the trial. The present study, however, did not evaluate the progeny chicks in term of ascites incidence. If the preventative effect of long-term maternal exposure to extra T_4 on the ascites incidence in progeny chicks (not evaluated here) is comparable to that of the short-term exposure (Akhlaghi et al., 2012), the long-term maternal administration of T_4 may then be a helpful alternative to decrease the incidence of ascites in the progeny of broiler breeder hens; however, application at the farm level requires further substantiation of the present findings, with an eye to immune- and growth-associated effects in progeny broilers.

ACKNOWLEDGMENTS

Grateful thanks are extended to Iran Hormone Co. for providing pure T_4 in the present study.

REFERENCES

- Akhlaghi, A., M. J. Zamiri, Y. Jafari Ahangari, H. Atashi, Z. Ansari Pirsaraei, H. Deldar, A. N. Eghbalian, A. A. Akhlaghi, B. Navidshad, and K. Yussefi Kelarikolaei. 2013. Oral exposure of broiler breeder hens to extra thyroxine modulates early adaptive immune responses in progeny chicks. *Poult. Sci.* 92:1040–1049.
- Akhlaghi, A., M. J. Zamiri, Y. Jafari Ahangari, M. R. Mollasalehi, H. Shojaee, H. Atashi, B. Navidshad, A. A. Akhlaghi, and M. Dadpasand. 2013. Growth performance and intestinal morphology in broiler chickens produced from hyperthyroid breeder hens. *Anim. Prod. Sci.* 53:1046–1051.
- Akhlaghi, A., M. J. Zamiri, A. Zare Shahneh, Y. Jafari Ahangari, A. Nejati Javaremi, G. Rahimi Mianji, M. R. Mollasalehi, H. Shojaie, A. A. Akhlaghi, H. Deldar, H. Atashi, Z. Ansari Pirsaraei, and M. Zhandi. 2012. Maternal hyperthyroidism is associated with a decreased incidence of cold-induced ascites in broiler chickens. *Poult. Sci.* 91:1165–1172.
- Bachman, S. E., and M. M. Mashaly. 1987. Relationship between circulating thyroid hormones and cell-mediated immunity in immature male chickens. *Dev. Comp. Immunol.* 11:203–213.

- Banovac, K., and E. Koren. 2000. Triiodothyronine stimulates the release of membrane-bound alkaline phosphatase in osteoblastic cells. *Calcif. Tissue Int.* 67:460–465.
- Black, B. L. 1988. Development of glucose active transport in embryonic chick intestine. Influence of thyroxine and hydrocortisone. *Comp. Biochem. Physiol.* 90:379–386.
- Brenta, G. 2010. Diabetes and thyroid disorders. *Br. J. Diabetes Vasc. Dis.* 10:172–177.
- Buitenhuis, A. J., J. B. Kjaer, R. Labouriau, and H. R. Juul-Madsen. 2006. Altered circulating levels of serotonin and immunological changes in laying hens divergently selected for feather pecking behavior. *Poult. Sci.* 85:1722–1728.
- Butler, L., and K. Laqua. 1995. Nomenclature, symbols, units and their usage in spectrochemical analysis-IX. Instrumentation for the spectral dispersion and isolation of optical radiation (IUPAC Recommendations 1995). *Pure Appl. Chem.* 67:1725–1744.
- Campbell, T., and E. Coles. 1986. Avian clinical pathology. *Vet. Clin. Pathol.* 4:279–300.
- Chatterjee, S., and A. S. Chandel. 1983. Immunomodulatory role of thyroid hormones: in vivo effect of thyroid hormones on the blastogenic response of lymphoid tissues. *Act. Endocrinol.* 103:95–100.
- Davey, F., R. Hutchison, and J. Henry. 2001. *Clinical Diagnosis and Management by Laboratory Methods*. WB Saunders Company, Philadelphia, PA.
- Decuypere, E., J. Buyse, C. Scanes, L. Huybrechts, and E. Kühn. 1987. Effects of hyper- or hypothyroid status on growth, adiposity and levels of growth hormone, somatomedin C and thyroid metabolism in broiler chickens. *Reprod. Nutr. Dev.* 27:555–564.
- Dhanwal, D. K. 2011. Thyroid disorders and bone mineral metabolism. *Indian J. Endocr. Metab.* 15:S107–S112.
- Duntas, L. H. 2002. Thyroid disease and lipids. *Thyroid* 12:287–293.
- Fabris, N. 1973. Immunodepression in thyroid-deprived animals. *Clin. Exp. Immunol.* 15:601–611.
- Geng, A., B. Li, and Y. Guo. 2007. Effects of dietary L-carnitine and coenzyme Q10 at different supplemental ages on growth performance and some immune response in ascites-susceptible broilers. *Arch. Anim. Nutr.* 61:50–60.
- Golde, D., N. Bersch, I. Chopra, and M. Cline. 1977. Thyroid hormones stimulate erythropoiesis in vitro. *Br. J. Haematol.* 37:173–177.
- Goodman, H. (2009) Principal of hormonal integration. Pages 91–99 in *Basic Medical Endocrinology*. Academic Press, Burlington.
- Harrison, G. J., and L. R. Harrison. 1986. *Clinical Avian Medicine and Surgery: Including Aviculture*. W B Saunders Co.
- Holness, M. J., and M. C. Sugden. 1987. Continued glucose output after re-feeding contributes to glucose intolerance in hyperthyroidism. *Biochem. J.* 247:801–804.
- Huang, M. J., and Y. F. Liaw. 1995. Clinical associations between thyroid and liver diseases. *J. Gastroenterol. Hepatol.* 10:344–350.
- Jowsey, J., and L. C. Detenbeck. 1969. Importance of thyroid hormones in bone metabolism and calcium homeostasis. *Endocrinology* 85:87–95.
- Julian, R. J. 1993. Ascites in poultry. *Avian Pathol.* 22:419–454.
- Klecha, A. J., A. M. Genaro, G. Gorelik, M. L. B. Arcos, D. M. Silberman, M. Schuman, S. I. Garcia, C. Pirola, and G. A. Cremaschi. 2006. Integrative study of hypothalamus–pituitary–thyroid–immune system interaction: thyroid hormone-mediated modulation of lymphocyte activity through the protein kinase C signaling pathway. *J. Endocrinol.* 189:45–55.
- Luger, D., D. Shinder, V. Rzepakovsky, M. Rusal, and S. Yahav. 2001. Association between weight gain, blood parameters, and thyroid hormones and the development of ascites syndrome in broiler chickens. *Poult. Sci.* 80:965–971.
- Luger, D., D. Shinder, and S. Yahav. 2002. Hyper- or hypothyroidism: its association with the development of ascites syndrome in fast-growing chickens. *Gen. Comp. Endocrinol.* 127:293–299.
- Maxon, H. R., M. Kreines, M. Goldsmith, and H. C. Knowles, Jr. 1975. Long-Term observations of glucose tolerance in thyrotoxic patients. *Arch. Intern. Med.* 135:1477–1480.
- Mosekilde, L., and M. S. Christensen. 1977. Decreased parathyroid function in hyperthyroidism: interrelationships between serum parathyroid hormone, calcium-phosphorus metabolism and thyroid function. *Act. Endocrinol.* 84:566–575.
- Mosekilde, L., E. F. Eriksen, and P. Charles. 1990. Effects of thyroid hormones on bone and mineral metabolism. *Endocrinol. Metab. Clin. North Am.* 19:35–63.
- Pantazi, H., and P. D. Papapetrou. 2000. Changes in parameters of bone and mineral metabolism during therapy for hyperthyroidism. *J. Clin. Endocrinol. Metab.* 85:1099–1106.
- Park, E. A., S. Song, C. Vinson, and W. J. Roesler. 1999. Role of CCAAT enhancer-binding protein β 2 in the thyroid hormone and cAMP induction of phosphoenolpyruvate carboxykinase gene transcription. *J. Biol. Chem.* 274:211–217.
- Rostami, S., Y. Jafari Ahangari, A. Akhlaghi, Z. Ansari Pirsaraei, S. R. Hashemi, and F. Saemi. 2019. Effects of long-term induced hyperthyroidism on egg quality traits in Cobb 500 broiler breeder hens. *Poult. Sci. J.* (Accepted).
- Saemi, F., A. Z. Shahneh, M. Zhandi, A. Akhlaghi, Z. Khaksar, and M. Dadpasand. 2018. Long-term effects of thyroxine-induced hyperthyroidism on the histological attributes of the oviduct in broiler breeder hens. *Comp. Clin. Pathol.* 27:605–609.
- Saemi, F., A. Z. Shahneh, M. Zhandi, A. Akhlaghi, and Z. Ansari Pirsaraei. 2018. TGF- β 2 and HSP70 responses in breeder hens treated with thyroxine. *Anim. Reprod. Sci.* 198:82–89.
- SAS. 2002. *User's Guide. Statistical Analysis Systems Institute Inc.*, Cary, NC, USA.
- Suh, J. H., D. H. Sieglaff, A. Zhang, X. Xia, A. Cvorov, G. E. Winnier, and P. Webb. 2013. SIRT1 is a direct coactivator of thyroid hormone receptor β 1 with gene-specific actions. *PLoS One* 8:e70097.
- Suvarna, S., F. McNabb, E. Dunnington, and P. Siegel. 1993. Intestinal 5' deiodinase activity of developing and adult chickens selected for high and low body weight. *Gen. Comp. Endocrinol.* 91:259–266.
- Ullmann, J. E., G. A. Hyman, and B. Calderz. 1963. The occurrence of lymphoma in patients with long-standing hyperthyroidism. *Blood* 21:282–297.