



The Open Rheumatology Journal

Content list available at: www.benthamopen.com/TORJ/

DOI: 10.2174/1874312901711010030



RESEARCH ARTICLE

Hip Abductor Strengthening Improves Physical Function Following Total Knee Replacement: One-Year Follow-Up of a Randomized Pilot Study

Karvannan Harikesavan^{1,*}, Raj D. Chakravarty², Arun G Maiya³, Sanjay P. Hegde² and Shivakumar Y. Shivanna²

¹Department of Physiotherapy, School of Allied Health sciences, Manipal University, Bangalore. India.

²Orthopaedic joint replacement surgeon. Manipal Hospital, Bangalore. India.

³Department of Physiotherapy, School of Allied Health Sciences, Manipal University, Manipal, India.

Received: September 09, 2016

Revised: December 07, 2016

Accepted: February 02, 2017

Abstract:

Background:

Total knee replacement (TKR) is the commonest surgical procedure for patients with severe pain and impaired physical function following end stage knee osteoarthritis. The hip abductors are well renowned in stabilization of the trunk and hip during walking, maintaining the lower limb position, and transferring the forces from the lower limbs to the pelvis.

Objective:

To assess the efficacy of hip abductor strengthening exercise on functional outcome using performance based outcome measures following total knee replacement.

Methods:

An observer blinded randomized pilot trial design was conducted at Manipal hospital, Bangalore, India. Participants designated for elective TKR were randomized to experimental group hip abductor strengthening along with standard rehabilitation (n=10) or control group standard rehabilitation alone (n=10). Participants followed for one year to assess physical function using performance based outcomes, such as timed up and go test, single leg stance test, six minute walk test, knee extensor strength and hip abductor strength.

Result:

Eighteen participants with a mean age of 63.1 ± 5.5 years (8 Males and 10 Females) completed the study. Improvement in hip abduction strength, single leg stand test was superior in hip abductor strengthening group at 3 months and 1 year when compared to standard rehabilitation alone.

Conclusion:

Hip abductor strengthening showed superior improvements in single leg stance test and six minute walk test. Hip abductor strengthening exercises has the potential to improve physical function following total knee replacement.

Keywords: Total knee replacement, Hip abductor strengthening, Knee exercises, Single leg stance, Osteoarthritis.

* Address correspondence to this author at the Department of Physiotherapy, No 98, Rustum Bagh, Old Airport Road, School of Allied Health Sciences, Manipal University, Bangalore, India; Tel: +91-9986565966; Fax: +91-8202571915; E-mails: karvannan.h@manipal.edu, karvannan.h@gmail.com

INTRODUCTION

Total knee replacement (TKR) is the most renowned surgical procedure for patients with severe knee pain and impaired function following end stage knee osteoarthritis [1, 2]. Significant improvements in pain reduction, improved function and patient satisfaction were the main expectations following TKR [3]. Studies reported that participants showed substantial improvements in arthritis pain, but a varied physical function following TKR [4, 5].

Despite more advanced and excellent surgical procedure, declined functional tasks were reported following TKR when compared to the healthy age matched controls with 15% reduced walking speed, 50% more time taken to complete stair climbing tasks and 20% less distance covered during the six minute walk test [6, 7]. Franklin *et al.* [3], suggested that demographic and clinical variables would predict the functional improvement especially the quadriceps strength. The strengthening of the quadriceps muscles was established by previous studies and proved its short-term improvements in physical function after TKR [8 - 10].

A systematic review of Minne lowis *et al.* [11], and Artz *et al.* [12], recommended that future trials should tailor the post-operative interventions for enhanced functional performance measures and long-term benefits. Arnold *et al.* in their recent systematic review found the long-term changes in physical activity following TKR, but with limited evidence and recommended to reiterating the approaches to improve the physical function and patients' expectations following TKR [13].

Reconnoitering the modifiable factors contributing to the functional performances following TKR might help to improve performance based outcomes, one such modifiable factor is the lower extremity muscle weakness following knee osteoarthritis (OA). A recent systematic review and meta-analysis of symptomatic knee OA patients revealed the weakness of isometric and isokinetic hip abductor strength [14]. Recent studies on total knee arthroplasty revealed that hip abductor strength was positively correlated to quadriceps strength for enhanced performance-based function measures [15, 16]. The hip abductors are well renowned as a primary muscle group for the stabilization of trunk and hip during walking, maintaining the femoropelvic alignment, femoral head stability and transferring the forces from the lower limbs to the pelvis [17 - 19]. Moreover, weakness of the hip abductor resulting in poorer functional performance in older adults and associated with reduced physical performance in patients with knee OA [20]. Hence, we hypothesized that the hip abductor strengthening might augment physical function in participants post TKR.

MATERIALS AND METHODOLOGY

Study Design

We conducted an observer blinded, randomized pilot study with outcome measurement taken before surgery and at 1 month, 3 months and 12 months following TKR. The study was conducted at College of Physiotherapy, School of Allied Health Sciences, Manipal hospital, Bangalore, India. The study protocol was approved by the Institutional Research Committee of School of Allied Health Sciences, Manipal University, Manipal, India.

Participants

Participants posted for the elective TKR screened for the inclusion and exclusion criteria. Participants were included if 50 years of age or older with osteoarthritis as diagnosis of the either knee or both knees was painful and they specified the most painful knee. All the study participants were screened by the orthopaedic surgeons for the diagnosis and staging of knee osteoarthritis based on the major elements of the diagnostic criteria like history, physical examination, imaging studies [21, 22]. The participants who were diagnosed with end stage knee osteoarthritis were posted for elective total knee replacement.

They were excluded, if diagnosed other than knee osteoarthritis, any neurological impairment that may alter the lower extremity performance, any other orthopedic surgery in either leg in the past year. All eligible participants signed an informed consent. Using block randomization, participants were randomly allocated to either experimental or control group. Prior to the enrollment of the participants, the Institutional research committee of Manipal University approved the study. The rights of the included participants have been reserved..

A total of 20 participants were included and randomly assigned to hip abductor strengthening group (n=10) or the control group (n=10) using block randomization prior to the surgery. A total of 10 males (6 in hip abductor strengthening group and 4 in the control group) and 10 females (4 in hip abductor strengthening group and 6 in the control group) were included in the study with a mean age of 63.3 (5.4) in hip abductor strengthening group and 62.8

(5.9) in the control group. The mean body mass index (BMI) of the hip abductor strengthening group was 26.5 (3.2) and the control group was 29.8 (3.2), respectively.

Rehabilitation Protocol

The rehabilitation protocol was established by the authors, physiotherapy department, Manipal University, Bangalore, India. All the participants in the control group underwent a standard rehabilitation protocol from day of operation until discharge. The standard rehabilitation focused on early mobilization, such as reducing pain and swelling, improving knee flexion and extension range of motion, and progressive quadriceps strengthening exercises to maximize function (Appendix-1). Hip abductor strengthening exercise group underwent hip abductor strengthening exercises along with the standard rehabilitation (Appendix-2). Both the groups underwent 30- 40 minutes of supervised physical therapy per session, twice a day until discharge. Following discharge the participants underwent supervised rehabilitation for 4-5 sessions in a week over 4 weeks duration, and thereafter 2-3 sessions per week for 12 weeks with each session lasted for about 40- 45 minutes. They were asked to continue their home based exercise and the exercise adherence was ensured through logbook and periodic telephone calls.

Variables

Baseline outcome measures of timed up and go test (TUG), six minute walk test (SMWT), single leg stance (SLS), numeric pain rating scale (NPRS), knee extensor strength and hip abductor strength were taken prior to the surgery. Post outcome measure of the same was recorded at 1 month, 3 months and 1 year following the surgery. An independent blinded observer with more than 5 years of experience in physical therapy collected the outcome measures.

Physical Function

Physical function allied to the person's ability to move around for performing the activities of daily living [23]. It can be measured using the performance based tests or self- reported measures using the questionnaires, both measures quarantine different forms of physical functioning [24 - 26]. The performance based measures were often assessed through observing the number of repetitions, timing or the totaling distance covered. These measures ideally evaluate the performance of the participants rather than the perception of what they can do [23, 25]. Performance based tests might evaluate distinguishing pain and function other than familiar self-reported questionnaires [25, 26]. However, studies have revealed that both the measure is seen as amenable, rather than challenging when appraising physical function for patients with hip and knee OA also follows total knee replacement [26 - 28].

Timed Up and Go Test

Timed up-and-go test (TUG) also termed as the ambulatory transitions test used for identifying problems in functional mobility [29 - 31]. The TUG comprises various activities, such as sit-to-stand from a standard chair with armrest (46 cm seat height), walking 3 m distance in their usual manner (customary walking aids if necessary), and turning at the end while walking and back to their chair to sit down again [32, 33]. With regular footwear, all the participants underwent one warm up trial prior to being timed with fastest of two trials [29]. TUG has an excellent inter-rater and intra-rater reliability in older adults and is responsive to changes after TKR and it has the ability to distinguish the physical functional performance of healthy subjects from patients with TKR [24, 32, 33].

Six Minute Walk Test

The six minute walk test (SMWT) assesses the physical function by totaling the distance covered maximally by the participant walking at their free speed in patients with knee osteoarthritis and people who are indicated for surgery [24, 34 - 37]. Participants asked to walk as quickly as they felt safe on a measured 46 meter uncarpeted rectangular indoor circuit during the 6-minute period. The participants walked as much distance as possible with an assistive device if required and the distance covered was measured to the nearest meter [35, 38].

Single Leg Stance

Single leg stance (SLS) used to assess the static postural control during standing on the operated leg. The participants were asked to balance on their stance leg with hands on the hips and the test duration was limited to 30 Sec, the longest duration of the three trials was recorded, followed by one warm up trial. The test was discontinued when the swing leg touches the stance limb or the floor for balance, the stance foot displaced from the ground, or when the participants moved his arm away from hips. This test has shown to be reliable, responsive to interventions and it is a

common clinical assessment tool for determining the functional ability of the participants of various ages and functional levels [39 - 42].

Muscle Strength Test

Participants underwent hip abductor and knee extensor maximal isometric strength assessment using hand held dynamometer (HHD) (Fabricatio enterprises incorporation, New York). They were positioned in supine lying with the tested limb in neutral position for hip abductor measurement. The HHD placed 5 cm above the knee joint line of the tested limb and the non-tested limb was positioned in neutral for stability. Participants asked to abduct their tested lower limb as hard as they can for 5 seconds with the therapist holding the force transducer of the HHD. The participants were encouraged verbally during the test for their maximal effort; and the maximum volitional force generated by the participant was quantified hip abductor muscle strength in pounds. Knee extensor maximal isometric strength (Quadriceps) of the participants was assessed in sitting position with hip in 90° and knee in 70° of flexion. They performed knee extension as hard as they can of the tested limb with pelvis and trunk stabilized [43 - 46].

Single blinded assessor conducted all the test and retest measurement with three trials performed per side, followed by one warm up trial. The maximal volitional force generated on hip abductors and knee extensors quantified muscle strength in pounds. This method provides the objective and quantitative measurements of the hip abductor and knee strength and showed to be a valid and reliable clinical tool to test muscle strength for athletes, patients with knee OA and those who underwent TKR [44 - 46].

Numeric Pain Rating Scale (NPRS)

The participants are asked to rate their pain intensity on an 11-point numeric pain rating scale (NPRS) where '0' indicates no pain and '10' indicates the worst pain imaginable. An independent blinded observer recorded the knee pain of the participant at baseline, 1 month, 3 months and 1 year following the surgery.

Statistical Analysis

Statistical analysis was conducted using SPSS software (version 16.0; SPSS. Chicago, IL, USA). Repeated measures ANOVA was used to determine the effects of hip abductor strengthening and standard rehabilitation post total knee replacement at baseline, 1 month, 3 months and 1 year, respectively. Post-hoc analysis using Bonferroni's test was used to determine the pairwise comparisons at different measurement levels amongst experimental and control groups. Equality of variance for the continuous variables was tested using Leven's test. The statistical significance was set at p value < 0.05 with 95% confidence intervals, and all the tests were 2-tailed.

RESULTS

Eighteen participants (8 males and 10 females) with a mean age of 63.1 ± 5.5 years completed the study. Both groups were comparable on clinical and demographic characteristics at baseline, and also for functional measures of TUG, SMWT, SLS, NPRS, and hip and knee strength measurements. The baseline characteristics of the included participants are mentioned in (Table 1). Two participants in the control group were not able to follow-up due to immediate post-operative complications. The pre-operative baseline score and changes within group at 1 month, 3 months, 1 year of knee extension strength, hip abductor strength, TUG, SMWT, SLS and NPRS are mentioned in (Table 2). The comparison between the groups of all the outcome variables is mentioned in (Table 3).

Table 1. Baseline characteristics of participants.

Variables	Hip abductor strengthening group Mean (SD) or N (%)	Control group Mean (SD) or N (%)	p value
Age in years	63.3 (5.4)	62.8 (5.9)	0.962
Sex M: F	6 (60%): 4 (40%)	4 (40%): 6(60%)	0.913
BMI	26.5 (3.2)	29.8 (3.2)	0.618
Knee strength in pounds	36.1 (8.4)	34.8 (4.9)	0.060
Hip Abductor strength in pounds	36.1 (6.0)	36 (7.0)	0.537
SMWT in meters	255.1 (79.2)	201.1(64.4)	0.597
TUG in meters	16.3 (4.2)	18(4.3)	0.538
SLS in sec	8.3 (4.5)	6.5 (2.5)	0.355
NPRS	7.1(.56)	6.5(1.5)	

Table 2. Within group analysis of outcome measures.

Outcome measures	Baseline	1 month	3 month	1 year	Baseline to 1 month Mean (95% CI)	p value	Baseline to 3 month Mean (95% CI)	p value	Baseline to 1 Year Mean (95% CI)	p value
Hip abductor strengthening group										
Knee strength	36.1(8.4)	36.4(8.8)	40.9(10.1)	41.7(7.5)	.3(-.71-1.3)	<0.520	4.8(2.6-6.9)	<0.001	5.6(2.7-8.4)	<0.002
Hip abductor strength	36.1(6.0)	40.3(7.2)	42.6(6.9)	45.7(6.7)	4.2(2.4-5.9)	<0.001	6.5(4.1-8.8)	<0.001	9.6(7.8-11.3)	<0.001
SMWT	255.1(79.2)	304.6(108.6)	387.6(119.5)	474.1(160.5)	49.5(-8.1-107.1)	<0.084	132.5(62.6-202.3)	<0.002	219(104.1-333.8)	<0.002
SLS	8.3(4.5)	6.6(3.4)	13.7(5.4)	15.2(5.9)	-1.7 (-4.1-.75)	<0.271	5.4(2.5 -8.2)	<0.001	6.9 (4.6-9.1)	<0.001
TUG	16.3(4.2)	17.5(3.9)	13.3(4.0)	11.3(2.2)	-1.2(-3.9-1.5)	<0.347	3(.78-5.2)	<0.013	5(2.1-7.8)	<0.003
NPRS	7.1(.56)	4.2(1.1)	1.7(.6)	1.4(.9)	2.9(1.9-3.8)	<0.001	5.4(4.7-6)	<0.001	5.7(4.9-6.4)	<0.001
Control group										
Knee strength	34.8(4.9)	37.5(4.4)	42.3(4.3)	42.8(3.1)	2.6(-.53-5.7)	<0.070	7.5(6.3-8.6)	<0.001	8(5.6-10.3)	<0.001
Hip abductor strength	36(7)	37.12(5.8)	37.3(6.4)	39.8(6.2)	1.1(-.89-3.1)	<0.229	1.3(.38-2.3)	<0.014	3.8(2.8-4.9)	<0.001
SMWT	201.1(64.4)	216.8(90.5)	319.3(134.8)	380.6(124.7)	15.7(-30.3-61.8)	<0.445	118.2(35.3-201.1)	<0.012	179.5(90.5-268.4)	<0.002
SLS	6.5(2.5)	5.3(1.1)	7.62(2.7)	9.5(2.3)	-1.1(-4.8-2.6)	<0.100	1.1(-2.3-4.5)	<0.100	3(.3-5.6)	<0.027
TUG	18(4.3)	18.7(5.3)	15.2(4)	13.8(2.6)	-.75(-6.1-4.6)	<0.754	2.7(-.58-6.0)	<0.092	4.1(1.5-6.7)	<0.007
NPRS	6.5(1.5)	3.6(1.5)	2.1(.8)	1.1(.8)	2.8(1.4-4.3)	<0.002	4.3(2.8-5.9)	<0.001	5.3(3.9-6.7)	<0.001

SMWT- Six minute walk test, measured in meters; TUG- Timed up and Go test, measured in seconds; SLS- Single leg stance test, measured in seconds; NPRS- Numeric pain rating scale; Knee strength (quadriceps strength) in pounds; Hip abductor strength in pounds.

Table 3. Between group analysis of outcome measures.

Outcome measures	Baseline to 1 month Mean (95% CI)	p value	Baseline to 3 month Mean (95% CI)	p value	Baseline to 1 Year Mean (95% CI)	p value
Between group analysis						
Knee strength	-2.3(-5.4)	<0.040	-2.9(-5.3 - -.45)	<0.023	-3.2(-6.5-.18)	<0.062
Hip abductor strength	3.0(.6-5.5)	<0.018	4.0(1.2-6.7)	<0.005	5.4(3.3-7.5)	<0.001
SMWT	19(-29.4-68.3)	<0.410	53.6(-10.4-117.7)	<0.095	88.3(36.5-140.2)	<0.002
SLS	-.82(-3.4-1.7)	<0.511	4.2(1.5-6.9)	<0.004	3.9(1.8-5.9)	<0.001
TUG	-.45(-5.6-4.7)	<0.857	.25(-3.2-3.7)	<0.883	.77(-2.9-4.4)	<0.661
NPRS	0.025(-1.4-1.5)	<0.972	1.02(-.4-2.4)	<0.148	-.72(-2.4-.86)	<0.330

SMWT- Six minute walk test, measured in meters; TUG- Timed up and Go test, measured in seconds; SLS- Single leg stance test, measured in seconds; NPRS- Numeric pain rating scale; Knee strength (quadriceps strength) in pounds; Hip abductor strength in pounds.

The Hip abductor strengthening group (HAS) has shown statistically significant changes in knee strength at 3 months when compared to baseline and the improvements were sustained during 1 year follow-up. The HAS group showed a significant change from 36.1 ± 8.4 to 40.9 ± 10.1 at 3 months and preserved till 1 year 41.7 ± 7.5 . Knee strengthening group or the control group (KS) showed statistical significant changes from 34.8 ± 4.9 to 42.3 ± 4.3 at 3 months and 42.8 ± 3.1 at 1 year respectively. Both the groups did not show statistical significant changes at 1 month following TKR. Between group comparison using post hoc analysis did not show a statistical significant at all the post outcome measures.

Hip abductor strength test at baseline in the HAS group was 36.1 ± 6.0 and control group was 36 ± 7 . The strength in HAS showed significant improvement at 1 month, 3 months and year with a value of 40.3 ± 7.2 , 42.6 ± 6.9 and 45.7 ± 6.7 , respectively. The control group showed a significant difference from baseline 36 ± 7 to 39.8 ± 6.2 at 1 year. Between groups comparison showed a statistically significant strength difference in the HAS group at 3 months and 1 year following TKR.

From the baseline to 1 month, both the groups had spent more duration while performing TUG. Both the groups showed the statistically significant improvements in one year following TKR. Between groups comparison on TUG, the

functional task at all the levels did not show a superior change. The time taken to complete the TUG was reduced from 16.3 ± 4.2 from baseline to 11.3 ± 2.2 at 1 year follow-up for HAS group and control group has improved from 18 ± 4.3 to 13.8 ± 2.6 .

The HAS group walked faster than the control group at 1 year duration during SMWT. The HAS group showed a significant change from a baseline value of 255.1 ± 79.2 meters to 387.6 ± 119.5 and to 474.1 ± 160.5 meters at 3 months and 1 year, respectively. The control group showed the improvement from a baseline value of 201.1 ± 64.4 to 380.6 ± 124.7 at 1-year duration. Between groups analysis showed a significant mean difference in SMWT at one year in HAS group with a p value of <0.002 .

The test for static postural control was assessed using SLS and it showed improvement in HAS group at 3 months and 1 year duration. However, it was not statistically significant at 1 month duration. The HAS showed changes in SLS from a baseline value of 8.3 ± 4.5 to 13.7 ± 5.4 at 3 months and 15.2 ± 5.9 Sec at 1 year duration. Between group analysis revealed a superior improvements in the HAS group at 3 months and 1 year as compared to the control group.

The NPRS showed a statistically significant difference in both the groups at all the periods. The HAS group showed a significant difference from a baseline value of $7.1 \pm .56$ to 4.2 ± 1.1 , 1.7 ± 0.6 , 1.4 ± 0.9 at 1 month, 3 months and 1 year, respectively. The control group showed a significant improvement with 3.6 ± 1.5 , $2.1 \pm .8$, $1.1 \pm .8$ at 1 month, 3 months and 1 year when compared to a baseline value of 6.5 ± 1.5 . Between groups analysis did not find a significant difference in NPRS during all the period.

DISCUSSION

The primary aim of this study is to determine whether hip abductor strengthening would improve physical function following total knee replacement when compared to standard rehabilitation. Hip abductor strengthening group showed a statistically significant improvement in physical functional measure, such as single leg stance at 3 months and 1 year and six-minute walk test at 1 year following TKR.

At one month and 3 months, the time taken to complete the TUG was not statistically significant and the participants in both groups showed an increased time to complete the task. Greater deficits were shown in the performance at 1 month and recovered to the preoperative levels at 3 months and it further showed significant gains at 1 year following surgery. Similarly, previous studies showed poor performances following TKR [7, 10] and this could possibly be due to the decreased quadriceps and hip abductor strength at one month following surgery.

The participants in the HAS group walked faster and longer than the KS participants at 1 year. The HAS group walked additional 132 meters at 3 months, 219 meters at 1 year and the KS group walked 118 meters more at 3 months and 179 meters at 1 year. The group's analysis has shown a significant difference at 1 year with a mean difference of 88.3 meters. A study targeting functional rehabilitation by Moffet *et al.* [10], found that their subjects walked 145 meters lesser at 12 months duration in SMWT following TKR. Petterson *et al.* [9], in their study found that the study patients walked 150 meters further 12 months post-operatively, our study participants performed additional 219 meters during SMWT, and these improvements might be attributed to the increased strength of hip abductors so as to provide the lateral stability of the pelvis during walking.

Single leg stance on the surgical side had decreased at one month in both the groups and the HAS group has shown a statistically and clinically significant differences at 3 months and 1 year duration. Piva *et al.* [15], observed a 25% difference in SLS between groups at 6 months following TKR on the surgical side in comparison our study between the group analysis has shown a mean difference of 3.9 Sec at 1 year. This could possibly due to the strength increments of the hip abductor muscle, augmentations of strength may enhance the stabilization of trunk and hip during standing by transferring the forces from the stance leg to the pelvis with an improved the femoral pelvic stability. In normal, healthy individuals, an average of 5° of hip adduction angle was reported from double leg stance to single leg stance [46] since the hip abductors are an important pelvic stabilizer by resisting the varus torque of femur and enhance the femoro-pelvic stability during SLS [17, 18].

The measurement of knee extension strength showed a significant difference for both the groups at 3 months and preserved until 1 year, and NPRS has shown significant changes at all periods in both the groups. The improvement in quadriceps strength and a reduction in pain intensity is imperative which could have possibly enhanced the better physical function following TKR.

Significant improvement in hip abductor strength at 3 months and 1 year in HAS group was seen when compared to

KS group. The participants with strong hip abductor strength walked further in SMWT at 1 year and performed better on SLS test at 3 months and 1-year duration. Considering the improvement observed in the HAS group it is most likely that the hip abductor would play a major role on the physical function measures following TKR. We recommend that the future randomized controlled trials should address the effectiveness of hip abductor strengthening following TKR with larger sample size.

CONCLUSION

Hip abductor strengthening showed superior improvements in single leg stance test at 3 months and one year duration. Participants with strong hip abductors walked further in six minute walk test at one year duration when compared to the knee strengthening alone. Thus hip abductor strengthening exercises could be the potential contributors for improving physical function following TKR.

APPENDIX-1

Rehabilitation Following Total Knee Replacement (Knee Strengthening -Control Group)

Postoperative Day 1

Bedside exercises: ankle pumps, quadriceps sets, gluteal sets, hip abduction (supine), short-arc quads, straight-leg raise (if able with knee brace on).

Bed mobility and transfer training (bed to/from chair) Gait training with assistive device on level surfaces and functional transfer training (eg, sit to/from stand, bed mobility).

Postoperative Day 2

Exercises for active ROM, active assisted ROM, Knee ROM heel slides. Strengthening exercises (eg, ankle pumps, quadriceps sets, gluteal sets, heel slides, straight-leg raises) 1-3 sets of 10 repetitions for all strengthening exercises, twice daily.

Postoperative Day 3- 7(or on Discharge to Rehabilitation Unit)

Progression of ROM with active assisted exercises as necessary. Progression of strengthening exercises to the patient's tolerance, 1-3 sets of 10 repetitions for all strengthening exercises twice/daily. Progression of ambulation distance. Progression of activities-of-daily-living training for discharge to home

Modalities

Ice 2-3 times per day, with lower extremity elevated for 20-30 min.

Day 8 – Week 4

Range of Motion

Active assisted ROM for knee flexion, sitting or supine, using other lower extremity to assist Knee extension stretch with manual pressure (in clinic) or weights (at home)

Strength

Quad sets, straight leg raises (without knee extension lag), hamstring curls (standing), sitting knee extension, terminal knee extensions from 45° to 0°, step-ups (5- to 15-cm block), wall slides to 45° knee flexion, 1-3 sets of 10 repetitions for all strengthening exercises.

Criteria for progression: exercises are to be progressed (eg, weights, step height, etc.) only when the patient can complete the exercise and maintain control through 3 sets of 10 repetitions.

Pain and Swelling

Ice and compression as needed.

Functional Activities

Ambulation training with assistive device, as appropriate, with emphasis on heel strike, push-off at toe-off, and

normal knee joint excursions.

Emphasis on heel strike, push-off at toe-off, and normal knee joint excursions when able to walk without assistive device.

Stair ascending and descending step over step when patient has sufficient concentric/eccentric strength.

Outpatient Physical Therapy Advanced Strengthening Program Semi-Independent Phase (week 5 - week 12).

**Supervised Physical Therapy Sessions of 2 Days a Week with Treatment Duration of 45-60 Minutes
Warm-up (10-15 min)**

Seated or supine knee AROM (flexion and extension). Alternated ankle dorsiflexion and plantar flexion. Passive knee extension stretch. Patellar mobilizations

Specific Strengthening (10-15 min), 1-3 sets of 10 Repetitions

Add 1 to 2-kg weights to the exercises for appropriate. Quadriceps sets. Straight-leg raises (goal to perform without a knee extension lag). Standing leg curls. Seated knee extension. Terminal knee extension from 45° to 0°

The specific strengthening exercises, performed in a supine or seated position, consisted of maximal isometric pain-free contractions (knee extensors and flexors), at different angles of knee flexion, and dynamic (concentric-eccentric) contractions against gravity. The isometric exercises were performed at multiple angles at 0, 45 and 90 degrees.

Functional Exercises (10-15 min)

Step-ups (forward and lateral), 5-15 cm, or climbing a flight of stairs with support. Sit-to-stands. Ambulation training with or without assistive device, as appropriate, with emphasis on heel strike, push-off at toe-off, and normal knee joint excursions. Stair ascending and descending step over step when patient has sufficient concentric/eccentric strength

Endurance Exercises (5-10 min)

Walking.

Cool-Down (5 min)

Ice and compression as needed, gentle stretching and ROM.

Incision Mobility

Soft tissue mobilization until incision moves freely over subcutaneous tissue.

Criteria for Progression

Voluntary quadriceps muscle control or 0° knee extension lag. AROM 0° to greater than 105° of knee flexion. Minimal to no pain and swelling. Exercises are to be progressed once the patient can complete 3 sets of 10 reps of the exercise correctly and feels maximally fatigued.

Participants performed exercises weekly once or twice (continue all exercises, 1-3 sets at 10-20 repetition) after 12 weeks till year. Exercise adherence was ensured through logbook and periodic telephone calls.

APPENDIX-2

Rehabilitation following Total Knee Replacement (Hip Abductor strengthening group)

Phase – I Inpatient Rehabilitation

Supervised Physical therapy sessions six days a week, twice a day each session last for 30- 45 minutes

Postoperative day 1

Bedsides exercises: ankle pumps, quadriceps sets, gluteal sets, hip abduction (supine), short-arc quads, straight-leg raise (if able with knee brace on).

Bed mobility and transfer training (bed to/from chair). Gait training with assistive device on level surfaces and functional transfer training (eg, sit to/from stand, bed mobility).

Postoperative day 2

Exercises for active ROM, active assisted ROM, Knee ROM heel slides.

Strengthening exercises (e.g., ankle pumps, quadriceps sets, gluteal sets, heel slides, straight-leg raises, supine hip abduction) 1-3 sets of 10 repetitions for all strengthening exercises, twice daily.

Postoperative day 3- 7(or on discharge to rehabilitation unit)

Progression of Hip abduction exercises in supine and standing. Progression of ROM with active assisted exercises as necessary. Progression of strengthening exercises to the patient's tolerance, 1-3 sets of 10 repetitions for all strengthening exercises twice/daily. Progression of ambulation distance. Progression of activities-of-daily-living training for discharge to home.

Modalities

Ice 2-3 times per day, with lower extremity elevated for 20-30 min.

Outpatient Physical Therapy Specific Strengthening Programme (Day 8- week 4)**Supervised Physical Therapy Sessions Four Days a Week, Single Session Last for 45-60 Minutes****Goals**

Increase range of motion (ROM). Decrease edema and pain. Gait training. Independence with activities of daily living (ADLs).

Exercises

Seated or supine knee active range of motion (AROM). Alternated ankle dorsiflexion and plantar flexion. Quadriceps sets, Straight-leg raise, Hamstring sets, seated knee extension. Supported single standing for balance. Repeated sit-to-stand transfer training. Ambulating with appropriate assistive device (walker).

Hip Abduction Exercises

Abduction in side lying, unilateral hip abduction performed in standing, isometric hip abductor strengthening exercise, Calm exercises for hip abductors.

Modalities

Ice 2-3 times per day, with lower extremity elevated for 20-30 min.

❖ Criteria for Progression to Phase III

AROM approaching 100-120° of knee flexion, minimal pain/swelling.

Independence in mobility in and out of the home.

Outpatient Physical Therapy Advanced Strengthening Program Semi-Independent Phase (week 5 - week 12)**Supervised Physical Therapy Sessions of 2 Days a Week with Treatment Duration of 45-60 Minutes Warm-up (10-15 min)**

Seated or supine knee AROM (flexion and extension). Alternated ankle dorsiflexion and plantar flexion. Passive knee extension stretch. Patellar mobilizations.

Specific Strengthening (10-15 min), 1-3 Sets of 10 Repetitions

Add 1 to 2-kg weights to the exercises for appropriate. Hip abductors with weights in supine and standing. Quadriceps sets. Straight-leg raises (goal to perform without a knee extension lag). Standing leg curls. Seated knee extension. Standing terminal knee extension from 45° to 0°.

The specific strengthening exercises, performed in a supine or seated position, consisted of maximal isometric pain-free contractions (knee extensors and flexors), at different angles of knee flexion, and dynamic (concentric-eccentric) contractions against gravity (hip abductors). The isometric exercises were performed at multiple angles, because strength gains are known to be specific to the trained positions.

Hip Abduction Exercises

Abduction in side lying, abduction in standing, side walking, isometric contraction of hip abductors. Unilateral hip abduction performed in side lying with the use of ankle cuff weights (3 sets of 10 at a 10 RM resistance). Unilateral hip abduction performed in standing with the use of ankle weights (3 sets of 10 at a 10 RM resistance). Standing wall isometric hip abduction. Calm exercises with resistance on theraband. Performed in unipedal stance with the opposite limb in 90 degrees of hip and knee flexion (3 sets of 10 with 5 second holds).

Functional Exercises (10-15 min)

Step-ups (forward and lateral), 5-15 cm, or climbing a flight of stairs with support. Sit-to-stands. Ambulation training with or without assistive device, as appropriate, with emphasis on heel strike, push-off at toe-off, and normal knee joint excursions. Emphasis on heel strike, push-off at toe-off, and normal knee joint excursions when able to walk without assistive device. Stair ascending and descending step over step when patient has sufficient concentric/eccentric strength.

Endurance Exercises (5-10 min)

Walking.

Cool-down (5 min)

Ice and compression as needed, Gentle stretching and ROM.

Incision Mobility

Soft tissue mobilization until incision moves freely over subcutaneous tissue.

Criteria for Progression

Voluntary quadriceps muscle control or 0° knee extension lag. AROM 0° to greater than 105° of knee flexion. Minimal to no pain and swelling. Exercises are to be progressed once the patient can complete 3 sets of 10 reps of the exercise correctly and feels maximally fatigued. Participants will perform exercises weekly once or twice (continue all exercises, 1-3 sets at 10-20 repetition) after 12 weeks. They were asked to continue their home based exercise and the exercise adherence was ensured through logbook and periodic telephone calls.

CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

Declared none.

REFERENCES

- [1] Dieppe P, Basler HD, Chard J, *et al.* Knee replacement surgery for osteoarthritis: Effectiveness, practice variations, indications and possible determinants of utilization. *Rheumatology (Oxford)* 1999; 38(1): 73-83. [<http://dx.doi.org/10.1093/rheumatology/38.1.73>] [PMID: 10334686]
- [2] Gossec L, Hawker G, Davis AM, *et al.* OMERACT/OARSI initiative to define states of severity and indication for joint replacement in hip and knee osteoarthritis. *J Rheumatol* 2007; 34(6): 1432-5. [PMID: 17552070]
- [3] Franklin PD, Li W, Ayers DC. The Chitranjan Ranawat Award: functional outcome after total knee replacement varies with patient attributes. *Clin Orthop Relat Res* 2008; 466(11): 2597-604. [<http://dx.doi.org/10.1007/s11999-008-0428-8>] [PMID: 18810570]
- [4] Farquhar S, Snyder-Mackler L. The Chitranjan Ranawat Award: The nonoperated knee predicts function 3 years after unilateral total knee arthroplasty. *Clin Orthop Relat Res* 2010; 468(1): 37-44. [<http://dx.doi.org/10.1007/s11999-009-0892-9>] [PMID: 19472024]
- [5] Noble PC, Gordon MJ, Weiss JM, Reddix RN, Condit MA, Mathis KB. Does total knee replacement restore normal knee function? *Clin*

- Orthop Relat Res 2005; (431): 157-65.
[<http://dx.doi.org/10.1097/01.blo.0000150130.03519.fb>] [PMID: 15685070]
- [6] Walsh M, Woodhouse LJ, Thomas SG, Finch E. Physical impairments and functional limitations: A comparison of individuals 1 year after total knee arthroplasty with control subjects. *Phys Ther* 1998; 78(3): 248-58.
[<http://dx.doi.org/10.1093/ptj/78.3.248>] [PMID: 9520970]
- [7] Bade MJ, Kohrt WM, Stevens-Lapsley JE. Outcomes before and after total knee arthroplasty compared to healthy adults. *J Orthop Sports Phys Ther* 2010; 40(9): 559-67.
[<http://dx.doi.org/10.2519/jospt.2010.3317>] [PMID: 20710093]
- [8] Mizner RL, Petterson SC, Snyder-Mackler L. Quadriceps strength and the time course of functional recovery after total knee arthroplasty. *J Orthop Sports Phys Ther* 2005; 35(7): 424-36.
[<http://dx.doi.org/10.2519/jospt.2005.35.7.424>] [PMID: 16108583]
- [9] Petterson SC, Mizner RL, Stevens JE, *et al.* Improved function from progressive strengthening interventions after total knee arthroplasty: a randomized clinical trial with an imbedded prospective cohort. *Arthritis Rheum* 2009; 61(2): 174-83.
[<http://dx.doi.org/10.1002/art.24167>] [PMID: 19177542]
- [10] Moffet H, Collet J-P, Shapiro SH, Paradis G, Marquis F, Roy L. Effectiveness of intensive rehabilitation on functional ability and quality of life after first total knee arthroplasty: A single-blind randomized controlled trial. *Arch Phys Med Rehabil* 2004; 85(4): 546-56.
[<http://dx.doi.org/10.1016/j.apmr.2003.08.080>] [PMID: 15083429]
- [11] Minns Lowe CJ, Barker KL, Dewey M, Sackley M. Effectiveness of physiotherapy exercise after knee arthroplasty for osteoarthritis: systematic review and meta-analysis of randomised controlled trials *BMJ* 2007; 335: 812.
- [12] Artz N, Elvers KT, Lowe CM, Sackley C, Jepson P, Beswick AD. Effectiveness of physiotherapy exercise following total knee replacement: systematic review and meta-analysis. *BMC Musculoskelet Disord* 2015; 16: 15.
[<http://dx.doi.org/10.1186/s12891-015-0469-6>] [PMID: 25886975]
- [13] Arnold JB, Walters JL, Ferrar KE. Does Physical Activity Increase After Total Hip or Knee Arthroplasty for Osteoarthritis? A Systematic Review. *J Orthop Sports Phys Ther* 2016; 46(6): 431-42.
[<http://dx.doi.org/10.2519/jospt.2016.6449>] [PMID: 27117726]
- [14] Deasy M, Leahy E, Semciw AI. Hip Strength Deficits in People With Symptomatic Knee Osteoarthritis: A Systematic Review With Meta-analysis. *J Orthop Sports Phys Ther* 2016; 46(8): 629-39.
[<http://dx.doi.org/10.2519/jospt.2016.6618>] [PMID: 27374011]
- [15] Piva SR, Teixeira PE, Almeida GJ, *et al.* Contribution of hip abductor strength to physical function in patients with total knee arthroplasty. *Phys Ther* 2011; 91(2): 225-33.
[<http://dx.doi.org/10.2522/ptj.20100122>] [PMID: 21212373]
- [16] Alnahdi AH, Zeni JA, Snyder-Mackler L. Hip abductor strength reliability and association with physical function after unilateral total knee arthroplasty: a cross-sectional study. *Phys Ther* 2014; 94(8): 1154-62.
[<http://dx.doi.org/10.2522/ptj.20130335>] [PMID: 24652473]
- [17] Neumann DA. Kinesiology of the hip: A focus on muscular actions. *J Orthop Sports Phys Ther* 2010; 40(2): 82-94.
[<http://dx.doi.org/10.2519/jospt.2010.3025>] [PMID: 20118525]
- [18] Neumann DA. *Kinesiology of the Musculoskeletal System: Foundations for Rehabilitation*. Philadelphia, PA: Elsevier Science, Health Science Division 2010.
- [19] Semciw AI, Pizzari T, Murley GS, Green RA. Gluteus medius: An intramuscular EMG investigation of anterior, middle and posterior segments during gait. *J Electromyogr Kinesiol* 2013; 23(4): 858-64.
[<http://dx.doi.org/10.1016/j.jelekin.2013.03.007>] [PMID: 23587766]
- [20] Brown M, Sinacore DR, Binder EF, Kohrt WM. Physical and performance measures for the identification of mild to moderate frailty. *J Gerontol A Biol Sci Med Sci* 2000; 55(6): M350-5.
[<http://dx.doi.org/10.1093/gerona/55.6.M350>] [PMID: 10843356]
- [21] Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthrosis. *Ann Rheum Dis* 1957; 16(4): 494-502.
[<http://dx.doi.org/10.1136/ard.16.4.494>] [PMID: 13498604]
- [22] Bellamy N, Kirwan J, Boers M, *et al.* Recommendations for a core set of outcome measures for future phase III clinical trials in knee, hip, and hand osteoarthritis. Consensus development at OMERACT III. *J Rheumatol* 1997; 24(4): 799-802.
[PMID: 9101522]
- [23] Terwee CB, Mokkink LB, Steultjens MP, Dekker J. Performance-based methods for measuring the physical function of patients with osteoarthritis of the hip or knee: a systematic review of measurement properties. *Rheumatology (Oxford)* 2006; 45(7): 890-902.
[<http://dx.doi.org/10.1093/rheumatology/kei267>] [PMID: 16461441]
- [24] Kennedy DM, Stratford PW, Wessel J, Gollish JD, Penney D. Assessing stability and change of four performance measures: a longitudinal study evaluating outcome following total hip and knee arthroplasty. *BMC Musculoskelet Disord* 2005; 6: 3.
[<http://dx.doi.org/10.1186/1471-2474-6-3>] [PMID: 15679884]
- [25] Stratford PW, Kennedy D, Pagura SM, Gollish JD. The relationship between self-report and performance-related measures: questioning the content validity of timed tests. *Arthritis Rheum* 2003; 49(4): 535-40.

- [http://dx.doi.org/10.1002/art.11196] [PMID: 12910560]
- [26] Stratford PW, Kennedy DM. Performance measures were necessary to obtain a complete picture of osteoarthritic patients. *J Clin Epidemiol* 2006; 59(2): 160-7.
[http://dx.doi.org/10.1016/j.jclinepi.2005.07.012] [PMID: 16426951]
- [27] Wright AA, Hegedus EJ, Baxter GD, Abbott JH. Measurement of function in hip osteoarthritis: developing a standardized approach for physical performance measures. *Physiother Theory Pract* 2011; 27(4): 253-62.
[http://dx.doi.org/10.3109/09593985.2010.491150] [PMID: 20649479]
- [28] Jordan KP, Wilkie R, Muller S, Myers H, Nicholls E. Measurement of change in function and disability in osteoarthritis: current approaches and future challenges. *Curr Opin Rheumatol* 2009; 21(5): 525-30.
[http://dx.doi.org/10.1097/BOR.0b013e32832e45fc] [PMID: 19525848]
- [29] Dobson F, Hinman RS, Roos EM, *et al.* OARSIS recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. *Osteoarthritis Cartilage* 2013; 21(8): 1042-52.
[http://dx.doi.org/10.1016/j.joca.2013.05.002] [PMID: 23680877]
- [30] Podsiadlo D, Richardson S. The timed Up & Go: A test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991; 39(2): 142-8.
[http://dx.doi.org/10.1111/j.1532-5415.1991.tb01616.x] [PMID: 1991946]
- [31] Kennedy D, Stratford PW, Pagura SM, Walsh M, Woodhouse LJ. Comparison of gender and group differences in self-report and physical performance measures in total hip and knee arthroplasty candidates. *J Arthroplasty* 2002; 17(1): 70-7.
[http://dx.doi.org/10.1054/arth.2002.29324] [PMID: 11805928]
- [32] Boonstra MC, De Waal Malefijt MC, Verdonchot N. How to quantify knee function after total knee arthroplasty? *Knee* 2008; 15(5): 390-5.
[http://dx.doi.org/10.1016/j.knee.2008.05.006] [PMID: 18620863]
- [33] Botolfsen P, Helbostad JL, Moe-Nilssen R, Wall JC. Reliability and concurrent validity of the Expanded Timed Up-and-Go test in older people with impaired mobility. *Physiother Res Int* 2008; 13(2): 94-106.
[http://dx.doi.org/10.1002/pri.394] [PMID: 18288773]
- [34] Parent E, Moffet H. Comparative responsiveness of locomotor tests and questionnaires used to follow early recovery after total knee arthroplasty. *Arch Phys Med Rehabil* 2002; 83(1): 70-80.
[http://dx.doi.org/10.1053/apmr.2002.27337] [PMID: 11782835]
- [35] Maly MR, Costigan PA, Olney SJ. Determinants of self-report outcome measures in people with knee osteoarthritis. *Arch Phys Med Rehabil* 2006; 87(1): 96-104.
[http://dx.doi.org/10.1016/j.apmr.2005.08.110] [PMID: 16401446]
- [36] Kreibich DN, Vaz M, Bourne RB, *et al.* What is the best way of assessing outcome after total knee replacement? *Clin Orthop Relat Res* 1996; (331): 221-5.
[http://dx.doi.org/10.1097/00003086-199610000-00031] [PMID: 8895642]
- [37] Ouellet D, Moffet H. Locomotor deficits before and two months after knee arthroplasty. *Arthritis Rheum* 2002; 47(5): 484-93.
[http://dx.doi.org/10.1002/art.10652] [PMID: 12382296]
- [38] Guyatt GH, Pugsley SO, Sullivan MJ, *et al.* Effect of encouragement on walking test performance. *Thorax* 1984; 39(11): 818-22.
[http://dx.doi.org/10.1136/thx.39.11.818] [PMID: 6505988]
- [39] Curb JD, Ceria-Ulep CD, Rodriguez BL, *et al.* Performance-based measures of physical function for high-function populations. *J Am Geriatr Soc* 2006; 54(5): 737-42.
[http://dx.doi.org/10.1111/j.1532-5415.2006.00700.x] [PMID: 16696737]
- [40] Cesari M, Kritchevsky SB, Newman AB, *et al.* Added value of physical performance measures in predicting adverse health-related events: results from the Health, Aging And Body Composition Study. *J Am Geriatr Soc* 2009; 57(2): 251-9.
[http://dx.doi.org/10.1111/j.1532-5415.2008.02126.x] [PMID: 19207142]
- [41] Simonsick EM, Newman AB, Nevitt MC, *et al.* Measuring higher level physical function in well-functioning older adults: expanding familiar approaches in the Health ABC study. *J Gerontol A Biol Sci Med Sci* 2001; 56(10): M644-9.
[http://dx.doi.org/10.1093/gerona/56.10.M644] [PMID: 11584038]
- [42] Piva SR, Gil AB, Almeida GJ, DiGioia AM III, Levison TJ, Fitzgerald GK. A balance exercise program appears to improve function for patients with total knee arthroplasty: a randomized clinical trial. *Phys Ther* 2010; 90(6): 880-94.
[http://dx.doi.org/10.2522/ptj.20090150] [PMID: 20378678]
- [43] Edwards RH, McDonnell M. Hand-held dynamometer for evaluating voluntary-muscle function. *Lancet* 1974; 2(7883): 757-8.
[http://dx.doi.org/10.1016/S0140-6736(74)90947-7] [PMID: 4143018]
- [44] Kawaguchi JK, Babcock G. Validity and reliability of handheld dynamometric strength assessment of hip extensor and abductor muscles. *Ath Train Sports Health Care* 2010; 1: 11-7.
[http://dx.doi.org/10.3928/19425864-20101221-04]
- [45] Staehli S, Glatthorn JF, Casartelli N, Maffiuletti NA. Test-retest reliability of quadriceps muscle function outcomes in patients with knee osteoarthritis. *J Electromyogr Kinesiol* 2010; 20(6): 1058-65.
[http://dx.doi.org/10.1016/j.jelekin.2010.07.006] [PMID: 20688529]

- [46] DiMattia M, Livengood A, Uhl T, Mattaclola C, Malone T. What are the validity of the single-leg-squat test and its relationship to hip-abduction strength? *J Sport Rehabil* 2005; 14: 108-23.
[<http://dx.doi.org/10.1123/jsr.14.2.108>]

© 2017 Harikesavan *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.