

Hyperbilirubinemia as a predictor of gangrenous/perforated appendicitis: a prospective study

Poras Chaudhary, Ajay Kumar, Neeraj Saxena, Upendra C. Biswal

Postgraduate Institute of Medical Education and Research, Dr Ram Manohar Lohia Hospital (Dr RMLH and PGIMER), New Delhi, India

Abstract

Background The aims of the study were: 1) to establish the role of hyperbilirubinemia as a new diagnostic tool to predict gangrenous/perforated appendicitis; and 2) to compare other variables such as age, duration of symptoms, clinical profile, white blood cell count, C-reactive protein and ultrasound in a similar role.

Methods This is a prospective study conducted on 50 consecutive cases of acute appendicitis admitted to the emergency ward. These were subjected to investigations to support the diagnosis. These cases were also subjected to liver function tests and clinical diagnosis was confirmed perioperatively and post-operatively by histopathological examination. Their clinical and investigative data were compiled and analyzed. Statistical analysis was performed using either chi square test or fisher's exact test. The level of significance was set at $P < 0.05$.

Results Total serum bilirubin including both direct and indirect was found to be significantly increased in case of acute suppurative appendicitis. Serum bilirubin was much higher ($P < 0.000$) in cases of gangrenous/perforated appendicitis.

Conclusion Serum bilirubin is an important adjunct in diagnosing the presence of gangrenous/perforated appendicitis.

Keywords Acute suppurative appendicitis, gangrenous/perforated appendicitis, hyperbilirubinemia
Ann Gastroenterol 2013; 26 (4): 325-331

Introduction

Appendicitis is one of the commonest causes of abdominal pain requiring emergency surgery. Often, it is difficult to reach a proper diagnosis. There may not be classical symptoms and signs of appendicitis. Different clinical signs and symptoms always mimic the diagnosis of acute appendicitis, as there are a number of causes leading to pain in right iliac fossa particularly in female patients. Diagnosing acute appendicitis clinically still remains a common surgical problem. Accurate diagnosis can be aided by additional testing or expectant management or

both. These might delay laparotomy and lead to appendiceal perforation with increased morbidity and hospital stay [1-3]. A safe alternative seems to be appendectomy as soon as the condition is suspected, a strategy that increases the number of unnecessary appendectomies [4,5]. A timelier and more accurate diagnosis has been attempted by the employment of additional laboratory tests [6-11], scoring systems [12-15], ultrasound imaging [16,17], computed tomography (CT) scan [18,19], scintigraphy [20], MRI [21], and laparoscopy [22-24]. None of these methods stands alone as they all come in support of, and are secondary to a primary clinical assessment.

Hyperbilirubinemia is a new diagnostic tool for perforation of appendix. Hyperbilirubinemia is the result of imbalance between production and excretion of bilirubin by the liver. It may be because of hepatocellular, cholestatic or hemolytic diseases. Liver receives blood mainly through the portal venous system, which receives blood from abdominal organs. Portal blood carries nutrients and other substances absorbed from gut including bacteria and its product (toxins). In a small percentage, even in normal healthy people, bacteria are found in portal blood. It is commonly cleared by detoxification and immunological action of the reticuloendothelial system of the liver that acts as first-line defense in clearing toxic substances, bacteria and its products. But when bacterial load

General Surgery, Postgraduate Institute of Medical Education and Research, Dr Ram Manohar Lohia Hospital (Dr RMLH and PGIMER), New Delhi, India

Funding: Financial support was provided by Dr Ram Manohar Lohia Hospital, New Delhi, India

Conflict of Interest: None

Correspondence to: Poras Chaudhary, M.S., General Surgery, Postgraduate Institute of Medical Education and Research, Dr Ram Manohar Lohia Hospital (Dr RMLH and PGIMER), New Delhi, India, Tel.: +98 914 47358, e-mail: drporaschaudhary@yahoo.com

Received 31 March 2013; accepted 13 June 2013

overwhelms the Kupffer cell function, it may cause dysfunction or damage to hepatocytes (liver parenchyma). It reflects a rise in serum bilirubin (SB) alone or in combination with liver enzymes depending upon the type, severity and site of the lesion. Recently, another substance known as cytokines e.g. interleukin (IL)-6, tumor necrosis factor (TNF), has also been considered to be responsible for depressed excretory function of the liver and may lead to increase in SB levels without a rise in liver enzymes.

The association between the elevated SB levels and the variety of infectious diseases has been noted in few studies [25-27]. This finding most commonly occurs in neonates with gram negative bacterial infection. It has also been described in patients with severe intra-abdominal infection. The pathogenesis is thought to be because of bacteremia or endotoxemia causing impaired excretion of bilirubin from the bile canaliculi. There are only a few reports in the literature that describe the finding of hyperbilirubinemia in patients with either severe post-operative infection after appendectomy or with complicated appendicitis.

The present study has been designed to evaluate the association between hyperbilirubinemia in cases of acute appendicitis and its complications. The significance of other parameters such as age, duration of symptoms, total leucocyte count (TLC), ultrasonography, Alvarado score and C-reactive protein (CRP) has also been evaluated in these cases.

The establishment of a possible role of hyperbilirubinemia as a predictor of gangrenous/perforated appendicitis has been stressed so that SB levels upon admission can be used in conjunction with other diagnostic tests such as ultrasonography and CT to help determine the presence of perforation and aid in proper clinical management.

Materials and methods

Study design

This was a single-center cohort, prospective study conducted by the department of Surgery from October 2008 to April 2010. 50 consecutive cases of acute appendicitis admitted in surgical unit I were recruited in the study. These were subjected to investigation to support the diagnosis. Investigation included total leukocytes count, differential leukocytes count, ultrasound, liver function tests (LFT) and CRP. The protocol was approved by the ethical review board of the Institute. According to the principles of the declaration of Helsinki 1975, written, informed consent was obtained from all participants.

Participants

The patients were selected from those attending the emergency department at the hospital. The age of patients varied from 15 to 64 years with most of the patients falling

within the age range from 15-24 years. Inclusion criteria were: patients of 15 years of age and above scheduled for appendectomy for acute appendicitis at the emergency unit of our hospital. Exclusion criteria were: appendectomy performed incidentally or for other indications; age below 15 years; patients with appendicular lump; history of alcoholic liver disease; hemolytic or liver diseases associated with hyperbilirubinemia; history of gastrointestinal or hepatopancreatobiliary malignancy in the past. 64 patients were enrolled, 14 excluded. Out of these 14 patients, 3 had presence of liver abscesses, 4 had multiple large mesenteric lymph nodes, 2 had the presence of peritoneal tubercles suggestive of tuberculosis and 5 patients had multiple liver nodules suggestive of cirrhosis. Patients were clinically evaluated by detailed history, routine examination on initial contact with patients and the following investigations were done; complete hemogram, liver function test (LFT) (Table 1), kidney function test (KFT), blood sugar, CRP, serum proteins, X-ray chest posteroanterior (PA) view these investigations blood samples were drawn within half an hour of presentation in the emergency department and radiological investigations were done within 2 h of admission. Determination of SB was done with photometric testing using 2,4-dichloroaniline. This method is based on the principle that in acidic solution, direct bilirubin forms a red colored azocompound with diazotized 2,4-dichloroaniline. A specific mixture of detergents enables a safe determination of the total bilirubin.

Routine LFT results were compared with laboratory reference values given in Table 1. Alvarado scoring criteria is given in Table 2. These cases were operated and clinical diagnosis was confirmed per-operatively and post-operatively by histopathological examination. Final histopathological examination was considered as a gold standard for diagnosing and categorizing patients as having normal appendix, acute appendicitis and acute appendicitis with perforation and/or gangrene. Their clinical and investigative data were compiled and analyzed. Based on histopathological examination patients were categorized as negative (acute appendicitis without perforation or gangrene) and positive (acute appendicitis with perforation and/or gangrene).

Statistical analysis

Table 1 Reference range of serum bilirubin and liver enzymes

Test	Normal range
Serum Bilirubin	
Total	1.2 mg/dL
Direct	0.2 mg/dL
Liver Enzymes	
ALT	≤50 U/L
AST	≤50 U/L
ALP	50-300 U/L

ALT, alanine aminotransferase; AST, aspartate aminotransferase ALP, alkaline phosphatase

Table 2 Alvarado scoring system

Features	Score
Symptoms	
Migratory right iliac fossa pain	1
Nausea / Vomiting	1
Anorexia	1
Signs	
Right iliac fossa tenderness	2
Fever >37.30 °C	1
Rebound tenderness in right iliac fossa	1
Laboratory tests	
Leukocytosis (>10000/ μ L)	2
Neutrophilic shift to the left >75%	1
Total score	10

Statistical analysis was performed by the SPSS program for Windows, version 17.0. Continuous variables are presented as mean \pm SD, and categorical variables are presented as absolute numbers and percentage. Data were checked for normality before statistical analysis. Normally distributed continuous variables were compared using the unpaired t test, whereas the Mann-Whitney U test was used for those variables that were not normally distributed. Categorical variables were analyzed using either the chi square test or Fisher's exact test. A receiver operating characteristics analysis was calculated to determine optimal cut-off values for SB, TLC, CRP, aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP) and Alvarado score. The area under the curve, the sensitivity, and the specificity was calculated to analyze the diagnostic value of all these markers. For all statistical tests, a P value less than 0.05 was considered statistically significant.

Results

General characteristics

Of 50, 40 cases were male and 10 female. Their age ranged from 15-60 years. The average was 27.1 years (Table 3). Duration of symptoms ranged from 1-5 days. Among 50 cases clinically diagnosed as acute appendicitis preoperatively, 42 had acute appendicitis, 3 had gangrenous appendix, 5 cases had perforated appendix and no patient had normal appendix.

Evaluation of LFT (Tables 4,5)

Among 50 cases, total SB (TSB) was raised in 38 (76%) cases whereas 12 (24%) cases had normal TSB level. TSB was more elevated in gangrenous and perforated appendix than

Table 3 Distribution of cases in different age groups (n = 50)

Age group	No. (%)
15-24	24 (48%)
25-34	17 (34%)
35-44	7 (14%)
45- 54	1 (2%)
55- 64	1 (2%)
Total	50 (100%)

those in acute appendicitis ($P < 0.001$). The mean of TSB in acute appendicitis cases without perforation or gangrene was 1.52 mg/dL and the mean of TSB in cases with perforated or gangrenous appendix was 3.62 mg/dL (ranged from 3.0- 5.0 mg/dL).

Liver enzyme ALT was within normal range in 40 (80%) cases, and minimally elevated (<2 times) in 10 (20%) cases. AST was within normal range in 45 (90%) cases, minimally elevated (<2 times) in 4(8%) cases, and moderately elevated (>2 times) in 1 (2%) case. ALP was within normal range in 46 (92%) cases, minimally elevated (<2 times) in 3 (6%) cases, moderately elevated (>2 times) in 1 (2%) case.

Table 4 Distribution of the cases according to level of total serum bilirubin (SB) (n= 50) and histological examination

Histopathology	Total SB		Total
	<1.2 mg/dL	>1.2 mg/dL	
	No. (%)	No. (%)	
Acute Appendicitis	12 (24%)	30 (60%)	42 (84%)
Gangrenous Appendix	Nil	3 (6%)	3 (6%)
Perforated Appendix	Nil	5 (10%)	5 (10%)
Normal Appendix	Nil	Nil	Nil
Total	12 (24%)	38 (76%)	50 (100%)
Histopathology	Frequency		%
Negative (acute appendicitis with no perforation or gangrene)	42		84%
Positive (acute appendicitis with perforation and/or gangrene)	8		16%
Total	50		100%

Table 5 Distribution of liver function test according to alanine aminotransferase (ALT) (n= 50), aspartate aminotransferase (AST) and alkaline phosphatase

	Acute appendicitis	Gangrenous appendix	Perforated appendix	Normal appendix	Total
Alaline aminotransferase (Normal range 15-50 U/L)					
Normal	35 (70%)	3 (6%)	2 (4%)	Nil	40 (80%)
Minimally elevated (<1 times - <2 times)	7 (14%)	Nil	3 (6%)	Nil	10 (20%)
Moderately elevated (>2 times)	Nil	Nil	Nil	Nil	Nil
Total	42 (84%)	3 (6%)	5 (10%)	Nil	50 (100%)
Aspartate aminotransferase (Normal range 15-50 U/L)					
Normal	39 (78%)	3 (6%)	3 (6%)	Nil	45 (90%)
Minimally elevated (<1 times - <2 times)	3 (6%)	Nil	1 (2%)	Nil	4 (8%)
Moderately elevated (>2 times)	Nil	Nil	1 (2%)	Nil	1 (2%)
Total	42 (84%)	3 (6%)	5 (10%)	Nil	50 (100%)
Alkaline phosphatase (Normal range 50-300 U/L)					
Normal	39 (78%)	5 (10%)	2 (4%)	Nil	46 (92%)
Minimally elevated (<1 times - <2 times)	2 (4%)	Nil	1 (2%)	Nil	3 (6%)
Moderately elevated (>2 times)	1 (2%)	Nil	Nil	Nil	1 (2%)
Total	42	03	05	Nil	50

Alvarado score (Table 6)

Alvarado score was calculated based on the criteria given in Table 2 and P value for Alvarado score was 0.01.

Comparing different parameters

Comparison of different parameters- SB, TLC, liver enzymes, Alvarado score, CRP and duration of symptoms and their sensitivity, specificity, positive and negative predictive value with the relevant P values are shown in Tables 7 and 8.

Table 6 Distribution of Alvarado score

Alvarado score	No of cases
10	1 (2%)
9	5 (10%)
8	10 (20%)
7	18 (36%)
6	7 (14%)
5	1 (2%)
4	2 (4%)
3	2 (4%)
2	4 (8%)

Discussion

In this study of 50 patients, hyperbilirubinemia was found in 30 of 42 patients with acute suppurative appendicitis and in all 8 patients with gangrenous/perforated appendicitis. This hyperbilirubinemia was mixed in type (both conjugated and unconjugated) in most of the patients and at the same time there was no elevation or minimal elevation (<100 U/L) in ALT and AST in most of the cases. Similarly, ALP was either within the normal range or was minimal to moderately elevated.

For gangrenous/perforated appendicitis, the P-value of SB was <0.001, specificity 92.9%, sensitivity 100%, positive predictive value 72.7% and negative predictive value was 100%.

The level of SB was higher than 3 mg/dL in cases of gangrenous/perforated appendicitis while in cases with acute appendicitis it was lower than 3 mg/dL (P<0.05). Broadly, we can say that it was predominantly isolated hyperbilirubinemia in the majority of cases. These findings are almost similar to another reported study [28]. Since these findings were documented at the time of admission, it is unlikely that liver injury because of anesthetic agents, blood transfusion, or medication was the cause of elevated bilirubin levels. Moreover, as per our exclusion criteria patients with alcoholic liver disease, viral hepatitis, hemolytic or congenital liver diseases were excluded from the study.

Table 7 P-value chart of different parameters

	Negative		Positive		P value
	Mean±SD	Min-Max	Mean±SD	Min-max	
Age	26.93±9.33	15-60	28.63±8.83	15-40	0.466
Hb	13.85±1.11	12-17	13.88±0.99	13-15	0.958
TLC	11690.48±4132.49	6200-24000	12162.50±4624.45	8100-22000	0.822
Serum bilirubin	1.52±0.59	0.6-3.0	3.62±0.70	3.0-5.0	<0.001
AST	28.02±15.88	10-90	37.13±28.83	16-102	0.551
ALT	33.83±17.21	14-83	46.38±18.90	25-72	0.047
ALP	178.52±137.97	50-847	189.13±164.58	48-584	0.0947
CRP	107.19±28.86	60-179	168.25±12.24	156-186	<0.001
Alvarado score	6.48±1.78	2-9	8.25±1.49	6-10	0.01
Duration	2.26±1.21	1-5	2.00±0.92	1-4	0.71

Hb, hemoglobin; TLC, total leukocyte count; AST, aspartate aminotransferase; ALT, alanine aminotransferase; ALP, alkaline phosphatase; CRP, C-reactive protein

The most likely explanation of the rise in SB is therefore circulating endotoxemia as a result of appendiceal infection. Utili *et al* [29-31] has shown with *in vitro* infusion of endotoxin into the isolated rat liver that there is a dose-dependent decrease in bile salt excretion from the liver and that it is possible that *Escherichia coli* endotoxin exerts direct damage at the cholangiolar level.

It was demonstrated by Sisson *et al* in 1971 [32] that in appendicitis mucosal ulceration occurs early and this facilitates invasion of bacteria into the muscularis propria of the appendix thereby causing classical acute suppurative appendicitis. Subsequent events lead to edema, elevated intraluminal pressure, and ischemic necrosis of mucosa, causing tissue gangrene and perforation [33,34]. This process is associated with progressive bacterial invasion most likely facilitated by bacterial cytotoxins. The number of organisms isolated from patients with gangrenous appendicitis is five times greater than those with acute suppurative appendicitis. Estrada *et al* [35] also found significantly higher peritoneal culture in patients with gangrenous/perforated appendicitis.

This elevated load of bacteria in appendicitis causes either direct invasion or translocation into the portal

venous system. Direct invasion of bacteria into the hepatic parenchyma interferes with the excretion of bilirubin into the bile canaliculi by a mechanism that is thought to be caused by the bacterial endotoxin and is biochemical in nature rather than obstructive.

Indirect evidence of bacterial translocation from inflamed gastrointestinal tract or peritonitis to the liver via the portal vein and the development of hepatitis and pyogenic liver abscess was observed by Dieulafoy [36]. Two classical findings were described: firstly, simultaneous inflammation of the intestine (e.g. appendix), peritoneum and development of pyogenic liver abscesses, and secondly, bacteriological similarities of the gastrointestinal tract and pyogenic liver abscesses. These bacteria commonly reach liver from intra-abdominal organs, commonly from the appendix.

Direct evidence of bacterial translocation from inflamed organs was observed in clinical and experimental studies. Recently, in one study, blood samples from the superior mesenteric vein in acute appendicitis showed bacteria in 38% of patients. These findings suggest that bacteria may transmigrate and produce portal bacteremia, hepatocellular dysfunction or pyogenic liver abscess. This low percentage of positive

Table 8 Sensitivity, specificity, positive and negative predictive value of different parameters, AUC-area under curve

	AUC	Cut off	Sensitivity	Specificity	PPV	NPV	P value
Serum bilirubin	0.997 (0.0-1.00)	2.1	100%	92.9%	72.7%	100%	<0.001
TLC	0.525 (0.307-0.704)	13500	37.5%	78.6%	25%	86.8%	0.379
CRP	0.973 (0.0-1.000)	149	100%	95.2%	80%	100%	<0.001
AST	0.567 (0.336-0.798)	28	50%	69%	23.5%	87.9%	0.419
ALT	0.723 (0.556-0.890)	35	75%	64.3%	28.6%	93.1%	0.056
ALP	0.507 (0.297-0.718)	172.5	87.5%	40.5%	21.9%	94.4%	0.231
Alvarado score	0.781 (0.556-1.000)	9.0	62.5%	95.2%	71.4%	93%	<0.001

AUC, area under the curve; PPV, positive predictive value; NPV, negative predictive value; TLC, total leukocyte count; CRP, C-reactive protein; AST, aspartate aminotransferase; ALT, alanine aminotransferase; ALP, alkaline phosphatase

blood cultures cannot explain hepatocellular dysfunction in the majority of cases. Thus, there must be other substances involved. It has been shown that liver dysfunction is caused by cytokines released from the gut due to injury/ inflammation. In a study [29,30], rats were subjected to intra-abdominal sepsis from cecal ligation and puncture and the following observations were made: 1) the small intestine is an important source of adrenomedullin release during poly microbial sepsis; 2) norepinephrine induced hepatocellular dysfunction in early sepsis, mediated by activation of α -2 adreno-receptors; and 3) TNF produces hepatocellular dysfunction despite normal cardiac output and hepatic microcirculation [37].

Thus, it is concluded that hepatocellular function is depressed during the early stage of sepsis despite the increased cardiac output and hepatic blood flow and decreased peripheral resistance. The depression of hepatocellular function in the early, hyper-dynamic stage of sepsis does not appear to be due to reduction in hepatic perfusion but is associated with elevated levels of circulating pro-inflammatory cytokines such as TNF and IL-6. Thus up regulation of TNF and/ or IL-6 may be responsible for producing hepatocellular dysfunction during the early hyper-dynamic stage of sepsis.

Our study shows that isolated hyperbilirubinemia without much elevation in the liver enzymes is a significant predictor of appendiceal perforation. This was demonstrated by a study by Estrada *et al* [35] and other studies [38,39] showing nearly a threefold risk of perforated appendicitis in patients with total bilirubin levels greater than 1 mg/dL. The other factors which we studied in this series were age, duration of symptoms, Alvarado score, total leukocyte count, ultrasonography, and CRP. P value was not significant in any of these criteria except CRP ($P < 0.001$).

The positive predictive value of SB and CRP was 72.7% and 80%, comparable to other published studies [40,41]. The negative predictive value of SB was 100% in our study as compared to 16% in the study by Khan *et al*. The negative predictive value of CRP was also 100% in our study. Therefore, in suspected cases of appendicitis elevation of SB/CRP can be used as a criterion to diagnose and manage acute appendicitis. Both sensitivity and specificity of elevated total SB level and CRP in acute appendicitis with perforation and/or gangrene is higher as compared to TLC and liver enzyme. This finding is similar to other reported studies [35,42,43].

This study also shows that the Alvarado scoring system is also of great value with a significant P value and comparable sensitivity and specificity.

Therefore, SB estimation, a simple cheap and easily available test in every laboratory, can be added to the routine investigation list of clinically suspected case of acute appendicitis for the confirmation of diagnosis. Since the rise in SB level was significantly higher in patients with appendiceal perforation, it has a definite predictive potential in these cases. Therefore, obtaining SB values upon admission can be used in conjunction with more modern diagnostic tests such as CT scan, ultrasonography to help determine the presence of perforation and thus aid in prompt clinical management.

Summary Box

What is already known:

- Diagnosing acute appendicitis clinically still remains a common surgical problem
- Accurate diagnosis can be aided by additional testing or expectant management or both

What the new findings are:

- Isolated hyperbilirubinemia without much elevation in liver enzymes has a predictive potential for the diagnosis of appendiceal perforation
- The risk of perforation is significantly higher in patients with acute appendicitis with total bilirubin levels greater than 2.1 mg/dL
- Serum bilirubin can be a significant addition to the list of routine investigations at the time of admission in cases of suspected appendicitis

References

1. Harrison MW, Lindner DJ, Campbell JR, Campbell TJ. Acute Appendicitis in children: factors affecting morbidity. *Am J Surg* 1984;**147**:605-610.
2. Savrin RA, Clatworthy HW Jr. Appendiceal rupture: a continuing diagnostic problem. *Paediatrics* 1979;**63**:36-43.
3. Scher KS, Coil JA. The continuing challenge of perforating appendicitis. *Surg Gynecol Obstet* 1980;**150**:535-538.
4. Karakas SP, Guelfguat M, Leonidas JC, Springer S, Singh SP. Acute appendicitis in children: comparison of clinical diagnosis with ultrasound and CT imaging. *Pediatr Radiol* 2000;**30**:94-98.
5. Hoffmann J, Rasmussen OO. Aids in the diagnosis of acute appendicitis. *Br J Surg* 1989;**76**:774-779.
6. Oruc MT, Kulah B, Ozozan O, et al. The value of 5-hydroxyl indole acetic acid measurement in spot urine in diagnosis of acute appendicitis. *East African Med J* 2004;**81**:40-41.
7. Taha AS, Grant V, Kelly RW. Urine analysis for interleukin-8 in the non-invasive diagnosis of acute and chronic inflammatory diseases. *Postgrad Med J* 2003;**79**:159-163.
8. Hallan S, Asberg A. The accuracy of C-reactive protein in diagnosing acute appendicitis – a meta analysis. *Scan J Clin Lab Invest* 1997;**57**:373-80.
9. Goodwin AT, Swift RI, Bartlett MJ, Fernando BS, Chadwick SJ. Can serum interleukin-6 levels predict the outcome of patients with right iliac fossa pain? *Ann R Coll Surg Engl* 1997;**79**:130-133.
10. Vermeulen B, Morabia A, Unger PF. Influence of white cell count on surgical decision making in patients with abdominal pain in the right lower quadrant. *Eur J Surg* 1995;**161**:483-486.
11. Albu E, Miller BM, Choi Y, Lakhnani S, Murthy RN, Gerst PH. Diagnostic value of C-reactive protein in acute appendicitis. *Dis Colon Rectum* 1994;**37**:49-51.
12. Christian F, Christian GP. A simple scoring system to reduce the negative appendectomy rate. *Ann R Coll Surg Engl* 1992;**74**:281-285.
13. Alvarado A. A practical score for the early diagnosis of acute

- appendicitis. *Ann Emerg Med* 1986;**15**:557-564.
14. Ohmann C, Franke C, Yang Q. Clinical benefit of a diagnostic score for appendicitis: results of a prospective interventional study. German Study Group of Acute Abdominal Pain. *Arch Surg* 1999;**134**:993-996.
 15. Zielke A, Sitter H, Rampp T, et al. Can diagnostic scoring systems help decision making in primary care of patients with suspected acute appendicitis? *Zielke Dtsch Med Wochenschr* 1999;**124**:545-550.
 16. Douglas CD, Macpherson NE, Davidson PM, Gani JS. Randomised controlled trial of ultrasonography in diagnosis of acute appendicitis, incorporating the Alvarado score. *BMJ* 2000;**321**:919-922.
 17. Rettenbacher T, Hollerweger A, Gritzmann N, et al. Appendicitis: should diagnostic imaging be performed if the clinical presentation is highly suggestive of the disease? *Gastroenterology* 2002;**123**:992-998.
 18. Walker S, Haun W, Clark J, McMillin K, Zeren F, Gilliland T. The value of limited computed tomography with rectal contrast in the diagnosis of acute appendicitis. *Am J Surg* 2000;**180**:450-455.
 19. Hong JJ, Cohn SM, Ekeh AP, Newman M, Salama M, Leblang SD. A prospective randomized study of clinical assessment versus computed tomography for the diagnosis of acute appendicitis. *Surg Infect (Larchmt)* 2003;**4**:231-239.
 20. Colak T, Gungor F, Ozugur S, et al. The value of 99mTc-HMPAO labelled white blood cell scintigraphy in acute appendicitis patients with an equivocal clinical presentation. *Eur J Nucl Med* 2001;**28**:575-580.
 21. Incesu L, Coskun A, Selcuk MB, Akan H, Sozubir S, Bernay F. Acute appendicitis: MR imaging and sonographic correlation. *Am J Roentgenol* 1997;**168**:316-318.
 22. Larsson PG, Henriksson G, Olsson M, et al. Laparoscopy reduces unnecessary appendectomies and improves diagnosis in fertile women. A randomized study. *Surg Endosc* 2001;**15**:200-202.
 23. Lamparelli MJ, Hoque HM, Pogson CJ, Ball AB. A prospective evaluation of the combined use of the modified Alvarado score with selective laparoscopy in adult females in the management of suspected appendicitis. *Ann R Coll Surg Engl* 2000;**82**:192-195.
 24. Bruwer F, Coetzer M, Warren BL. Laparoscopic versus open surgical exploration in premenopausal women with suspected acute appendicitis. *Afr J Surg* 2003;**41**:82-85.
 25. Johnson AM. Sepsis and jaundice. *Pediatrics* 1993;**91**:10-18.
 26. Miller DJ, Keeton DJ, Webber BL, Pathol FF, Saunders SJ. Jaundice in severe bacterial infection. *Gastroenterology* 1976;**71**:94-97.
 27. Whitehead MW, Hainsworth I, Kingham JG. The causes of obvious jaundice in South West Wales: perceptions versus reality. *Gut* 2001;**48**:409-413.
 28. Khan S. Evaluation of hyperbilirubinemia in acute inflammation of appendix: a prospective study of 45 cases. *KUMJ* 2008;**4**:281-289.
 29. Utili R, Abernathy CO, Zimmerman HJ. Cholestatic effects of *Escherichia coli* endotoxin on the isolated perfused rat liver. *Gastroenterology* 1976;**70**:248-253.
 30. Utili R, Abernathy CO, Zimmerman HJ. Studies on the effects of *E coli* endotoxin on canalicular bile formation in the isolated perfused rat liver. *J Lab Clin Med* 1977;**89**:471-478.
 31. Utili R, Abernathy CO, Zimmerman HJ. Endotoxin effects on the liver. *Life Sci* 1977;**20**:553-568.
 32. Sisson RG, Ahlvin RC, Harlow MC. Superficial mucosal ulceration and the pathogenesis of acute appendicitis. *Am J Surg* 1971;**122**:378-380.
 33. Bennion RS, Wilson SE, Serota AI, Williams RA. The role of gastrointestinal microflora in the pathogenesis of complications of mesenteric ischemia. *Rev Infect Dis* 1984;**6**:S132-S138.
 34. Bennion RS, Wilson SE, Williams RA. Early portal bacteremia in mesenteric ischemia. *Arch Surg* 1984;**119**:151-155.
 35. Estrada JJ, Petrosyan M, Barnhart J, et al. Hyperbilirubinemia in appendicitis: a new predictor of perforation. 2007;**11**:714-718.
 36. Dieulafoy G. Le foie appendiculaire: abcès du foi concutifs à l'appendicite. *Semin Med (Paris)* 1898;**18**:449.
 37. Franström TR, Hierholzer WJ Jr, LaBrecque DR. Frequency and characteristics of hyperbilirubinemia associated with bacteremia. *Rev Infect Dis* 1985;**7**:1-9.
 38. Bennion RS, Baron EJ, Thompson JE. The bacteriology of gangrenous and perforated appendicitis- revisited. *Ann Surg* 1990;**211**:165-171.
 39. Bennion RS, Thompson JE, Baron EJ, Finegold SM. Gangrenous and perforated appendicitis with peritonitis: treatment and bacteriology. *Clin Ther* 1990;**12**:31-44.
 40. Khan S. Elevated serum bilirubin in acute appendicitis: a new diagnostic tool. *KUMJ* 2008;**6**:161-165.
 41. Beltran MA, Mendez PE, Barrera RE, et al. Is hyperbilirubinemia in appendicitis a better predictor of perforation than C-reactive protein?- a prospective study. *Indian J Surg* 2009;**71**:265-272.
 42. Sand M, Bechara F, Holland-Letz T, Sand D, Mehnert G, Mann B. Diagnostic value of hyperbilirubinemia as a predictive factor for appendiceal perforation in acute appendicitis. *Am J Surg* 2009;**198**:193-198.
 43. Khan S. The diagnostic value of hyperbilirubinemia and total leukocyte count in the evaluation of acute appendicitis. *J Clin Diagn Res* 2009;**3**:1647-1652.