

Poblap

SEROLOGICAL STUDY OF HANTAVIRUS IN THE RODENT POPULATION OF NAKHON PATHOM AND NAKHON RATCHASIMA PROVINCES THAILAND

Narong Nitatpattana¹, Gilles/Chauvancy^{1,4}, Julien Dardaine^{1,4}, Thaval Poblap³, Kanittha Jumronsawat², Waraluk Tangkanakul⁵, Duangporn Poonsuksombat⁶, Sutee Yoksan¹ and Jean-Paul/Gonzalez^{1,4}

¹Center for Vaccine Development, Institute of Science and Technology for Research and Development, Mahidol University, Thailand; ²ASEAN Institute for Health Development, Mahidol University, Thailand; ³Nakhon Pathom Provincial Health Office, Nakhon Pathom Province, Thailand; ⁴Institut de Recherche pour le Développement (IRD), France; ⁵Department of Communicable Diseases Control, Ministry of Public Health, Nonthaburi, Thailand; ⁶Armed Forces Research Institute of Medical Sciences, Bangkok, Thailand

Abstract. A serological survey has been carried out to detect evidence of hantavirus infection in rodents from two provinces of Thailand. This study aimed to examine virus antibody in 354 rodents trapped among 6 different villages of Nakhon Pathom Province (February-March, 1998) and in 326 rodents trapped among 14 villages of Nakhon Ratchasima Province (August-October, 1998). Seroprevalence among rodents from Nakhon Pathom Province (2.3%), was mostly found in *Rattus norvegicus* (3.8%) and *Bandicota indica* (2.6%). In Nakhon Ratchasima Province seroprevalence (4.0%) was mostly in *Bandicota indica* (19.1%) and *Rattus exulans* (3.5%).

INTRODUCTION

Hantaviruses are single-stranded, negative-sense RNA viruses, and members of the Bunyaviridae family. They cause zoonotic diseases which are transmitted to humans by contact with infected rodents. Hantaviruses are associated to rodents of the Muroides superfamily. Muroids are vectors and reservoirs of virus, with one species as a primary reservoir for one virus type. Rodent infection occurs through inhalation of infectious aerosols generated from rodent secretions and excretions, leading to a short, transient viremia (Nuzum *et al*, 1988). Virus persists in rodents despite high-titer antibodies, and disseminates to various tissues, particularly lung, salivary gland and kidney, resulting in a lifelong but asymptomatic infection (Lee *et al*, 1981; Lee, 1982; Yanagihara *et al*, 1985). Horizontal spread of virus occurs through exposure to contaminated nesting material, grooming behavior; interspecies wounding and biting are also important factors in virus maintenance within the enzootic cycle (Glass *et al*, 1988; Yanagihara and Gadjusek, 1988). Vertical transmission probably does not occur (Yanagihara, 1990). The human disease presents two major clinical pictures of severe hantavirus infections: a hemorrhagic fever with renal syndrome (HFRS), which is typically defined by fever, acute interstitial nephropathy with transient renal dysfunction and less commonly hemorrhage (Chun

et al, 1989) and the more recently described hantavirus group which is responsible of an acute pulmonary syndrome (HPS) (Dohmae *et al*, 1993).

Hantavirus antibodies were previously found in 33% of the population living in slum areas of Bangkok (Elwell *et al*, 1985). Some studies have been conducted in Thailand to determine the presence of hantavirus antibodies in various rodent species. High antibody prevalence by IFA and plaque reduction neutralization test (PRNT) was found in *Bandicota indica* (24.0%) and *Rattus norvegicus* (5.7%) (Elwell *et al*, 1985). Those studies did not clearly identify the risk of infection by hantavirus regarding specific potential vector and environment. In the present paper a preliminary approach to identify risk factors associated with rodent species and environmental issues is presented.

MATERIALS AND METHODS

Study site

Nakhon Pathom Province (2,168.327 km²) is located in the western part of Bangkok metropolitan area, (13° 49' 43''N; 100° 04' 35''E) (Fig 1) in a paddy field area. A rainy season lasts 6 to 7 months when the temperature increases to a maximum of 40°C and a dry season lasts for the other remaining 5 to 6 months. Land use characteristics




Fig 1—Study sites of Nakhon Pathom Province.




Fig 2—Study sites of Nakhon Ratchasima Province.

are classified as urban area, municipal area, agricultural area (rice and sugar cane fields) and animal farms (especially pigs). The population size of Nakhon Pathom Province in 1997 was 753,599 persons.

Nakhon Ratchasima Province with population of 2,510,839 as of 1997 (20,493.964 km²) is located in the north eastern part of Bangkok metropolitan area (14° 38'N; 1002° 26'E) (Fig 2). The geographical characteristics are classified as urban area, municipal area, agricultural area (mostly rice, corn fields and sugar cane fields).

Rodent sampling

Rodents from Nakhon Pathom Province were trapped in six villages (February-March, 1998). After carbon dioxide anesthesia, whole blood and organs (liver, kidney, spleen and heart) were collected. Specimens were labeled, kept in cryotube and stored at -70°C. The study sample were classified as follows by their different geographical origin:

- Maung district: Ban Wang Yen village where most of the land occupation is residential and rice fields.
- Sam Phran district: Ban Talad Jinda village where most of the areas are gardens.
- Don Tum district: Ban Samkoen village, mostly an area covered by rice fields.
- Kamphaeng Saen District: Ban Thung Look Nok, Ban Thung Kwang and Ban Thung Bau villages in a sugar cane field.

Rodents from Nakhon Ratchasima Province were trapped in two villages (August-October, 1998) and sampled as described above. The study areas were classified as follows:

- Nong Bunnak district: villages including moo bahn 1, 4, 5, 6, 7, 9, 10, 12, 15 in Nong Bunnak subdistrict where most of the areas are rice fields.
- Chalem Prakiat district: villages including moo bahn 1, 5, 6, 7, 9 in Nong Jan subdistrict where most of the areas are rice fields, and gardens.

Serological tests

The ELISA plates (MaxiSorp; Nunc, Roskilde, Denmark) were coated with either hantavirus antigen (Hantaan ROK 84/105 SMRV kindly provided by Professor Ho Wang Lee) dilution 1:1,000 in sterile PBS 7.2 overnight at 4°C. The next morning the plates were washed with PBS/tween 0.5%. Then 100 µl of rodent serum diluted 1:100 in PBS/ tween 0.5%/ skim milk 3% was added and incubated for 1 hour at 37°C after three times washing with PBS/

tween 0.5%, 100 µl of HRP/Anti-Rodent IgG (HRP Conjugate Goat Anti-Rodent IgG, ZYMED) was added and incubated for 1 hour at 37°C. After three times washing, the reaction was stopped with 50 µl 4M H₂SO₄ and the plates were read in a Metertech Σ490 spectrophotometer at 490 nm. The cutoff value for presence of antibody was defined as 3 times of the negative control absorbance.

RESULTS

Rodent characteristics and seroprevalence

354 rodents were trapped from different geographical areas of Nakhon Pathom Province. Specimens were identified as *Bandicota indica*, *Bandicota savilei*, *Mus caroli*, *Mus castaneus*, *Rattus argentiventer*, *Rattus exulans*, *Rattus losea*, *Rattus norvegicus* and *Rattus rattus*. The overall seroprevalence against hantavirus antigen was 2.3%. Mostly it was found among paddy field dike rodents, especially *Bandicota indica* (2.6%), while among the indoor-dwelling rodents, the higher seroprevalence was found in *Rattus norvegicus* (3.8%), *Rattus exulans* (2.2%) and *Rattus rattus* (1.5%) (Table 1). Among the different geographical areas of the province, the highest seroprevalence against hantavirus was found in the rodent popu-

lation collected from Don Tum district (4.2%), among *Bandicota indica* (2/65), *Rattus norvegicus* (1/7) and *Rattus exulans* (1/11) (Table 2).

In Nakhon Ratchasima Province, we trapped 326 rodents from different geographical areas. Only six species were present: *Bandicota indica*, *Mus caroli*, *Bandicota savilei*, *Bandicota* sp, *Rattus exulans* and *Rattus rattus*. The overall seroprevalence against hantavirus antigen was 4.0%. Most of the seroprevalence was found in *Bandicota indica* 19.1% and *Rattus exulans* 3.5% (Table 1).

DISCUSSION

All rodents collected in Nakhon Pathom and Nakhon Ratchasima provinces, were members of the Muridae family. *Bandicota indica* (44.4%) was the most frequent species trapped from Nakhon Pathom Province, while in Nakhon Ratchasima Province most of the rodent specimens caught were *Rattus exulans* (78.8%).

The highest hantavirus seroprevalence in Nakhon Pathom was found essentially among *Rattus norvegicus* (3.8%) and *Bandicota indica* (2.6%). In Nakhon Ratchasima only *Bandicota indica* (19.0%) and *Rattus exulans* (3.5%) had hantavirus reacting

Table 1
Hantavirus antibody reacting sera from rodent species trapped in Nakhon Pathom and Nakhon Ratchasima Province, Thailand, 1998.

Species	Nakhon Pathom	Nakhon Ratchasima	Total
Paddy field dike rodents			
<i>Bandicota indica</i>	4/157 (2.6) ^a	4/21 (19.1)	8/178 (4.5)
<i>Bandicota savilei</i>	0/14 (0.0)	0/6 (0.0)	0/20 (0.0)
<i>Bandicota</i> sp	-	0/5 (0.0)	0/5 (0.0)
<i>Mus caroli</i>	0/3 (0.0)	0/1 (0.0)	0/4 (0.0)
<i>Rattus argentiventer</i>	0/9 (0.0)	-	0/9 (0.0)
<i>Rattus losea</i>	0/2 (0.0)	-	0/2 (0.0)
Total	4/185 (2.2)	4/33 (12.1)	8/218 (3.7)
Indoor rodents			
<i>Mus castaneus</i>	0/3 (0.0)	-	0/3 (0.0)
<i>Rattus exulans</i>	1/45 (2.2)	9/257 (3.5)	10/302 (3.3)
<i>Rattus norvegicus</i>	2/53 (3.8)	-	2/53 (3.8)
<i>Rattus rattus</i>	1/68 (1.5)	0/36 (0.0)	1/104 (1.0)
Total	4/169 (2.4)	9/293 (3.1)	13/462 (2.8)
Total	8/354 (2.3)	13/326 (4.0)	21/680 (3.1)

^aNumber of positive/Total tested (percentage)

Table 2
Hantaanvirus antigen antibody reacting sera from rodent species of Nakhon Pathom Province
by district of origin.

Species	District				Total
	Muang	Sam Phran	Don Tum	Kamphaeng Saen	
<i>Bandicota indica</i>	2/41 (4.9)*	0/8 (0.0)	2/65 (3.1)	0/43 (0.0)	4/157 (2.5)
<i>Bandicota savilei</i>	0/11 (0.0)	0/1 (0.0)	0/2 (0.0)	-	0/14 (0.0)
<i>Mus caroli</i>	-	-	-	0/3 (0.0)	0/3 (0.0)
<i>Rattus argentiventer</i>	0/2 (0.0)	0/4 (0.0)	0/1 (0.0)	0/2 (0.0)	0/9 (0.0)
<i>Rattus losea</i>	0/1 (0.0)	0/1 (0.0)	-	-	0/2 (0.0)
<i>Mus castaneus</i>	-	0/3 (0.0)	-	-	0/3 (0.0)
<i>Rattus exulans</i>	0/4 (0.0)	0/22 (0.0)	1/11 (9.1)	0/8 (0.0)	1/45 (2.2)
<i>Rattus norvegicus</i>	1/32 (3.1)	0/7 (0.0)	1/7 (14.3)	0/7 (0.0)	2/46 (4.3)
<i>Rattus rattus</i>	0/7 (0.0)	1/47 (2.1)	0/10 (0.0)	0/4 (0.0)	1/68 (1.5)
Total	3/98 (3.1)	1/93 (1.1)	4/96 (4.2)	0/67 (0.0)	8/354 (2.3)

*positive/total tested

Table 3
Hantavirus antigen antibody reacting sera from rodent species of Nakhon Ratchasima Province by
district of origin.

Species	District		Total
	Nong Bunnak	Chalerm Prakiet	
<i>Bandicota indica</i>	3/18 ^a (16.7)	1/3 (33.3)	4/21 (19.0)
<i>Bandicota savilei</i>	0/3 (0.0)	0/3 (0.0)	0/6 (0.0)
<i>Bandicota sp</i>	0/4 (0.0)	0/1 (0.0)	0/5 (0.0)
<i>Mus caroli</i>	0/1 (0.0)	0/0 (0.0)	0/1 (0.0)
<i>Rattus exulans</i>	5/156 (3.2)	4/101 (4.0)	9/257 (3.5)
<i>Rattus rattus</i>	0/18 (0.0)	0/18 (0.0)	0/36 (0.0)
Total	8/200 (4.0)	5/126 (4.0)	13/326 (4.0)

*positive/total tested

antibodies. These findings show an active hantavirus circulation in the provinces with *Bandicota indica* as a primary virus reservoir candidate as previously reported by others (Elwell *et al*, 1985; Sawadikol *et al*, 1989; Tantivanich *et al*, 1988, 1992). In some instances farmers and other low-income groups eat rodent meat trapped from rice fields or sugar cane areas. Some rodents like *Bandicota indica* can be found also at the local market. This is a risk factor which needs to be evaluated.

Rattus exulans can be another potential hantavirus vector as well as *Rattus norvegicus* and *Rattus rattus*. *Rattus exulans* have been reported as infected for the first time in those areas. Indoor

rodents which have positive hantavirus serology such as *Rattus exulans* can potentially transmit virus to human because of their domestic habits.

Outdoor rodents appear to be more infected, as judged by the rate of antibody prevalence, than indoor rodents. When we compared hantavirus circulation in both provinces we found more paddy field dike rodents with higher hantavirus seropositivity than indoor rodents in Nakhon Ratchasima but no difference in Nakhon Pathom. However the hantavirus prevalence by district of Nakhon Pathom showed higher positivity in Muang and Don Tum areas which are more included in the rice field environment than the two other sites.

From previous studies and our observation (Elwell *et al*, 1985) *Bandicoda indica* and *Rattus exulans* appear as a primary potential reservoir/vector of hantavirus in the central plain of Thailand. *Bandicoda indica* lives outside the housing and its domain overlaps with that of *Rattus exulans* which is the most abundant domestic rodent in Thailand living in close contact with humans. *Rattus exulans* is the smallest rat in Thailand and originally comes from the Pacific Islands where it is known to be an outdoor rodent. From our study we can hypothesize that in Thailand *Bandicoda indica* can play a role as a wild reservoir of virus and *Rattus exulans* will be more likely a vector of hantavirus to humans.

It appears from a previous study done in the human population that 10 years ago 1.2% of the population from the Bangkok area had hantavirus antibody (Sawasdikol *et al*, 1989). We found similar seroprevalence in humans from Nakhon Pathom Province (1.9%) however, a high density population living in slum areas of Bangkok showed an unexplained, much higher, prevalence of 31.4% (Elwell *et al*, 1985).

Those findings of seroprevalence of antibodies against hantavirus among rodents will lead to other serological surveys among human populations who have experienced contact with rodents. Also it is important to address a warning to physicians to emphasize laboratory investigations and differentiate the causes of acute respiratory syndrome (Kiatboonsri *et al*, 1995), renal failure and leptospirosis cases in Thailand (Hindrichsen *et al*, 1993).

ACKNOWLEDGEMENTS

This study was supported by the Department of Technical and Economic Cooperation (DTEC) Thailand; the Center for Vaccine Development (CVD) Thailand; CDC/MOPH Thailand and the Institut de Recherche pour le Developpement (IRD), France.

We would like to thank all staff in Environmental and Sanitation Section, Nakhon Pathom Provincial Health Office, Dr Darika Kingnate Department of Communicable Diseases Control, Ministry of Public Health, Nonthaburi, Thailand. Miss Yuvaluk Khoprasert Agricultural Zoology Research Group, Division of Entomology and Zoology, Department of Agriculture. Mr Greangsak Hamarit, Agricultural Zoology Research Group, Division of Entomology

and Zoology, Department of Agriculture. Mr Noppadol Sanngjun, Armed Forces Research Institute of Medical Sciences. Dr Yupha Rongsriyam, Department of Medical Entomology, Faculty Tropical Medicine, Mahidol University, University and Office for their Cooperation, CVD-RCEVD in the program of emerging viral diseases.

We are also deeply indebted to Professor Natth Bhamarapavati for his constant support to the project.

REFERENCES

- Chun CH, Lahdevirta J, Lee HW. Clinical manifestations of HFRS. In: Lee HW, Darlymple JM, eds. Manual of hemorrhagic fever with renal syndrome. World Health Organization Collaborating Center for Virus Reference And Research (HFRS), Institute for Viral Diseases, Korea University, Seoul. 1989: 19-38.
- Dohmae K, Koshimizu U, Nishimune. In utero and mammary transfer of hantavirus antibody from dams to infant rats. *Lab Anim Sci* 1993; 43: 557-61.
- Elwell MR, Ward GS, Tingpalapong M, LeDuc JW. Serologic evidence of Hantaan-like virus in rodents and man in Thailand. *Southeast Asian J Trop Med Public Health* 1985; 16: 349-54.
- Glass GE, Childs JE, Korch GW, LeDuc JW. Association of intraspecific wounding with hantaviral infection in wild rats (*Rattus norvegicus*). *Epidemiol Infect* 1988; 101: 459-72.
- Hindrichsen S, Medeiros A, Clement J, McKenna P, Matthys P, Neild GH. Hantavirus infection in Brazilian patients from Recife with suspected leptospirosis. 1993; 341: 50.
- Kiatboonsri S, Vathesatogit P, Charoenpan P. Adult respiratory distress syndrome in Thai medical patients. *Southeast Asian J Trop Med Public Health* 1995; 26: 774-80.
- Lee HW, French GR, Lee PW, Baek LJ, Tsuchiya K, Foulke RS. Observations on natural and laboratory infection of rodents with the etiologic agent of Korean hemorrhagic fever. *Am J Trop Med Hyg* 1981; 30: 477-82.
- Lee HW, Baek LJ, Johnson KM. Isolation of Hantaan virus, the etiologic agent of Korean hemorrhagic fever from wild urban rats. *J Infect Dis* 1982; 146: 638-44.
- Lee HW. Korean hemorrhagic fever. In Melnick JL, ed: Progress in medical virology. Basel: S Karger AG, 1982: 96-111.
- Nuzum EO, Rossi CA, Stephenson EH, LeDuc JW. Aerosol transmission of Hantaan and related viruses to

- laboratory rats. *Am J Trop Med Hyg* 1988; 38: 636-40.
- Sawasdikol S, Tamura M, Jamjit P. Antibody to hemorrhagic fever with renal syndrome in man and rat in Thailand. *Bull Dept Med Sci* 1989; 31: 125-30.
- Tantivanich S, Chongsa-nguan M, Impand P, Potha U, Implaps S. Serological studies of Hantaan virus among Thai people and urban rats. *J Parasit Trop Med Assoc Thai* 1988; 11: 76.
- Tantivanich S, Ayuthaya PI, Usawattanakul W, Imphand P. Hantaanvirus among urban rats from a slum area in Bangkok. *Southeast Asian J Trop Med Public Health* 1992; 23: 504-9.
- Yanagihara R, Amyx HL, Gajdusek DC. Experimental infection with Puumala virus, the etiologic agent of nephropathia epidemica, in bank voles (*Clethrionomys glareolus*). *J Virol* 1985; 55: 34-8.
- Yanagihara R, Gajdusek DC. Hemorrhagic fever with renal syndrome; a historical perspective and review of recent advances. In: Gear JHS, ed. *CRC handbook of viral and rickettsial hemorrhagic fever*. Boca Raton: CRC Press 1988; 151-88.
- Yanagihara R. Hantavirus infection in the United States: Epizootology and epidemiology. *Rev Infect Dis* 1990; 12: 449-57.

**THE SOUTHEAST ASIAN JOURNAL OF TROPICAL MEDICINE
AND PUBLIC HEALTH**
ISSN : 0038-3619

Volume 31

Number 2

June 2000

CONTENTS

Page

Editorial	C Kidson	201
Prognostic significance of skin and subcutaneous fat sequestration of parasites in severe falciparum malaria	P Wilairatana, M Riganti, P Puchadapirom, B Punpoowong, S Vannaphan, R Udomsangpetch, S Krudsood, GM Brittenham, Sornchai Looareesuwan	203
<i>Plasmodium falciparum</i> : Detection and strain identification of Indian isolates by polymerase chain reaction	ABS Sidhu, R Madhubala	213
Malaria outbreak in a tribal area of Gujarat State, India	HC Srivastava, RS Yadav	219
Review : Status of malaria in Thailand	T Chareonviriyaphap, MJ Bangs, S Ratanatham	225
Treatment of uncomplicated <i>Plasmodium falciparum</i> malaria in Myanmar : A clinical decision analysis	Cho-Min-Naing, A Saul	238
Autogenous <i>Aedes togoi</i> sub-colony (Chanthaburi, Thailand strain), an efficient laboratory vector in study of filariasis	D Riyong, W Choochote, A Jitpakdi, S Suvannadabba, S Leemingsawat, U Chaithong	246
A preliminary analysis of <i>Wuchereria bancrofti</i> microfilarial antigens for potential use in diagnosis	JK Casinader, Saverimuttu, K Berzins, P Perlmann, EH Karunanayake, MM Ismail	252
A comparison of the pattern of liver involvement in dengue hemorrhagic fever with classic dengue fever	SFSA Walid, S Sanusi MM Zawawi, RA Ali	259
Case report : A dengue shock patient with negative serology and polymerase chain reaction (PCR) tests	C Pancharoen, U Thisyakorn, A Nisalak	264
Molecular evolution, distribution and genetic relationship among the dengue 2 viruses isolated from different clinical severity.....	BD Pandey, K Morita, F Hasebe, M del Carmen Parquet, A Igarashi	266
Clinico-epidemiology of hepatitis C viral infection in northeastern Thailand.....	K Chunlertrith, W Sukeepaisarnjaroen, P Mairiang, Y Urwijitaroon, K Takase, T Yamauchi, H Yoshimura, Y Tameda	273
Serological study of hantavirus in the rodent population of Nakhon Pathom and Nakhon Ratchasima Provinces, Thailand.....	N Nitatpattana, G Chauvancy, J Dardaine, T Poblap, K Jumronsawat, W Tangkanakul, D Poonsuksombat, S Yoksan, J-P Gonzalez	277 ✕
Human immunodeficiency virus type 1 subtypes among Malaysian intravenous drug users.....	TS Saraswathy, KP Ng, M Sinniah	283
Immunogenicity and safety of purified Vero-cell rabies vaccine in severely rabies-exposed patients in China	XJ Wang, J Lang, XR Tao, JD Shu, V Le Mener, SC Wood, JT Huang, SL Zhao	287
Blood lead levels of a population group not occupationally exposed to lead in Singapore.....	KS Neo, KT Goh, CT Sam	295
Risk factors for Nipah virus transmission, Port Dickson, Negeri Sembilan, Malaysia: results from a hospital-based case-control study	NM Amal, MS Lye, TG Ksiazek, PD Kitsutani, KS Hanjeet, MA Kamaluddin, F Ong, S Devi, PC Stockton, O Ghazali, R Zainab, MA Taha	301
Assessment of Nipah virus transmission among pork sellers in Seremban, Malaysia	GD Premalatha, MS Lye, J Ariokasamy, UD Parashar, R Rahmat, BY Lee, TG Ksiazek	307
Assessment of community maternal care performance of public health midwives of a province in Sri Lanka : A multi-method approach	W Gunathunga, DN Fernando	310
Fluoride in water consumed by children in remote areas of Thailand.....	S Chuckpaiwong, S Nakornchai, R Surarit, S Soo-ampon, R Kasetsuwan	319

