

# Multi-strange baryon elliptic flow in Au + Au collisions at $\sqrt{s_{NN}}=200$ GeV

J. Adams,<sup>3</sup> M.M. Aggarwal,<sup>29</sup> Z. Ahammed,<sup>43</sup> J. Amonett,<sup>20</sup> B.D. Anderson,<sup>20</sup> D. Arkhipkin,<sup>13</sup> G.S. Averichev,<sup>12</sup> S.K. Badyal,<sup>19</sup> Y. Bai,<sup>27</sup> J. Balewski,<sup>17</sup> O. Barannikova,<sup>32</sup> L.S. Barnby,<sup>3</sup> J. Baudot,<sup>18</sup> S. Bekele,<sup>28</sup> V.V. Belaga,<sup>12</sup> A. Bellingeri-Laurikainen,<sup>38</sup> R. Bellwied,<sup>46</sup> J. Berger,<sup>14</sup> B.I. Bezverkhny,<sup>48</sup> S. Bharadwaj,<sup>33</sup> A. Bhasin,<sup>19</sup> A.K. Bhati,<sup>29</sup> V.S. Bhatia,<sup>29</sup> H. Bichsel,<sup>45</sup> J. Bielcik,<sup>48</sup> J. Bielcikova,<sup>48</sup> A. Billmeier,<sup>46</sup> L.C. Bland,<sup>4</sup> C.O. Blyth,<sup>3</sup> B.E. Bonner,<sup>34</sup> M. Botje,<sup>27</sup> A. Boucham,<sup>38</sup> J. Bouchet,<sup>38</sup> A.V. Brandin,<sup>25</sup> A. Bravar,<sup>4</sup> M. Bystersky,<sup>11</sup> R.V. Cadman,<sup>1</sup> X.Z. Cai,<sup>37</sup> H. Caines,<sup>48</sup> M. Calderón de la Barca Sánchez,<sup>17</sup> J. Castillo,<sup>21</sup> O. Catu,<sup>48</sup> D. Cebra,<sup>7</sup> Z. Chajecski,<sup>28</sup> P. Chaloupka,<sup>11</sup> S. Chattopadhyay,<sup>43</sup> H.F. Chen,<sup>36</sup> Y. Chen,<sup>8</sup> J. Cheng,<sup>41</sup> M. Cherney,<sup>10</sup> A. Chikanian,<sup>48</sup> W. Christie,<sup>4</sup> J.P. Coffin,<sup>18</sup> T.M. Cormier,<sup>46</sup> J.G. Cramer,<sup>45</sup> H.J. Crawford,<sup>6</sup> D. Das,<sup>43</sup> S. Das,<sup>43</sup> M. Daugherty,<sup>40</sup> M.M. de Moura,<sup>35</sup> T.G. Dedovich,<sup>12</sup> A.A. Derevschikov,<sup>31</sup> L. Didenko,<sup>4</sup> T. Dietel,<sup>14</sup> S.M. Dogra,<sup>19</sup> W.J. Dong,<sup>8</sup> X. Dong,<sup>36</sup> J.E. Draper,<sup>7</sup> F. Du,<sup>48</sup> A.K. Dubey,<sup>15</sup> V.B. Dunin,<sup>12</sup> J.C. Dunlop,<sup>4</sup> M.R. Dutta Mazumdar,<sup>43</sup> V. Eckardt,<sup>23</sup> W.R. Edwards,<sup>21</sup> L.G. Efimov,<sup>12</sup> V. Emelianov,<sup>25</sup> J. Engelage,<sup>6</sup> G. Eppley,<sup>34</sup> B. Erasmus,<sup>38</sup> M. Estienne,<sup>38</sup> P. Fachini,<sup>4</sup> J. Faivre,<sup>18</sup> R. Fatemi,<sup>17</sup> J. Fedorisin,<sup>12</sup> K. Filimonov,<sup>21</sup> P. Filip,<sup>11</sup> E. Finch,<sup>48</sup> V. Fine,<sup>4</sup> Y. Fisyak,<sup>4</sup> J. Fu,<sup>41</sup> C.A. Gagliardi,<sup>39</sup> L. Gaillard,<sup>3</sup> J. Gans,<sup>48</sup> M.S. Ganti,<sup>43</sup> F. Geurts,<sup>34</sup> V. Ghazikhanian,<sup>8</sup> P. Ghosh,<sup>43</sup> J.E. Gonzalez,<sup>8</sup> H. Gos,<sup>44</sup> O. Grachov,<sup>46</sup> O. Grebenyuk,<sup>27</sup> D. Grosnick,<sup>42</sup> S.M. Guertin,<sup>8</sup> Y. Guo,<sup>46</sup> A. Gupta,<sup>19</sup> T.D. Gutierrez,<sup>7</sup> T.J. Hallman,<sup>4</sup> A. Hamed,<sup>46</sup> D. Hardtke,<sup>21</sup> J.W. Harris,<sup>48</sup> M. Heinz,<sup>2</sup> T.W. Henry,<sup>39</sup> S. Heppelmann,<sup>30</sup> B. Hippolyte,<sup>18</sup> A. Hirsch,<sup>32</sup> E. Hjort,<sup>21</sup> G.W. Hoffmann,<sup>40</sup> H.Z. Huang,<sup>8</sup> S.L. Huang,<sup>36</sup> E.W. Hughes,<sup>5</sup> T.J. Humanic,<sup>28</sup> G. Igo,<sup>8</sup> A. Ishihara,<sup>40</sup> P. Jacobs,<sup>21</sup> W.W. Jacobs,<sup>17</sup> M. Jedynek,<sup>44</sup> H. Jiang,<sup>8</sup> P.G. Jones,<sup>3</sup> E.G. Judd,<sup>6</sup> S. Kabana,<sup>2</sup> K. Kang,<sup>41</sup> M. Kaplan,<sup>9</sup> D. Keane,<sup>20</sup> A. Kechechyan,<sup>12</sup> V.Yu. Khodyrev,<sup>31</sup> J. Kiryluk,<sup>22</sup> A. Kisiel,<sup>44</sup> E.M. Kislov,<sup>12</sup> J. Klay,<sup>21</sup> S.R. Klein,<sup>21</sup> D.D. Koetke,<sup>42</sup> T. Kollegger,<sup>14</sup> M. Kopytine,<sup>20</sup> L. Kotchenda,<sup>25</sup> K.L. Kowalik,<sup>21</sup> M. Kramer,<sup>26</sup> P. Kravtsov,<sup>25</sup> V.I. Kravtsov,<sup>31</sup> K. Krueger,<sup>1</sup> C. Kuhn,<sup>18</sup> A.I. Kulikov,<sup>12</sup> A. Kumar,<sup>29</sup> R.Kh. Kutuev,<sup>13</sup> A.A. Kuznetsov,<sup>12</sup> M.A.C. Lamont,<sup>48</sup> J.M. Landgraf,<sup>4</sup> S. Lange,<sup>14</sup> F. Laue,<sup>4</sup> J. Lauret,<sup>4</sup> A. Lebedev,<sup>4</sup> R. Lednicky,<sup>12</sup> S. Lehocka,<sup>12</sup> M.J. LeVine,<sup>4</sup> C. Li,<sup>36</sup> Q. Li,<sup>46</sup> Y. Li,<sup>41</sup> G. Lin,<sup>48</sup> S.J. Lindenbaum,<sup>26</sup> M.A. Lisa,<sup>28</sup> F. Liu,<sup>47</sup> H. Liu,<sup>36</sup> L. Liu,<sup>47</sup> Q.J. Liu,<sup>45</sup> Z. Liu,<sup>47</sup> T. Ljubicic,<sup>4</sup> W.J. Llope,<sup>34</sup> H. Long,<sup>8</sup> R.S. Longacre,<sup>4</sup> M. Lopez-Noriega,<sup>28</sup> W.A. Love,<sup>4</sup> Y. Lu,<sup>47</sup> T. Ludlam,<sup>4</sup> D. Lynn,<sup>4</sup> G.L. Ma,<sup>37</sup> J.G. Ma,<sup>8</sup> Y.G. Ma,<sup>37</sup> D. Magestro,<sup>28</sup> S. Mahajan,<sup>19</sup> D.P. Mahapatra,<sup>15</sup> R. Majka,<sup>48</sup> L.K. Mangotra,<sup>19</sup> R. Manweiler,<sup>42</sup> S. Margetis,<sup>20</sup> C. Markert,<sup>20</sup> L. Martin,<sup>38</sup> J.N. Marx,<sup>21</sup> H.S. Matis,<sup>21</sup> Yu.A. Matulenko,<sup>31</sup> C.J. McClain,<sup>1</sup> T.S. McShane,<sup>10</sup> F. Meissner,<sup>21</sup> Yu. Melnick,<sup>31</sup> A. Meschanin,<sup>31</sup> M.L. Miller,<sup>22</sup> N.G. Minaev,<sup>31</sup> C. Mironov,<sup>20</sup> A. Mischke,<sup>27</sup> D.K. Mishra,<sup>15</sup> J. Mitchell,<sup>34</sup> B. Mohanty,<sup>43</sup> L. Molnar,<sup>32</sup> C.F. Moore,<sup>40</sup> D.A. Morozov,<sup>31</sup> M.G. Munhoz,<sup>35</sup> B.K. Nandi,<sup>43</sup> S.K. Nayak,<sup>19</sup> T.K. Nayak,<sup>43</sup> J.M. Nelson,<sup>3</sup> P.K. Netrakanti,<sup>43</sup> V.A. Nikitin,<sup>13</sup> L.V. Nogach,<sup>31</sup> S.B. Nurushev,<sup>31</sup> G. Odyniec,<sup>21</sup> A. Ogawa,<sup>4</sup> V. Okorokov,<sup>25</sup> M. Oldenburg,<sup>21</sup> D. Olson,<sup>21</sup> S.K. Pal,<sup>43</sup> Y. Panebratsev,<sup>12</sup> S.Y. Panitkin,<sup>4</sup> A.I. Pavlinov,<sup>46</sup> T. Pawlak,<sup>44</sup> T. Peitzmann,<sup>27</sup> V. Perevoztchikov,<sup>4</sup> C. Perkins,<sup>6</sup> W. Peryt,<sup>44</sup> V.A. Petrov,<sup>46</sup> S.C. Phatak,<sup>15</sup> R. Picha,<sup>7</sup> M. Planinic,<sup>49</sup> J. Pluta,<sup>44</sup> N. Porile,<sup>32</sup> J. Porter,<sup>45</sup> A.M. Poskanzer,<sup>21</sup> M. Potekhin,<sup>4</sup> E. Potrebenikova,<sup>12</sup> B.V.K.S. Potukuchi,<sup>19</sup> D. Prindle,<sup>45</sup> C. Pruneau,<sup>46</sup> J. Putschke,<sup>21</sup> G. Rakness,<sup>30</sup> R. Raniwala,<sup>33</sup> S. Raniwala,<sup>33</sup> O. Ravel,<sup>38</sup> R.L. Ray,<sup>40</sup> S.V. Razin,<sup>12</sup> D. Reichhold,<sup>32</sup> J.G. Reid,<sup>45</sup> J. Reinnarth,<sup>38</sup> G. Renault,<sup>38</sup> F. Retiere,<sup>21</sup> A. Ridiger,<sup>25</sup> H.G. Ritter,<sup>21</sup> J.B. Roberts,<sup>34</sup> O.V. Rogachevskiy,<sup>12</sup> J.L. Romero,<sup>7</sup> A. Rose,<sup>21</sup> C. Roy,<sup>38</sup> L. Ruan,<sup>36</sup> M. Russcher,<sup>27</sup> R. Sahoo,<sup>15</sup> I. Sakrejda,<sup>21</sup> S. Salur,<sup>48</sup> J. Sandweiss,<sup>48</sup> M. Sarsour,<sup>17</sup> I. Savin,<sup>13</sup> P.S. Sazhin,<sup>12</sup> J. Schambach,<sup>40</sup> R.P. Scharenberg,<sup>32</sup> N. Schmitz,<sup>23</sup> K. Schweda,<sup>21</sup> J. Seger,<sup>10</sup> P. Seyboth,<sup>23</sup> E. Shahaliev,<sup>12</sup> M. Shao,<sup>36</sup> W. Shao,<sup>5</sup> M. Sharma,<sup>29</sup> W.Q. Shen,<sup>37</sup> K.E. Shestermanov,<sup>31</sup> S.S. Shimanskiy,<sup>12</sup> E. Sichtermann,<sup>21</sup> F. Simon,<sup>23</sup> R.N. Singaraju,<sup>43</sup> N. Smirnov,<sup>48</sup> R. Snellings,<sup>27</sup> G. Sood,<sup>42</sup> P. Sorensen,<sup>21</sup> J. Sowinski,<sup>17</sup> J. Speltz,<sup>18</sup> H.M. Spinka,<sup>1</sup> B. Srivastava,<sup>32</sup> A. Stadnik,<sup>12</sup> T.D.S. Stanislaus,<sup>42</sup> R. Stock,<sup>14</sup> A. Stolpovsky,<sup>46</sup> M. Strikhanov,<sup>25</sup> B. Stringfellow,<sup>32</sup> A.A.P. Suaide,<sup>35</sup> E. Sugarbaker,<sup>28</sup> C. Suire,<sup>4</sup> M. Sumner,<sup>11</sup> B. Surrow,<sup>22</sup> M. Swanger,<sup>10</sup> T.J.M. Symons,<sup>21</sup> A. Szanto de Toledo,<sup>35</sup> A. Tai,<sup>8</sup> J. Takahashi,<sup>35</sup> A.H. Tang,<sup>27</sup> T. Tarnowsky,<sup>32</sup> D. Thein,<sup>8</sup> J.H. Thomas,<sup>21</sup> S. Timoshenko,<sup>25</sup> M. Tokarev,<sup>12</sup> T.A. Trainor,<sup>45</sup> S. Trentalange,<sup>8</sup> R.E. Tribble,<sup>39</sup> O.D. Tsai,<sup>8</sup> J. Ulery,<sup>32</sup> T. Ullrich,<sup>4</sup> D.G. Underwood,<sup>1</sup> G. Van Buren,<sup>4</sup> M. van Leeuwen,<sup>21</sup> A.M. Vander Molen,<sup>24</sup> R. Varma,<sup>16</sup> I.M. Vasilievski,<sup>13</sup> A.N. Vasiliev,<sup>31</sup> R. Vernet,<sup>18</sup> S.E. Vigdor,<sup>17</sup> Y.P. Viyogi,<sup>43</sup> S. Vokal,<sup>12</sup> S.A. Voloshin,<sup>46</sup> W.T. Waggoner,<sup>10</sup> F. Wang,<sup>32</sup> G. Wang,<sup>20</sup> G. Wang,<sup>5</sup> X.L. Wang,<sup>36</sup> Y. Wang,<sup>40</sup> Y. Wang,<sup>41</sup> Z.M. Wang,<sup>36</sup> H. Ward,<sup>40</sup> J.W. Watson,<sup>20</sup> J.C. Webb,<sup>17</sup> G.D. Westfall,<sup>24</sup> A. Wetzler,<sup>21</sup> C. Whitten Jr.,<sup>8</sup> H. Wieman,<sup>21</sup> S.W. Wissink,<sup>17</sup> R. Witt,<sup>2</sup> J. Wood,<sup>8</sup> J. Wu,<sup>36</sup> N. Xu,<sup>21</sup> Z. Xu,<sup>4</sup> Z.Z. Xu,<sup>36</sup> E. Yamamoto,<sup>21</sup> P. Yepes,<sup>34</sup> V.I. Yurevich,<sup>12</sup> I. Zborovsky,<sup>11</sup> H. Zhang,<sup>4</sup> W.M. Zhang,<sup>20</sup> Y. Zhang,<sup>36</sup> Z.P. Zhang,<sup>36</sup> R. Zoukarneev,<sup>13</sup> Y. Zoukarneeva,<sup>13</sup> and A.N. Zubarev<sup>12</sup>

(STAR Collaboration)

<sup>1</sup>Argonne National Laboratory, Argonne, Illinois 60439

<sup>2</sup>University of Bern, 3012 Bern, Switzerland

<sup>3</sup>University of Birmingham, Birmingham, United Kingdom

<sup>4</sup>Brookhaven National Laboratory, Upton, New York 11973

- <sup>5</sup>California Institute of Technology, Pasadena, California 91125  
<sup>6</sup>University of California, Berkeley, California 94720  
<sup>7</sup>University of California, Davis, California 95616  
<sup>8</sup>University of California, Los Angeles, California 90095  
<sup>9</sup>Carnegie Mellon University, Pittsburgh, Pennsylvania 15213  
<sup>10</sup>Creighton University, Omaha, Nebraska 68178  
<sup>11</sup>Nuclear Physics Institute AS CR, 250 68 Řež/Prague, Czech Republic  
<sup>12</sup>Laboratory for High Energy (JINR), Dubna, Russia  
<sup>13</sup>Particle Physics Laboratory (JINR), Dubna, Russia  
<sup>14</sup>University of Frankfurt, Frankfurt, Germany  
<sup>15</sup>Institute of Physics, Bhubaneswar 751005, India  
<sup>16</sup>Indian Institute of Technology, Mumbai, India  
<sup>17</sup>Indiana University, Bloomington, Indiana 47408  
<sup>18</sup>Institut de Recherches Subatomiques, Strasbourg, France  
<sup>19</sup>University of Jammu, Jammu 180001, India  
<sup>20</sup>Kent State University, Kent, Ohio 44242  
<sup>21</sup>Lawrence Berkeley National Laboratory, Berkeley, California 94720  
<sup>22</sup>Massachusetts Institute of Technology, Cambridge, MA 02139-4307  
<sup>23</sup>Max-Planck-Institut für Physik, Munich, Germany  
<sup>24</sup>Michigan State University, East Lansing, Michigan 48824  
<sup>25</sup>Moscow Engineering Physics Institute, Moscow Russia  
<sup>26</sup>City College of New York, New York City, New York 10031  
<sup>27</sup>NIKHEF and Utrecht University, Amsterdam, The Netherlands  
<sup>28</sup>Ohio State University, Columbus, Ohio 43210  
<sup>29</sup>Panjab University, Chandigarh 160014, India  
<sup>30</sup>Pennsylvania State University, University Park, Pennsylvania 16802  
<sup>31</sup>Institute of High Energy Physics, Protvino, Russia  
<sup>32</sup>Purdue University, West Lafayette, Indiana 47907  
<sup>33</sup>University of Rajasthan, Jaipur 302004, India  
<sup>34</sup>Rice University, Houston, Texas 77251  
<sup>35</sup>Universidade de Sao Paulo, Sao Paulo, Brazil  
<sup>36</sup>University of Science & Technology of China, Anhui 230027, China  
<sup>37</sup>Shanghai Institute of Applied Physics, Shanghai 201800, China  
<sup>38</sup>SUBATECH, Nantes, France  
<sup>39</sup>Texas A&M University, College Station, Texas 77843  
<sup>40</sup>University of Texas, Austin, Texas 78712  
<sup>41</sup>Tsinghua University, Beijing 100084, China  
<sup>42</sup>Valparaiso University, Valparaiso, Indiana 46383  
<sup>43</sup>Variable Energy Cyclotron Centre, Kolkata 700064, India  
<sup>44</sup>Warsaw University of Technology, Warsaw, Poland  
<sup>45</sup>University of Washington, Seattle, Washington 98195  
<sup>46</sup>Wayne State University, Detroit, Michigan 48201  
<sup>47</sup>Institute of Particle Physics, CCNU (HZNU), Wuhan 430079, China  
<sup>48</sup>Yale University, New Haven, Connecticut 06520  
<sup>49</sup>University of Zagreb, Zagreb, HR-10002, Croatia

We report on the first measurement of elliptic flow  $v_2(p_T)$  of multi-strange baryons  $\Xi^- + \bar{\Xi}^+$  and  $\Omega^- + \bar{\Omega}^+$  in heavy-ion collisions. In minimum bias Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV, a significant amount of elliptic flow, comparable to other non-strange baryons, is observed for multi-strange baryons which are expected to be particularly sensitive to the dynamics of the partonic stage of heavy-ion collisions. The  $p_T$  dependence of  $v_2$  of the multi-strange baryons confirms the number of constituent quark scaling previously observed for lighter hadrons. These results support the idea that a substantial fraction of the observed collective motion is developed at the early partonic stage in ultra-relativistic nuclear collisions at RHIC.

PACS numbers: 25.75.Ld

Lattice QCD calculations, at vanishing or finite net-baryon density, predict a transition from the deconfined thermalized partonic matter Quark Gluon Plasma (QGP) to ordinary hadronic matter at a critical temperature  $T_c \approx 150 - 180$  MeV [1, 2]. Measurements of hadron yields in the intermediate ( $2 \lesssim p_T \lesssim 6$  GeV/c) and high ( $p_T \gtrsim 6 - 8$  GeV/c) transverse

momentum  $p_T$  region indicate that dense matter has been produced in Au + Au collisions at RHIC [3, 4, 5, 6, 7, 8, 9, 10]. Furthermore, previous measurements of elliptic flow of hadrons indicate that the matter created at RHIC is also strongly interacting [11, 12]. Thus, in the early stage of the collision, dense and strongly interacting matter will lead to

collective effects among constituents such as transverse collective motion. If these interactions occur frequently enough, the system will finally reach thermalization. Due to the initial spatial anisotropy of the system in non-central collisions, an elliptic component of the collective transverse motion should also be present. Collectivity is cumulative throughout the whole collision and should survive the hadronization process [13, 14]; therefore, the amount of transverse flow observed in the final state will have a contribution from the pre-hadronic, i.e. partonic, stage.

Early dynamic information might be masked by later hadronic rescatterings. Multi-strange baryons with their large mass and presumably small hadronic cross sections [15, 16, 17, 18, 19], should be less sensitive to hadronic rescattering in the later stages of the collision and therefore a good probe of the early stage of the collision [20]. Indeed, a systematic study of hadron  $p_T$  spectra from high-energy heavy ion collisions, using a hydrodynamically inspired model, shows that multi-strange baryons thermally freeze-out close to the point where chemical freeze-out occurs with  $T_{\text{ch}} \sim 160$  MeV [20, 21] which at these collision energies coincides with the critical temperature  $T_c$  [1, 2]. This may mean that multi-strange baryons are not, or much less, affected by hadronic rescatterings during the later stage of heavy ion collisions [15, 16]. Their observed transverse flow would then primarily reflect the partonic flow. Moreover, elliptic flow is in itself considered to be a good tool for understanding the properties of the early stage of the collisions [22, 23], primarily due to its self-quenching nature. Elliptic flow is generated from the initial spatial anisotropy of the system created in non-central collisions by rescatterings among the constituents of the system. The generated elliptic flow will reduce the spatial anisotropy of the system and quench its own origin. Thus multi-strange baryon elliptic flow could be a valuable probe of the initial partonic system.

In this Letter we present the first results on elliptic flow of multi-strange baryons  $\Xi^- + \Xi^+$  and  $\Omega^- + \Omega^+$  from Au + Au collisions at  $\sqrt{s_{\text{NN}}} = 200$  GeV, as measured with the STAR detector [24]. About 2 million events from Au + Au collisions collected with a minimum bias trigger are used in this analysis. Multi-strange baryons are reconstructed via their decay topology:  $\Xi \rightarrow \Lambda + \pi$  and  $\Omega \rightarrow \Lambda + K$  with the subsequent decay of  $\Lambda \rightarrow p + \pi$  as described in [20]. Charged tracks were reconstructed in the STAR Time Projection Chamber (TPC) [25]. Simple cuts on geometry, kinematics and particle identification via specific ionization are applied to reduce the combinatorial background. A detailed description of the analysis procedure can be found in [20, 26].

Figure 1 shows the invariant mass distribution for (a)  $\Xi^- + \Xi^+$  and (b)  $\Omega^- + \Omega^+$  candidates from minimum bias collisions (0–80% of the total hadronic cross-section). The  $\Xi^- + \Xi^+$  and  $\Omega^- + \Omega^+$  signals appear as clear peaks around the rest masses (indicated by the vertical arrows) in the invariant mass distribution, above a combinatorial background. The combinatorial background of uncorrelated decay candi-




FIG. 1: (color online). (a)  $\Xi^- + \Xi^+$  and (b)  $\Omega^- + \Omega^+$  invariant mass distribution from minimum bias (0–80%) Au + Au collisions at  $\sqrt{s_{\text{NN}}} = 200$  GeV. The solid line shows the combinatorial background as estimated from a same event rotating method (see text for details). Azimuthal distributions with respect to the event plane of the (c)  $\Xi^- + \Xi^+$  and (d)  $\Omega^- + \Omega^+$  raw yields. Dashed lines represent the fit results. All plots shown include  $\Xi^- + \Xi^+$  and  $\Omega^- + \Omega^+$  in the transverse momentum range  $1 < p_T < 4$  GeV/c.

dates under the peak can be determined by sampling the regions on both sides of the peak. It can also be reproduced by rotating the  $\Lambda$  candidates by  $180^\circ$  in the transverse plane and then reconstructing the  $\Xi$  and  $\Omega$  candidates. The rotation of the  $\Lambda$  breaks the correlation in the invariant mass and therefore mimics the background of uncorrelated decay pairs. Both background determination methods provide consistent results. In Fig. 1 (a) and (b), the combinatorial background as calculated from the rotation method is shown as solid lines. Outside the region of the corresponding mass peak, the rotation method describes the background well. The residual *bump* at lower invariant mass than the peak in Fig. 1 (a) can be understood as fake  $\Xi$  candidates being reconstructed as  $\Xi_{\text{fake}}(\pi_\Lambda, \Lambda_{\text{fake}}(\pi_{\text{random}}, p_\Lambda))$ , where  $\pi_\Lambda$  and  $p_\Lambda$  are the daughters of a real  $\Lambda$  and  $\pi_{\text{random}}$  is a random  $\pi$ . The real correlation between  $\pi_\Lambda$  and  $p_\Lambda$  remains in the  $\Xi_{\text{fake}}$  reconstruction resulting in the observed bump in the  $\Xi$  invariant mass distribution. A similar mis-association happens in the  $\Omega$  case with the addition of the  $\pi_\Lambda$  being misidentified as a kaon. Our studies have shown that this residual correlation does not affect the signal peak. The raw yields are then extracted from the invariant mass distribution by counting the number of entries in the mass peak above the estimated background.

The elliptic flow  $v_2$  is calculated from the distribution of particle raw yields as a function of azimuthal angle  $\phi$  with respect to the event plane angle  $\Psi$ . The  $\Xi$  and  $\Omega$  candidates are divided in  $\phi - \Psi$  bins, and the raw yields for each bin are

extracted from the invariant mass distributions as described above. The event plane angle  $\Psi$  is used as an estimate of the reaction plane angle [27, 28]. Here, the event plane is determined from the azimuthal distribution of charged primary tracks with  $0.2 < p_T < 2.0$  GeV/c and pseudo-rapidity  $|\eta| < 1.0$ . To avoid autocorrelations, tracks associated with a  $\Xi$  or  $\Omega$  candidate are explicitly excluded from the event plane calculation. Figure 1 shows the azimuthal distributions of raw yields for (c)  $\Xi^- + \Xi^+$  and (d)  $\Omega^- + \Omega^+$  with respect to the event plane from the minimum bias collisions in the  $1 < p_T < 4$  GeV/c range. To reduce the statistical uncertainties in the  $\Xi$  and  $\Omega$  signal extraction and because of the  $\cos 2(\phi - \Psi)$  dependence of  $v_2$ , we have folded around  $\pi/2$  the candidates in the  $\pi/2 < \phi - \Psi < \pi$  range into the  $\pi/2 > \phi - \Psi > 0$  range. The distributions exhibit a clear oscillation with azimuthal angle  $\phi - \Psi$  for both  $\Xi$  and  $\Omega$  particles indicating the presence of significant elliptic flow. Dashed lines are results from fitting a function  $\frac{dN}{d(\phi - \Psi)} = A[1 + 2v_2 \cos 2(\phi - \Psi)]$ , where  $A$  is the normalization constant. Furthermore, we note that the amplitude of the oscillation for the  $\Xi$  and  $\Omega$  are of similar magnitude indicating that their  $v_2$  is similar, as will be discussed later. The finite resolution in the event plane determination smears out the azimuthal distributions and leads to a lower signal in the apparent anisotropy [28]. We determine the event plane resolution by dividing each event into random sub-events and determine the correction factor to be  $1/0.72$  for minimum-bias collisions. In the following, all numbers reported on  $v_2$  are corrected for this resolution. Systematic uncertainties in  $v_2$  were studied by comparing the background determination methods described above and by changing the cuts used in the  $\Xi$  and  $\Omega$  reconstruction. For the  $\Xi$ , the estimated absolute systematic uncertainties are 0.02 for the lowest  $p_T$  bin and smaller than 0.01 for all other  $p_T$  bins. For the  $\Omega$ , the absolute systematic uncertainty is 0.04 for both measured transverse momentum bins. Correlations unrelated to the reaction plane (non-flow effects) can modify the apparent  $v_2$  [11]. Non-flow contributions for multi-strange baryons were not studied yet, but are expected to be similar to those calculated for  $\Lambda$  ( $\sim -0.01$  at  $p_T=1$  GeV/c and  $\sim -0.04$  at  $p_T=2.5$  and 4.0 GeV/c) [11].

Figure 2 shows the results of the elliptic flow parameter  $v_2(p_T)$  for multi-strange baryons (a)  $\Xi^- + \Xi^+$  and (b)  $\Omega^- + \Omega^+$  from minimum bias (0–80%) Au + Au collisions. As a reference, the open symbols represent the published [11]  $K_S^0$  and  $\Lambda$   $v_2(p_T)$  from the same event class. As guideline, results of the fit [29] to  $v_2(p_T)$  of  $K_S^0$  and  $\Lambda$  are shown as dot-dashed-lines. Hydrodynamic model calculations using an Equation Of State (EOS) with a phase transition at  $T_c = 165$  MeV and a thermal freeze-out at  $T_{fo} \sim 130$  MeV [30] are shown as dotted lines, from top to bottom, for  $\pi$ ,  $K$ ,  $p$ ,  $\Lambda$ ,  $\Xi$ , and  $\Omega$ , respectively. The expected mass ordering in hydrodynamics of  $v_2(p_T)$  is observed with lighter particles having larger  $v_2(p_T)$  than heavier particles.

First, we observe in Fig. 2 (a) that for  $\Xi$ , the  $v_2$  increases with  $p_T$  reaching a saturation value of  $\sim 18\%$  at  $p_T \sim 3.0$




FIG. 2: (color online).  $v_2(p_T)$  of (a)  $\Xi^- + \Xi^+$  and (b)  $\Omega^- + \Omega^+$  from 200 GeV Au + Au minimum bias collisions. The  $v_2$  of  $K_S^0$  and  $\Lambda$  [11] are also shown as open symbols and the results of the fits [29] are shown as dashed-dot lines. Hydrodynamic model calculations [30] are shown as dotted curves for  $\pi$ ,  $K$ ,  $p$ ,  $\Lambda$ ,  $\Xi^-$ , and  $\Omega^-$  masses, from top to bottom, respectively.

GeV/c. This is similar to the result for  $\Lambda$  baryons [11]. In the lower  $p_T$  region ( $p_T < 2.5$  GeV/c), the  $\Xi$  results are in agreement with the hydrodynamic model prediction [30]. Second, we observe in Fig. 2 (b) that the values of  $v_2$  for the  $\Omega$ , are clearly non-vanishing although they have larger statistical uncertainties due to their smaller abundance. Over the measured  $p_T$  range and considering the statistical uncertainties, the  $v_2$  of the  $\Omega$  is non-zero with 99.73% confidence level ( $3\sigma$  effect). The  $\Omega$   $v_2$  values are, within uncertainties, consistent with those measured for the  $\Xi$ , indicating that even the triply-strange baryon  $\Omega$  has developed significant elliptic flow in Au + Au collisions at RHIC. In the scenario where multi-strange baryons are less affected by the hadronic stage [20] and where  $v_2$  develops primarily at the early stage of the collision [22, 23], the large  $v_2$  of multi-strange baryons reported in this paper shows that partonic collectivity is generated at RHIC.

Previously, a particle type (baryon versus meson) difference in  $v_2(p_T)$  was observed for  $\pi$  and  $p$  [31] as well as for  $K_S^0$  and  $\Lambda$  [11] at the intermediate  $p_T$  region. The present results




FIG. 3: (color online). Number of quark ( $n_q$ ) scaled  $v_2$  as a function of scaled  $p_T$  for  $\Xi^- + \Xi^+$  (filled circles) and  $\Omega^- + \Omega^+$  (filled squares). Same distributions also shown for  $\pi^+ + \pi^-$  (open diamonds),  $p + \bar{p}$  (open triangles) [31],  $K_S^0$  (open circles),  $\Lambda + \bar{\Lambda}$  (open squares) [11]. All data are from 200 GeV Au + Au minimum bias collisions. The dot-dashed-line is the scaled result of the fit to  $K_S^0$  and  $\Lambda$  [29].

on the  $\Xi$   $v_2(p_T)$  follow closely the ones for  $\Lambda$  confirming that this observed particle type difference is a meson-baryon effect rather than a mass effect. This particle type dependence of the  $v_2(p_T)$  is naturally accounted for by quark coalescence or recombination models [32, 33, 34]. In these hadronization models, hadrons are formed dominantly by coalescing massive quarks from a partonic system with the underlying assumption of collectivity among these quarks. Should there be no difference in collectivity among  $u$ -,  $d$ -, and  $s$ -quarks near hadronization, these models predict a universal scaling of  $v_2$  and the hadron transverse momentum  $p_T$  with the number of constituent quarks ( $n_q$ ). This scaling has previously been observed to hold within experimental uncertainties for the  $K_S^0$  and the  $\Lambda$  when  $p_T/n_q \geq 0.7$  GeV/c [11].

The  $n_q$ -scaled  $v_2$  versus the  $n_q$ -scaled  $p_T$  are shown in Fig. 3 for  $\pi^- + \pi^+$  (open diamonds),  $p + \bar{p}$  (open circles) [31],  $K_S^0$  (open triangles),  $\Lambda + \bar{\Lambda}$  (open squares) [11],  $\Xi^- + \Xi^+$  (filled circles) and  $\Omega^- + \Omega^+$  (filled squares). Except for pions, all hadrons including  $\Xi$  and  $\Omega$  scale well within statistics. The discrepancy in the pion  $v_2$  may in part be attributed to its Goldstone boson nature (its mass is smaller than the sum of its constituent quark masses) or to the effects of resonance decays (a large fraction of the measured pions will come from the decays of resonances at higher  $p_T$ ) [29, 35]. This further success of the coalescence models in describing the multi-strange baryon  $v_2(p_T)$  also lends strong support to the finding that collectivity developed in the partonic stage at RHIC. In addition, the good agreement of  $v_2(p_T/n_q)/n_q$  for  $p(uud)$ ,  $\Lambda(uds)$ ,  $\Xi(dss)$  and  $\Omega(sss)$  further supports the idea that the partonic flow of  $s$  quarks is similar to that of  $u, d$  quarks. Future measurements with higher statistics, specially for the  $\Omega$ , will allow for a more quantitative comparison.

In summary, we reported the STAR results on multi-strange baryon,  $\Xi^- + \Xi^+$  and  $\Omega^- + \Omega^+$ , elliptic flow  $v_2$  from minimum bias Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. The observations of sizable elliptic flow and the constituent quark scaling behavior for the multi-strange baryons suggest that substantial collective motion has been developed prior to hadronization in the high-energy nuclear collisions at RHIC.

We thank the RHIC Operations Group and RCF at BNL, and the NERSC Center at LBNL for their support. This work was supported in part by the HENP Divisions of the Office of Science of the U.S. DOE; the U.S. NSF; the BMBF of Germany; IN2P3, RA, RPL, and EMN of France; EPSRC of the United Kingdom; FAPESP of Brazil; the Russian Ministry of Science and Technology; the Ministry of Education and the NNSFC of China; IRP and GA of the Czech Republic, FOM of the Netherlands, DAE, DST, and CSIR of the Government of India; Swiss NSF; the Polish State Committee for Scientific Research; and the STAA of Slovakia.

- 
- [1] F. Karsch, Nucl. Phys. **A698**, 199c (2002).
  - [2] Z. Fodor, Nucl. Phys. **A715**, 319c (2003) and references therein; Z. Fodor and S.D. Katz, Phys. Lett. **B534**, 87 (2002).
  - [3] C. Adler *et al.* (STAR Collaboration), Phys. Rev. Lett. **89**, 202301 (2002).
  - [4] J. Adams *et al.* (STAR Collaboration), Phys. Rev. Lett. **91**, 172302 (2003).
  - [5] J. Adams *et al.* (STAR Collaboration), Phys. Rev. Lett. **91**, 072304 (2003).
  - [6] K. Adcox *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **88**, 022301(2002).
  - [7] S.S. Adler *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **91**, 072301 (2003).
  - [8] S. S. Adler *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **91**, 072303 (2003).
  - [9] B. B. Back *et al.* (PHOBOS Collaboration), Phys. Rev. Lett. **91**, 072302 (2003).
  - [10] I. Arsene *et al.* (BRAHMS Collaboration), Phys. Rev. Lett. **91**, 072305 (2003).
  - [11] J. Adams *et al.* (STAR Collaboration), Phys. Rev. Lett. **92**, 052302 (2004).
  - [12] C. Adler *et al.* (STAR Collaboration), Phys. Rev. Lett. **87**, 182301 (2001).
  - [13] P. F. Kolb and U. Heinz, arXiv:nucl-th/0305084.
  - [14] D. Teaney, J. Lauret and E. V. Shuryak, arXiv:nucl-th/0110037.
  - [15] H. van Hecke, H. Sorge and N. Xu, Phys. Rev. Lett. **81**, 5764 (1998).
  - [16] S.A. Bass *et al.*, Phys. Rev. C **60**, 021902 (1999); A. Dumitru, S.A. Bass, M. Bleicher, H. Stöcker, W. Greiner, Phys. Lett. **B460**, 411 (1999); S.A. Bass and A. Dumitru, Phys. Rev. C **60**, 064909 (2000).
  - [17] Y. Cheng *et al.*, Phys. Rev. C **68**, 034910 (2003).
  - [18] S. F. Biagi *et al.*, Nucl. Phys. **B186**, 1 (1981).
  - [19] R. A. Muller, Phys. Lett. B **38**, 123 (1972).
  - [20] J. Adams *et al.* (STAR Collaboration), Phys. Rev. Lett. **92**, 182301 (2004).
  - [21] C. Alt *et al.* (NA49 Collaboration), submitted to Phys. Rev. Lett. nucl-ex/0409004.
  - [22] H. Sorge, Phys. Rev. Lett. **82**, 2048 (1999).

- [23] J. Y. Ollitrault, Phys. Rev. D **46**, 229 (1992).
- [24] K. H. Ackermann *et al.* (STAR Collaboration), Nucl. Instrum. Meth. A **499**, 624 (2003).
- [25] M. Anderson *et al.*, Nucl. Instrum. Meth. A **499**, 659 (2003).
- [26] C. Adler *et al.* (STAR Collaboration), Phys. Rev. Lett. **89**, 092301(2002).
- [27] K. H. Ackermann *et al.* (STAR Collaboration), Phys. Rev. Lett. **86**, 402 (2001).
- [28] A. Poskanzer and S. A. Voloshin, Phys. Rev. C **58**, 1671 (1998).
- [29] X. Dong, S. Esumi, P. Sorensen, N. Xu , Z. Xu, Phys. Lett. **B597**, 328 (2004).
- [30] P. Huovinen, P.F. Kolb, U. Heinz, P.V. Ruuskanen, and S. Voloshin, Phys. Lett. **B503**, 58 (2001).
- [31] S. S. Adler *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **91**, 182301 (2003).
- [32] S. Voloshin, Nucl. Phys. **A715**, 379c (2003); D. Molnar and S. Voloshin, Phys. Rev. Lett. **91**, 092301 (2003).
- [33] R.J. Fries, B. Müller, C. Nonaka, S.A. Bass, Phys. Rev. Lett. **90**, 202303 (2003).
- [34] Z. Lin and C. Ko, Phys. Rev. Lett. **89**, 202302 (2002).
- [35] V. Greco and C. Ko, Phys. Rev. C **70**, 024901 (2004).