

# Changes in Volatile Profiles and Activity of Hydroperoxide Lyase and Alcohol Dehydrogenase During the Development of Cabernet Sauvignon Grapes (*Vitis vinifera* L.)

Ya Nan' OuYang<sup>1</sup>, Jin Shan Gao<sup>1</sup>, Ruo Lan Li<sup>1</sup>, Mei Rong Zhu<sup>1</sup>, Xin Hao Hu<sup>1</sup>, Zhuo Min<sup>1</sup>, Shu Xia Chen<sup>2</sup>, Zhen Wen Zhang<sup>1,3</sup>, Yu Lin Fang<sup>1,3\*</sup>

(1) College of Oenology, Northwest A&F University, Yangling, Shaanxi 712100, China

(2) College of Horticulture, Northwest A&F University, Yangling, Shaanxi 712100, China

(3) Grape and Wine Engineering Research Centre in Shaanxi Province, Yangling, Shaanxi 712100, China

Submitted for publication: April 2015

Accepted for publication: October 2015

Key words: Alcohol dehydrogenase, Cabernet Sauvignon, hydroperoxide lyase, volatiles

**In this study we focused on the development of Cabernet Sauvignon grapes and investigated changes in the activity of alcohol dehydrogenase (ADH) and hydroperoxide lyase (HPL) in different tissues. We sampled grape skin at four, six, seven, eight, nine, 10, 12, 14 and 16 weeks after anthesis; developing flowers when blooming at 0%, 5%, 50%, and 90%; and leaves at two and four weeks before anthesis and at two, four, six, eight, nine, and 10 weeks after anthesis. We also examined the type and fluctuation of volatile contents. ADH activity increased with the development of flowers and grape skins, which led to the increasing of types and concentration of alcohols. Low levels of 9-HPL led to low concentrations of C9 compounds. According to this paper, C6 compounds became abundant with the development of grape berries, while the activity of 13-HPL kept at a low level in the flowers and grape skins. There might have been a high level of 13-HPL activity from the end of flowering until fruit setting that we did not detect. Furthermore, similar C6 and C5 compounds were detected across all tissues, including hexanal, (E)-2-hexenal, (Z)-3-hexenal, (Z)-2-penten-1-ol, (Z)-3-hexen-1-ol, 1-hexanol and 3-hexen-1-ol. Generally speaking, the concentrations of C6 and C5 compounds could be used as the criterion of maturation of the three grape tissues.**

## INTRODUCTION

Green leaf volatiles (GLVs) are short-chain acyclic aldehydes, alcohols and esters that form as a result of the catalysis by hydroperoxide lyases (HPLs) and alcohol dehydrogenase (ADH), and they are the main source of the green aroma in grapes. GLVs are known for the aroma of crushed or injured leaves (Hatanaka, 1993; Matsui *et al.*, 2006), and plants produce GLVs after damage and biotic or abiotic stress. GLVs are not only the main source of volatiles in fruit and vegetables – they also have important physiological functions. They are critical signals that help plants recognise and compete with other organisms (Matsui *et al.*, 2006) and also impart resistance to pathogenic bacteria and insects (Nakamura & Hatanaka, 2002; Tufariello *et al.*, 2012). Furthermore, GLVs are similar to the aroma from spices in the food and beverage industry (Kalua & Boss, 2009).

The presence of short-chain volatiles in grape juice and wine, as well as their contribution to the final flavour and aromatic composition of wine, has been well documented

(Baumes *et al.*, 1988; Terrier *et al.*, 1996). The major components of these short-chain volatiles are aldehydes, alcohols and esters. They are primarily synthesised via the metabolic pathway of aliphatic acid (Fig. 1S) (Cynthia & Kenji, 2014). During the maturation of the grape berry, the metabolism of aliphatic acid can be catalysed directly by lipoxygenase (LOX) and the following steps of metabolism are similar to the GLV synthesis pathway described above.

In a previous study, the volatiles formed during the development of three types of non-aromatic grape cultivars revealed that only C6 volatile compounds were present in high concentrations; therefore, changes in the proportion of C6 volatile compounds can be used to confirm optimum maturity of the fruit (Gomez *et al.*, 1995). Other studies revealed that the proportion of volatiles increased significantly after grape colouring (Park *et al.*, 1991; Coelho, *et al.*, 2007). The major C6 volatiles found during the early, middle and late periods of grape development are acetates, aldehydes and alcohols respectively. The ratio of alcohols to aldehydes influences the aroma of grapes and wines. Thus,

\*Corresponding author: Yu Lin Fang, E-mail address: fangyulin@nwsuaf.edu.cn

Acknowledgements: The authors would like to express their gratitude to the Programme of State Forestry Administration, 948 Project [2009-4-09]; The National Eleventh Five-year Research Programme of China [2012BAD31B07]; and the National Technology System for Grape Industry [nycytx-30-2p-04], for the generous financial support provided for this work.

changes in volatiles during grape development indicate greater dependence on enzyme activity and specificity than on the unsaturated fatty acids.

LOX-HPL is the confirmed pathway for GLV formation and, in many other species, it is dependent on the activity of ADH and HPLs. The dependence on LOX, HPLs, ADH, acetyltransferase (AAT) and aldehyde isomerase (AH) in the LOX pathway during grape development allows for the control of grape and wine aroma (Kalua & Boss, 2009). Moreover, several studies have suggested that most unsaturated fatty acids are metabolised through the 13-HPL pathway during the maceration of grape must (Hashizume & Samuta, 1997; Coelho *et al.*, 2007; Jiang *et al.*, 2011). Moreover, the pleasant flavours in cucumbers (*Cucumis sativus*) result from the enzymatic action of LOX on linolenic acid and linoleic acid. LOX introduces molecular oxygen at the C13 or C9 of linoleic acid or linolenic acid, thereby forming 13-hydroperoxylinolenic acid (13-HPOT) or 9-hydroperoxylinolenic acid (9-HPOT). HPL cleaves 13-HPO (13-hydroperoxide) and 9-HPO (9-hydroperoxide), producing C6 and C9 aldehydes, which are responsible for the flavours in cucumber (Matsui *et al.*, 2000).

In contrast to other plants, very few studies have examined the formation and regulation of GLVs in grapes, especially for Cabernet Sauvignon, which is one of the most important *Vitis vinifera* varieties cultivated in China. Our study focused on the changes in the activity of ADH and HPLs, as well as on the development of the characteristic aromatic compounds, to investigate the relationship between key enzyme activity and volatile contents. Our research will enhance our understanding of the enzymes involved in the formation of GLVs during the development of characteristic grape aromas. It will lead to the establishment of links between changes in grapevine physiology and wine quality.

## MATERIALS AND METHODS

### Plant material and sampling

The Cabernet Sauvignon (*V. vinifera* L.) grapes grown at the Caoxinzhuang vineyard (Yangling, Shaanxi Province, China) were planted in March 2008. Samples were collected during anthesis and grape development from April to May and July to September 2012 respectively. Grape berries were randomly sampled four to 16 weeks after anthesis, and flowers were collected randomly during inflorescence (0, 5, 50 and 90% florets flowering per cluster). The third to fifth leaves from the bottom of the shoots were collected two and four weeks before anthesis, as well as two, four, six, eight, nine and 10 weeks after anthesis. The five-point random sampling method was used (picking four points with the same distance from the centre point of diagonal lines) for each replicate and there were three replicates. Approximately 30 standard clusters, 30 leaves and 15 inflorescence were sampled at each stage or time point, and the grape berries were separated from the bunches. The grape berries were snap-frozen in liquid nitrogen, and the grape skins were separated from the berries and transferred to a tube. Leaves and flowers were immediately snap-frozen in liquid nitrogen and stored at -80°C.

### Analysis of enzyme activity

For enzyme extraction from the HPLs, the sample (1 g) was pulverised in liquid nitrogen and dissolved in a sodium acetate buffer (3 mL; 50 mM, pH 5.0, containing 0.21% vitamin C and 0.5% [w/v] Triton X-100). The sample was then centrifuged at  $14\,000 \times g$  and 4°C for 20 min. The supernatant liquid was the crude extract of the HPL enzyme.

To extract the potato 9-LOX protein, a slice of potato was ground into slurry using liquid nitrogen and dissolved in a 50 mM potassium phosphate buffer (pH 7.0 with 2 mM sodium hydrogen sulphite). The solution was then centrifuged at  $12\,000 \times g$  for 30 min, and ammonium sulphate was added to the resultant supernatant over a 2 h stirring period to obtain 50% saturation. After centrifugation, the pellet was re-suspended in a 50 mM potassium phosphate buffer (pH 7.0) and dialysed overnight using a 50 mM potassium phosphate buffer (pH 7.0) and water. All work was performed at 4°C (Suurmeijer *et al.*, 1998) and the protein levels were measured by the Coomassie brilliant blue method (Smith *et al.*, 1985).

The activity of HPLs was determined by adding 50 µL of 10 mM NADH, 50 µL of 5 U/µL ADH, 420 µL of substrate (product of linoleic acid catalysed by soybean 13-LOX [Sigma, Aldrich] or potato 9-LOX protein), and 250 µL of crude HPLs enzyme extract to 4.5 mL of 0.1 M potassium phosphate buffer (pH 6.0). The reaction proceeded for 4 min at 20°C, and then the absorbance was recorded at 340 nm for 15 s (Anthon & Barrett, 2003).

For ADH enzyme extraction, the samples were ground using liquid nitrogen and added to 100 mM MES buffer (pH 6.5) containing 2 mM DTT and 1% w/v PVPP in a ratio of 1:3 (sample:buffer). The sample was homogenised, filtered by gauze and then centrifuged at  $14\,000 \times g$  and 4°C for 20 min. The resulting supernatant was the crude extract of ADHs.

The ADHs activity was determined using 0.05 mL of 1.6 mM NADH, 0.05 mL of 80 mM acetaldehyde, and 0.1 mL of crude extract ADHs in 0.8 mL of 100 mM MES buffer (pH 6.5). The absorbance was recorded at 340 nm for 15 s (Defilippi *et al.*, 2005). Three biological replicates were analysed in three technical replicates

### Analysis of volatile compounds

The volatile compounds were extracted by headspace–solid-phase micro-extraction (HS-SPME) and analysed using a Gas Chromatograph-Mass Spectrometer (GC-MS, TRACE DSQ, Finnigan, USA) according to the methods of Song and Barros (Perestrelo *et al.*, 2011; Song *et al.*, 2012). The grape skin, flowers and leaf samples were ground under liquid nitrogen and the resulting powder (20 g) was suspended in 20 mL of water and allowed to soak in the dark for 24 h at 4°C in a nitrogen atmosphere. The mixture was then centrifuged at  $8\,000 \times g$  for 30 min, and the supernatant was used for the volatile analysis.

The extracted liquid (8 mL) was added to a 15 mL vial containing 2 g of NaCl and 5 µL of 4-methyl-2-pentanol (0.93 g/L, internal standards). The vial was sealed and heated to 30°C for 10 min. Then, a fibre (50/30µm PSMS/CAR/DVB, Supelco, USA) that was pre-conditioned at 250°C for 1 h was inserted into the headspace of the vial

for 40 min. The fibre was subsequently desorbed in the GC injector at 250°C for 5 min. Three independent biological replicates were analysed for each sample. The GC-MS was performed with a DB-WAX elastic quartz capillary column, an Xcalibur workstation, and the 2002 NIST Standard Reference Database. The oven temperature was maintained at 40°C for 2.5 min, increased to 230°C at a rate of 6°C/min and then held at 230°C for 7 min. The MS transfer line and ion source temperatures were 230 and 200°C respectively. A constant helium flow of 1 mL/min was used, and the electron impact mass spectrometric data from the *m/z* 35 to 400 range were collected with an ionisation voltage of 70 eV, an emission current of 100  $\mu$ A, and a detection voltage of 1.4 kV. Pentanal, butanoic acid, ethyl ester, hexanal, (E)-2-pentenal, heptanal, (E)-2-hexenal, 1-pentanol, (Z)-2-penten-1-ol, 1-hexanol, (E)-2-hexen-1-ol, 1-nonanol and (E/Z)-6-nonen-1-ol were used as individual standards to make calibration curves. The individual standards were dissolved in ethanol and then diluted with water to obtain a set of serial dilutions to establish the calibration curve. The concentration of volatile compounds for which it was not possible to establish calibration curves were estimated on the basis of the equations of compounds with the same functional group and/or similar numbers of C atoms. GC data was normalised by the area normalisation method to calculate the component concentration.

### Statistical analysis

Mean values, data normalisation (Z-score), PCA analysis and cluster (TwoStep cluster) analysis were calculated and performed using SPSS (Version 19.0). Heatmap and hierarchical clustering analysis were performed to display the change in concentration and types of volatiles in three different tissues by MeV-4-9-0 using normalised and mean-centred data. All data presented were the mean values.

## RESULTS AND DISCUSSION

### Changes in volatile compounds in different tissues

Calibration curves of 12 individual standards are shown in Table S1. In total, 36 volatile compounds were detected in the developing grape skin from the 2012 harvest, including 14 aldehydes, 18 alcohols and 10 C6 compounds (Table 1). A heatmap representation of the volatile profiles of grape skins is shown in Fig. 1, revealing that large numbers of volatiles were highly responsive to the sampling dates. Hexanal, benzaldehyde, furfural, 3-methyl-butanal, 2-methyl-butanal, propanal, 1-heptanol, 1-hexanol, (E)/(Z)-2-hexen-1-ol, etc. were the most abundant volatiles present in the grape skins at 16 weeks after anthesis.

All of the volatile profiles clustered into three categories. The first cluster refers to those whose concentrations were abundant at a later development stage of the grape berries. The concentration of the second cluster was higher at the early stage and included 3,7-dimethyl-1,6-octadien-3-ol, 2-hexanol, borneol, (Z)/(E)-2-heptenal, etc. The third cluster of volatiles contained some C5 and C6 compounds and the concentrations of C5, C6, C9 aldehyde were abundant at 12 weeks. Many studies have shown that the typical aroma of wine from Cabernet Sauvignon grapes comes from C6 compounds (Xu *et al.*, 2015). Our results reveal

that different structures of C6 and C5 compounds occurred at different development stages. Most alcohol compounds were detected at 16 weeks and this was consistent with the pathway of LOX-HPL-ADH. This indicated that the fruity and flowery aromas of grape berries increased with grape development. Conversely, the content of eucalyptol (minty odour) declined with berry development, with the highest amounts occurring in the earlier stages. D-limonene (fresh citrus odour) was only identified during some of the berry development stages and reached its highest content 10 weeks after anthesis. Eucalyptol was the predominant monoterpene in the early stage of berry development, which indicates that it was probably left by the flowers (Delphia *et al.*, 2007; Köllner *et al.*, 2008).

Principle component analysis was applied to interpret the results, and the components are shown in Fig. 2. The first three components explained 70.58% of the variance, which was caused by the difference in volatile components of the grape skins from different sampling dates. According to the loading plot and the PCA score plot, the concentrations of (E)-2-pentenal, 2,6-dimethyl-4-heptanone, and (E,E)-2,4-hexadienal were different in the grape skins sampled at 12 weeks after anthesis than those from other development stages. The compounds 3-hexen-1-ol, borneol, (S)-3-ethyl-4-methyl-pentanol and 2-ethyl-furan were associated with samples collected at 14 weeks after anthesis. In addition, 3-methyl-butanal, furfural, benzaldehyde, (E)/(Z)-2-hexen-1-ol, 1-heptanol and 1-octanol contributed more to the separation of samples at 16 weeks after anthesis (i.e., at harvest). The results of the PCA analysis are consistent with the heatmap analysis.

Forty volatile compounds were identified in the developing flowers, including six aldehydes and 20 alcohols (Table 2). Two different clusters of volatile profiles are displayed in Fig. 3. One of these contained more aldehydes and alcohols. Fig. 3 shows that the concentrations of aldehydes and alcohols were abundant when the flowers were at the bud stage. The other cluster contained more ketones and terpenes. The concentrations of caryophyllene,  $\alpha$ -caryophyllene,  $\alpha$ -farnesene, caryophyllene oxide, 2-tridecanone, cyclopentadecanone, etc. increased at the 50% blooming stage of the flowers. There also were some C5, C6 and C9 alcohols in the second cluster, such as 1-penten-3-ol, 2-nonanol, 4-methyl-5-hexen-2-ol, cyclopentanol, (Z)-2-hexen-1-ol, (E)-2-hexen-1-ol and 1-hexanol. All of the volatiles from the second cluster became abundant at the stage of 5% flower blooming until full blooming. The contents of the aldehydes and alcohols as important components of green leaf odour generally declined during flower development. Various terpenes were identified in the blooming flowers and reached their highest contents in the 50% flowering stage (Table 2). Bestmann *et al.* (1997) showed that limonene and the  $\alpha$ -terpenes were the monoterpenes typically found in flowers, while eucalyptol and caryophyllene were the highest oxyterpene and sesquiterpene respectively.

PCA analysis revealed that the first three components explained 100% of the variance, with the first component explaining 50.29% of the total variability. Fig. 4 shows that caryophyllene,  $\alpha$ -caryophyllene,  $\alpha$ -farnesene, caryophyllene oxide, 2-tridecanone, 2-pentadecanone, 8-heptadecene,

TABLE 1  
Volatile compounds in grape skin during development in 2012.

Compound	symbol	Weeks after anthesis/w								
		4	6	7	8	9	10	12	14	16
		Concentration (µg/kg)								
<b>Aldehydes</b>										
Propanal	bs 1	nd	0.37	0.50	0.80	1.05	1.50	1.15	1.41	1.65
2-methyl-butanal	bs 2	nd	0.42	0.65	0.82	1.05	1.29	2.19	2.44	2.60
Pentanal	bs 3	4.22	5.24	4.67	6.64	40.22	16.61	16.66	9.22	9.65
Hexanal	bs 4	17.46	12.45	9.28	17.93	162.27	347.32	317.09	526.22	703.26
(E)-2-pentenal	bs 5	nd	nd	nd	nd	nd	nd	1.19	nd	nd
(Z)-3-hexenal	bs 6	nd	0.52	0.94	1.50	2.27	5.29	10.71	11.89	2.04
2-methyl-4-pentenal	bs 7	nd	nd	nd	nd	105.74	nd	nd	nd	nd
3-methyl-butanal	bs 8	nd	0.42	0.88	1.91	2.03	1.29	1.89	2.44	6.03
(E)-2-hexenal	bs 9	12.06	11.76	11.11	15.21	115.51	413.02	1964.01	762.67	377.38
(Z)/(E)-2-heptenal	bs 10	nd	0.29	nd	nd	nd	nd	nd	nd	nd
Nonanal	bs 11	nd	nd	nd	nd	0.78	0.99	1.68	1.36	nd
Furfural	bs 12	nd	nd	nd	nd	nd	nd	nd	nd	1.48
(E,E)-2,4-hexadienal	bs 13	1.61	nd	nd	nd	1.35	2.98	8.93	1.11	1.95
Benzaldehyde	bs 14	nd	nd	nd	nd	nd	nd	nd	nd	29.24
<b>Alcohols</b>										
1-Penten-3-ol	bs 15	3.25	3.18	1.40	2.98	6.85	11.12	11.61	6.44	5.01
2-Hexanol	bs 16	185.97	186.00	nd	nd	1.31	6.78	118.00	98.78	90.40
3-Methyl-1-butanol	bs 17	0.80	1.02	0.51	0.63	1.48	6.10	5.95	9.79	6.22
3-Methyl-3-buten-1-ol	bs 18	nd	nd	nd	nd	0.96	4.00	2.38	4.44	nd
1-Pentanol	bs 19	0.89	1.09	0.68	0.62	1.83	6.10	5.95	9.78	6.05
(Z)-2-Penten-1-ol	bs 20	nd	6.32	nd	4.02	9.86	15.12	17.41	11.56	9.38
1-Hexanol	bs 21	13.01	4.15	6.01	7.34	16.80	29.77	34.67	98.78	90.40
3-Hexen-1-ol	bs 22	14.62	6.61	6.62	6.85	10.95	14.24	19.64	26.11	7.89
(E)/(Z)-2-hexen-1-ol	bs 23	6.17	1.09	2.46	2.74	7.59	19.26	41.07	nd	54.67
1-Heptanol	bs 24	nd	nd	nd	nd	nd	1.29	1.19	nd	1.48
1-Octen-3-ol	bs 25	0.72	0.46	0.71	nd	1.09	3.39	2.53	4.00	3.90
3,7-Dimethyl-1,6-octadien-3-ol	bs 26	1.44	1.49	1.09	0.73	nd	nd	0.89	2.22	1.30
1-Octanol	bs 27	nd	nd	nd	nd	nd	2.30	2.98	4.00	4.94
1-Nonanol	bs 28	nd	0.40	0.61	nd	nd	nd	1.49	1.56	1.30
Benzyl alcohol	bs 29	nd	nd	nd	nd	nd	2.92	3.27	5.89	4.73
Phenylethyl alcohol	bs 30	nd	nd	nd	nd	0.70	4.82	5.21	8.44	6.12
Borneol	bs 31	2.81	3.44	2.46	4.29	3.14	2.78	2.38	3.67	2.23
(S)-3-Ethyl-4-methylpentanol	bs 32	nd	nd	nd	nd	nd	nd	nd	14.44	nd
<b>Others</b>										
2-Ethyl-furan	bs 33	1.84	4.01	2.03	1.31	2.18	1.76	3.87	5.67	1.95
Eucalyptol	bs 34	20.18	15.63	3.86	1.52	nd	nd	nd	nd	nd
2,6-Dimethyl-4-heptanone	bs 35	4.64	3.24	4.39	3.56	5.41	2.85	5.66	2.67	3.71
D-Limonene	bs 36	0.78	nd	nd	nd	nd	4.34	nd	1.56	nd

nd, not detected




FIGURE 1  
Heatmap of the concentration of volatiles in grape skin.  
Note: W (weeks after anthesis)




FIGURE 2  
Principle component analysis of volatile compounds of grape skin.  
A. PCA score plot. B. Loading plot of components 1, 2 and 3.  
Note: “bs” refers to volatiles in grape skins. For details, refer to the Table 1.

cyclopentadecanone and cyclotridecanone were major contributors to the variation among flower stages, and could discriminate the 50% blooming stage from others (0%, 5%, 90%). (E)-2-hexenal, (E,E)-2,4-hexadienal, 2-butyl-2-octenal, 2-heptanol, 1-pentanol, 1-heptanol, (R)-3,7-dimethyl-6-octen-1-ol, 3,7-dimethyl-2,6-octadien-1-ol and

benzyl alcohol contributed more to the volatile composition of the inflorescences, whereas the concentration of these compounds decreased or even declined to zero with the development of flowers.

In the developing leaves, 53 volatile compounds were detected, including 17 aldehydes and 25 alcohols (Table 3).

TABLE 2  
Volatile compounds in flowers during development in 2012.

Compound	Symbol	Percentage of flowering/%			
		0	5	50	90
		Concentration ( $\mu\text{g}/\text{kg}$ )			
<b>Aldehydes</b>					
Pentanal	F1	20.67	20.20	nd	nd
Hexanal	F2	2082.61	184.96	187.09	235.13
(Z)-3-Hexenal	F3	17.12	21.60	8.70	8.91
(E)-2-Hexenal	F4	944.76	824.46	392.12	349.05
(E,E)-2,4-hexadienal	F5	94.48	5.57	nd	nd
2-Butyl-2-octenal	F6	164.15	nd	nd	nd
<b>Alcohols</b>					
Cyclopentanol	F7	nd	nd	29.91	20.25
1-Penten-3-ol	F8	31.30	23.68	34.26	25.38
1-Pentanol	F9	19.48	nd	nd	nd
2-Heptanol	F10	180.10	118.08	121.28	96.64
(E)-2-Hexen-1-ol	F11	nd	142.46	296.40	121.48
(Z)-2-Hexen-1-ol	F12	nd	nd	12.51	nd
1-Hexanol	F13	124.00	271.34	494.37	298.30
(Z)-3-Hexen-1-ol	F14	97.43	84.29	84.30	37.52
1-Heptanol	F15	7.08	nd	nd	nd
2-Nonanol	F16	20.08	16.37	20.67	15.66
(1R)-1,7,7-trimethyl-bicyclo[2.2.1]heptan-2-ol	F17	91.52	nd	nd	nd
3,7-Dimethyl-1,6-octadien-3-ol	F18	33.66	5.92	nd	nd
1-Octanol	F19	9.45	nd	nd	nd
4-Methyl-5-hexen-2-ol	F20	nd	12.89	10.33	11.07
1-Nonanol	F21	14.17	nd	nd	nd
(Z)-3-nonen-1-ol	F22	21.85	nd	nd	nd
(R)-3,7-dimethyl-6-octen-1-ol	F23	42.52	nd	17.40	9.18
3,7-Dimethyl-2,6-octadien-1-ol	F24	205.48	36.22	48.95	25.10
Benzyl alcohol	F25	21.85	8.01	11.42	13.23
Phenylethyl alcohol	F26	18.30	nd	nd	4.86
<b>Others</b>					
2-Ethyl-furan	F27	7.68	nd	nd	6.75
2,6-Dimethyl-4-heptanone	F28	nd	nd	nd	5.67
4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3-buten-2-one	F29	10.04	nd	nd	nd
6-Methyl-5-hepten-2-one	F30	nd	79.42	76.14	133.09
Caryophyllene	F31	nd	154.30	343.72	130.39
$\alpha$ -Caryophyllene	F32	nd	20.20	46.23	17.82
$\alpha$ -Farnesene	F33	nd	45.63	306.74	32.12
Caryophyllene oxide	F34	nd	9.06	14.14	5.67
2-Tridecanone	F35	nd	372.35	722.24	277.24
2-Pentadecanone	F36	nd	36.92	204.49	19.44
8-Heptadecene	F37	nd	0.00	83.21	0.00
Cyclopentadecanone	F38	nd	117.03	440.53	54.53
Cyclotridecanone	F39	nd	0.00	26.65	9.72
(1R)-1,7,7-trimethyl-Bicyclo[2.2.1]heptan-2-one	F40	nd	24.03	46.77	23.76

nd, not detected




FIGURE 3

Heatmap of the concentration of volatiles in developing flowers

Note: 0, 5, 50, 90 (percentage of blooming)

Alcohols were the most abundant class of compounds present in the leaves. Volatile profiles in the leaves were clustered into four groups (Fig. 5), and the four groups of volatiles separated leaves into two different development stages. Fig. 5 reveals that leaves at four weeks before flowering had high concentrations of polymers or more substituent aldehydes and alcohols. The concentrations of 1-heptanol, 2-methyl-bicyclo[2.2.2]octan-1-ol, 3,7-dimethyl-1,6-octadien-3-ol, 1-octanol, 3,3-dimethyl-4-heptanol, 1-undecanol, 2-pentanol and 1-nonanol were abundant at this stage. C5 and C6 compounds such as 1-hexanol, 1-pentanol, (E)-2-hexenal, (E)-2-pentenal and (Z)-3-hexen-1-ol became obviously abundant from six to 10 weeks after anthesis, although there also were some polymeric compounds including 2,3-dimethyl-1-butanol and (E)-3,7-dimethyl-2,6-octadienal.

PCA analysis revealed that the most clear separation was noted for samples collected at four and two weeks before anthesis, whereas other sampling points are grouped quite closely together. As above, polymer or more substituent aldehydes and alcohols separated leaves sampled four weeks before anthesis from other samples. Leaves sampled

at two weeks before anthesis contained more unsaturated alcohols and some unsaturated oxygen heterocyclic compounds. (Z)-3-hexen-1-ol, 1-hexanol, (Z)-2-hexen-1-ol, 1-octen-3-ol, 1-pentanol, 2,3-dimethyl-1-butanol, benzyl alcohol, (Z/E)-3/6-nonen-1-ol, (s)-3-ethyl-4-methyl-pentanol, 2-methyl-4-pentenal and (E)-2-hexenal contributed more to the composition of volatiles in leaves sampled at 10 weeks after anthesis (Fig. 6). This result was consistent with the hypothesis that C6 compounds were major contributors to the composition of volatiles in mature leaves (i.e., 10 weeks after anthesis).

#### Activity of ADH and HPL in different tissues

ADH and HPL activity in the leaves, flowers and grape skins is shown in Fig. 7. In the grape skin there was high ADH activity at the early stage of berry development, and it peaked at eight weeks after anthesis, which corroborated several previous studies (Terrier *et al.*, 1996; Tesnière & Verriès, 2000). In contrast, the activity of both 13-HPL and 9-HPL was low, but they demonstrated a similar trend, with a slight increase (peaking 10 and 12 weeks after flowering) followed

TABLE 3  
Volatile compounds in leaves during development in 2012.

Compounds	Symbol	Weeks after anthesis/w								
		-4	-2	2	4	6	8	9	10	
		Concentration (µg/kg)								
Aldehydes	Propanal	L1	nd	nd	3.94	4.35	13.93	15.06	10.03	nd
	Pentanal	L2	nd	16.87	nd	nd	nd	12.51	11.34	nd
	Hexanal	L3	4343.40	2557.09	305.36	97.61	230.98	232.08	229.61	206.75
	(E)-2-Pentenal	L4	nd	5.76	3.03	4.35	16.25	16.51	7.63	18.49
	2-Methyl-4-pentenal	L5	22.20	nd						
	(Z)-3-hexenal	L6	22.20	65.43	38.47	37.24	nd	69.48	56.04	nd
	(Z/E)-2-heptenal	L7	nd	nd	4.24	nd	nd	nd	nd	nd
	(E)-2-hexenal	L8	1335.60	3432.77	1443.92	1920.38	3459.42	3595.26	2318.57	4474.56
	4-Ethyl-2-hexynal	L9	56.40	15.22	nd	nd	nd	nd	nd	nd
	2-Ethyl-2-hexenal	L10	88.80	nd						
	(E,E)-2,4-hexadienal	L11	48.60	95.88	35.44	42.98	104.75	120.52	133.88	27.16
	(E,E)-2,5-hexadienal	L12	42.60	nd						
	(E,E)-2,4-heptadienal	L13	15.60	nd	14.09	nd	18.57	18.95	11.34	4.90
	2-Butyl-2-octenal	L14	950.40	nd	nd	nd	nd	14.88	7.41	nd
	Benzaldehyde	L15	nd	30.45	22.57	30.62	nd	31.37	nd	26.79
	2-Butyl-2-octenal	L16	85.80	nd						
	(E)-3,7-dimethyl-2,6-octadienal	L17	18.60	6.17	nd	nd	12.77	14.51	nd	16.60
Alcohols	Cyclopentanol	L18	nd	nd	25.75	14.96	nd	nd	nd	nd
	1-Penten-3-ol	L19	36.60	61.72	16.81	12.00	19.15	20.51	13.52	27.54
	2-Hexanol	L20	186.00	nd	nd	189.65	nd	153.68	157.87	nd
	1-Pentanol	L21	nd	8.23	5.76	14.09	11.03	8.30	12.43	26.03
	3-Methyl-1-butanol	L22	nd	nd	nd	5.22	nd	5.72	6.54	nd
	2,3-Dimethyl-1-butanol	L23	nd	4.53						
	(Z)-2-Penten-1-ol	L24	nd	158.84	nd	nd	nd	nd	nd	nd
	(Z)-2-Hexen-1-ol	L25	165.60	156.37	nd	nd	nd	nd	nd	136.58
	1-Hexanol	L26	61.80	88.06	48.62	54.29	68.77	61.67	51.02	117.71
	(Z)-3-Hexen-1-ol	L27	107.40	216.04	94.06	86.13	493.87	323.66	241.60	561.77
	1-Octen-3-ol	L28	nd	7.82	6.51	11.48	8.12	9.04	9.38	18.49
	1-Heptanol	L29	35.40	nd	nd	5.92	6.38	6.88	8.72	10.56
	2-Methyl-bicyclo[2.2.2]octan-1-ol	L30	20.40	nd						
	(S)-3-Ethyl-4-methylpentanol	L31	nd	nd	nd	91.17	42.94	135.04	190.80	469.34
	3,7-Dimethyl-1,6-octadien-3-ol	L32	26.40	42.80	19.39	32.36	10.74	8.28	nd	nd
	1-Octanol	L33	24.00	6.17	3.94	5.04	4.64	5.24	5.89	7.17
	2-Pentanol	L34	22.80	nd						
	3,3-Dimethyl-4-heptanol	L35	22.80	nd	nd	nd	3.48	4.08	nd	nd
	1-Undecanol	L36	24.00	nd						
	1-Nonanol	L37	23.40	5.76	3.03	3.13	nd	nd	nd	nd
	(Z/E)-3/6-nonen-1-ol	L38	nd	14.82	15.45	9.74	4.64	6.07	7.63	8.68
	3,7-Dimethyl-6-octen-1-ol	L39	14.40	nd	6.06	4.00	nd	nd	nd	nd
	3,7-dimethyl-2,6-Octadien-1-ol	L40	561.00	248.14	39.38	41.58	nd	16.28	13.74	15.09
	Benzyl alcohol	L41	58.20	15.22	10.00	35.67	11.32	14.88	27.04	81.87
	Phenylethyl alcohol	L42	15.00	nd	10.60	17.75	5.80	11.08	15.48	29.05
Others	2-Methyl-3-hexanone	L43	nd	nd	2.42	nd	nd	nd	nd	nd
	3,4-Dihydro-2H-pyran	L44	nd	6.58	nd	nd	nd	nd	nd	nd
	2,3-Dimethyl-1-pentene	L45	nd	nd	37.87	33.76	84.44	98.92	56.70	80.36
	2,3-Dimethyl-1-pentene	L46	21.60	nd						
	2,6-Dimethyl-4-heptanone	L47	14.40	nd	nd	nd	nd	5.44	4.80	nd
	2-Ethyl-furan	L48	0.00	116.04	10.15	12.88	23.50	24.04	nd	24.90
	Cyclopentanecarboxaldehyde	L49	57.60	27.57	nd	nd	nd	nd	nd	nd
	2-methyl-3-methylene-									
	3,3,5-Trimethyl-cyclohexene	L50	201.00	nd						
	2,5-Dimethyl-3-ethyl-1,3-hexadiene	L51	16.80	nd						
	5-Ethyl-2(5H)-furanone	L52	nd	nd	nd	2.44	5.51	5.64	4.36	nd
Eugenol	L53	nd	nd	3.03	3.65	nd	nd	nd	nd	

nd, not detected

by a decline to the initial level during the maturation of the berry (Fig. 7a).

The activity of these enzymes in flowers displayed a different trend. The activity of ADH and 9-HPL was dynamic

until the full-bloom stage, with a continuous increase (peaking at 90% blooming), whereas 13-HPL activity maintained a low but stable level during the development of the flowers (Fig. 7b). In the leaves, the activity of three




FIGURE 4

Principle component analysis of volatile compounds of grape flowers.

A. PCA score plot. B. Loading plot of components 1, 2 and 3. Note: “F” refers to volatiles in flowers. For details, refer to Table 2.




FIGURE 5

Heatmap of volatiles in developing leaves. Note: W (weeks after anthesis)

enzymes followed an interesting trend. ADH and 9-HPL were more active than 13-HPL, but they all decreased with the development of the leaves (Fig. 7c).

The highest activities of 13-HPL, 9-HPL and ADH occurred in the grape skin, flowers and grape skin respectively. The activity of 13-HPL did not change much in the developing flowers and leaves. The activity of 9-HPL was much higher in the flowers and leaves than in the skins. The activity of ADH was obviously high in these three tissues compared to that of the HPLs. In this paper, we report that the activity of ADH and 9-HPL decreased in the leaves during anthesis (-2 to 2 weeks after anthesis), while they increased in the flowers during the entire process of anthesis. The activity of ADH increased until eight weeks after anthesis in the grape skins. ADH catalysed the conversion of aldehydes to alcohols (Chen & Chase Jr, 1993; Speirs *et al.*, 1998), and exhibited high activity during berry development (Tesnière *et al.*, 1993; 2006; Köllner *et al.*, 2008). The results on ADH activity were in accordance with the results above. Furthermore, Chen *et al.* (2015) reported that 9-HPL activity was approximately twice as high as that of 13-HPL in cucumber fruit and that 13-HPL activity decreased with the development of fruit, although 9-HPL activity increased. In this paper, the same trend of 9-HPL and 13-HPL activity as in Chen *et al.*'s report was tracked in the grape flowers, but was not in the grape skins. We supposed that 13-HPL and 9-HPL activity first increased and then decreased during blossoming and fruit setting.

#### Correlation analysis of enzyme activity and volatiles

Fluctuations in the concentrations of C5 and C6 are shown in Fig. 8. Generally speaking, the content of C5 maintained a low level in all three tissues, which may be a result of the weak activity of 13-HPL. C5 volatiles were synthesised by the homolytic mechanism of 13-HPL (Grechkin, 1998; Salch *et al.*, 1995.), while C5 was more abundant in the leaves than in the grape skins and flowers. The concentration of C5 decreased rapidly at the first three sampling dates, and increased slightly later on in the leaves, whereas it kept increasing in the flowers and even in grape skins until nine weeks after anthesis. The concentration of C5 stayed at a stable level in the grape skins from nine to 12 weeks after

anthesis, while the concentration of C6 started increasing sharply from nine weeks.

There was quite a significant increase in C6 compounds in the grape skin from nine to 12 weeks, whereas the concentrations decreased later. The concentration of C6 decreased obviously in the flowers and leaves at the first three sampling dates, and they all increased slightly later (Fig. 8). According to previous studies, the C6 content has a direct relationship with the activity of 13-HPL in other species. C6 aldehydes and C12 oxyacids are formed through the heterolytic mechanism of 13-HPL (Blee, 1998; Grechkin, 1998; Mita *et al.*, 2005). Our study showed that the concentration of C6 increased (peaking at 12 weeks) with increasing 13-HPL activity (peaking at 10 weeks) in the grape skins. It seems that both C6 and C5 were formed by the catalysis of 13-HPL, while they were not formed at the same time in the grape skins.

The principal alcohols gradually increased in the grape skin with grape development (peaking at 16 weeks) (Fig. 1), which is consistent with high ADH activity, but they may also be synthesised by other metabolic pathways, such as those involved in amino acid metabolism (Sanz *et al.*, 1996). The trend for ADH is consistent with other reports (Van Der Straeten *et al.*, 1991; Echeverría *et al.*, 2004).

A high terpene content was detected in the flowers, which is consistent with the synthesis of terpenes in *Arabidopsis* (Echeverría *et al.* 2004) and *Antirrhinum majus* (Dudareva *et al.*, 2003); therefore, the formation of terpenes may exhibit tissue specificity. The correlation of volatiles in the leaves, flowers and berries and the conjecture of molecule transfer among tissues necessitate further study on the transfer mechanism.

Pearson's correlation analysis was performed on the concentration of volatile components and the activity of 13-HPL, 9-HPL and ADHs. The results are shown in Table 4. More kinds of volatile compounds in the leaves correlated with the activity of the three enzymes than in the flowers and grape skins. Furthermore, the concentration of 3,7-dimethyl-1,6-octadien-3-ol was positively correlated with the activity of 13-HPL ( $p < 0.01$ ) and 9-HPL ( $p < 0.05$ ), and negatively correlated with activity of ADHs ( $P < 0.05$ ) in the leaves, whereas the concentration was negatively correlated with




FIGURE 6

Principle component analysis of volatile compounds of leaves. A. PCA score plot. B. Loading plot of components 1, 2 and 3.

Note: "L" refers to volatiles in leaves. For detail, refer to Table 3.




FIGURE 7

Changes in HPLs, ADHs activity during tissues development. A: Changes of activity in grape skin; B: Changes of enzyme activity in developing flowers; C: Changes of enzyme activity in developing leaves.

TABLE 4

Results of Pearson's  $r$  correlation analysis between volatile concentration and activity of 13-HPL, 9-HPL and ADHs enzyme.

Tissue	Compounds	13-HPL	9-HPL	ADHs
Leaf	Propanal	-0.449	-0.168	<b>0.747*</b>
	Hexanal	0.493	<b>0.707*</b>	-0.631
	(E)-2-Pentenal	-0.328	-0.42	<b>0.899**</b>
	(Z)-2-Penten-1-ol	<b>0.820*</b>	0.49	-0.409
	(Z)-3-Hexen-1-ol	-0.217	-0.407	<b>0.755*</b>
	(S)-3-Ethyl-4-methylpentanol	-0.529	<b>-0.744*</b>	0.573
	3,7-Dimethyl-1,6-octadien-3-ol	<b>0.877**</b>	<b>0.810*</b>	<b>-0.789*</b>
	3,4-Dihydro-2H-pyran	<b>0.820*</b>	0.49	-0.409
	2,3-Dimethyl-1-pentene	-0.638	-0.569	<b>0.950**</b>
	2-Ethyl-furan	<b>0.803*</b>	0.446	-0.23
	5-Ethyl-2(5H)-furanone	-0.418	-0.147	<b>0.767*</b>
Flower	Cyclopentanol	0.57	0.6	<b>0.96*</b>
	2-Heptanol	<b>-0.983*</b>	-0.8	-0.686
	(Z)-3-Hexen-1-ol	-0.876	<b>-0.983*</b>	-0.413
	Caryophyllene	<b>1**</b>	0.894	0.641
Berry skin	1-Penten-3-ol	0.654	<b>0.704**</b>	-0.18
	2-Hexanol	-0.651	-0.527	<b>-0.762**</b>
	(Z)-2-Penten-1-ol	0.559	<b>0.702*</b>	-0.206
	3,7-Dimethyl-1,6-octadien-3-ol	<b>-0.746*</b>	<b>-0.689*</b>	-0.632
	D-limonene	<b>0.706*</b>	0.244	0.013

\*\* Values in bold are significantly correlated at  $p < 0.01$ . \* Values in bold are significantly correlated at  $p < 0.05$ . The absence of '\*' indicates  $P > 0.05$ .




FIGURE 8

Concentration of six-carbon compounds and five-carbon compounds in (a) grape skin; (b) flower; (c) leaves




FIGURE 1S  
Schematic drawing of the GLV pathway in plants

TABLE S1  
Standard curves of characteristic volatiles.

No.	Retention time/min	English name	Standard curve equation	R <sup>2</sup>
1	8.94	Pentanal	$y = 40.89x - 12.92$	0.994
2	10.49	Butanoic acid, ethyl ester	$y = 2.443x + 13.16$	0.998
3	11.71	Hexanal	$y = 46.15x - 603.0$	0.995
4	13.09	(E)-2-Pentenal	$y = 20.85x + 24.13$	0.996
5	14.72	Heptanal	$y = 4.673x + 37.45$	0.998
6	15.79	(E)-2-Hexenal	$y = 12.70x + 161.7$	0.997
7	16.58	1-Pentanol	$y = 71.10x - 8.869$	0.998
8	18.56	(Z)-2-Penten-1-ol	$y = 79.66x + 22.38$	0.997
9	19.62	1-Hexanol	$y = 13.95x + 82.38$	0.997
10	21.1	(E)-2-Hexen-1-ol	$y = 14.16x + 82.27$	0.998
11	28.47	1-Nonanol	$y = 0.916x + 8.604$	0.997
12	30.02	(E/Z)-6-Nonen-1-ol	$y = 3.436x + 19.65$	0.997

the activity of both 13-HPL and 9-HPL ( $P < 0.05$ ) in the grape skins. It appears that the correlation of the activity of the enzyme and the concentration of volatile compounds exhibited tissue specificity. The concentrations of 1-penten-3-ol and (Z)-2-penten-1-ol were positively correlated with the activity of 9-HPL in the berry skin. In addition, the concentration of 2-hexanol was negatively correlated with the activity of ADHs. The concentration of (Z)-3-hexen-1-ol in the flowers was negatively correlated with the activity of 9-HPL, but it was positively correlated with the activity of ADHs in the leaves.

#### CONCLUSIONS

The multivariate analysis of the concentrations of volatiles in Cabernet Sauvignon grapes revealed that the volatile

profiles were dramatically different among the grapes, leaves and flowers. The volatile profiles of the grape skins were found to be more complex, with not only more types of volatile compounds detected, but also higher concentrations, especially of C6 compounds, resulting in a higher total overall at 16 weeks after anthesis. C6 compounds in grapes are also known to be direct precursors of the formation of some pleasant esters in wines. In the leaves, polymer alcohols and ketones were gradually broken down, whereas some C6 and C5 alcohols accumulated in old leaves. In the flowers, aldehydes and alcohols were major contributors at the bud stage, but most of these compounds were not retained in the full blooming flowers. The accumulation of alcohols in the grape skin was apparent, which correlated with the high activity of ADHs. This is consistent with the documented

pathway of fatty acid oxidation, LOX-HPL. In addition, the decreasing concentration of volatiles in the leaves might be caused by transportation from the leaves to the grapes, which is in need of further study.

The activity of 13-HPL was low during the growing season, whereas the content of C6 was noticeably high. As we hypothesised, the activity of 13-HPL occurred at a stage we did not examine. Generally, the activity of 9-HPL in the leaves and flowers was high, but few C9 compounds were detected in these two kinds of tissue. This might be because the C9 compounds were unstable and were degraded quickly.

## LITERATURE CITED

- Anthon, G.E. & Barrett, D.M., 2003. Thermal inactivation of lipoxygenase and hydroperoxytrienoic acid lyase in tomatoes. *Food Chem.* 81, 275-279.
- Baumes, R., Bayonove, C., Barillère, J. M., Escudie, R. & Cordonnier, R., 1988. La macération pelliculaire dans la vinification en blanc. Incidence sur la composante volatile des moûts. *Connaissance de la Vigne et du Vin. Journal International des sciences de la vigne*, 32, 209-223.
- Bestmann, H.J., Winkler, L. & Von, H.O., 1997. Headspace analysis of volatile flower scent constituents of bat-pollinated plants. *Phytochem.* 46, 1169-1172.
- Blee, E., 1998. Phytooxylipins and plant defense reactions. *Prog. Lipid Res.* 37, 33-72.
- Cynthia, M.M. & Kenji M., 2014. The importance of lipoxygenase control in the production of green leaf volatiles by lipase-dependent and independent pathways. *Plant Biotechnol.* 31, 445-452.
- Chen, A.R. & Chase Jr., T., 1993. Alcohol dehydrogenase 2 and pyruvate decarboxylase induction in ripening and hypoxic tomato fruit. *Plant Physiol. Bioch.* 31, 875-885.
- Chen, S.X., Zhang, R.R., Hao, L.N., Chen, W.F. & Cheng, S.Q., 2015. Profiling of volatile compounds and associated gene expression and enzyme activity during fruit development in two cucumber cultivars. *Plos One* 3, 1-22.
- Coelho, E., Rocha, S.M., Barros, A.S., Delgadillo, I. & Coimbra, M.A., 2007. Screening of variety and pre-fermentation-related volatile compounds during ripening of white grapes to define their evolution profile. *Anal. Chim. Acta* 597, 257-264.
- Defilippi, B.G., Kader, A.A. & Dandekar, A.M., 2005. Apple aroma: Alcohol acyltransferase, a rate limiting step for ester biosynthesis, is regulated by ethylene. *Plant Sci.* 168, 1199-1210.
- Delphia, C.M., Mescher, M.C. & De Moraes, C.M., 2007. Induction of plant volatiles by herbivores with different feeding habits and the effects of induced defenses on host-plant selection by thrips. *J. Chem. Ecol.* 33, 997-1012.
- Dudareva, N., Martin, D., Kish, C.M., Kolosova, N., Gorenstein, N., Faldt, J., Miller, B. & Bohlmann, J., 2003. (E)-beta-ocimene and myrcene synthase genes of floral scent biosynthesis in snapdragon: Function and expression of three terpene synthase genes of a new terpene synthase subfamily. *Plant Cell* 15, 1227-1241.
- Echeverria, G., Graell, J., López, M. & Lara, I., 2004. Volatile production, quality and aroma-related enzyme activities during maturation of 'Fuji' apples. *Postharvest Biol. Tec.* 31, 217-227.
- Gomez, E., Martinez, A. & Laencina, J., 1995. Changes in volatile compounds during maturation of some grape varieties. *J. Sci. Food Agr.* 67, 229-233.
- Grechkin, A., 1998. Recent developments in biochemistry of the plant lipoxygenase pathway. *Prog. Lipid Res.* 37, 317-352.
- Hashizume, K. & Samuta, T., 1997. Green odorants of grape cluster stem and their ability to cause a wine stemmy flavor. *J. Agr. Food Chem.* 45, 1333-1337.
- Hatanaka, A., 1993. The biogenesis of green odour by green leaves. *Phytochem.* 34, 1201-1218.
- Jiang, W.G., Li, J.M., Xu, Y., Fan, W. & Yu, Y., 2011. Analysis of aroma components in four red grape varieties. *Food Sci.* 32(6), 225-229.
- Köllner, T.G., Held, M., Lenk, C., Hiltbold, I., Turlings, T.C., Gershenzon, J. & Degenhardt, J., 2008. A maize (E)-beta-caryophyllene synthase implicated in indirect defense responses against herbivores is not expressed in most American maize varieties. *Plant Cell* 20, 482-494.
- Kalua, C.M. & Boss, P.K., 2009. Evolution of volatile compounds during the development of Cabernet Sauvignon grapes (*Vitis vinifera* L.). *J. Agr. Food Chem.* 57, 3818-3830.
- Matsui, K., Minami, A., Hornung, E., Shibata, H., Kishimoto, K., Ahnert, V., Kindl, H., Kajiwara, T. & Feussner, I., 2006. Biosynthesis of fatty acid derived aldehydes is induced upon mechanical wounding and its products show fungicidal activities in cucumber. *Phytochem.* 67, 649-657.
- Matsui, K., Ujita, C., Fujimoto, S.H., Wilkinson, J., Hiatt, B., Knauf, V., Kajiwara, T. & Feussner, I., 2000. Fatty acid 9- and 13-hydroperoxide lyases from cucumber. *FEBS Lett.* 481, 183-188.
- Mita, G., Quarta, A., Fasano, P., De Paolis, A., Di Sansebastiano, G.P., Perrotta, C., Iannacone, R., Belfield, E., Hughes, R. & Tsesmetzis, N., 2005. Molecular cloning and characterization of an almond 9-hydroperoxide lyase, a new CYP74 targeted to lipid bodies. *J. Exp. Bot.* 56, 2321-2333.
- Nakamura, S. & Hatanaka, A., 2002. Green-leaf-derived C6-aroma compounds with potent antibacterial action that act on both gram-negative and gram-positive bacteria. *J. Agr. Food Chem.* 50, 7639-7644.
- Park, S.K., Morrison, J.C., Adams, D.O. & Noble, A.C., 1991. Distribution of free and glycosidically bound monoterpenes in the skin and mesocarp of Muscat of Alexandria grapes during development. *J. Agr. Food Chem.* 39, 514-518.
- Perestrelo, R., Barros, A.S., Rocha, S.M. & Câmara, J.S., 2011. Optimisation of solid-phase microextraction combined with gas chromatography-mass spectrometry based methodology to establish the global volatile signature in pulp and skin of *Vitis vinifera* L. grape varieties. *Talanta* 85, 1483-1493.
- Salch, Y. P., Grove, M. J., Takamura, H. & Gardner, H. W., 1995. Characterization of a C-5,13-cleaving enzyme of 13(S)-Hydroperoxide of linolenic acid by soybean seed. *Plant Physiol.* 108, 1211-1218.
- Sanz, C., Olias, J.M. & Perez, A., 1996. Aroma biochemistry of fruits and vegetables. *Pr. Phyt. Soc.* 41, 125-156.
- Smith, P., Krohn, R.I., Hermanson, G., Mallia, A., Gartner, F., Provenzano, M., Fujimoto, E., Goeke, N., Olson, B. & Klenk, D., 1985. Measurement of protein using bicinchoninic acid. *Anal. Biochem.* 150, 76-85.
- Song, J., Shellie, K.C., Wang, H. & Qian, M.C., 2012. Influence of deficit irrigation and kaolin particle film on grape composition and volatile compounds in Merlot grape (*Vitis vinifera* L.). *Food Chem.* 134, 841-850.
- Speirs, J., Lee, E., Holt, K., Yong-Duk, K., Scott, N.S., Loveys, B. & Schuch, W., 1998. Genetic manipulation of alcohol dehydrogenase levels in ripening tomato fruit affects the balance of some flavor aldehydes and alcohols. *Plant Physiol.* 117, 1047-1058.
- Suurmeijer, C.N., Pérez-Gilbert, M., Van der Hijden, H.T., Veldink, G.A. & Vliegthart, J.F., 1998. Purification, product characterization and kinetic properties of soluble tomato lipoxygenase. *Plant Physiol. Bioch.* 36, 657-663.

- Terrier, N., Sauvage, F. & Romieu, C., 1996. Absence de crise respiratoire, induction de l'activité alcool deshydrogénase et diminution de l'acidité vacuolaire lors de la maturation du raisin In Œnologie 95, Compte Rendu 5eme Symposium International d'Œnologie. Tec & Doc Lavoisier, Paris. pp. 24 – 28.
- Tesnière, C. & Verriès, C. 2000. Molecular cloning and expression of cDNAs encoding alcohol dehydrogenases from *Vitis vinifera* L. during berry development. *Plant Sci.* 157, 77-88.
- Tesnière, C.M., Romieu, C. & Vayda, M.E., 1993. Changes in the gene expression of grapes in response to hypoxia. *Am. J. Enol. Vitic.* 44, 445-451.
- Tesnière, C., Davies, C., Sreekantan, L., Bogs, J., Thomas, M. & Torregrosa, L., 2006. Analysis of the transcript levels of VvAdh1, VvAdh2 and VvGrip4, three genes highly expressed during *Vitis vinifera* L. berry development. *Vitis* 45, 75-79.
- Tufariello, M., Capone, S. & Siciliano, P., 2012. Volatile components of Negroamaro red wines produced in Apulian Salento area. *Food Chem.* 132, 2155-2164.
- Van Der Straeten, D., Pousada, R.A.R., Gielen, J. & Van Montagu, M., 1991. Tomato alcohol dehydrogenase: Expression during fruit ripening and under hypoxic conditions. *FEBS Lett.* 295, 39-42.
- Xiao qing, X., Guo, C., Ling ling, D., Jiang, R., Qiu hong, P., Chang qing, D., & Jun, W., 2015. Effect of training systems on fatty acids and their derived volatiles in Cabernet Sauvignon grapes and wines of the north foot of Mt. Tianshan. *Food Chem.* 181, 198-206.