

Review Article

Depression: Abnormality in Neural Circuits

Chanyi Lu^{1,2,5#}, Yao-Yao Li^{1,2#}, Hua-Zhen Lin^{3,4#}, and Yun-Feng Zhang^{1,2,6*}

¹School of Ophthalmology & Optometry, The Eye Hospital, Wenzhou Medical University, China

²State Key Laboratory Cultivation Base and Key Laboratory of Vision Science, Ministry of Health P. R. China and Zhejiang Provincial Key Laboratory of Ophthalmology and Optometry, China

³The 2nd Affiliated Hospital and Yuying Children's Hospital, Wenzhou Medical University, China

⁴The 2nd School of Medicine, Wenzhou Medical University, China

⁵University of Pennsylvania School of Dental Medicine, USA

⁶Department of Neuroscience, University of Pennsylvania Perelman School of Medicine, USA

Chanyi Lu, Yao-Yao Li and Hua-Zhen Lin contributed equally to this work

***Corresponding author**

Yun-Feng Zhang, Department of Neuroscience, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA; Tel: (215) 746-2791; Fax: 215-573-9050; Email: zhangyunfeng1983@163.com

Submitted: 08 August 2017

Accepted: 18 September 2017

Published: 19 September 2017

Copyright © 2017 Zhang et al.

ISSN: 2374-0124

OPEN ACCESS

Keywords

- Depression
- Mental illness
- Pathophysiology

Abstract

Depression as a kind of mental illness affects huge number of population worldwide and is ranked as the fourth leading cause of disability worldwide, and is predicted to increase to the second place by 2020. Depression leads to great social burden because of its substantial impairment and disability in everyday activities. Though great efforts in previous decades have been made to understand this mental disease, to date, much still remains to be known about its pathophysiology, especially for the underlying neural circuit mechanisms. To better improve the diagnosis, treatment and prevention of this mental disorder, it is imperative and also very important to make more clarifications on the underlying neural circuitry mechanisms responsible for the occurrence of depression. Accordingly, here we briefly summarized some key studies pertaining to the neural circuits responsible for depression, and hopefully to shed some lights on the future studies concerning this annoying mental disorder.

INTRODUCTION

Depression as a kind of mental illness affects around 20% of the world's population [1]. It is ranked the fourth leading cause of disability worldwide, and is predicted to increase to the second place by 2020 [2]. Depression causes great social burden because of its substantial impairment and disability in everyday activities, which makes imperative investigations on the mechanisms underlying its pathophysiology so as to improve the diagnosis, treatment and prevention of this mental disorder. A growing body of studies emerges aiming to make a better understanding of depression [3-11]. Our recent work demonstrates that alterations of motor cortical neural microcircuits contribute greatly to the depressive-like behaviors in a light-deprivation induced mouse model of depression [8], which emphasizes the potentially robust changes in neural circuits during the occurrence of depression. Generally, neural circuits between distinct brain areas are responsible for the depressive-like phenotypes [9, 12-14]. The occurrence of depression is potentially associated with abnormalities of these specific neural circuits in/among distinct brain regions [15]. Here, we concisely summarized some key literatures pertaining to the neural circuits responsible for depression, hopefully to provide some insights into the future studies concerning this annoying mental disorder.

Multiple lines of evidence implicated the important role of the ventral pallidum (VP) in depression because of its special anatomical location at the interface of the motivational and reward circuitry [16-18]. A recent study indicated that neural circuits between the ventral pallidum (VP) and lateral habenula (LHb) as well as ventral tegmental area (VTA) mediated separate core symptoms of depression [9]. The VP parvalbumin-positive (PV) neurons projecting to LHb subserve the behavioral despair and those projecting to VTA contribute the social withdrawal of depressive behaviors. This work by Knowland et al. established a systematic approach that could be taken to explore how distinct neural circuits may subserve the heterogeneous symptoms seen in depression. It also implies that VP neural circuits endowed with distinct components could contribute to related, yet different depressive-like phenotypes in mice, which provides some insights into symptom-specific treatments of depression.

The amygdala is another key brain structure closely associated with depression [19-22]. A new clinical study documented that maternal depressive symptoms during pregnancy were closely related to the amygdala hyper responsivity in their children [23]. Moreover, a historic study has reported a critical role of amygdala-cingulate feedback circuit in depression [21]. Using the functional connectivity density (FCD) mapping method, it was found a decreased FCD in the mid-cingulate cortex (MCC)

while an increased FCD in the occipital cortex (OCC) in patients with depression [13]. In addition, evidence from functional magnetic resonance imaging (fMRI) showed that the amygdala-left rostral PFC (rPFC) functional connectivity was decreased in response to negative emotional stimuli in medication-naive subjects with major depressive disorder (MDD) [22], suggesting that abnormalities in neural circuitry between the amygdala and left rPFC potentially play a key role in the pathophysiology of MDD. A recently work by Young et al. found that the increased amygdala hemodynamic response to positive memories dependent on the real-time functional MRI neurofeedback (rtfMRI-nf) training could dramatically decreased depressive symptoms of patients. In addition, using the resting-state fMRI, another recent study reported that the resting-state amygdala-cortical functional connectivity with the prefrontal-cingulate-temporal circuit is abnormal chronic tinnitus patients with depressive mood [24]. Collectively, it suggests that abnormalities of neural circuits between amygdala and other brain areas are potentially responsible for the core symptoms of depression, and the amygdala as a potential target for the treatment depression should be paid more attention.

In a chronic mild stress (CMS) rat model of depression, it was found that the dopamine DA neuron population activity was remarkably decreased, which could be restored by attenuating the activity of either the VP or the basolateral nucleus of the amygdala (BLA). It highlighted that the CMS rat depression model was closely related to the BLA-VP-VTA inhibition of dopamine neuron activity, and shed some lights on the important role of neural circuits of the BLA-VP-VTA in depression [25]. In fact, VTA is also a very important brain area closely associated with depression [5, 26], and the inhibition of VTA dopaminergic neurons could lead to several depression-like phenotypes in mice [27]. The optogenetic stimulation of VTA projecting dopamine neurons in the nucleus accumbens (NAc) could greatly change the neural encoding of depression-related behaviors of freely moving rodents [27], which underscored the crucial role of the VTA-NAc neural circuitry in depression. The importance of the VTA-NAc neural circuitry in depression has been strengthened by a recent study depending on the chronic bilateral high frequency deep brain stimulation (DBS) of the medial forebrain bundle (MFB) [28]. It was documented that the lesion of VTA dopamine neurons could lead to the dopamine depletion in the NAc and, in the end, produced depressive-like behaviors in the rats, which was, to some extent, reversed by MFB-DBS. This study also highlights the DBS as a potentially efficient way in relieving depressive symptoms especially resulting from abnormalities of VTA-NAc neural circuitry.

In summary, future studies should make much more room for the important roles of neural circuits in depression [29], which will definitely supply some informative evidence for the clinical treatment of depression. Though the crucial roles of neural circuitry between the VP and LHb as well as VTA, the amygdala and PFC, the BLA-VP-VTA, and also the VTA-NAc, we should keep in mind that some other local neural microcircuits mentioned elsewhere also should not be neglected [29]. From this point, much more efforts in future studies should be made for the search of new potential brain regions and neural circuits related to depression. The rapid development of optogenetics makes

it more convenient for the analysis of the role of specific neural circuits in specific behaviors in living animals [30,31], including depression-like behaviors, which subserves the investigations of new potential brain regions and neural circuits responsible for depression.

ACKNOWLEDGEMENTS

This study was supported by the grants from the Wenzhou Science and Technology Plans (No.Y20140139), the Foundation of School of Ophthalmology & Optometry, The Eye Hospital, Wenzhou Medical University (YNCX201311), the Natural Science Foundation of Zhejiang Province of China (No. LQ14H120003) and the National Natural Science Foundation for Young Scientists of China (No. 81301117).

REFERENCES

- Stepanichev M, Dygalo NN, Grigoryan G, Shishkina GT, Gulyaeva N. Rodent Models of Depression: Neurotrophic and Neuroinflammatory Biomarkers. *Biomed Res Int.* 2014; 2014: 932757.
- Bromet E, Andrade LH, Hwang I, Sampson NA, Alonso J, de Girolamo G, et al. Cross-national epidemiology of DSM-IV major depressive episode. *BMC Med.* 2011; 9: 90.
- Seo JS, Zhong P, Liu A, Yan Z, Greengard P. Elevation of p11 in lateral habenula mediates depression-like behavior. *Mol Psychiatry.* 2017; 16.
- Lu C, Wang Y, Zhang YF. Light deprivation produces a sexual dimorphic effect on neural excitability and depression-like behavior in mice. *Neurosci Lett.* 2016; 633: 69-76.
- Pena CJ, Kronman HG, Walker DM, Cates HM, Bagot RC, Purushothaman I, et al. Early life stress confers lifelong stress susceptibility in mice via ventral tegmental area OTX2. *Science.* 2017; 356: 1185-1188.
- Svenningsson P, Kim Y, Warner-Schmidt J, Oh YS, Greengard P. p11 and its role in depression and therapeutic responses to antidepressants. *Nat Rev Neurosci.* 2013; 14: 673-680.
- Li K, Zhou T, Liao L, Yang Z, Wong C, Henn F, et al. β CaMKII in lateral habenula mediates core symptoms of depression. *Science.* 2013; 341: 1016-1020.
- Zhang Y-F, Li Q-Q, Qu J, Sun C-M, Wang Y. Alterations of motor cortical microcircuit in a depressive-like mouse model produced by light deprivation. *Neuroscience.* 2016; 341: 79-94.
- Knowland D, Lilascharoen V, Pacia CP, Shin S, Wang EH, Lim BK. Distinct Ventral Pallidal Neural Populations Mediate Separate Symptoms of Depression. *Cell.* 2017; 170: 284-297.
- Gonzalez MM, Aston-Jones G. Light deprivation damages monoamine neurons and produces a depressive behavioral phenotype in rats. *Proc Natl Acad Sci USA.* 2008; 105: 4898-4903.
- Ramirez S, Liu X, MacDonald CJ, Moffa A, Zhou J, Redondo RL, et al. Activating positive memory engrams suppresses depression-like behaviour. *Nature.* 2015; 522: 335-339.
- Pezawas L, Meyer-Lindenberg A, Drabant EM, Verchinski BA, Munoz KE, Kolachana BS, et al. 5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: a genetic susceptibility mechanism for depression. *Nat Neurosci.* 2005; 8: 828-834.
- Zhang B, Li M, Qin W, Demenescu LR, Metzger CD, Bogerts B, et al. Altered functional connectivity density in major depressive disorder at rest. *Eur Arch Psychiatry Clin Neurosci.* 2016; 266: 239-248.
- Warden MR, Selimbeyoglu A, Mirzabekov JJ, Lo M, Thompson KR,

- Kim S-Y, et al. A prefrontal cortex-brainstem neuronal projection that controls response to behavioural challenge. *Nature*. 2012; 492: 428-432.
15. Price JL, Drevets WC. Neural circuits underlying the pathophysiology of mood disorders. *Trends Cogn Sci*. 2012; 16: 61-71.
 16. Smith KS, Tindell AJ, Aldridge JW, Berridge KC. Ventral pallidum roles in reward and motivation. *Behav Brain Res*. 2009; 196: 155-167.
 17. Heimer L, Switzer RD, Vanhoesen GW. Ventral Striatum and Ventral Pallidum - Components of the Motor System. *Trends in Neurosciences*. 1982; 5: 83-87.
 18. Haber SN, Groenewegen HJ, Grove EA, Nauta WJ. Efferent connections of the ventral pallidum: evidence of a dual striato pallidofugal pathway. *J Comp Neurol*. 1985; 235: 322-335.
 19. Young KD, Siegle GJ, Zotev V, Phillips R, Misaki M, Yuan H, et al. Randomized Clinical Trial of Real-Time fMRI Amygdala Neurofeedback for Major Depressive Disorder: Effects on Symptoms and Autobiographical Memory Recall. *Am J Psychiatry*. 2017; 174: 748-755.
 20. Klimek V, Schenck JE, Han H, Stockmeier CA, Ordway GA. Dopaminergic abnormalities in amygdaloid nuclei in major depression: a postmortem study. *Biol Psychiatry*. 2002; 52: 740-748.
 21. Pezawas L, Meyer-Lindenberg A, Drabant EM, Verchinski BA, Munoz KE, Kolachana BS, et al. 5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: a genetic susceptibility mechanism for depression. *Nat Neurosci*. 2005; 8: 828-834.
 22. Kong LT, Chen KY, Tang YQ, Wu F, Driesen N, Womer F, et al. Functional connectivity between the amygdala and prefrontal cortex in medication-naive individuals with major depressive disorder. *J Psychiatry Neurosci*. 2013; 38: 417-422.
 23. Van der Knaap NJF, Klumpers F, El Marroun H, Mous S, Schubert D, Jaddoe V, et al. Maternal depressive symptoms during pregnancy are associated with amygdala hyperresponsivity in children. *Eur Child Adolesc Psychiatry*. 2017; 30.
 24. Chen YC, Bo F, Xia W, Liu S, Wang P, Su W, et al. Amygdala functional disconnection with the prefrontal-cingulate-temporal circuit in chronic tinnitus patients with depressive mood. *Prog Neuropsychopharmacol Biol Psychiatry*. 2017; 79: 249-257.
 25. Chang CH, Grace AA. Amygdala-ventral pallidum pathway decreases dopamine activity after chronic mild stress in rats. *Biol Psychiatry*. 2014; 76: 223-320.
 26. Small KM, Nunes E, Hughley S, Addy NA. Ventral tegmental area muscarinic receptors modulate depression and anxiety-related behaviors in rats. *Neurosci Lett*. 2016; 616: 80-85.
 27. Tye KM, Mirzabekov JJ, Warden MR, Ferenczi EA, Tsai HC, Finkelstein J, et al. Dopamine neurons modulate neural encoding and expression of depression-related behaviour. *Nature*. 2013; 493: 537-541.
 28. Furlanetti LL, Coenen VA, Dobrossy MD. Ventral tegmental area dopaminergic lesion-induced depressive phenotype in the rat is reversed by deep brain stimulation of the medial forebrain bundle. *Behav Brain Res*. 2016; 299: 132-140.
 29. Lu CY, Li QQ, Li YY, Wang Y, Zhang YF. A Short Glance at the Neural Circuitry Mechanism Underlying Depression. *World Journal of Neuroscience*. 2016; 6: 184-192.
 30. Boyden ES, Zhang F, Bamberg E, Nagel G, Deisseroth K. Millisecond-timescale, genetically targeted optical control of neural activity. *Nat Neurosci*. 2005; 8: 1263-1268.
 31. Zhang YF. Times for the Neuroscience: Patch-Clamp Recordings and Optogenetics in Decoding Neural Circuits. *EC Ophthalmology*. 2016; 4: 589-591.

Cite this article

Lu C, Li YY, Lin HZ, Zhang YF (2017) Depression: Abnormality in Neural Circuits. *Ann Psychiatry Ment Health* 5(5): 1115.