

Phenotypic and Whole Cell Protein Analysis by SDS-PAGE for Identification of Dominant Lactic Acid Bacteria Isolated from Algerian Raw Milk

Fatima Ghazi, Djamel Eddine Henni, Zineb Benmechernene and Mebrouk Kihal

Laboratory of Applied Microbiology, Department of Biology,
Faculty of Sciences, Oran University, B.P. 16, Es-Senia, Oran 31100, Algeria

Abstract: Phenotypic identification of 21 representative strains revealed the presence of the following species: *Leuconostoc mesenteroides* subsp. *dextranicum*, *Weissella paramesenteroides*, *Lactococcus lactis* biovar. *diacetylactis*, *Pediococcus acidilactici*, *Enterococcus faecium*, *Lactobacillus rhamnosus*, *Lactobacillus paracasei* subsp. *paracasei* and *Lactobacillus plantarum*. In order to validate the previous results, whole cell protein patterns were obtained using Sodium Dodecyl Sulphate Gel Electrophoresis (SDS-PAGE) of these lactic acid bacteria strains, were analysed by calculating the coefficients of similarity (>100) for each two strains which were 80.7% and 78% for next couple of strains (6, 13) and (43, L₄) that were identified phenotypically as *Weissella paramesenteroides* and *Leuconostoc mesenteroides* subsp. *dextranicum*, respectively. The coefficients of similarity (>100) between one strain of *Leuconostoc mesenteroides* and one strain of *Weissella paramesenteroides* were 36%, 48%, 44.4%, 48.6%, 44.4% and 48.6%, for the following couples (L₄, 13), (L₄, 6), (43, 13), (43, 6), (27, 13) and (27, 6), respectively, in agreement with distant phylogenetic relationship between *Leuconostoc mesenteroides* and *Weissella paramesenteroides*. The SDS-PAGE method allowed clarifying some ambiguous points in phenotypic identification. It had corroborate, complete and correct phenotypic identification, although, further study is required to accurate it.

Key words: Lactic acid bacteria • Identification • Phenotypic method • SDS-PAGE • raw milk • *Leuconostoc*

INTRODUCTION

During the last decade studies concerning development of new dairy products have focused on selection of new strains isolated from wild niches that are able to increase biodiversity and restore the unique characteristics of traditional dairy products [1].

Lactic acid bacteria (LAB) are Gram positive and usually catalase negative, grow under anaerobic conditions but they are aerotolerant, non-spore forming and have a fermentative sugar metabolism with lactic acid as a major final product. They have a great economic importance in dairy and other fermented food industries [2-4].

Lactic acid bacteria consist of a wide range of genera including a considerable number of species. Traditionally, they were divided into four genera, *Streptococcus*, *Lactobacillus*, *Leuconostoc* and *Pediococcus*. However, a considerable change in the taxonomy of lactic acid bacteria was observed during the last years [5].

Currently, the group of LAB in foods exist in the following genera *Carnobacterium*, *Enterococcus*, *Lactococcus*, *Streptococcus*, *Leuconostoc*, *Oenococcus*, *Lactobacillus*, *Pediococcus*, *Tetragenococcus*, *Vagococcus*, *Weissella*, *Aerococcus*, *Alloiococcus*, *Globicatella* and *Dolosigranulum* [6].

Phenotypic identification of LAB in dairy products are based mainly on morphological, physiological and biochemical characteristics, which are useful and indispensable tools that have been commonly used [7].

Although, phenotypic tests provide some evidence of metabolic capabilities, there are some problems, such as a lack of reproducibility and a lack of discriminatory power. Designation of certain neotype strains based only on phenotypic characteristics gave confused results which were resolved only by using molecular technique [8]. Recently, genetic techniques such as the mol % G+C contents of the DNA, DNA: DNA hybridization studies and structures and sequence of rRNA have been developed in order to get more consistent and accurate

identification of LAB [9-11]. Ogier *et al.* [9] observed a spatial distribution of bacteria species in cheese, based on metabolite availability and competition between bacteria. The bacteria of the cheese surface generally correspond to high G+C content genomes, whereas bacteria present in the core generally have low G+C content genomes.

Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) of whole cell protein is widely used for identification, since it offers the advantage to have a good level of taxonomic resolution at species and subspecies. Molecular methods have been used successfully to identify lactic acid bacteria isolated from different fermented foods [12-14]. The interest in LAB occurring in milk is primarily due to the biotechnological potential of new species.

The use of starters improved the technological quality of dairy products, but at the same time limited their biodiversity as well as the organoleptic variation of the end products. Therefore, an increasing demand exists for new strains that show desirable effects on the product characteristics [15].

The objective of this study is to identify the LAB isolated from Algerian raw milk using phenotypic methods and whole cell proteins fingerprinting.

MATERIALS AND METHODS

Sampling: Raw milk samples (cow, goat and camel) were collected from Oran, Mascara and Tindouf regions between June 2001 and January 2002. The Roquefort French cheese was also used to isolate LAB.

Isolation, Purification and Growth Conditions of Bacterial Strains: In order to select the *Leuconostoc* species, the strains have been isolated on MRS agar at pH 6.5, supplemented with 25 µg/ml of vancomycin [16] (MRSv) and were incubated at 28°C for 48h. Other LAB has been isolated on MRS agar without the presence of vancomycin at pH 6.5 and was incubated at 28°C and 45°C for 24h-72h. One reference strain of *Leuconostoc mesenteroides* subsp. *dextranicum* (L₄) obtained from Biotechnology-Microbiology Laboratory of ENSBANA, Dijon (France), has been used in this study. Morphologically, different colonies were selected and purified in the same previous conditions. A total of 62 isolates which were Gram positive and catalase negative, isolated from raw milks and Roquefort French cheese, have been retained.

Stock cultures were routinely maintained in MRS agar slants at 4°C and subcultured every 15 days. For long-term storage, frozen samples (-20°C) were prepared in sterile skim milk with 30% (v/v) glycerol. Enumeration of colonies of LAB in raw coagulate milk samples was performed.

Strains Identification: On the basis of morphological criteria, 21 dominant strains were chosen for this study. They were tested for: The morphological aspect, the proteolytic activity on PCA at 2% of skim milk, the production of CO₂ from glucose, the growth at different temperatures 10, 15, 37 and 45°C, the thermotolerance at 60°C for 30min, the growth at 6.5% NaCl and at pH 9.6, the hydrolysis of arginine, the dextran production which was detected at 10 % sucrose agar medium and the hydrolysis of aesculin was tested with 0.5% (w/v) aesculin. The citrate utilization, in the presence of glucose, was performed on the KMK medium [17].

The production of acetoin from glucose was determinate using Voges-Proskauer test on Clak and Lubs broth. All isolates were tested for the fermentation of the following carbohydrates: L(+) arabinose, D(-) ribose, D(+) xylose, dulcitol, glucose, galactose, fructose, D-mannose, D-mannitol, sorbitol, lactose, maltose, sucrose, D(+) cellobiose, D(+) raffinose and L(+) rhamnose (4, 5).

The identification of LAB isolates was referred to the following authors: Garvie [18], Collins *et al.* [19] for genus *Leuconostoc* and *Weissella*. Garvie [20] for *Pediococcus*. Farrow and Collins [21], Schleifer and Kilpper-Bälz [22], Schleifer *et al.* [13], for *Streptococcus*, *Lactococcus* and *Enterococcus*. Garvie [20] for *Lactobacillus* Samelis *et al.*, [12].

Electrophoresis SDS-PAGE of Whole Cell Protein Extracts: Only 14 representative species have been chosen for closer phenotypic profiles.

Sample Preparation: Cultures were grown in MRS broth, for 18h-48h and were streaked on buffered MRS agar (phosphate buffer 0.1M, pH 6.8) then were incubated for 18h to 48h. Cells were collected and resuspended in distilled water, then, harvested by centrifugation at 10000 rpm for 10min. The pellet was washed twice with distilled water and 50mg from the pellet was recuperated for each sample. Cells were sonicated at 30W for 30 sec and then incubated on ice for 20 sec. This performance was repeated twice. Subsequently, samples were resuspended in 0.9ml phosphate buffer 0.1M (pH 6.8), containing 5%

β -mercaptoethanol plus 5% sucrose, then an 0.1ml SDS (20%) was added to the suspension and the mixture was heated at 100°C for 2min. After cooling the mixture on ice and centrifuging at (10000 rpm for 10 min), the supernatant was recuperated and preserved at -20°C [23].

Protein concentration in the supernatant was measured by Bradford assay, using bovin serum albumin as standard. A standard protein solution was prepared with serum albumin (68kDa), casein (24kDa) and β -lactoglobulin (18kDa).

SDS-PAGE Electrophoresis: Cell-free extracts were subjected to SDS-PAGE electrophoresis on vertical slabs gel. The polyacrylamide gels were consisted of resolving gel 12% and stacking gel 5%. The electrode buffer was 0.3% (w/v) Tris, 1.44% (w/v) glycine and 0.1% (w/v) SDS (pH 8.3). A volume of sample containing 50 μ g of proteins was layered on top of gel and 10-20 μ l of loading sample buffer [0.01% (w/v) Bromophenol blue, 2% (w/v) SDS in buffer phosphate 1M (pH 6.8)] was added to each layer as a visible marker. Electrophoresis was performed at current of 50 mA and 120V. Proteins were fixed by immersing the gels in an aqueous of 10% (v/v) trichloroacetic solution for 1h and stained overnight in coomassie blue stain [0.25% (w/v) coomassie blue R-250, 50% (v/v) methanol and 10% (v/v) acetic acid]. The excess stain was washed out by destaining the gel with a solution of 25% (v/v) methanol and 10% (v/v) acetic acid [23].

Analysis of Protein Patterns: Molecular weight of proteins in electrophoregrams and coefficients of similarity (Jaccard coefficient) were calculated.

$$Sab = Ns / Ns + Nd$$

Sab: coefficients of similarity between two strains (a) and (b). Ns: Number of proteins having similar molecular weight between two strains (a) and (b) in electrophoregram. Nd: Number of proteins having different molecular weight between two strains (a) and (b) in electrophoregram.

RESULTS AND DISCUSSION

Results of the enumeration of lactic acid bacteria on MRS and MRSv have shown that the proportion of mesophilic lactic acid bacteria resisting to vancomycin are respectively: 7.3%, 0.265% and 1.7% from raw milk

of goat, cow and camel. Mathot *et al.* [16] had mentioned the resistance of *Leuconostoc* strains and the sensitivity of *Lactococcus* strains towards vancomycin at this concentration. However resistance to this antibiotic is among the general features of *Leuconostoc*, although it has been shown that *Pediococcus* and *Lactobacillus* are also vancomycin resistant. The 21 isolates were identified as belonging to the genus *Leuconostoc-Weissella* (10), *Lactobacillus* (8), *Lactococcus* (1), *Pediococcus* (1) and *Enterococcus* (1).

Phenotypic Identification of Lactic Acid Bacteria: Firstly, we have limited our research to the group of LAB that can be found in milk and dairy products. Strains numbering 17, 2, 13, 6, 43 and 27 isolated on MRSv and strains 19, 39 on MRS without vancomycin have the coccoid rod cells and occurred in pairs, in short chains or little mass. They were heterofermentative mesophilic, they did not hydrolyse arginine and they were considered as *Leuconostoc* (Table 1). Production of dextran from sucrose is an important character [18] to differentiate *Leuconostoc* species. Only two strains L₄ (*Leuconostoc mesenteroides* subsp. *dextranicum*) and 43 had shown this property. A weak dextran production was detected with the 43 strain. Devoyod and Poullain [24] had also isolated the species of *Leuconostoc mesenteroides* subsp. *mesenteroides* from dairy products which had also shown a weak dextran production. The *Leuconostoc* dextran (+) can be differentiated by the ability to ferment L-arabinose [25]. This reaction was negative with the strains L₄, 43, which were identified as *Leuconostoc mesenteroides* subsp. *dextranicum* (Table 1). On the basis of sugar fermentation patterns, we could differentiate them into 43 (xylose⁻, mannitol⁻, raffinose⁺) and L₄ (xyl⁺, mannit⁺, raff⁻).

The *Leuconostoc cremoris* has a limited sugar fermentation profile and does not grow at 37°C. This is not the case of strains 17, 2, 13, 6, 27, 19 and 39 which doesn't produce dextran and doesn't ferment fructose, sucrose, maltose and mannose (Table 1). It is difficult to distinguish between *Leuconostoc lactis* and *Leuconostoc paramesenteroides* by their sugar fermentation patterns [25]. However, our leuconostocs were assigned to *Leuconostoc paramesenteroides* (currently *Weissella paramesenteroides*; Collins *et al.* [19]. Furthermore, they can be divided into four groups: Group 1: 27 (arabinose⁺, xylose⁻, Cellobiose⁻, raffinose⁻, mannitol⁺). Group 2 : 6, 1, 2, 17, 39 (ara⁻, xyl⁺, Celb⁺, raff⁺, mannit⁻). Group 3 : 13 (ara⁻, xyl⁻, Celb⁻, raff⁺, mannit⁻). Group 4 : 19 (ara⁻, xyl⁻, Celb⁺, raff⁺, mannit⁻).

Table 1 : Physiological and biochemical properties of coccoid and lactic acid bacteria.

Tests	Strains												
	17	39	2	19	13	6	38	21	43	27	1	L ₄	20
Fermentative metabolism	ht	ht	ht	ht	ht	ht	hm	hm	ht	ht	ht	ht	hm
Growth in NaCl 6,5%	-	-	-	-	-	-	-	-	-	-	-	-	+
Growth at 45°C	-	-	-	-	-	-	+	-	-	-	-	-	+
Growth at 37°C	±	±	+	+	+	+	+	+	±	+	+	+	+
Growth at 10°C	+	+	+	+	+	+	-	+	+	+	+	+	+
Growth at 50°C	Nt	Nt	Nt	Nt	Nt	Nt	+	Nt	Nt	Nt	Nt	Nt	Nt
Thermotolerance	-	-	-	-	-	-	+	-	-	-	-	-	+
Growth in pH 9,6	-	-	-	-	-	-	-	-	-	-	-	-	+
Hydrolysis of arginine	-	-	-	-	-	-	+	+	-	-	-	-	+
Citrate (KMK)	+	+	+	+	+	+	+	+	+	-	+	+	+
VP	-	-	-	-	-	-	+	+	+	-	-	+	-
Dextran	-	-	-	-	-	-	Nt	Nt	±	-	-	+	Nt
Hydrolysis of aesculin	-	-	-	-	-	+	-	-	-	+	+	-	+
Proteolytic activity	+	+	+	+	+	+	+	+	+	+	+	+	+
Sugar fermentation :													
Fructose	+	+	+	+	+	+	+	+	+	+	+	+	+
Dulcitol	-	-	-	-	-	-	-	-	-	-	-	-	-
Galactose	+	+	+	+	+	+	+	+	±	+	+	+	+
Glucose	+	+	+	+	+	+	+	+	+	+	+	+	+
D(+) xylose	+	+	+	-	-	+	±	-	-	-	+	+	-
L(+) arabinose	-	-	-	-	-	-	+	+	-	+	-	-	+
Maltose	+	+	+	+	+	+	-	+	+	+	+	+	+
Lactose	+	+	+	+	+	+	+	+	+	±	+	+	+
L(+) rhamnose	-	-	-	-	-	-	-	-	-	-	-	-	-
Cellobiose	±	±	+	±	-	+	+	+	±	-	±	+	+
Sucrose	+	+	+	+	±	+	-	+	+	+	+	+	+
D mannose	±	±	±	+	±	+	+	+	±	+	+	+	+
D mannitol	-	-	-	-	-	-	-	+	-	±	-	+	±
D(-) ribose	+	+	+	±	±	+	+	+	+	+	+	+	+
D(+) raffinose	+	+	+	+	+	+	-	±	+	-	+	-	-
Sorbitol	-	-	-	-	-	-	-	-	-	-	-	-	-

Symbols : +; positive reaction at 24h or 48h , -; negative reaction, ±; weak and / or delayed reaction (the time of reaction exceed 48h), ht; heterofermentative metabolism, hm; homofermentative metabolism, Nt; not tested, KMK; Kempler and Mac Kay media contained 3g/l of yeast extract, VP; Voges Proskauer test (positive result detected after incubation 2-3 days, except, strain 21 at 6h)

Since the Ribose fermentation is a distinguishable character between *Leuconostoc lactis* and *Weissella paramesenteroides* (8). All the strains, mentioned previously, are characterised by ribose fermentation and they belong to *Weissella paramesenteroides*. Moreover, hence The strains 27, 13, 19 and all strains from group 2 are a variable sugar (Cellobiose, xylose, mannitol, arabinose and raffinose) (Table 1) fermentation, therefore, they cannot belong to the species of *Leuconostoc pseudomesenteroides*.

It should be noted that one *Weissella paramesenteroides* number 27, was citrate defective variant. The instability of plasmid encoded citrate metabolism is known in the case of *Leuconostoc* sp. [26]. Strains 20, 21 and 38 were homofermentative cocci.

On the basis of the growth at both 10°C and 45°C, in 6.5% NaCl, pH 9.6, surviving at 60°C for 30min and the hydrolysis of arginine, the strain 20 was identified as *Enterococcus* (Table 1). Notably, the ability of *Enterococcus faecium* to ferment arabinose led to distinct it from *En. faecalis* and *En. durans* [27,28].

Except, the growth at 10°C and arginine hydrolysis, the above other tests were negative for strain 21, therefore, it is assigned to the genus *Lactococcus*. It is arginine(+) and citrate(+) and VP(+). The latter properties had suggested the identification of this strain as *Lc. lactis* subsp. *lactis* biovar. *diacetylactis*.

Strain 38 had spherical cells, formed tetrads and was assigned to genus *Pediococcus*. Strain 38 was heat resisting, able to grow at 45°C but did not grow at

Table 2: Physiological and biochemical properties of lactobacilli

Tests	Strains							
	23	24	52	58	61	25	50	32
Fermentative metabolism	hm	hm	hm	hm	hm	hm	hm	hm
Growth at 45°C	+	+	+	-	+	+	+	-
Growth at 15°C	+	+	-	+	+	+	-	+
Arginine hydrolysis	-	-	-	-	-	-	-	-
Citrate (KMK)	-	-	+	+	+	-	-	-
Vp	+	+	+	+	+	-	+	+
Aesculin hydrolysis	-	-	+	+	+	-	-	-
Proteolytic activity	+	+	+	+	+	+	+	+
Sugar fermentation :								
Fructose	+	+	+	+	+	+	+	+
Dulcitol	-	-	-	-	-	-	-	-
Galactose	+	+	+	+	+	+	+	+
Glucose	+	+	+	+	+	+	+	+
D(+) xylose	-	-	-	-	-	-	-	-
L(+) arabinose	-	-	+	+	+	+	-	-
Maltose	+	+	+	+	+	+	+	+
Lactose	+	+	+	+	+	+	+	+
L(+) rhamnose	+	+	-	-	-	-	±	-
Cellobiose	+	+	+	+	+	-	±	+
Sucrose	±	+	+	+	+	+	+	+
D mannose	+	+	+	+	+	+	+	+
D mannitol	+	+	+	+	+	+	+	+
D(-) ribose	+	+	+	+	+	±	+	+
D(+) raffinose	-	-	+	+	+	-	+	-
Sorbitol	+	+	+	+	+	-	+	±

Symbols : +; positive reaction at 24h or 48h, - ; negative ; reaction, ±; weak and / or delayed reaction (the time of reaction exceed 48h), ht; heterofermentative metabolism, hm; homofermentative metabolism, KMK; Kempler and Mac Kay media contained 3g/l of yeast extract, VP ; Voges Proskauer test (positive result detected after incubation 2-3 days)

temperature of 10°C, 6.5% of NaCl and pH 9.6. Contrary to *Streptococcus thermophilus*, it was arginine positive, fermented arabinose, cellobiose and did not fermented sucrose, consequently, the previous definition was confirmed, unlike to *Pc. damnosus*, which is able to grow at pH 7.0, at 37°C and at 45°C. On the other hand, *Pc. acidilactici* and *Pc. pentosaceus* are closely related species that may not be clearly differentiated by phenotypic characteristics, but they are differentiated by ADN-AND homology. On the basis of growth at 50°C, the strain 38 was identified as *Pc. acidilactici* [5,25,28]. The latter was the only *Pediococcus* (isolated from raw milk and dairy products) that tolerate this temperature.

All strains 23, 24, 52, 58, 61, 25, 50 and 32 (Table 2) were characterised as mesophilic homofermentative lactobacilli belong to the group [12,25]. These strains do not hydrolyse arginine.

The rod-shaped cells from strain 25 had a tendency to a curve. Strains 52 and 61 had shown thickest rod cells. Colonies from the strains 58, 61, 52 and 25 were easily distinguished from all the other strains. Strains 58, 61 and 52 were larger, more convex and cheesy white, however, 25 were smaller colonies, lesser convex and transparency white.

The strains 23, 24, 52, 58, 61, 50 and 32 (Table 2) were lactose positive, ribose positive, sucrose positive and sorbitol positive, that is why it cannot belong to *Lactobacillus casei*.

The fact that all those strains were sorbitol positive and able to grow at 15°C, their identification was restricted to *Lb. paracasei* subsp. *paracasei*, *Lb. plantarum*, *Lb. rhamnosus*. Only *Lb. pentosus* doesn't grow at 15°C. Moreover, fermenting arabinose by strains 52, 58 and 61, it's the other argument in favour. These strains were

Table 3: Coefficients of similarity (>100) between each two strains of LAB

Strains	L ₄	13	6	43	27	52	58	32	50	23	25	38	20
13	36.0												
6	48.6	80.7											
43	78.0	44.4	48.6										
27	78.0	44.4	48.6	100.0									
52	27.1	30.2	37.2	30.6	30.6								
58	26.8	22.5	25.0	30.9	30.9	55.5							
32	29.5	18.2	27.9	30.4	30.4	77.8	52.0						
50	35.9	23.0	27.5	33.3	33.3	36.4	37.8	36.6					
23	27.5	21.0	19.5	28.6	28.6	31.8	36.0	31.7	85.2				
25	35.2	9.3	11.7	7.5	7.5	7.0	8.6	10.2	8.3	8.8			
38	19.1	18.6	22.7	11.3	11.3	31.6	22.7	22.9	14.6	6.5	2.4		
20	25.7	25.8	23.5	20.5	20.5	18.6	16.6	20.5	22.8	17.1	11.11	17.9	
21	21.1	20.8	22.0	22.2	22.2	29.6	32.6	26.9	21.5	22.4	4.2	35.4	26.2

grown at 15°C and, except 58, did not grow at 45°C. They did not ferment rhamnose, therefore, were assigned to *Lb. plantarum* or *Lb. paracasei* subsp. *paracasei* and strain 58 to *Lb. paracasei* subsp. *paracasei*.

Strains 23, 24, 50 and 32 were arabinose negative and rhamnose positive except strain 32 which was rhamnose negative. Also, the ability of their growth on 15°C and 45°C was observed with all the strains except with strain 32 which did not show a grow on 45°C. Thus, 23, 24, 50 were assigned to *Lb. rhamnosus* and 32 (resistant to 25µg/ml to vancomycin) to *Lb. plantarum* or *Lb. paracasei* subsp. *paracasei*. Furthermore, The fermentation of raffinose can differentiate strains of *Lb. rhamnosus* into groups; (1) containing strain 50 was raffinose positive and (2) containing strains 23, 24 were raffinose negative.

The following characteristics sorbitol negative and cell-shaped indicated that strain 25 (resistant to 25µg/ml to vancomycin) do not belong to any of the above species of lactobacilli. Sugar fermentation pattern of this strain from it did not perfectly match coincided with the other lactobacilli in group II. Therefore, it was remain to be unidentified.

Identification by Whole Cell Protein Analysis: The results obtained by SDS-PAGE of whole-cell proteins discriminates as much as the DNA-DNA hybridization [29,30] and present a very good correlation between the results of numerical analysis of protein patterns and 16 rRNA based on oligonucleotide probe hybridization has been reported [14,31,32].

Our phenotypic identification is a little ambiguous. In order to resolve this problem and improve it, protein fingerprinting by SDS-PAGE method as complementary analysis to phenotypic characterization has been used. The analysis of cell protein extracts by SDS-PAGE of 14 representative strains was studied. It was undertaken by visual comparison of the electrophoretic patterns




Fig. 1: Whole cell protein profiles of *Leuconostoc* (L4, 43, 27) and *Weissella*




Fig. 2: Whole cell protein profiles of *Lactobacillus* (52, 58, 32, 50, 23) and *Enterococcus faecium* (20)

(Fig. 1, Fig. 2). That was done by the coefficients of similarity (>100) calculated by using the molecular weight of patterns in electrophoregrams. The coefficients of similarity (>100) were obtained for each two strains of the studied LAB (Table 3).

Sanchez *et al.* [13] have observed that the SDS-PAGE technique generated complex and stable patterns that were easy to interpret and compare with the reference strains of lactic acid bacteria. After numerical analysis of the resulting electrophoretic protein patterns the clusters

were discriminated with some exceptions and correlated well with different groups phenotypically found on Almagro Spain cheese.

Leuconostoc and Weissella paramesenteroides: The results of SDS-PAGE confirmed the assignment of strain 43 (isolated from goat milk, xyl⁻, mannit⁻, raff⁺) to *Ln. mesenteroides* subsp. *dextranicum* by coefficient of similarity of 78% (Table 3) with strain L₄ (ENSBANA-Dijon, xyl⁺, mannit⁺, raff⁻). This result clarifies the identity of the *Leuconostoc mesenteroides* subsp. *dextranicum* strain 43. The slight dissimilarities between strains 43 and L₄ may be due to the different origin of the compared strains, as it was indicated by Samelis *et al.* [12]; Pérez *et al.* [33].

The two strains 6 (isolated from camel milk, ara⁻, xyl⁺, Celb⁺, raff⁻, mannit⁻) and 13 (isolated from cow milk, ara⁻, xyl⁻, Celb⁻, raff⁺, mannit⁻) have shown an 80.7% of similarity. That corroborated the assignment of them to the same identity, phenotypically defined as *Weissella paramesenteroides*.

Protein fingerprinting of strains 43 and 27 (isolated from the same goat milk sample) were very similar with a coefficient of 100%. Nevertheless, they were phenotypically identified as *Ln. mesenteroides* subsp. *dextranicum* for the first strain and *Weissella paramesenteroides* for the second. The same phenomenon has been found by Perez *et al.* [33]. The difficulty of the electrophoresis of cell-free extracts is to distinguish between the *Leuconostoc* subspecies which has been reported by Tsakalidou *et al.* [34] and Villani *et al.* [35] and it is probably because they are very close phylogenetic subspecies which produce very similar protein patterns [31,36]. These data and the sugar fermentation pattern indicated that it is quite possible to identify strain 27 as *Ln. mesenteroides* subsp. *mesenteroides*, which has lost dextran production and citrate utilisation. Several authors have mentioned this specificity in *Leuconostoc mesenteroides* [24,26].

The difference from two characters (arabinose, dextran) between strains 43 and 27 could not be demonstrated by the SDS-PAGE protein profile analysis, the same fact has been reported by Sanchez *et al.* [13].

The strain 27 and the strains 13 and 6, which were previously identified as *Weissella paramesenteroides*, had shown a low coefficient of similarity (Table 3). Hence, the protein profile revealed the atypical character of the strain 27.

The strain couples (L₄, 13), (L₄, 6), (43,13), (43,6), (27,13) and (27, 6) had given the following coefficients of

similarity 36, 48, 44.4, 48.6, 44.4 and 48.6%, respectively. These low level of similarity are in agreement with distant phylogenetic relationship between the genus of *Leuconostoc* sp. and *Weissella* [19].

Lactobacilli: The two strains 23 and 50 (respectively raff⁻ and raff⁺, isolated from the same sample of goat milk) have shown a high level of similarity with 85.2% (Table 3) and were defined as *Lb. rhamnosus*.

The strains 58, isolated from goat milk and 52, isolated from camel milk, except, the growth at 45°C for the first strain, they presented similar phenotypic profile. Strain 58 was previously identified as *Lb. paracasei* subsp. *paracasei*. The coefficient of similarity 55.5% between the two strains, have shown different identity of them. Consequently, if strain 58 had previously identified as *Lb. paracasei* subsp. *paracasei*, the strain 52 had assigned as *Lb. plantarum* rather than *Lb. paracasei* subsp. *paracasei*. On the other hand, the coefficient of similarity between strains 52 (isolated from camel milk, ara⁺, raff⁺) and 32 (isolated from roquefort cheese, ara⁻, raff⁻) was 77.8%. In regard to the level of similarity 52% between strains 58 and 32, the latter strain is closely related to 52 rather than 58 and could be named *Lb. plantarum*.

The low coefficients of similarity between strains 50 and 52, 50 and 58, 50 and 32, 23 and 52, 23 and 58, 23 and 32 (Table 3) are in agreement with the precedent propositions.

Therefore, protein electrophoresis SDS-PAGE had allowed the separation of strains possessing very high or similar phenotypic profiles. On the other hand, the possible mistake to the assignment of strain 52 to the genus *Lactobacillus* or *Lactococcus*, which can be caused by one's rod cell shaped (making the morphological determination difficult), had been excluded by the above result. The very low coefficients of similarity between *Lactobacillus* 25 and other *Lactobacillus* (Table 3) confirmed the phenotypic results, so, it is clearly that strain 25 is distant to the group of *Lb. plantarum*, *Lb. paracasei*, *Lb. rhamnosus*.

Simova *et al.*, [37] described the clustering of kefir lactic acid bacteria strains by using amino acid profiles.

The SDS-PAGE technique generated complex and stable patterns that were easy to interpret and compare the profiles of the strains used in this study. After numerical analysis of the resulting electrophoretic protein pattern, the clusters were discriminated and correlated well with the different groups phenotypically found [13].

Homofermentative Cocci: The very low coefficient of similarity 17.9% (Table 3) between strains 38 and 20 indicated that strain 38 can not be identified as *Enterococcus faecium*.

The microscopic observation of strain 52 had shown the thickest rod-shaped cells, which is easy to be confused with cocci. If this strain was assigned to the genus *Lactococcus*, therefore, it could be identified phenotypically as *Lactococcus lactis* biovar. *diacetylactis*. However, the low coefficient of similarity 29.6% between strains 52 and 21 showed that the previous proposition was not correct. The reason of this problem is that lactobacilli may produce very short or ellipsoid cells, under certain growth conditions. Pérez *et al.* [33] had indicated that the major problem encountered in the classical identification of LAB was the assignment to the genera *Lactococcus* or *Lactobacillus* which was based on the microscopic observation of the strains.

All strains occurring coccoid rod cells isolated from MRS supplemented with 25µg/ml of vancomycin were identified as *Leuconostoc* or *Weissella paramesenteroides*. *Weissella paramesenteroides* was dominante heterofermentative cocci in raw milk samples.

The biochemical and physiological tests are unsatisfactory for the identification of isolated LAB. The use of SDS-PAGE method had allowed the clarification of some ambiguous points in phenotypic identification, for example, it allowed the correction of the misidentification which is due to the presence of atypical characters, separation between strains which have closer phenotypic profiles, resolve the problem of microscopic determination of cell shape. Consequently, our results show that, protein fingerprinting analysis corroborate, complete and confirm the phenotypic identification. De Vuyst et Vancanneyt [38] shows that the isolation of novel taxa mainly depends on the cultivation approach used selective incubation media and conditions. The identification of isolated strains needs a polyphasic approach, including a combination of phenotypic and genotypic methods.

REFERENCES

1. De Palencia, P.F., M. Plaza, F. Amarita, T. Requena and C. Pelaez, 2006. Diversity of amino acid converting enzymes in wild lactic acid bacteria. *Enz. Microbial. Technol.*, 38: 88-93.
2. Badis, A., N. Laouabdia-Sellami, D. Guetarni, M. Kihal and R. Ouzrout, 2005. Caractérisation phenotypique des bactéries lactiques isolées à partir de lait cru de chèvre de Deux populations caprines locales Arrabia et Kabyle. *Sci. Technol.*, 23: 30-37.
3. Guessas and M. Kihal, 2004. Characterization of lactic acid bacteria isolated from Algerian arid zone: Raw goats' milk. *African J. Biotechnol.*, 3(6): 339-342.
4. Thapa, N., J. Pal and J.P. Tamang, 2006. Phenotypic identification and technological properties of lactic acid bacteria isolated from traditionally processed fish products of eastern Himalayas. *Int. J. Food. Microbial.*, 107: 33-38.
5. Stiles, M.E., H. Wilhelm and W.H. Holzapfel, 1997. Lactic acid bacteria of foods and their current taxonomy. *Int. J. Food Microbiol.*, 36: 1-29.
6. Axelsson, L., 1998. Lactic acid bacteria: classification and physiology. In: S. Salminen, A. Von wright and M. Dekker (Eds.), *Lactic Acid Bacteria: Microbiology and Functional Aspects*, 2nd edition, Inc., New York, pp: 1-72.
7. Badis, A., D. Guetarni, B. Moussa-Boudjemaa, D.E. Henni and M. Kihal, 2004b. Identification and technological properties of lactic acid bacteria isolated from raw goat milk of four Algerian races. *Food Microbiol.*, 21: 579-588.
8. De Angelis, M., A. Corsetti, N. Tosti, J. Rossi, M.R. Corbo and M. Gobbetti, 2001. Characterization of non starter lactic acid bacteria from Italian ewe cheeses based on phenotypic, genotypic and cell wall protein analysis. *Appl. Environ. Microbiol.*, 67(5): 2011-2020.
9. Ogier, J.C., V. Lafarge, V. Girard, A. Rault, V. Maladen, A. Gruss, J.Y. Ieveau and A. Delacroix-Buchet A., 2004. Molecular fingerprinting of dairy microbial ecosystems by use of temporal temperature and denaturing gradient gel electrophoresis. *Appl. Environ. Microbiol.*, 70(9): 5628-5643.
10. Catzeddu, P., E. Mura, E. Parents, M. Sanna and G.A. Farris, 2006. Molecular characterization of lactic acid bacteria from sourdough breads produced in Sardinia (Italy) and multivariate statistical analyses of results. *Syst. Appl. Microbiol.*, 29: 138-144.
11. Bensalah, F., M.J. Flores and A. Mouats, 2006. A rapid PCR based method to distinguish between *Enterococcus* species by using degenerate and species-specific *sodA* gene primers. *African J. Biotechnol.*, 5(9): 697-702.

12. Samelis, J., Tsakalidou, E., Metaxopoulos, J., Kalantzopoulos, G., 1995. Differentiation of *Lactobacillus sake* and *Lactobacillus curvatus* isolated from naturally fermented Greek dry salami by SDS-PAGE of whole cell proteins. J. Appl. Bacteriol. 78, 157-163.
13. Sanchez, I., Sesena, S., Palop, L., 2003. Identification of lactic acid bacteria from spontaneous fermentation of 'Almagro' eggplant by SDS-PAGE whole cell protein fingerprinting. Int. J. Food Microbiol., 2555: 181-189.
14. Sacilik S.C., Osmanagaoglo O., Gunduz U., Cokmus C., 2000. Availability of use of total extracellular proteins in SDS-PAGE for characterization of gram positive cocci. Turk. J. Biol. 24 : 817-823 .
15. Zamfir, M., M. Vancanneyt, L. Makras, F. Vaningelgem, K. Lefebvre, B. Pot, J. Swings and L. Vuyst, 2006. Biodiversity of lactic acid bacteria in Romanian dairy products. Syst. Appl. Microbiol., 29: 487-495.
16. Mathot, A.G., M. Kihal, H. Prevost and C. Diviès, 1994. Selective enumeration of *Leuconostoc* on vancomycin agar Media. Int. Dairy. J., 4: 459-469.
17. Kempler, G.M. and L.L. Mac Kay, 1980. Improved medium for detection of citrate fermenting *Streptococcus lactis* subsp. *diacetylactis*. Appl. Environ. Microbiol., 39 (4): 926-927.
18. Garvie, E.I., 1986b. Genus *Leuconostoc* Van Tieghem 1978. In: Sneath, P.H.A., Mair, N.S., Sharpe, M.E., Holt J.C. (Eds.), Bergey's Manual of systematic Bacteriology Vol 2. Williams and Wilkins, Baltimore, MD, pp: 1071-1075.
19. Collins, M.D., J. Samelis, J. Metaxopoulos and S. Wallbanks, 1993. Taxonomic studies on some *Leuconostoc*-like organism from fermented sausages: Description of a new genus *Weissella* for the *Leuconostoc paramesenteroides* group of specie. J. Appl. Bacteriol., 75: 595-603.
20. Garvie, E.I., 1986a. Genus *Pediococcus* Claussen 1903. In: Sneath, P.H.A., Mair, N.S., Sharpe, M.E., Holt, J.C. (Eds.), Bergey's Manual of systematic Bacteriology, vol. 2. Williams and Wilkins, Baltimore, MD, pp. 1075-1079.
21. Farrow, J.A.E. and M.D. Collins, 1984. DNA base composition, DNA-DNA homology and long-chain fatty acid studies on *Streptococcus thermophilus* et *Streptococcus salivarius*. J. Gen. Microbiol., 130: 357-362.
22. Schleifer, K.H. and R. Kilpper-Bälz, 1984. Transfer of *Streptococcus faecalis* and *Streptococcus faecium* to the genus *Enterococcus* NOM. rev. as *Enterococcus faecalis* Comb. nov and *Enterococcus faecium* Comb. nov. Int. J. Syst. Bacteriol., 34,:31-34.
23. Kersters, K., 1980. Polyacrylamide Gel electrophoresis of bacterial protein. In: Clement, Z., Rudolph, K., Sands, D.C. (Eds.), Methods in phytobacteriology. Akademia Kiado, Budapest, pp: 191-197.
24. Devoyod, J.J. and F. Poullain, 1988. Les leuconostocs propriétés: Leur rôle en technologie laitière. Le lait, 68 (3), 249-280.
25. Singh, S.K., S.U. Ahmed and A. Pandey, 2006. Metabolic engineering approaches for lactic acid production. Process. Biochem., 41: 991-1000.
26. Kihal, M., H. Prevost, M.E. Lhotte, D.Q. Huang and C. Diviès, 1996. Instability of plasmid encoded citrate permease in *Leuconostoc*. Lett. Appl. Microbiol., 22: 219-223.
27. Schleifer, K.H., J. Kraus, C. Dvorak, R. Kilpper-Bälz, M.D. Collins and W. Fischer, 1985. Transfer of *Streptococcus lactis* and related streptococci to the genus *Lactococcus* gen. Nov. System. Appl. Microbiol., 6: 183-195.
28. Carr, F.J., D. Chill and N. Maida, 2002. The lactic acid bacteria: A literature Survey. Cur. Rev. Microbiol., 28(4): 281-370.
29. Ferraz, P.N., C.C.P. Menezes, G.M. Danelli, J.O.P. Lizeu, E.R. Nascimento and M.D. Lucchesi, 2000. protein profile analysis by SDS-PAGE of mycoplasma gallisepticum strains S6(208) and F-K810 growth in hayflick's and frey's media. Brazilian. J. Microbiol., 31(2): 11-17.
30. Priest, F.G., Austin, B., 1993. Modern bacterial taxonomy (2nd ed) Chapman and Hall, London.
31. Pot, B., C. Hertel, W. Ludwig, P. Descheemaeker, K. Kersters and K.H. Schleifer, 1993. Identification and classification of *Lactobacillus acidophilus*, *L. Gasseri* and *L. Johnsoni*. Strains by SDS-PAGE and rRNA-targeted oligonucleotide probe hybridisation. J. Gen. Microbiol., 139: 513-517.
32. Booyesen, C., L.M.T. Dicks, I. Meijering and A. Ackermann, 2002. Isolation, identification and changes in the composition of lactic acid bacteria during the malting of two different barley cultivars. Int. J. Food Microbiol., 76: 63-73.

33. Pérez, G., E. Cardell and V. Zarate, 2000. Protein fingerprinting as a complementary analysis to classical phenotyping for the identification of lactic acid bacteria from Tenerife cheese. *Lait*, 80: 589-600.
34. Tsakalidou, E., E. Manolopoulou, E. Kabarak, E. Zoidou, B. Pot, K. Kersters and G. Kalantzopoulos, 1994. The combined use of whole-cell protein extracts for the identification (SDS-PAGE) and enzyme activity screening of lactic acid bacteria isolated from traditional Greek dairy products. *Syst. Appl. Microbiol.*, 17: 444-458.
35. Villani, F., G. Moschetti, G. Blaiotta and S. Coppola, 1997. Characterization of strains of *Leuconostoc mesenteroides* by analysis of soluble Whole-Cell protein pattern, DNA fingerprintings and restriction ribosomal DNA. *J. Appl. Microbiol.*, 82: 578-588.
36. Moschetti, G., G. Blaiotta, F. Villani and S. Coppola, S., 2000. Specific detection of *Leuconostoc mesenteroides* subsp. *mesenteroides* with DNA primers identified by randomly amplified polymorphic DNA analysis. *Appl. Environ. Microbiol.*, 66: 422-424.
37. Simova, E., Z. Simov, D. Beshkova, G. Frengova, Z. Dimitrov and Z. Spasov, 2006. Amino acid profiles of lactic acid bacteria, isolated from Kefir grains and kefir starter made from them. *Int. J. Food Microbiol.*, 107: 112-123.
38. De Vuyst, L. and M. Vancanneyt, 2006. Biodiversity and identification of sourdough lactic acid bacteria. *Food. Microbiol.*, 24(2): 120-127.