

Cytological investigations and new chromosome number reports in yarrow (*Achillea millefolium* Linnaeus, 1753) accessions from Iran

Fatemeh Afshari¹, Mohsen Ebrahimi², Mohammad Akbari³, Mostafa Farajpour²

1 Biology Department, Faculty of Science, Payam Noor University, Tehran, Iran **2** Department of Agronomy and Plant Breeding, College of Abouraihan, University of Tehran, Tehran, Iran **3** Department of Horticultural Sciences, Faculty of Agriculture, University of Tabriz, Tabriz, Iran

Corresponding author: *Mostafa Farajpour* (Farajpour_m@ut.ac.ir)

Academic editor: *J.R. Daviña* | Received 9 August 2013 | Accepted 7 October 2013 | Published 24 October 2013

Citation: Afshari F, Ebrahimi M, Akbari M, Farajpour M (2013) Cytological investigations and new chromosome number reports in yarrow (*Achillea millefolium* Linnaeus, 1753) accessions from Iran. *Comparative Cytogenetics* 7(4): 271–277. doi: 10.3897/CompCytogen.v7i4.6075

Abstract

In this study, a new chromosome number for Iranian yarrow (*Achillea millefolium* L.) accessions was reported. Cytological analyses on four *A. millefolium* accessions, indicated that two accessions were diploids ($2n=2x=18$) and two tetraploids ($2n=4x=36$). Cluster analysis based on chromosomal characteristics and karyotype asymmetry, categorized the four accessions separated into two groups. In terms of the Stebbins' system, the karyotype of diploid accessions grouped in 2A class. The average value of the total form percentage (TF%) in the group one (diploid accessions) and two (tetraploid accessions) were 40.85 and 41.15, respectively. The group one had the highest mean value for the symmetry index ($S\%=57.5$). Consequently, it can be inferred that diploids belonging to the group one are the earlier evolutionary forms.

Keywords

Achillea millefolium, karyotype, chromosome number, ploidy level

Introduction

Achillea is one of the most recent genera of the Asteraceae family which exists throughout the world (Rechinger 1963). More than 100 species have been identified in this genus. Many of those who used these plants reported properties such as anti-inflammatory, anti-rheumatic, antiseptic, antispasmodic, analgesic, astringent, carminative, diaphoretic, digestive, expectorant, hypotensive, stomachic and etc. (Balbir et al. 2012). These plants are native to Europe and Western Asia but are also found in Australia, New Zealand, and North America (Rechinger 1963).

Achillea millefolium has a high genetic and phenotypic variation in Iran (Farajpour et al. 2012, Ebrahimi et al. 2012). The basic chromosome number is often reported in different species of *Achillea* is $x = 9$; however, the diversity in chromosome numbers and ploidy levels are frequently occurring in the genus (Ebrahim et al. 2012). Polyploid taxa have originated in many clades including $4x$, $6x$ and $8x$ species, and as a result, several *Achillea* species show high morphological variability (Sheidai et al. 2009). Biste (1987) explained worthy diversity in leaf width, height, shoot number, and stomata length in different populations of the same species.

In most of the chemotypes in *Achillea* sp, camphor, borneol (Rohloff et al. 2000 and Mockute and Judzentiene 2003) and 1.8-Cineole (Saeidnia et al. 2004; Barghamadi et al. 2009 and Azizi et al. 2010) have been detected. Among a number of data that can be obtained by chromosome studies: karyotype structure, karyotype asymmetry, chromosome banding, FISH, GISH and chromosome painting (Stace 2000, Levin 2002, Graphodatsky et al. 2011, Guerra 2012), the most popular, cheap and widely used approaches is that concerning karyotype asymmetry (Peruzzi and Eroğlu 2013).

Achillea millefolium has been cytologically analyzed extensively in different regions of the world (Felfoldy 1947, Mizianty and Frey 1973, Pireh and Tyril 1980, Lavrenko et al. 1991, Guo et al. 2012, Bala and Gupta 2013). Three cytological studies have been reported in Iran and showed the following ploidy levels: tetraploid $2n = 4x = 36$, hexaploid $2n = 6x = 54$ and octoploid $2n = 8x = 72$ (Farsi et al. 2000, Ebrahim et al. 2012, Sheidai et al. 2009). The aims of this study were (1) to determine the chromosome numbers of four *A. millefolium* accessions and (2) to find any relationship between the karyotype characteristics and asymmetrical index with ploidy levels.

Material and methods

The aerial parts of the four *Achillea millefolium* accessions were collected from three provinces in north, west and south of Iran (Table 1). Voucher samples were deposited at the herbarium of Research Institute of Forests and Rangeland (RIFR) of Tehran, Iran. Seeds were germinated on moist filter paper in Petri dishes. Actively growing root tips, 1 to 2 cm length were cut from the germinating seeds and pretreated with 8-hydroxyquinoline (0.002M) for 2 to 4 hours and fixed in Farmer (1:3, glacial acetic acid : ethanol 95%) for 24 hours at 4° C. Thereafter, the root tips were hydrolyzed in

Table 1. Accessions of *Achillea millefolium* studied.

Accessions no.	Location	Latitude	Longitude	Elevation (m.a.s.l.)
Am1	Iran, Ardabil, Ardabil	38°15'N	48°17'E	1332
Am2	Iran, Ardabil, Meshkin-Shahr	37°58'N	48°58'E	1723
Am3 [†]	Iran, Ilam, Ilam	34°27'N	46°25'E	1387
Am4	Iran, Fars, Estahban	53°04'N	29°12'E	1767

[†]some of the characteristics of this accession were reported by Farajpour et al. (2012) in table 1 (Am30)

1 N NaOH at 60° C for 5-10 minutes, stained for 45 minutes in esterase stain at 30° C, and squashed in 45% glacial acetic acid. Finally, the chromosome images were obtained with photomicroscope.

Karyotypic characteristics such as differences of range relative length (DRL), mean chromosome length (MCL), and mean arm ratio (MAR) were calculated using MICROMEASURE (Version 3.3) Software (Reeves 2001). Stebbin's classification was calculated (Stebbins 1971). Cluster analysis was performed to differentiate the accessions according to the Ward's method SPSS software for Windows 20.0 (SPSS Inc., Chicago, IL, USA).

Results and discussion

Karyological data

Am1 and Am2 accessions were diploids ($2n = 2x = 18$) whereas the two other accessions (Am3 and Am4) showed tetraploid ($2n = 4x = 36$) level (Figure 1). According to previous studies, Farsi et al. (2009) and Khaniki (1995), reported $2n = 4x = 36$ chromosomes, while Sheidai et al. (2009) and Ebrahim et al. (2012) reported hexaploid and octoploid cytotypes. In our findings, we have observed a new ploidy level ($2n = 2x = 18$) for two Iranian accessions of *A. millefolium* (Am3, Am4) that were collected in northern parts of Iran.

Karyotypic analysis indicated asymmetrical pattern in the four accessions of *A. millefolium* (Table 2). Mitotic metaphases and karyograms of the four accessions are shown in Figure 1. The highest TCL value was found in Am3 (60.9 μ m) and the lowest was found in Am2 (24.5 μ m) (Table 3). The lowest and the highest DRL values were found in Am3 and Am2 accessions, respectively (Table 3). High DRL value leads to more changes in the construction of chromosomes. DRL values in the two diploid accessions were higher than the tetraploid ones; it can be a relationship between ploidy level and DRL value. The tetraploid accessions had the most symmetric karyotypes.

Other parameters that indicate karyotype asymmetry are total form percentage (TF %; Huziwara 1962) and symmetrical index (S% or S/L%; Battaglia 1955) (Table 2). Group one (Am1 and Am2) had the highest mean value for the symmetrical index (S%=57.5) than the group two (S%= 50) (Table 2). It can be inferred that the group one, as diploids,




Figure 1. A–B Mitotic metaphases (A) and karyograms (B) of four *Achillea millefolium* accessions (Am1–Am4). Bars = 5µm.

Table 2. Karyotype features of four *Achillea millefolium* accessions.

Accession	2n	Ploidy level	TF%	S%	Karyotype formulae
Am1	18	2x	40.9	64	1M+8m
Am2	18	2x	40.8	51	1M+6m+2sm
Am3	36	4x	40.5	63	16M+2sm
Am4	36	4x	41.8	37	17M+1sm
Mean of group1	-	-	40.85	57.5	-
Mean of group2	-	-	41.15	50	-

‡ TF%=total form percentage (sum of short arm lengths of individual/total haploid length of the complement chromosomes), S% -symmetry index (shorter chromosome length / longer chromosome length), karyotype formula (m, median region; sm, submedian region; M, median point).

Table 3. Total chromosome length (TCL), mean chromosome length (MCL), mean arm ratio (MAR), difference of range relative length (DRL), chromosome length range (CLR), Symmetry Classes of Stebbins (SC) of four *Achillea millefolium* accessions.

Accession	TCL(µm)	MCL(µm) (±SD)	MAR(µm)	DRL(µm)	CLR(µm)	SC
Am1	26.4	2.93(±0.13)	0.71	4.92	2.4-3.7	2A
Am2	24.5	2.72(±0.17)	0.69	6.9	1.8-3.5	2A
Am3	60.8	3.37(±0.09)	0.68	2.2	2.6-4.1	1A
Am4	41	2.27(±0.11)	0.73	4.49	1.1-2.9	1B

are the earlier evolutionary form. The average value of the total form percentage (TF%) in the group one and two were 40.85 and 41.15, respectively. The TF% index has frequently been used to explain karyotype asymmetry (Mercado-Ruaro and Delgado-Salinas 1998, Ruas et al. 2000). In terms of the Stebbins’ system, the karyotype of Am1 and Am2 grouped in 2A class, and it can be ancient evolutionary origin of *A. millefolium* species.

Cluster Analysis

Cluster analysis was done based on karyotypic characteristics (TCL, MCL, MAR and DRL) and karyotype asymmetry (TF% and S%) (Figure 2) and agrees with Ebrahim




Figure 2. Dendrogram of four *Achillea millefolium* accessions based on the karyotype characteristics and asymmetry.

et al. (2012). The results of cluster analyses divided the four accessions in two groups (Figure 2); based on ploidy levels. The first group included diploid accessions (Am1 and Am2), while the second group comprised tetraploid accessions (Am3 and Am4). In the dendrogram, distance between diploid accessions is lower than tetraploid accessions that confirm the result of Stebbins' system that both of diploid accessions grouped in 2A class.

Conclusion

The results of the present study illustrated a new ploidy level ($2n = 2x = 18$) in Iranian *Achillea millefolium* accessions. Cluster analysis indicated that accessions can be classified based on ploidy levels.

References

- Azizi M, Chizzola R, Ghani A, Oroojalian F (2010) Composition at different development stages of the essential oil of four *Achillea* species grown in Iran. *Natural product communications* 5(2): 283–290.
- Bala S, Gupta RS (2013) Male meiosis and chromosome number in Asteraceae family from district Kangra of H. P. *International Journal Botanical Research* 3(1): 43–58.
- Balbir S, Kaur P, Kumari N, Kumar R (2012) Review on *Achillea millefolium*, important medicinal plant. *International Journal of Natural Product Science Spl. Issue* 1: 176.
- Barghamadi A, Mehrdad M, Sefidkon F, Yamini Y, Khajeh M (2009) Comparison of the volatiles of *Achillea millefolium* L. obtained by supercritical carbon dioxide extraction and hydrodistillation methods. *Journal of Essential Oil Research* 21: 259–263. doi: 10.1080/10412905.2009.9700164
- Battaglia E (1955) Chromosome morphology and terminology. *Caryologia* 8: 179–187.

- Biste C (1987) Cytotaxonomische Untersuchungen des Formenkreises *Achillea millefolium* (Asteraceae in der DDR. Feddes Repertorium 88: 533–613. doi: 10.1002/fedr.19780880902
- Ebrahim F, Pakniyat H, Arzani A, Rahimmalek M (2012) Karyotype analysis and new chromosome number reports in *Achillea* species. Biologia 67(2): 284–288. doi: 10.2478/s11756-012-0011-3
- Ebrahimi M, Farajpour M, Beigmohammadi M, Ebrahimi M (2012) Genetic relationships among yarrow based on random amplified polymorphic DNA markers. Journal of Biotechnology and Pharmaceutical Research 3(4): 69–73. doi: 10.1016/j.bse.2012.02.017
- Farajpour M, Ebrahimi M, Amiri R, Golzari S, Sanjari S (2012) Assessment of genetic diversity in *Achillea millefolium* accessions from Iran using ISSR marker. Biochemical Systematic and Ecology 37: 73–79.
- Farsi M, Alhoseini-Goreishi J, Jafari E (2000) Cytological studies of some Iranian *Achillea* species. Journal Agricultural Knowledge 11: 18–37. [in Persian]
- Felföldy LJM (1947) Chromosome numbers of certain Hungarian plants. Acta Biologica Hungarica 17: 101–103.
- Graphodatsky AS, Trifonov VA, Stanyon R (2011) The genome diversity and karyotype evolution of mammals. Molecular Cytogenetics 4: 22. doi: 10.1186/1755-8166-4-22
- Guerra M (2012) Cytotaxonomy: the end of childhood. Plant Biosystems 146: 703–710.
- Guo YP, Wang SZ, Vogl C, Ehrendorfer F (2012) Nuclear and plastid haplotypes suggest rapid diploid and polyploidy speciation in the N Hemisphere *Achillea millefolium* complex (Asteraceae). BMC Evolutionary Biology 12: 1–18. doi: 10.1186/1471-2148-12-2
- Huziwara Y (1962) Karyotype analysis in some genera of Compositae. VIII. Further studies on the chromosome of Aster. American Journal of Botany 49: 116–119. doi: 10.2307/2439026
- Khaniki GB (1995) Chromosome number and morphometry in *Achillea* (Anthemideae, compositae). Nucleus 38: 104–111.
- Lavrenko AN, Serditov NP, Ulle LG (1991) Chromosome numbers in some of vascular plants from the Pechoro-Ilychsky Reservation. Botanicheskii Zhurnal 76: 473–476 (in Russian).
- Levin DA (2002) The role of chromosome change in plant evolution. New York: Oxford University Press.
- Mercado-Ruaro P, Delgado-Salinas A (1998) Karyotypic studies on species of *Phaseolus* (Fabaceae: Phaseolinae). American Journal of Botany 85: 1–9. doi: 10.2307/2446547
- Mizianty M, Frey L (1973) Chromosome numbers of some vascular plants in the Western Bieszczady Mts. (South-eastern Poland). Polish Botanical Journal 28: 363–369.
- Mockute D, Judzentiene A (2003) Variability of the essential oils composition of *Achillea millefolium* ssp. *millefolium* growing wild in Lithuania. Biochemical Systematics and Ecology 31: 1033–1045. doi: 10.1016/S0305-1978(03)00066-8
- Peruzzi L, Eroğlu HE (2013) Karyotype asymmetry: again, how to measure and what to measure? Comparative Cytogenetics 7(1): 1–9. doi: 10.3897/compcytogen.v7i1.4431
- Pireh W, Tyrl RJ (1980) Cytogeography of *Achillea millefolium* in Oklahoma and adjacent States. Rhodora 80: 361–367.
- Rechinger KH (1963) Flora Iranica. No. 158. Akademische Druck-Verlagsanstalt, Wien, Austria, 49–71.

- Reeves A (2001) Micromeasure: a new computer program for the collection and analysis of cytogenetic data. *Genome* 44: 439–443. doi: 10.1139/g01-037
- Rohloff J, Skagen EB, Steen AH, Iversen TH (2000) Production of yarrow (*Achillea millefolium* L) in Norway: essential oil content and quality. *Journal of Agricultural and Food Chemistry* 48: 6205–6209. doi: 10.1021/jf000720p
- Ruas PM, Ruas CF, Maffei EMD, Marin-Morales MA (2000) Chromosome studies in the genus *Mikania* (Asteraceae). *Genetics and Molecular Biology* 23: 79–98. doi: 10.1590/S1415-47572000000400042
- Saeidnia S, Yassa N, Rezaeipoor R (2004) Comparative investigation of the essential oils of *Achillea talagonica* Boiss and *A. millefolium*, chemical composition and immunological studies. *Journal of Essential Oil Research* 16: 262–265. doi: 10.1080/10412905.2004.9698716
- Sheidai M, Azanei N, Attar F (2009) New chromosome number and unreduced pollen formation in *Achillea* species (Asteraceae). *Acta Biologica Szegediensis* 53: 39–43.
- Stace CA (2000) Cytology and cytogenetics as a fundamental taxonomic resource for the 20th and 21st centuries. *Taxon* 49: 451–477. doi: 10.2307/1224344
- Stebbins G L (1971) Chromosomal evolution in higher plants. London, UK: Edward Arnold (Publishers) Ltd.
- Ziaei-Nasab M, Hesamzadeh-Hejazi SM, Bihamta MR, Mirza M, Naderi-Shahab MA (2012) Assessment of karyotypical variation among 16 populations of *Thymus daenensis* Celak and *Thymus Kotschyanus* Boiss. Species in Iran. *African Journal of Biotechnology* 11(5): 1028–1036.