

Review

Correspondence

Richard E. Randall
rer@st-andrews.ac.uk

Stephen Goodbourn
s.goodbourn@sgul.ac.uk

Interferons and viruses: an interplay between induction, signalling, antiviral responses and virus countermeasures

Richard E. Randall¹ and Stephen Goodbourn²

¹School of Biology, University of St Andrews, The North Haugh, St Andrews KY16 9ST, UK

²Division of Basic Medical Sciences, St George's, University of London, London SW17 0RE, UK

The interferon (IFN) system is an extremely powerful antiviral response that is capable of controlling most, if not all, virus infections in the absence of adaptive immunity. However, viruses can still replicate and cause disease *in vivo*, because they have some strategy for at least partially circumventing the IFN response. We reviewed this topic in 2000 [Goodbourn, S., Didcock, L. & Randall, R. E. (2000). *J Gen Virol* **81**, 2341–2364] but, since then, a great deal has been discovered about the molecular mechanisms of the IFN response and how different viruses circumvent it. This information is of fundamental interest, but may also have practical application in the design and manufacture of attenuated virus vaccines and the development of novel antiviral drugs. In the first part of this review, we describe how viruses activate the IFN system, how IFNs induce transcription of their target genes and the mechanism of action of IFN-induced proteins with antiviral action. In the second part, we describe how viruses circumvent the IFN response. Here, we reflect upon possible consequences for both the virus and host of the different strategies that viruses have evolved and discuss whether certain viruses have exploited the IFN response to modulate their life cycle (e.g. to establish and maintain persistent/latent infections), whether perturbation of the IFN response by persistent infections can lead to chronic disease, and the importance of the IFN system as a species barrier to virus infections. Lastly, we briefly describe applied aspects that arise from an increase in our knowledge in this area, including vaccine design and manufacture, the development of novel antiviral drugs and the use of IFN-sensitive oncolytic viruses in the treatment of cancer.

Biology of the interferon system

The interferons (IFNs) are a group of secreted cytokines that elicit distinct antiviral effects. They are grouped into three classes called type I, II and III IFNs, according to their amino acid sequence. Type I IFNs (discovered in 1957; Isaacs & Lindenmann, 1957) comprise a large group of molecules; mammals have multiple distinct IFN- α genes (13 in man), one to three IFN- β genes (one in man) and other genes, such as IFN- ω , - ϵ , - τ , - δ and - κ . The IFN- α and - β genes are induced directly in response to viral infection, whereas IFN- ω , - ϵ , - δ and - κ play less well-defined roles, such as regulators of maternal recognition in pregnancy. Thus, rather than use the term 'type I IFN', we will use IFN- α/β when referring to the virally induced cytokines. Although the multigenic nature of IFN- α has been known for over 20 years, the significance of this is still debated – i.e. whether these genes are expressed differentially in distinct cell types, whether they are inducible by different

types of viruses or whether they are functionally specialized (Brideau-Andersen *et al.*, 2007). For the rest of this review, we will not distinguish between IFN- α subtypes. Type III IFNs have been described more recently and comprise IFN- λ 1, - λ 2 and - λ 3, also referred to as IL-29, IL-28A and IL-28B, respectively (reviewed by Ank *et al.*, 2006; Uze & Monneron, 2007). These cytokines are also induced in direct response to viral infection and appear to use the same pathway as the IFN- α/β genes to sense viral infection (Onoguchi *et al.*, 2007). Type II IFN has a single member, also called IFN- γ or 'immune IFN', and is secreted by mitogenically activated T cells and natural killer (NK) cells, rather than in direct response to viral infection, and will not be considered in depth in this review.

IFN- α/β acts through a common heterodimeric receptor (see below), which appears to be expressed ubiquitously, to activate a signal-transduction pathway that triggers the transcription of a diverse set of genes that, in total, establish an antiviral response in target cells. These genes are referred to as IFN-inducible genes or IFN-stimulated genes (ISGs). A subset of ISGs can also be induced directly

(i.e. in an IFN-independent manner) by viral infection, perhaps offering a degree of protection in the primary infected cells, although the dramatic viral sensitivity of IFN- α/β receptor-knockout mice suggests that this is much less effective than the IFN response itself. In addition to the cell-autonomous activities of IFN- α/β , these cytokines modulate the immune system by activating effector-cell function and promoting the development of the acquired immune response. The IFN- α/β system is summarized in Fig. 1.

Type III IFNs are also secreted and bind to receptors on cells (the IL-28 receptor, which comprises a heterodimer of IL10R2 and IFNLR1) and elicit an equivalent antiviral response to IFN- α/β . Like IFN- α/β , type III IFNs can be induced in many cells but, unlike IFN- α/β , the type III IFN receptor shows a limited tissue distribution (Meager *et al.*, 2005; Mennechet & Uze, 2006; Zhou *et al.*, 2007). The role of the type III IFNs remains to be established but, to date, there is little evidence to support an essential role for host survival in response to infection.

Induction of IFN by viruses

In recent years, our understanding of the nature of the signalling processes in response to viral infection has evolved significantly. Studies on synthetic inducers of IFN throughout the late 1960s and 1970s demonstrated that double-stranded RNAs (dsRNAs), especially poly(rI).poly(rC), were extremely efficient inducers (reviewed by De Clercq, 2006). Seminal studies by Marcus and

colleagues in the 1970s and 1980s generated a paradigm in which both RNA and DNA viruses induced IFN through the production of viral dsRNA (Marcus & Sekellick, 1977; Marcus, 1983). Thus, negative-stranded RNA viruses were proposed to generate a dsRNA molecule dependent upon transcription, positive-stranded RNA viruses to generate a dsRNA molecule via replication, and even DNA viruses were proposed to generate dsRNA as a result of convergent transcription.

However, whilst dsRNA, either viral or synthetic, is an efficient inducer of IFN- α/β , it is not the only inducer. There are several ways in which a cell can recognize the presence of an invading micro-organism, and multiple distinct routes by which hosts can recognize viruses and signal the induction of IFN through the recognition of pathogen-associated molecular patterns (PAMPs) by pattern-recognition receptors. The importance of any individual route of induction depends upon the specific virus, the cell being infected or the stage of infection, and requires a lot more investigation. The known routes of activation are discussed in detail below.

Before considering the specifics of each route of induction, it is worth reviewing our knowledge of the downstream components, namely the transcription factors, as these may be common to many of the induction pathways. The best-studied model for IFN- α/β induction is the production of IFN- β in fibroblastoid cells in response to either the synthetic dsRNA poly(rI).poly(rC) or to simple RNA viruses, such as Sendai virus (SeV; see Table 1 for a list




Fig. 1. Overview of the IFN- α/β system. Cells that secrete IFN- α/β have pattern-recognition receptors (PRRs) to detect molecules associated with infection. These molecules include viral nucleic acids such as dsRNA. These PRRs, once stimulated by their appropriate ligands, activate intracellular signalling cascades leading to transcription of IFN- α/β genes. Once secreted, IFN- α/β binds to the IFN- α/β receptor on neighbouring uninfected cells (as well as on the initial infected cell) and activates an intracellular signalling cascade leading to upregulation of several hundred IFN- α/β -responsive genes, many of which have direct or indirect antiviral action. Viruses released from the primary infected cell replicate inefficiently in cells that are in the antiviral state. The image shows a monolayer of cells infected at 0.01 p.f.u. per cell with PIV5 and, 24 h later, the cells were stained with antibody to the viral nucleocapsid protein (virus antigen) and DAPI (4,6-diamidino-2-phenylindole) to stain the nuclei.

Table 1. Virus abbreviations

Virus	Abbreviation
Adenovirus	AdV
African swine fever virus	ASFV
Borna disease virus	BDV
Bovine herpesvirus 1	BHV-1
Bovine respiratory syncytial virus	BRSV
Bovine viral diarrhea virus	BVDV
Bunyamwera virus	BUNV
Camelpox virus	CMLV
Canine distemper virus	CDV
Classical swine fever virus	CSFV
Cowpox virus	CPXV
Crimean Congo hemorrhagic virus	CCHV
Dengue virus	DV
Eastern equine encephalitis virus	EEEV
Ebola virus	EBOV
Ectromelia virus	ECTV
Encephalomyocarditis virus	EMCV
Epstein-Barr virus	EBV
Foot-and-mouth-disease virus	FMDV
Hantavirus	HTNV
Hendra virus	HeV
Hepatitis A virus	HAV
Hepatitis B virus	HBV
Hepatitis C virus	HCV
Herpes simplex virus	HSV
Human cytomegalovirus	HCMV
Human herpesvirus	HHV
Human immunodeficiency virus	HIV
Human papillomavirus	HPV
Human respiratory syncytial virus	HRSV
Japanese encephalitis virus	JEV
Kunjin virus	KUNV
La Crosse virus	LACV
Langat virus	LGTV
Lymphocytic choriomeningitis virus	LCMV
Mapuera virus	MPRV
Measles virus	MeV
Mumps virus	MuV
Murine cytomegalovirus virus	MCMV
Murine polyomavirus	MPyV
Myxoma virus	MYXV
Newcastle disease virus	NDV
Nipah virus	NiV
Parainfluenza virus	PIV
Poliovirus	PV
Rabies virus	RABV
Rift Valley fever virus	RVFV
Rinderpest virus	RPV
Sendai virus	SeV
Severe acute respiratory syndrome coronavirus	SARS Co-V
Simain virus	SV
Sindbis virus	SINV
Theiler's virus	TMEV
Thogoto virus	THOV
Vaccinia virus	VACV
Vesicular stomatitis virus	VSV
West Nile virus	WNV
Yaba-like disease virus	YLDV
Yellow fever virus	YFV

of virus abbreviations) or Newcastle disease virus (NDV). In this case, the induction is at the level of transcription and requires no new cellular protein synthesis, indicating that cells contain all of the required signalling components and transcription factors. In its simplest form, induction of IFN- β requires the activation of nuclear factor kappa B (NF- κ B) and IFN regulatory factor-3 (IRF-3) although, as discussed below, certain other members of the ten-member IRF family (reviewed by Honda & Taniguchi, 2006; Paun & Pitha, 2007) may play a role under some circumstances.

Prior to induction, IRF-3 and NF- κ B are both cytoplasmic. Upon receipt of an appropriate signal (see below), the C terminus of IRF-3 is phosphorylated, causing a conformational change leading to dimerization and the unveiling of a nuclear-localization signal (NLS) (Lin *et al.*, 1998; Dragan *et al.*, 2007; Panne *et al.*, 2007a). Translocated IRF-3 is retained in the nucleus until it is dephosphorylated (Kumar *et al.*, 2000). NF- κ B is held in the cytoplasm by association with the inhibitor molecule named inhibitor of NF- κ B (I κ B). Signals generated during viral infection cause phosphorylation of I κ B and its subsequent ubiquitination and degradation by proteasomes; freed of association with its inhibitor, the NLS of the p65 subunit of NF- κ B becomes accessible and NF- κ B is translocated to the nucleus (reviewed by Hayden & Ghosh, 2004; Wullaert *et al.*, 2006).

Optimal induction of the IFN- β gene also requires binding of a c-jun/ATF-2 heterodimer to the promoter. The IRF-3, NF- κ B and c-jun/ATF-2 complexes assemble on the promoter in a co-operative manner to form the so-called enhanceosome, and formation of this complex is aided by the high-mobility group (HMG) chromatin-associated protein HMGI(Y) (also known as HMGA) (reviewed by Merika & Thanos, 2001). The assembled enhanceosome components aid the recruitment of CREB-binding protein (CBP)/p300 that, in turn, promote the assembly of the basal transcriptional machinery and RNA polymerase II. Although the proposed model for the IFN- β enhanceosome features HMGI(Y) as a structural component, NF- κ B, IRF-3, and c-jun/ATF-2 can each form a stable structure with the promoter without HMGI(Y) (Berkowitz *et al.*, 2002; Panne *et al.*, 2004). Indeed, although HMGI(Y) could aid crystal formation, it could not be positioned within the resolved structure of NF- κ B bound to the IFN- β promoter. A revision of the enhanceosome model thus suggests that any role for HMGI(Y) is limited to enhancing complex assembly through modulation of the DNA structure.

A feature of the enhanceosome model is that each of the transcription factors binds the IFN- β promoter with limited affinity and that cooperativity between factors is required for optimal induction. However, promoter-mapping studies have shown that the individual binding sites from the IFN- β promoter can respond independently to inducers and that a subset of binding sites will confer some degree of response to inducers. The consensus view is that binding of IRF-3 and/or IRF-7 (see below) is indispensable for induction, but that activation of both

NF- κ B and c-jun/ATF-2 may not be essential; indeed, IFN induction has been reported under conditions where NF- κ B or c-jun/ATF-2 are not activated or their binding sites are not required (Goodbourn *et al.*, 1985; Ellis & Goodbourn, 1994; King & Goodbourn, 1994; Peters *et al.*, 2002; Poole *et al.*, 2002). Presumably, the relative contributions that the individual transcription factors make to IFN- β induction will vary depending on cell type and inducer, and there might be alternative enhanceosome structures.

An analysis of IFN- β induction is complicated by the fact that pre-treatment of cells with IFN can sometimes enhance the IFN yield (an effect called priming) and the distinct possibility of positive autoregulation, where IFN produced during the induction cycle can enhance ongoing production by inducing the synthesis of alternative signalling or transcriptional components. The production of IFN during viral infection leads to the induction of at least three transcription factors (IRF-1, IRF-7 and IRF-9) that, under some circumstances, play a role in the induction process (Matsuyama *et al.*, 1993; Kawakami *et al.*, 1995; Sato *et al.*, 1998, 2000; Honda *et al.*, 2005b). The precise role of each of these factors is unclear. IRF-7 expression levels are undetectable or very low in tissues other than plasmacytoid dendritic cells (pDCs – see below) or cells exposed to IFN. Thus, positive-feedback models have been proposed in which direct induction of the ‘primary’ IFN genes (IFN- β and murine IFN- α 4) (Erlandsson *et al.*, 1998; Marie *et al.*, 1998) takes place, utilizing IRF-3 as the IRF family member; the IFN thus produced feeds back onto cells and induces the synthesis of IRF-7, which, in the presence of a continued infection, then enhances the transcription of the primary IFN genes and allows transcription of the ‘secondary’ IFN genes (the remaining IFN- α genes) (Marie *et al.*, 1998; Sato *et al.*, 1998; Prakash *et al.*, 2005). In support of this model, IRF-7 can bind to the IFN- β promoter (Wathelet *et al.*, 1998; Panne *et al.*, 2007b) and can enhance transcription dramatically (Yang *et al.*, 2004). However, in addition to this feedback role, it remains possible that, even at very low levels, IRF-7 may function as a primary transcription factor in IFN- β induction in fibroblasts. Mouse embryonic fibroblasts (MEFs) from both IRF-3^{0/0} and IRF-7^{0/0} mice show impaired induction of IFN- β (Sato *et al.*, 2000; Honda *et al.*, 2005b), indicating that, although neither factor is essential, both appear important in induction in these cells, although it remains uncertain whether the IRF-7 effect is direct or indirect.

Induction of the IFN- α genes is less well understood (reviewed by Civas *et al.*, 2002). In fibroblastoid cells, these genes do not display a primary induction profile, with the exception of murine IFN- α 4. Unlike the IFN- β promoter, the IFN- α genes lack NF- κ B sites in their promoters, but contain several binding sites for members of the IRF family, including some sites that fit poorly to the consensus. The identity of the IRF family member that stimulates IFN- α gene transcription is uncertain, but it is

notable that IRF-7 has a more relaxed DNA-binding specificity for IRF sites than other members of the IRF family (Morin *et al.*, 2002), and there is considerable evidence that IRF-7 stimulates IFN- α gene transcription preferentially (Au *et al.*, 1998; Lin *et al.*, 2000), probably in association with IRF-3 (Morin *et al.*, 2002). The apparent requirement for IRF-7 is consistent with the lack of primary IFN- α gene induction in fibroblasts, where induction would be dependent upon the feedback induction of the IRF-7 gene (Yeow *et al.*, 2000).

Whilst the general applicability of this model remains to be seen, there are clearly many situations where different profiles are observed. In lymphocytes, viral infection leads to the production of IFN- α without the need to pre-produce IFN- β , indicating that these cells are somehow primed constitutively (Hata *et al.*, 2001). This may correlate with the pre-existence of IRF-7 in these cells, although the evidence for this is equivocal. However, in the case of pDCs, there is constitutive IRF-7 expression, and these cells respond to IFN inducers by making massive amounts of IFN- α/β (hence their alternative name of IFN-producing cells or IPCs). Recent work has also suggested a role for IRF-1 in the induction of IFN- β in myeloid-derived dendritic cells (mDCs) in response to the Toll-like receptor 9 (TLR9) ligand CpG (Negishi *et al.*, 2006; Schmitz *et al.*, 2007).

We will next consider the known individual routes of activation in more detail.

(i) Extracellular dsRNA or dsRNA delivered through endosomes. It has been known for over 30 years that many cell lines and cell types respond to the addition of synthetic dsRNA to the culture medium (referred to here as extracellular dsRNA). Given the size of this molecule, it seemed likely that cells either have a surface-bound dsRNA receptor or contain such a receptor in endosomes, which is activated by internalizing dsRNA by endocytosis. As the kinetics of response to extracellular dsRNA can vary widely between cells, it is probable that a dsRNA receptor can be localized in both compartments. The nature of this receptor remained elusive until 2001, when TLR3 was identified as a molecule that permitted signal responses to added dsRNA (Alexopoulou *et al.*, 2001). TLR3 shows a relatively wide tissue distribution, but is expressed at a high level in mDCs, especially the CD4⁻CD8⁺ subset. TLR3 relocalizes from the endoplasmic reticulum to endosomes in mDCs (Johnsen *et al.*, 2006), to lysosomes in bone marrow-derived macrophages (de Bouteiller *et al.*, 2005; Lee *et al.*, 2006) and to both endosomes and the cell surface in fibroblasts (Matsumoto *et al.*, 2002). The location of TLR3 (and, indeed of TLR7 and TLR9 – see below) enables the detection of viral nucleic acid present in the extracellular environment or produced by uncoating or degradation of entering viral particles; these features permit the development of an IFN response without the need for viral replication. Furthermore, under some circumstances, pDCs can respond through TLR3 to the dsRNA presented

in the phagocytosed apoptotic cells derived from viral infection (Schulz *et al.*, 2005).

The importance of TLR3 in antiviral defence has been probed by using knockout mice; TLR3-deficient mice succumb to infection by murine cytomegalovirus (MCMV) due to reduced IFN production (Tabeta *et al.*, 2004), but remain resistant to some other viruses, such as lymphocytic choriomeningitis virus (LCMV), vesicular stomatitis virus (VSV) and reovirus (Edelmann *et al.*, 2004). Engagement of TLR3 by dsRNA triggers a complex signal-transduction pathway (summarized in Fig. 2), starting with the dimerization of TLR3 and its tyrosine phosphorylation (Sarkar *et al.*, 2004) and recruitment of an adaptor called

Toll-interleukin (IL)-1-resistance (TIR) domain-containing adaptor inducing IFN- β (TRIF) (Hoebe *et al.*, 2003; Yamamoto *et al.*, 2003a), as well as phosphatidylinositol 3 (PI3) kinase (Sarkar *et al.*, 2004). Engagement of TRIF signals the activation of both the NF- κ B and the IRF-3 'arms' of the IFN-induction pathway (Jiang *et al.*, 2004). In the NF- κ B 'arm', TRIF activation leads to the recruitment of the tumour necrosis factor (TNF) receptor-associated factor 6 (TRAF6) (Sato *et al.*, 2003; Jiang *et al.*, 2004) and receptor-interacting protein 1 (RIP1) (Meylan *et al.*, 2004; Cusson-Hermance *et al.*, 2005). TRAF6 recruitment and oligomerization activate its lysine 63-linked ubiquitin E3 ligase activity, leading to polyubiquitination of itself and RIP1 (reviewed by Chen, 2005). The polyubiquitin chains




Fig. 2. TLR3-dependent signalling in response to dsRNA. dsRNA, presented to the outside of the cell or presented to endosomes by endocytosis of extracellular dsRNA, uncoating of endocytosed viral particles or degradation of engulfed apoptotic cells, binds to TLR3. Activated TLR3 recruits the adaptor TRIF that, in turn, acts as a scaffold to recruit signalling components that feed into either the IRF-3 or the NF- κ B pathways. NF- κ B activation requires TRAF6 and RIP1 recruitment to TRIF and their co-operation in recruiting the IKK complex and TAK1. TAK1 phosphorylates the IKK β subunit of the IKK complex, leading to its activation and phosphorylation of I κ B. Phosphorylated I κ B is ubiquitinated and subsequently degraded by proteasomes, releasing NF- κ B for migration to the nucleus (green arrow) and assembly on the IFN- β promoter. IRF-3 activation requires recruitment of TRAF3 to TRIF. TRAF3 binds to TANK, which then binds to TBK-1 and/or IKK ϵ , which are activated in an uncharacterized manner and can phosphorylate IRF-3 directly. The related proteins NAP1 and SINTBAD may function in a non-redundant manner at the same level as TANK (indicated as TANK etc.). IRF-7, where present due to the feedback action of IFN, is activated by TBK-1 and IKK ϵ in a similar manner (NB this is distinct from the TLR7- and TLR9-dependent pathway described in Fig. 3). The activated IRFs also migrate to the nucleus (green arrows) and assemble on the IFN- β promoter with NF- κ B and ATF-2/c-jun, leading to the recruitment of co-factors such as CBP/p300 and RNA polymerase II and, ultimately, stimulation of transcription. See text for more details and references.

are recognized by TAK1-binding proteins 2 and 3 (TAB2 and TAB3) (Kanayama *et al.*, 2004), which chaperone transforming growth factor β -activated kinase 1 (TAK1) to the complex (Deng *et al.*, 2000; Wang *et al.*, 2001). Polyubiquitinated RIP1 is recognized by the NF- κ B essential modifier (NEMO) [also known as the γ subunit of the I κ B kinase (IKK) complex; Ea *et al.*, 2006; Li *et al.*, 2006a; Wu *et al.*, 2006] and thus the IKK complex is recruited to the TRIF–RIP1–TRAF6–TAB–TAK1 complex; as a result of this juxtaposition, the IKK β subunit of the IKK complex is phosphorylated directly by TAK1 (Wang *et al.*, 2001), leading to the downstream phosphorylation of I κ B, its subsequent ubiquitination and degradation and the eventual nuclear uptake of NF- κ B. To emphasize the importance of the ubiquitination profile of this complex in innate immune responses, the de-ubiquitination enzyme A20 is required for the termination of TLR-dependent NF- κ B activation, in a manner that requires the removal of ubiquitin from TRAF6 (Boone *et al.*, 2004).

The IRF-3 'arm' is less well understood. Activated TRIF recruits TRAF3, a molecule that is essential for IFN induction in response to TLR stimulation (Hacker *et al.*, 2006; Oganessian *et al.*, 2006). TRAF3 interacts directly with TRAF family member-associated NF- κ B activator (TANK) (Li *et al.*, 2002), a protein that also associates directly with TANK-binding kinase 1 (TBK-1) (Pomerantz & Baltimore, 1999). TBK-1 and the related IKK ϵ (also known as IKKi) are the kinases that interact with and phosphorylate IRF-3 (Fitzgerald *et al.*, 2003; Sharma *et al.*, 2003). The associations offer a means to explain the observed recruitment of TBK-1 to TRIF in response to TLR3 stimulation (Sato *et al.*, 2003). The precise role of TANK remains uncertain, as two related proteins, NF- κ B-activating kinase-associated protein 1 (NAP1) and similar to NAP1 TBK1 adaptor (SINTBAD), have also been implicated in TLR3-mediated responses (Sasai *et al.*, 2005, 2006; Guo & Cheng, 2007; Ryzhakov & Randow, 2007) and appear to have many of the same properties as TANK without apparently being redundant (Ryzhakov & Randow, 2007). It is also interesting to note that TANK interacts specifically with NEMO, suggesting a potential means to co-regulate the IRF-3 and NF- κ B arms of the signalling response (Chariot *et al.*, 2002; Zhao *et al.*, 2007).

(ii) ssRNA delivered through endosomes. During some viral infections, pDCs can make up to half of the circulating IFN (reviewed by Cao & Liu, 2007). One key feature of pDCs is their TLR-expression profile. They are one of the few cell types that express TLR7, and this is expressed exclusively in endosomes. The ligands for TLR7 include immunomodulatory compounds such as imiquimod and R848 (Hemmi *et al.*, 2002; Jurk *et al.*, 2002; Heil *et al.*, 2003) and single-stranded RNA (ssRNA) molecules, either synthetic (Diebold *et al.*, 2004) or derived from viruses such as human immunodeficiency virus (HIV), influenza and VSV (Diebold *et al.*, 2004; Heil

et al., 2004; Lund *et al.*, 2004). TLR7 shows no sequence specificity for the ssRNA, requiring only the presence of several uridines in close proximity (Diebold *et al.*, 2006). Interestingly, RNA modifications such as those found in mammalian tRNA and rRNA suppress the TLR7-agonist properties, suggesting that these modifications may have evolved to enable discrimination between host and pathogen RNA (Kariko *et al.*, 2005).

Thus, pDCs can mount robust IFN responses when exposed to ssRNA viruses. Uniquely, these cells are dependent on TLRs for IFN induction by NDV, with the RNA helicases melanoma differentiation-associated gene-5 (mda-5) and retinoic acid-inducible gene-I (RIG-I) (discussed below) playing no part (Kato *et al.*, 2005). There is a functional diversity within RNA viruses, in that some (such as influenza A virus; Diebold *et al.*, 2004) can activate TLR7 signalling without the need for replication, and presumably do so as a consequence of uncoating and RNA release within the endosome, whilst others (such as VSV and SeV) require replication to activate TLR7 signalling. This observation was puzzling, because the replication would be cytoplasmic and therefore in a different subcellular compartment from TLR7. A recent observation has offered a resolution of this issue by showing that cytoplasmic material (including replicated viruses) can be engulfed by autophagy and the resultant vesicles can fuse back to endosomes to present nucleic acids to TLR7 (Iwasaki, 2007; Lee *et al.*, 2007). It remains to be seen how this system prevents the sensing and response to cellular RNAs.

The mechanism of induction of IFN follows a different profile from dsRNA-induced activation of TLR3, in that ssRNA-activated TLR7 recruits a distinct adaptor called myeloid differentiation factor 88 (MyD88) that, in turn, recruits a complex containing the kinases interleukin-1 receptor-associated kinase 4 (IRAK-4), IRAK-1 and TRAF6 (Fig. 3); there is also an involvement of TRAF3 in this process (Hacker *et al.*, 2006). TRAF6 recruitment can activate NF- κ B through TAK1–TAB2–TAB3 and the canonical IKK complex, and this route is followed for NF- κ B activation in response to TLR7 engagement. However, in contrast to the activation of IRF-3 seen in response to TLR3 engagement (discussed above), the unique presence of significant constitutive levels of IRF-7 in pDCs offers an alternative route for induction of IFN. In these cells, the MyD88–IRAK-1–IRAK-4–TRAF6 complex binds directly to IRF-7 (Honda *et al.*, 2004; Kawai *et al.*, 2004; Uematsu *et al.*, 2005) and TRAF6 uses its ubiquitin E3 ligase function to polyubiquitinate IRF-7 (Kawai *et al.*, 2004). This process also requires RIP1, and IRF-7 interacts preferentially with polyubiquitinated RIP1 (Huye *et al.*, 2007). Recruited IRF-7 is phosphorylated by IRAK-1 in a TBK/IKK ϵ -independent manner and translocates into the nucleus (still in association with MyD88, TRAF6 and IRAK-1), where it can bind to DNA and stimulate transcription. Recent results also suggest a role for IKK α (Hoshino *et al.*, 2006) and osteopontin (Shinohara *et al.*, 2006) in IRF-7 activation.




Fig. 3. TLR7- and TLR9-dependent signalling. In pDCs, ssRNA or CpG DNA is presented to TLR7 or TLR9, respectively, in endosomes by endocytosis of extracellular nucleic acids or uncoating of endocytosed viral particles, or by degradation of engulfed apoptotic cells. TLR7 is also stimulated by viral PAMPs taken into the endosomes from the cytoplasm by autophagy. Activated TLRs recruit the adaptor MyD88 that recruits IRAK-4 and IRAK-1. This complex acts as a scaffold to recruit signalling components that feed into either the IRF-7 or NF- κ B pathways. NF- κ B activation follows a route similar to that described in Fig. 2, although the role for RIP1 remains to be clarified. IRF-7 recruitment to the MyD88 adaptor complex requires polyubiquitination by TRAF6 in a RIP1-dependent manner. IRF-7 is phosphorylated by IRAK-1 and a complex containing IRF-7, MyD88, TRAF6, IRAK-1 and possibly IRAK-4 is released and migrates to the nucleus (green arrows). Here, it assembles on the IFN- β promoter with NF- κ B and other factors, leading to the stimulation of transcription. IRF-7 can also stimulate IFN- α promoters strongly. See text for more details and references.

It should be emphasized that this mode of activation is distinct from the ‘feedback’ activation of IRF-7 discussed above. In the latter situation, the TBK-1 and IKK ϵ kinases can phosphorylate the C terminus of IRF-7 in the same way as IRF-3 in the TLR3 pathway (Caillaud *et al.*, 2005). Recent evidence indicates that IRF-7 activation by this TLR-independent pathway requires the activity of the Fas-associated death domain (FADD) protein (Balachandran *et al.*, 2007). Additionally, IRF-7 can also interact with small/mothers against decapentaplegic 3 (SMAD3) to enhance induction of IFN- β in a TGF- β -dependent manner (Qing *et al.*, 2004).

(iii) DNA delivered through endosomes. Certain peripheral blood mononuclear cells (PBMCs) show responses to engulfed ‘foreign’ DNA and can activate IFN production. In this case, the receptor is TLR9, again localized on the endosomes, and the DNA is recognized as foreign because it is usually unmethylated (CpG), unlike host genomic DNA. The importance of TLR9 in innate

antiviral immunity is illustrated by the susceptibility of TLR9-deficient mice to MCMV, as a result of reduced IFN production (Krug *et al.*, 2004; Tabeta *et al.*, 2004; Delale *et al.*, 2005).

pDCs, mDCs and macrophages each mount responses to TLR9 agonists, but these cell types use distinct signalling pathways. In pDCs, IFN induction requires endosomal retention of the CpG ligand (Honda *et al.*, 2005a) and involves a MyD88-dependent signalling pathway that is identical to the TLR7-mediated activation of IRF-7 and NF- κ B, as discussed above (Fig. 3). However, mDCs are unable to retain CpG ligands in their early endosomes and are unable to use this IRF-7 pathway, although they still activate NF- κ B; rather, they use IRF-1 (Negishi *et al.*, 2006; Schmitz *et al.*, 2007), which is also activated by TLR9-dependent engagement of the MyD88 adaptor and IRAK kinases.

IRF-5 is also involved in the production of IFN by pDCs in response to TLR7 and TLR9 agonists (Barnes *et al.*, 2004;

Schoenemeyer *et al.*, 2005; Takaoka *et al.*, 2005; Yasuda *et al.*, 2007). IRF-5 is not involved in IFN production by fibroblasts, but makes a contribution to serum IFN levels during viral infection (Yanai *et al.*, 2007). TLR7 or TLR9 engagement activates IRF-5 via the MyD88 pathway, as discussed above for IRF-7. Mutations in the IRF-5 promoter or allelic variations in the structure of the IRF-5 protein are associated with a predisposition to systemic lupus erythematosus (reviewed by Graham *et al.*, 2007; Kozyrev *et al.*, 2007; Kyogoku & Tsuchiya, 2007) and arthritis (Sigurdsson *et al.*, 2007).

(iv) Intracellular viral RNA. Cells possess TLR-independent pathways that respond to viral nucleic acids generated in the cytoplasm by viral replication. The activities of two widely expressed RNA helicase molecules (RIG-I and mda-5) are linked to these responses. Although the nature of the activators of RIG-I and mda-5 remain uncertain (see below), RIG-I was identified initially as an essential regulator for poly(rI).poly(rC)-induced signalling in a functional screen (Yoneyama *et al.*, 2004) and binds to poly(rI).poly(rC) *in vitro* (Rothenfusser *et al.*, 2005;

Yoneyama *et al.*, 2005). Ectopic expression of RIG-I enhances poly(rI).poly(rC) responses, and small interfering RNA (siRNA) knockdowns limit IFN- β induction by poly(rI).poly(rC) (Yoneyama *et al.*, 2004, 2005). The related protein mda-5 was identified initially as a binding partner for the IFN-induction antagonist protein of PIV5 (Andrejeva *et al.*, 2004) and has similar properties to RIG-I (Andrejeva *et al.*, 2004; Yoneyama *et al.*, 2005), although it binds less avidly to poly(rI).poly(rC). When activated, these molecules recruit and activate a mitochondrion-associated adaptor called CARD adaptor inducing IFN- β (Cardif)/virus-induced signalling adaptor (VISA)/mitochondrial antiviral signalling protein (MAVS)/IFN- β promoter stimulator protein 1 (IPS-1) (Kawai *et al.*, 2005; Meylan *et al.*, 2005; Seth *et al.*, 2005; Xu *et al.*, 2005; reviewed by Hiscott *et al.*, 2006a; Johnson & Gale, 2006; summarized in Fig. 4). Experiments with knockout mice indicate that this adaptor is essential for responses to both helicases (Kumar *et al.*, 2006; Sun *et al.*, 2006). Strikingly, although these mice have functionally intact TLR-signalling systems, they are extremely susceptible to RNA viruses, indicating that IFN production by the TLR and




Fig. 4. mda-5- and RIG-I-dependent signalling. Viral RNA, generated in the cytoplasm by uncoating, transcription or replication, activates the RNA helicases mda-5 and RIG-I. mda-5 and RIG-I are both activated by dsRNA, whilst RIG-I can also be activated by RNA molecules with 5' triphosphates. Both helicases have N-terminal CARD domains that recruit the adaptor Cardif/VISA/MAVS/IPS-1. This adaptor, in turn, acts as a scaffold to recruit signalling components that feed into either the IRF-3 or the NF- κ B pathways. Although the details of these downstream signalling pathways remain incomplete, for Cardif/VISA/MAVS/IPS-1 activation, they seem very similar to those events described in Fig. 2 downstream of TRIF. The assembly of an enhanceosome complex on the IFN- β promoter is also equivalent to that described in Fig. 2. See text for more details and references.

helicase pathways is not redundant and that IFN production by the helicase pathway is essential for controlling infection by these viruses.

RIG-I, but not mda-5, requires ubiquitination by TRIM25 to interact with Cardif/VISA/MAVS/IPS-1 (Gack *et al.*, 2007), and mda-5, but not RIG-I, interacts with dihydroacetone kinase in a manner that suggests its negative regulation (Diao *et al.*, 2007), although the significance of these findings is unknown. Both RIG-I and mda-5, as well as the Cardif/VISA/MAVS/IPS-1 adaptor, interact with the atg5/atg12 components of the autophagy machinery (Jounai *et al.*, 2007). This interaction inhibits signalling downstream of Cardif/VISA/MAVS/IPS-1. It remains to be seen whether this is a means to ensure that the induction of autophagy and IFN production are mutually exclusive in infected cells.

A third member of the helicase family (LGP2) has also been described; this protein lacks the CARD domains present in mda-5 and RIG-I and appears to function as a negative regulator of IFN production (Rothenfusser *et al.*, 2005; Yoneyama *et al.*, 2005; Saito *et al.*, 2007). However, this simple model has been challenged by the phenotypes of LGP2^{0/0} mice, which show unpredictable antiviral phenotypes (Venkataraman *et al.*, 2007). Engagement of Cardif/VISA/MAVS/IPS-1 leads to independent signalling to both the NF- κ B and IRF-3 'arms' of the IFN-induction pathway, as discussed above for the TLR3 adaptor TRIF. Cardif/VISA/MAVS/IPS-1 interacts directly with TRAF6 (Xu *et al.*, 2005) and probably with TRAF3, and the activation pathways are very similar. Specifically, in the context of intracellular RNA signalling, the IKK component NEMO acts as an essential adaptor both for NF- κ B activation and TBK-1/IRF-3 activation, through its interaction with TANK (Zhao *et al.*, 2007).

The relative roles of mda-5 and RIG-I have been studied intensively to determine whether the helicases are redundant parallel sensors or whether they sense different virus-derived signals. In cell-culture systems, both mda-5 and RIG-I signal responses to transfected poly(rI).poly(rC) (Andrejeva *et al.*, 2004; Yoneyama *et al.*, 2004, 2005; Yamashita *et al.*, 2005; Cardenas *et al.*, 2006; Taima *et al.*, 2006). Furthermore, helicase domain-swap experiments have revealed that there are no detectable differences between mda-5, RIG-I or either chimera in their ability to respond to poly(rI).poly(rC) (Childs *et al.*, 2007). However, studies on knockout mice indicate that mda-5 plays a much more important role than RIG-I in the regulation of total serum IFN- α/β levels in response to injected poly(rI).poly(rC) (Kato *et al.*, 2006), in IFN- β production in poly(rI).poly(rC)-treated dendritic cells or macrophages (Gitlin *et al.*, 2006; Kato *et al.*, 2006) and in poly(rI).poly(rC)-transfected embryo-derived fibroblasts (Kato *et al.*, 2006). Curiously, RIG-I plays a limited role in poly(rI).poly(rC) responses in these systems (Kato *et al.*, 2006).

Despite the ability of both helicases to mediate responses to poly(rI).poly(rC), the two molecules differ in their abilities

to recognize other types of RNA. The observation that RIG-I was much more efficient than mda-5 at responding to dsRNA generated by *in vitro* transcription raised the possibility that the two helicases recognized different types of structures within dsRNA. However, RIG-I can bind and respond to ssRNA molecules that contain a 5' triphosphate (Hornung *et al.*, 2006; Pichlmair *et al.*, 2006). Such a triphosphate moiety is present on the RIG-I-activating *in vitro*-transcribed dsRNA and on many RNA molecules generated during viral replication, but absent from most cellular cytosolic RNAs. For example, cellular mRNAs have 5' cap structures, and RNA polymerase I and most RNA polymerase III transcripts have 5' monophosphates. Specifically, mda-5 was not activated by 5' triphosphate-containing RNAs. The well-documented response of RIG-I to poly(rI).poly(rC) is not explained by these findings, because the bulk of poly(rI).poly(rC) contains little or no 5' triphosphate. Probably, RIG-I can be activated by both 5' triphosphate RNAs and dsRNAs or, alternatively, the 5' triphosphate RNAs examined may contain some secondary structure.

Intriguingly, the magnitude of IFN induction is enhanced by the activation of RNase L (Malathi *et al.*, 2007), an endonuclease that degrades both cellular and viral RNAs to generate short fragments with 3' monophosphates. The induction of IFN by RNase L utilized both mda-5 and RIG-I, and the activation of both was limited by removal of the 3' monophosphates. The authors speculate that the degradation products contain both ssRNA and dsRNA components, but these would both lack 5' triphosphates. Overall, knowledge of the ligands for both helicases remains incomplete, and a full understanding may require the structures of the helicases to be solved.

The above properties offer some explanation for the relative importance of mda-5 and RIG-I in inducing IFNs in response to different RNA viral infections. Experiments on knockout mice show that mda-5 is essential for IFN production in response to picornaviruses (Gitlin *et al.*, 2006; Kato *et al.*, 2006), but is less important in responses to other types of RNA virus, such as influenza A virus, where RIG-I is critical (Kato *et al.*, 2006). These results are consistent with the properties of mda-5 and RIG-I discussed above and the reported status of viral RNA structure generated during the viral infection. Thus, the picornaviruses poliovirus (PV) and encephalomyocarditis virus (EMCV) depend on mda-5 for IFN- β induction and generate detectable levels of dsRNA during infection, whilst influenza A virus and paramyxoviruses do not generate dsRNA and depend on RIG-I (Pichlmair *et al.*, 2006). However, results from cell culture suggest a less clear-cut situation. Ectopic overexpression or gene knockout achieved in cell culture by using siRNA or dominant interfering forms of factors indicate a role for both RIG-I and mda-5 in IFN induction by NDV and SeV (Andrejeva *et al.*, 2004; Yoneyama *et al.*, 2004, 2005; Melchjorsen *et al.*, 2005), whilst mda-5, but not RIG-I, is required for IFN- β induction by measles virus (MeV) (Berghall *et al.*, 2006)

and picornaviruses (Kato *et al.*, 2006). In all of these experiments, it should be remembered that the viruses will often encode inhibitors of IFN induction. For example, most paramyxoviruses (including NDV, SeV and MeV) encode an inhibitor of mda-5 (the V protein) that would help to explain why mda-5 gives the impression of making only a limited contribution to IFN induction. It is also important to stress that different preparations of some viruses might produce distinct types of IFN-inducing signal and, if so, that this might influence the apparent contribution of RIG-I and mda-5 to IFN induction. For example, the optimal induction of IFN by SeV was achieved with a mixture of live virus and defective interfering (DI) particles (Johnston, 1981), due to the presence of copyback DI genomes that could self-anneal and form dsRNA (Strahle *et al.*, 2006). It is probable that, during a viral infection, cells may be exposed to and need to block signals emanating from more than one type of PAMP.

(v) Cytoplasmic DNA. Some mammalian cells, most notably macrophages and DCs, respond to 'foreign' DNA present in the cytoplasm (Ishii *et al.*, 2006; Stetson & Medzhitov, 2006a). Such DNA might be distinguished from self DNA by its lack of methylation, higher A+T content (perhaps on the basis of minor-groove topology) or cytoplasmic localization. There seems a considerable degree of disagreement in the literature about the nature of the foreign DNA-recognition system, for example over the involvement of Cardif/VISA/MAVS/IPS-1 (Stetson & Medzhitov, 2006a; Cheng *et al.*, 2007; Takaoka *et al.*, 2007) or whether NF- κ B is activated (Ishii *et al.*, 2006; Stetson & Medzhitov, 2006a; Cheng *et al.*, 2007). It is possible that there are several distinct novel components, perhaps expressed in a tissue-specific manner. Studies utilizing transfected DNA generally do not investigate whether the DNA remains cytoplasmic, or indeed whether it is transcribed into RNA.

The receptor for cytoplasmic DNA is probably distinct from the TLRs and RIG-I/mda-5, and a candidate receptor molecule called DAI/DLM-1/ZBP1 was identified recently (Takaoka *et al.*, 2007). Consistent with a role in signalling responses to foreign DNA, depletion of DAI/DLM-1/ZBP1 led to a decrease in IFN- β mRNA production during herpes simplex type 1 (HSV-1), but not NDV, infection (Takaoka *et al.*, 2007). The induction of IFN- β by DAI/DLM-1/ZBP1 in response to cytoplasmic DNA involves the activation of TBK1 and IRF-3, but the potential involvement of NF- κ B remains to be resolved.

(vi) Induction of IFN by viral proteins. In addition to viral nucleic acids, several viral envelopes or particles can induce IFN- α/β , although this is not a general feature of viruses. For example, HIV gp120 is a potent inducer of IFN- α/β in PBMCs (Capobianchi *et al.*, 1992), but the best-studied systems in this regard are herpesviruses. In response to exposure to either human cytomegalovirus (HCMV) or HSV-1, human fibroblasts activate IRF-3 complex

formation without new protein synthesis (Navarro *et al.*, 1998; Mossman *et al.*, 2001; Preston *et al.*, 2001). Moreover, for herpesviruses, the activation of ISGs by IRF-3 seems sufficient to establish an antiviral state without NF- κ B activation or IFN production (Mossman *et al.*, 2001). Importantly, although both HCMV and HSV-1 activate IRF-3 rapidly, these viruses encode powerful antagonists of IFN production to counteract activation. Whether viral recognition at the cell surface or viral entry is required to initiate IRF-3 activation is uncertain. However, HSV-1 virions deficient in glycoprotein D (gD), gH, gB or gI were unable to activate IRF-3, suggesting that viral entry is required (Mossman *et al.*, 2001; Preston *et al.*, 2001). Similarly, HCMV-mediated ISG induction was blocked by fusion inhibitors (Netterwald *et al.*, 2004). On the other hand, soluble HSV-1 gD (Ankel *et al.*, 1998) and soluble HCMV gB (Boehme & Compton, 2004) induced IFN- β or IRF-3 activation in some cell types. In contrast to HCMV or HSV-1, Kaposi's sarcoma-associated herpesvirus (KSHV)/human herpesvirus 8 (HHV-8) virions do not activate IRF-3 or IFN- β , despite soluble envelope glycoprotein gpK8.1 doing so (Perry & Compton, 2006), suggesting that HHV-8 has evolved a mechanism to avoid this effect.

In addition to IRF-3 activation, efficient induction of IFN- β usually requires the activation of NF- κ B. At low m.o.i., HSV-1 and HCMV particles do not induce NF- κ B in the absence of viral replication (Paladino *et al.*, 2006), implying the existence of a signalling mechanism that activates IRF-3, but not NF- κ B. Interestingly, at higher m.o.i. with HCMV, NF- κ B activation and IFN- α/β production were observed (Paladino *et al.*, 2006).

Expression of MeV nucleocapsid can activate IRF-3 and cause IFN induction (tenOever *et al.*, 2002), but the generality of this phenomenon remains uncertain. Moreover, the nucleocapsid is an avid RNA-binding protein and could function simply to present RNA to the signalling system.

Macrophages and conventional DCs respond to bacterial pathogens and lipopolysaccharide (LPS) by making IFN. This activation is TLR4-dependent and, as for TLR3, TLR7/8 and TLR9 (discussed above), TLR4 activates IRF-3 in addition to NF- κ B. However, TLR4 is unique among these TLRs in responding entirely to an extracellular signal. Whilst the biological significance of induction of IFN in response to LPS remains unclear, there is evidence that respiratory syncytial virus (RSV) can induce IFN synthesis through TLR4 via recognition of its F protein (Kurt-Jones *et al.*, 2000; Haynes *et al.*, 2001). Furthermore, VSV gG can activate IFN- α/β production through a TLR4-dependent pathway (Georgel *et al.*, 2007).

Although TLR4 utilizes MyD88 to activate NF- κ B, it uses the same adaptor as TLR3 (namely TRIF) to activate IRF-3 (Yamamoto *et al.*, 2002; Oshiumi *et al.*, 2003). Both MyD88 and TRIF are recruited indirectly to TLR4, via the TIRAP/MyD88 adaptor-like (Mal) (Fitzgerald *et al.*,

2001; Horng *et al.*, 2001) and TRIF-related adaptor molecule (TRAM) (Yamamoto *et al.*, 2003b) proteins, respectively.

Signalling responses to IFN

The basic signalling pathway activated in response to IFN- α/β has been characterized in detail and reviewed comprehensively (Platanias, 2005; Fig. 5). Key players in this signalling pathway are the Janus/just another kinase (JAK) family (cytoplasmic tyrosine kinases that are recruited to a range of receptors in response to ligand binding) and the signal transducers and activators of transcription (STATs) (members of a family of transcription factors that are phosphorylated by JAKs and bind to DNA; reviewed by Murray, 2007).

In outline, all type I IFNs binds to a common heterodimeric receptor (composed of the products of the *IFNAR1* and *IFNAR2* genes), inducing oligomerization. Prior to activation, the cytoplasmic tail of IFNAR1 is associated with tyrosine kinase 2 (Tyk2) and the IFNAR2 subunit is associated with the tyrosine kinase JAK1. STAT2 is also bound to IFNAR2 before induction and is associated weakly with STAT1 (Stancato *et al.*, 1996; Precious *et al.*, 2005a; Tang *et al.*, 2007). Ligand-induced dimerization of the receptor causes a conformational change, such that Tyk2 phosphorylates tyrosine 466 on IFNAR1, creating a strong docking site for STAT2. Tyk2 then phosphorylates STAT2 on tyrosine 690, and STAT1 is phosphorylated by JAK1 on tyrosine 701; the phosphorylated STAT1 and STAT2 now form a stable heterodimer. Phosphorylation of STAT1 and its subsequent dimerization with STAT2 create a novel NLS (Banninger & Reich, 2004) and, simultaneously, phosphorylation of STAT2 inactivates the

dominant constitutive nuclear export of STAT2 (Frahm *et al.*, 2006) so that the dimers become transported into and retained in the nucleus until their dephosphorylation (reviewed by Reich & Liu, 2006). The STAT1–STAT2 complex associates with a monomer of IRF-9 to form the ISGF3 heterotrimer that binds to the IFN-stimulated response element (ISRE), present in the promoters of most IFN-responsive genes, and enhances transcription. Until recently, it was thought that the assembly of ISGF3 took place in the nucleus, but a recent study suggested that assembly may be co-ordinated at the receptor (Tang *et al.*, 2007). In response to IFN stimulation, the transcriptional co-factor CBP is recruited to the IFNAR2 chain of the receptor and catalyses IFNAR2 acetylation; the acetylation of IFNAR2 creates a docking site for IRF-9 that, in turn, also gets acetylated, as do the receptor-bound STAT1 and STAT2. Acetylation of IRF-9 is required for DNA binding, and acetylation of the STAT factors may aid ISGF3 complex assembly (Tang *et al.*, 2007).

Signalling in response to type III IFN follows a very similar pattern to that in response to type I IFN (Zhou *et al.*, 2007), and signalling in response to type II IFN also shares some features of these pathways. The IFN- γ receptor is also a heterodimeric glycoprotein, comprising two species of major subunit, IFNGR1 and IFNGR2, that pre-associate weakly in unstimulated cells. The cytoplasmic domains of the IFNGR1 and IFNGR2 receptor subunits are associated with a different set of JAKs, JAK1 and JAK2, respectively. When dimeric IFN- γ binds to the receptor subunits, the signalling pathway is initiated by triggering receptor dimerization. This brings JAK1 and JAK2 into close proximity, resulting in the activation of JAK2 which, in turn, *trans*-phosphorylates JAK1, thereby activating it. Subsequently, the activated JAKs phosphorylate a




Fig. 5. Signalling pathway activated by IFN- α/β . The biological activities of IFN- α/β are initiated by binding to the type I IFN receptor. This leads to the activation of the receptor-associated tyrosine kinases JAK1 and Tyk2, which phosphorylate STAT1 on tyrosine 701 and STAT2 on tyrosine 690. Phosphorylated STAT1 and STAT2 interact strongly with each other by recognizing SH2 domains, and the stable STAT1–STAT2 heterodimer is translocated into the nucleus, where it interacts with the DNA-binding protein IRF-9. The IRF-9–STAT1–STAT2 heterotrimer is called ISGF3 and it binds to a sequence motif (the IFN-stimulated response element or ISRE) in target promoters and brings about transcriptional activation. In addition to the phosphorylation of tyrosine, STAT1 also requires phosphorylation on serine 727 for function. See text for more details and references.

tyrosine-containing region in the C terminus of IFNGR1 (between tyrosine 440 and tyrosine 444), creating a pair of binding sites for STAT1. Two STAT1 molecules then interact via their SH2 domains with IFNGR1 and are phosphorylated at tyrosine 701, resulting in their activation and dissociation from the receptor. The activated STAT1 molecules dimerize through SH2 domain–tyrosine phosphate recognition, forming a STAT1–STAT1 homodimer. This homodimer translocates to the nucleus and binds to the unique element of ISGs, the gamma-activation sequence (GAS), and stimulates transcription. Importantly, this sequence is distinct from the ISRE, and the binding of the STAT1–STAT1 homodimer does not require IRF-9.

Although not a major component in IFN- α/β signalling, STAT1 homodimers can also be activated by IFN- α/β (the so-called AAF complex) and trigger the transcription of some genes (Decker *et al.*, 1991). Furthermore, the STAT1–STAT2 heterodimers formed by IFN- α/β stimulation bind to some promoters and stimulate transcription (Li *et al.*, 1996), but the importance of this in the overall antiviral response is unknown.

Additional signalling pathways that can influence the outcome of IFN-induced transcription are activated by IFN (reviewed by Plataniias, 2005). Transcriptional activation by STAT1, whether activated by IFN- α/β or IFN- γ , also requires phosphorylation on serine 727 for full activity and for mounting a full antiviral response. Several kinases catalyse this phosphorylation, including the protein kinase C isoform PKC- δ , which interacts directly with STAT1 (Uddin *et al.*, 2002; Deb *et al.*, 2003). PKC- δ is activated by the PI3 kinase pathway that is also activated by IFN- α/β (reviewed by Kaur *et al.*, 2005). Serine 727 phosphorylation facilitates the interaction of STAT1 with the basal transcription machinery and with other adaptor proteins. STAT2 is not serine-phosphorylated in response to IFN, but it also binds CBP/p300 (see above) and facilitates interaction with the basal transcriptional machinery (Bhattacharya *et al.*, 1996). Treatment of cells with IFN- α/β also activates the p38 MAP kinase (mitogen-activated protein kinase) pathway, which can have important effects on the antiviral programme, although the molecular basis of this is unknown (reviewed by Katsoulidis *et al.*, 2005).

Whilst the activation of an IRF-9–STAT1–STAT2 heterotrimer is the canonical mode of ISRE activation, there are alternative transcription-factor arrangements that activate ISGs through ISREs, and which may show promoter-specific effects or distinct kinetic profiles (reviewed by van Boxel-Dezaire *et al.*, 2006). Activation of a subset of ISRE-dependent genes by IFN- α/β depends on the phosphorylation of STAT1 on tyrosine 708 by IKK ϵ (which is activated in infected cells) and this phosphorylation may stabilize the association of ISGF3 with these ISREs (tenOever *et al.*, 2007). A heterotrimer containing IRF-9 and two molecules of STAT1 (as opposed to a STAT1–STAT2 pair) is activated by IFN- γ (Bluyssen *et al.*, 1995) and activates several ISRE-containing promoters, including the CXCL10

promoter (Majumder *et al.*, 1998). This type of complex, activated differentially by IFN- γ , may mediate the observed subtle differences in transcript profile induced by IFN- γ and IFN- α/β . STAT2–STAT2 homodimers can form a complex with IRF-9 (Bluyssen & Levy, 1997), although the biological role of this is unclear.

Importantly, many ISGs show either sustained induction in response to IFN or biphasic induction kinetics. These responses are dependent on new protein synthesis and, in many cases, depend on IRF-1, which binds to most, if not all, ISREs. In several cases, IRF-1 interacts with STAT1 for optimal transcriptional activation (e.g. Chatterjee-Kishore *et al.*, 2000a, b). Although IRF-1 is induced by IFN- α/β and IFN- γ , it is more responsive to IFN- γ and this is probably a major reason for the difference in gene expression between the two types of IFN. Notably, all of the components of ISGF3 are themselves IFN-inducible, and the increase in subunit concentration may permit the formation of an ISGF3-like complex that does not require STAT tyrosine phosphorylation. In this context, IRF-9 induction by IFN- β requires IRF-9 and STAT1 without a requirement for STAT1 phosphorylation (Rani & Ransohoff, 2005).

Nature of the IFN-induced antiviral state

Treatment of cells with IFN- α/β upregulates the expression of several hundred genes and, in combination, these specify the antiviral state (Fig. 6). No single gene is pivotal and, for any given virus, a subset of genes is probably required to limit viral replication. Several of the upregulated genes encode enzymes that have been studied intensively and reviewed comprehensively, e.g. dsRNA-dependent protein kinase R (PKR) (Garcia *et al.*, 2006), 2'5'-oligoadenylate synthetase (OAS) (Clemens, 2005; Silverman, 2007) and Mx (Martens & Howard, 2006; Haller *et al.*, 2007).

(i) Protein kinase R (PKR). PKR is synthesized in an inactive form and, in response to the cofactor dsRNA, produced during viral replication, undergoes dimerization and activation. In addition to activation by dsRNA, PKR is activated by a stress-activated protein called protein kinase R (PKR)-activating protein (PACT) (Ito *et al.*, 1999; Patel *et al.*, 2000). The best-characterized substrate for PKR is the α subunit of the eukaryotic translational initiation factor 2 (eIF2 α). Phosphorylation by PKR prevents recycling of eIF2 α such that initiation is halted. Additionally, eIF2 α phosphorylation can activate autophagy, by which the contents of a cell can be degraded (Taloczy *et al.*, 2002; reviewed by Espert *et al.*, 2007) and, for HSV-1, this limits viral replication (Taloczy *et al.*, 2006). As discussed below, the involvement of PKR has been invoked in a number of other antiviral mechanisms, including the induction of apoptosis and cell-cycle arrest.

(ii) 2'5' OAS. Like PKR, OAS is synthesized in an inactive form and utilizes dsRNA as a cofactor. Activation allows

OAS to oligomerize ATP through an unusual 2'5' phosphodiester linkage, and the 2'5'-oligoadenylates bind to and activate RNase L, which degrades cellular and viral RNAs. As discussed above, the OAS/RNase L system has been linked to the induction of IFN and, as discussed below, may also affect apoptosis.

(iii) **Mx.** Mx and the Mx family of genes encode large GTPases related to dynamin; the precise functions of Mx and the superfamily of GTPases are unknown, but they show antiviral activity against a wide range of RNA viruses. The Mx proteins act by recognizing nucleocapsid-like structures and restricting their localization within the cell, thereby restricting virus replication. For example, human MxA recognizes the viral nucleoprotein of Thogoto virus (THOV) and prevents transport of the incoming viral nucleocapsids into the nucleus (Kochs *et al.*, 1998; Weber *et al.*, 2000).

(iv) **ISG15, ISG54 and ISG56.** In addition to PKR, OAS and Mx, other enzymes have important roles in the antiviral response. Early studies showed that several genes were

upregulated dramatically in response to IFN, and these were named according to the molecular mass of the protein (e.g. ISG54 is a 54 kDa protein induced by IFN). Of all the genes upregulated by IFN- α/β , ISG15 is one of the most inducible, and yet little is known about its function (reviewed by Kerscher *et al.*, 2006). ISG15 is related to ubiquitin and, like ubiquitin, it becomes joined covalently to many cellular proteins (Giannakopoulos *et al.*, 2005; Zhao *et al.*, 2005). Interestingly, ISG15 is also released from IFN-treated cells and acts as a cytokine. Evidence for the importance of ISG15 in the antiviral response comes from studies of knockout mice that are deficient in Ubp43 (also called Usp18), a deconjugating enzyme that removes ISG15 from substrates. These mice are hypersensitive to IFN and poly(rI).poly(rC) (Malakhova *et al.*, 2003) and resistant to LCMV and VSV (Ritchie *et al.*, 2004), and MEFs derived from these mice show restricted replication of VSV and Sindbis virus (SINV) (Ritchie *et al.*, 2004). Similarly, siRNA directed against Ubp43 enhances the ability of IFN to inhibit hepatitis C virus (HCV) replication and infectious particle production (Randall *et al.*, 2006). Mice lacking the *ISG15* gene are more susceptible to infection by influenza A and B viruses, SINV




Fig. 6. Biological properties of IFN- α/β . IFN- α/β binds to its receptor and initiates the signalling programme outlined in Fig. 5. The IFN-induced transcripts encode proteins that mediate the antiviral response. Some of these proteins (e.g. PKR and OAS) are enzymes whose activities are dependent upon viral co-factors (e.g. dsRNA) and, when such co-factors are provided, the enzymes can bring about dramatic changes in cellular function (such as translational arrest). Other IFN-inducible factors trigger cell-cycle arrest (e.g. the G₁/S phase-specific cyclin-dependent kinase inhibitor p21) and others promote the presentation of viral antigens to the adaptive immune response (e.g. by upregulating MHC class I and the antigen-processing machinery). IFN- α/β also has immunomodulatory functions, by promoting the maturation of DCs, upregulating the activities of NK cells and CD8⁺ T cells and inducing the synthesis of IL-15, a factor that promotes the division of memory CD8⁺ T cells.

and several herpesviruses although, curiously, these mice show no alteration in sensitivity to VSV infection (Lenschow *et al.*, 2007). It seems likely that ISG15 acts to upregulate the efficiency of the IFN response globally; thus, macrophages from Ubp43^{0/0} mice show an increase in the strength and duration of the IFN-inducible transcription response (Zou *et al.*, 2007) and MEFs from these mice show prolonged IFN-signalling responses (Malakhova *et al.*, 2003). Interestingly, many of the defined substrates for ISG15 modification play important roles in the innate immune response [including JAK1, Erk1 and STAT1 (Malakhov *et al.*, 2003), PKR, MxA and RIG-I (Zhao *et al.*, 2005) and IRF-3 (Lu *et al.*, 2006)], and the consequences of ISG15 conjugation include stabilizing proteins against degradation (Lu *et al.*, 2006) and downregulating cap-dependent translation (Okumura *et al.*, 2007). The importance of ISG15 modification is underscored by the fact that the influenza B virus non-structural protein 1 (NS1) protein interacts with ISG15 and prevents its activation and conjugation to its substrates (Yuan & Krug, 2001).

Two other highly IFN-inducible genes have attracted attention. ISG54 and ISG56 are related proteins that inhibit translation by interacting with the multiprotein translation initiation factor eIF3 complex. Human ISG56 interacts with the eIF3e subunit (Guo *et al.*, 2000) and inhibits its ability to stabilize the eIF2-GTP-tRNAi^{Met} complex (Hui *et al.*, 2003). Curiously, whilst murine ISG56 also targets eIF3, it does not recognize the eIF3e subunit, but rather interacts with the eIF3c subunit, a property that it shares with murine ISG54 (Terenzi *et al.*, 2005). Murine ISG56 inhibits the ability of eIF3 to promote the formation of the eIF2-GTP-tRNAi^{Met}-mRNA-40S ribosomal subunit translational initiation complex (Hui *et al.*, 2005).

(v) Promyelocytic leukaemia (PML) nuclear bodies. The role of PML nuclear bodies in the antiviral response has been of considerable interest recently (reviewed by Everett & Chelbi-Alix, 2007). These bodies are heterogeneous in size and composition, and so are functionally poorly defined, but contain, amongst other proteins, the IFN-inducible protein PML and other IFN-inducible proteins, such as Sp100. They play roles in transcriptional responses to stress and may regulate chromatin structure and promoter accessibility (e.g. Kumar *et al.*, 2007). Overexpression of certain isoforms of PML impairs the replication of both RNA and DNA viruses, and depletion of key PML nuclear-body components enhances the growth of herpesviruses, indicating that the PML nuclear bodies do indeed play a role in restricting viral replication, although the details of their involvement remain to be determined. The best evidence that these functions are antiviral is that some viruses [e.g. HSV and adenovirus (AdV)] encode factors to disrupt these bodies (discussed under 'PML and PML nuclear bodies' in section on 'Viral evasion strategies').

(vi) APOBECs and TRIMs. Extensive studies on cellular genes that restrict the replication of retroviruses have revealed the existence of 'restriction factors'. These factors are members of protein families that control normal cellular functions, and the antiviral family members are evolving rapidly, suggesting that they are under positive selection pressure. The best characterized of these restriction factors are the apolipoprotein B mRNA-editing enzyme-catalytic polypeptide-like (APOBEC) 3F and 3G genes (APOBEC3F and 3G), and the unrelated tripartite motif-5 α (TRIM5 α) gene. Although expressed constitutively (as are many of the 'classical' IFN-response genes, such as PKR), these proteins are upregulated strongly by IFN- α/β in certain cell types (Asaoka *et al.*, 2005; Bonvin *et al.*, 2006; Chen *et al.*, 2006; Peng *et al.*, 2006; Sarkis *et al.*, 2006; Argyris *et al.*, 2007; Sakuma *et al.*, 2007; Ying *et al.*, 2007), indicative of a role in the innate immune response. APOBEC3F and 3G show antiviral activity against a range of retroviruses (reviewed by Malim, 2006; Soros & Greene, 2007) and against hepatitis B virus (HBV) (Baumert *et al.*, 2007). The mechanism of action of these enzymes involves both cytidine deamination and subsequent mutation of template, and inhibition of reverse transcription. Both APOBEC3F and 3G can be incorporated into retroviral virions; the HIV Vif protein prevents this, and also targets these two proteins for degradation by cellular proteasomes. TRIM5 α shows species-specific antiviral activity against retroviruses. For example, TRIM5 α restricts HIV-1 strongly in rhesus monkeys, but only weakly in humans (Hatzioannou *et al.*, 2004; Keckesova *et al.*, 2004; Stremlau *et al.*, 2004; Yap *et al.*, 2004; Song *et al.*, 2005). TRIM5 α interacts with incoming viral capsids and forms a complex in which the virion is rendered non-infectious, at least in part by being targeted for destruction by proteasomes (reviewed by Towers, 2007).

(vii) Adenosine deaminases. Members of another class of RNA-editing enzyme, the adenosine deaminases, also function in antiviral defence (reviewed by Toth *et al.*, 2006). Specifically, the large isoform of 'adenosine deaminase acting on RNA' (ADAR1-L) is cytoplasmic and upregulated by IFN. *In vitro*, this enzyme deaminates adenosines within double-stranded regions of RNA to generate inosines and this activity might thereby mutate viral RNA, although experimental evidence for this is limited. Thus, hypermutation has been observed in the genomes of distinct RNA viruses (reviewed by Cattaneo, 1994; Zahn *et al.*, 2007), whilst the enhanced hepatitis delta virus RNA editing observed in IFN-treated cells is associated with increased ADAR1-L activity (Hartwig *et al.*, 2006).

(viii) Viperin. Recent work has indicated that a protein referred to as viperin (also known as CMV-inducible gene 5, cig5) may play an important role in the IFN-induced antiviral response. Expression of viperin in cell-culture systems demonstrated potent antiviral activity against

HCMV (Chin & Cresswell, 2001), HCV (Helbig *et al.*, 2005), HIV (Rivieccio *et al.*, 2006), alphaviruses (Zhang *et al.*, 2007) and influenza A virus (Wang *et al.*, 2007). The mode of action of viperin is unclear, but it has recently been shown that viperin expression disrupts the formation of lipid rafts (which are important in the budding process of some viruses) on the plasma membrane by binding to and inhibiting an enzyme involved in the biosynthesis of isoprenoid-derived lipids (Wang *et al.*, 2007).

(ix) IFN-inducible microRNAs (miRNAs). Recently, it has been reported that IFN induces the expression of several cellular miRNAs and that some of these have sequence-predicted targets within HCV genomic RNA (Pedersen *et al.*, 2007). Neutralization of these miRNAs with anti-miRNAs reduced the antiviral effects of IFN- β against HCV, whilst the introduction of synthetic miRNA mimics that corresponded to these miRNAs reduced HCV replication, suggesting that IFN-inducible miRNAs may have activity against HCV (and thus potentially other viruses). However, given the high mutation rates of RNA viruses, it might be expected that viruses such as HCV would rapidly evolve resistance to such miRNAs. Furthermore, as it is not clear that HCV has played a significant role in human evolution, it is debatable as to how anti-HCV IFN-inducible miRNAs might have evolved. Nevertheless, this is a potentially important area of research in which progress into the possible roles, and importance, of IFN-induced miRNAs is likely over the coming years.

(x) IFN and apoptosis. Another important function of IFNs is to establish a pro-apoptotic state in target cells (reviewed by Clemens, 2003; Maher *et al.*, 2007); this helps to remove infected cells, especially in circumstances where apoptosis depends on viral co-factors such as dsRNA. The molecular basis for the role of IFN remains unclear and there are several distinct pathways by which IFN can act. For instance, procaspase genes are IFN-inducible (Tamura *et al.*, 1996; Chin *et al.*, 1997; Dai & Krantz, 1999), and both PKR (reviewed by Garcia *et al.*, 2006) and OAS (mediated through the actions of RNase L; Castelli *et al.*, 1998) affect apoptosis and are activated by viral dsRNA. Indeed, several viruses induce apoptosis in a PKR-dependent manner (Balachandran *et al.*, 2000a; Gaddy & Lyles, 2007; Zhang & Samuel, 2007), although this is not universal (Zhang & Samuel, 2007). The apoptotic functions of PKR are sometimes mediated by eIF2 α -dependent pathways (Srivastava *et al.*, 1998; Gil *et al.*, 1999; Scheuner *et al.*, 2006). dsRNA can probably activate a 'classical' apoptosis pathway similar to that activated by TNF. Consistent with this, dsRNA treatment leads to the formation of a complex containing both FADD and procaspase-8 (Iordanov *et al.*, 2005), with subsequent cleavage of procaspase-8 to the active form (Iordanov *et al.*, 2005; Takahashi *et al.*, 2006). The formation of this complex may involve PKR (Balachandran *et al.*, 1998, 2000a). Finally, the apoptosis-activating gene p53 is also IFN-inducible and can be activated indirectly in response

to dsRNA or virus infection to trigger an apoptotic response (Takaoka *et al.*, 2003). Mice lacking PKR are especially sensitive to VSV (Takaoka *et al.*, 2003), whereas mice expressing an extra copy of p53 show increased resistance to VSV, due to enhanced apoptosis (Munoz-Fontela *et al.*, 2005). The mechanism linking viral infection to p53 activation remains obscure, but p53 can be phosphorylated directly by PKR, and p53 activity is enhanced by PKR (Cuddihy *et al.*, 1999a, b). However, it is unlikely that p53-mediated apoptosis is essential for clearing all types of viral infection, and one report shows that p53 actually protects cells against dsRNA-induced apoptosis (Marques *et al.*, 2005).

(xi) IFN and cell-cycle arrest. In addition to establishing an antiviral state, IFNs are often profoundly cytostatic, triggering a growth arrest at the G₁/S transition point in many cell types, and this has led to IFNs being used against a variety of cancers (reviewed by Ferrantini *et al.*, 2007). As discussed above for apoptosis, there are several mechanisms by which this is achieved. The best understood is the IFN-induced upregulation of the G₁/S cyclin-dependent kinase (CDK) inhibitor p21 (also known as pWAF, CIP), and other CDK inhibitors can also be upregulated. IFNs also upregulate the levels of a group of IFN-inducible proteins (the 'p200' family) that affect cell-cycle progression (reviewed by Asefa *et al.*, 2004). The best characterized of this group is p202 which binds to, amongst other targets, the transcription factors pRB, p53 and E2F. It also inhibits E2F-dependent gene expression of S-phase-specific genes, and it seems probable that other family members have similar activities. p53 activation may also play a role because its induction can be associated with either apoptosis or cell-cycle arrest, depending on the extent of DNA damage or the concentration of other cellular factors.

(xii) IFN- α/β in immunomodulation. Although IFN- γ plays a major role in promoting the transition from innate to adaptive immune responses, IFN- α/β is also important in this regard and has profound immunomodulatory activity (reviewed by Le Bon & Tough, 2002). The most obvious example is the upregulation of class I major histocompatibility complex (MHC) molecules and components of the antigen-presenting machinery [for example, transporter associated with antigen processing 1 (TAP-1); Epperson *et al.*, 1992]. This mechanism helps to counter the frequently observed downregulation of class I MHC associated with specific viruses, which is beyond the scope of this review. Downregulation of class I MHC leaves infected cells open to attack by NK cells. IFNs also help to activate NK cells by complex processes including the upregulation of perforin and granzymes (reviewed by Bolitho *et al.*, 2007).

IFN- α/β also has less obvious immunomodulatory functions. Production of IFN by infected cells can promote the maturation of DCs (Le Bon *et al.*, 2003) and, whilst IFN

inhibits the proliferation of naïve CD8⁺ T cells (Petricoin *et al.*, 1997; Sun *et al.*, 1998), it can also sustain the proliferation of antigen-specific CD8⁺ T cells (Tough *et al.*, 1996; Marrack *et al.*, 1999; Honda *et al.*, 2005b; Kolumam *et al.*, 2005; Le Bon *et al.*, 2006). Thus, it seems likely that, once a CD8⁺ T cell has been engaged by an antigen-presenting cell (APC), the production of cytokines by that APC can play a crucial role in stimulating the expansion of the CD8⁺ T-cell clone. During viral infections, the presentation of processed antigens on MHC class I (either through direct cytoplasmic infection of the DC or, more likely, through the degradation of an engulfed infected cell, a process called cross-priming) would be accompanied by the stimulation of TLRs by the engulfed infected cell and the induction of IFN- α/β , hence ensuring the delivery of a high local concentration of cytokine to the selected CD8⁺ T cell. Interestingly, there are specialized types of CD8⁺ effector T cell that require IFN for their expansion (Honda *et al.*, 2005b; Stetson & Medzhitov, 2006b). In addition to this mitogenic role, IFNs upregulate the effector mechanisms of cytotoxic T cells and promote NK-cell maturation. Finally, IFNs can trigger the production of IL-15 by APCs, and this cytokine plays an important role in stimulating the division of memory cytotoxic T cells (reviewed by Surh *et al.*, 2006; Boyman *et al.*, 2007) and the maintenance of NK-cell populations (Nguyen *et al.*, 2002).

Viral evasion strategies

General considerations of how viruses evade the IFN response

There are five main ways by which viruses circumvent the IFN response, namely by: (i) interfering globally with host-cell gene expression and/or protein synthesis; (ii) minimizing IFN induction by limiting the production of viral PAMPs and/or by specifically blocking IFN-induction cascades; (iii) inhibiting IFN signalling; (iv) blocking the action of IFN-induced enzymes with antiviral activity; and (v) having a replication strategy that is (largely) insensitive to the action of IFN. Within each of these strategies, different viruses have evolved a great diversity of molecular mechanisms to achieve similar ends. Furthermore, to circumvent the IFN response, it may be necessary for viruses to use combinations of these strategies. Thus, if a virus blocks IFN signalling but fails to limit IFN production, the virus may not spread rapidly from the infected cell to neighbouring cells, because the latter would be in an IFN-induced antiviral state. Probably because of this, viruses that block IFN signalling often limit IFN production specifically, and some even inactivate specific IFN-induced proteins with antiviral activity, e.g. PKR.

Despite the production of some extremely powerful IFN antagonists, there must be a balance between the ability of viruses to block the IFN response and the cells' ability to produce and respond to IFN. Consistent with this, many wild-type viruses form bigger plaques on cells engineered to be non-responsive to IFN than on the parental cells

(Young *et al.*, 2003). Consequently, until viruses can synthesize enough of the viral IFN antagonist(s) to limit the IFN response, it is also important for viruses to keep production of any potential inducer(s) of IFN to a minimum. This may manifest itself in the mode and kinetics of virus transcription and replication. Indeed, any viral defect, be it in attachment, entry, RNA synthesis or packaging, has the potential to tip the balance of this race against the virus. This implies that, in a normal acute infection, the IFN response probably constitutes a constant selective pressure that keeps viruses maximally fit in terms of replication speed and competence, and in the maintenance of mechanisms to circumvent the IFN response. It also suggests that the efficiency by which a given virus overcomes the IFN response may be extremely important in terms of its pathogenesis and host range (Young *et al.*, 2003).

Viral IFN antagonists are often multifunctional proteins and their different properties may vary in importance at different stages of the virus replication cycle. For example, the NS1 protein of most strains of influenza A virus blocks expression of cellular genes by interfering with stability, processing or export of mRNA from the nucleus (Fortes *et al.*, 1994; Chen *et al.*, 1999; Li *et al.*, 2001b; Kim *et al.*, 2002; Noah *et al.*, 2003; Satterly *et al.*, 2007). However, the NS1 protein of some influenza virus strains also inhibits RIG-I-mediated inhibition of the IFN pathway (Guo *et al.*, 2006; Opitz *et al.*, 2006; Pichlmair *et al.*, 2006; Mibayashi *et al.*, 2007). This may be because there is a window of opportunity for an infected cell to respond to influenza virus infection before the NS1 protein accumulates to sufficient levels to block cellular mRNA processing and export from the nucleus completely. However, influenza viruses may also be able to exploit this window to upregulate specific cellular genes to their own advantage, e.g. by the NS1 protein activating PI3 kinase (Hale *et al.*, 2006; Ehrhardt *et al.*, 2007; Shin *et al.*, 2007). In addition to these properties, NS1 also binds dsRNA (and thus blocks OAS activity) and interacts with PKR to block its activity (all of these activities are described in more detail below). Thus, the relative affinity of NS1 for its cellular targets, its cellular location and the relative abundance of free NS1 will all influence the biological properties of NS1 at any given point in the virus replication cycle. However, it should be noted that although a great many viral IFN antagonists with multiple functions have been identified *in vitro* by using genetic approaches, often there is no formal proof that the supposed targets have relevance *in vivo*. One approach to address such questions is to infect animals with null mutations in host defence genes with (recombinant/mutant) viruses with specific defects in their IFN antagonist(s) in order to provide a formal genetic test for identifying *in vivo* mechanisms and targets of virus virulence (Leib *et al.*, 2000).

Whilst over the last few years we have learnt a great deal about the molecular mechanisms of how different viruses block the IFN response, it is less clear what all the biological

consequences may be, both for the virus and host. Undoubtedly the mechanism employed by a given virus to circumvent the IFN response will be a major factor that will influence the molecular pathogenesis of virus infections. This may be especially true for simple RNA viruses, where most of the virus-encoded proteins will be involved in virus replication rather than in 'luxury' functions.

(i) Interfering with host gene expression and/or protein synthesis globally. Many viruses that cause acute infections have developed a variety of mechanisms to inhibit cellular protein expression globally, including inhibition of cellular gene transcription, mRNA processing or export or cellular protein synthesis (specific examples of how viruses inhibit cellular protein expression are illustrated in Table 2). Whilst many host-cell functions will be affected by general inhibition of cellular gene expression, there is good evidence that one major reason that viruses do this is to circumvent the IFN response. For example, the NSs protein of Bunyamwera virus (BUNV) inhibits cellular mRNA transcription by blocking the activity of RNA polymerase II (Thomas *et al.*, 2004). However, a recombinant BUNV that does not encode a NSs protein is pathogenic in IFN- α/β receptor-knockout mice and grows to high titres in cells that do not produce IFN (Weber *et al.*, 2002; Young *et al.*, 2003). Similarly, the

NSs protein of La Crosse virus (LACV) serves primarily to suppress the IFN response of mammalian hosts (Blakqori *et al.*, 2007). A mutation in the VSV matrix (M) protein (which, in wild-type virus, causes a general inhibition of host-cell transcription and translation) also leads to an attenuated virus with efficient IFN- β -inducing properties (Ferran & Lucas-Lenard, 1997). Similarly, foot-and-mouth-disease virus (FMDV) L proteinase shuts off host-cell protein synthesis, and mutation of this protein generated an attenuated strain that induced elevated levels of IFN- α/β (Chinsangaram *et al.*, 1999).

Whilst there are many potential advantages in having a mechanism that prevents host-cell gene expression in some way, there are also many disadvantages. On the upside, by inhibiting cellular gene expression, the cell will not produce IFN (or other cytokines) or respond to IFN by upregulating the production of IFN-inducible proteins. On the downside, if a virus inhibits host-gene expression, then the cell will die fairly rapidly and thus the virus will have only a limited time to replicate. Also, by inhibiting global expression of cellular proteins, the virus will be unable to manipulate the cell to its own advantage, e.g. by forcing it to enter the cell cycle to provide the enzymes that might be necessary for virus replication. Furthermore, a virus cannot establish a persistent or latent infection in cells in which

Table 2. Examples of viruses that inhibit host-cell gene expression globally

Mode of action	Virus	References
Virus-encoded proteases cleave translation initiation factor eIF4-G, poly(A)-binding protein, TATA box-binding protein (TBP) and other cellular factors, including nucleoporins	PV	Etchison <i>et al.</i> (1982); Clark <i>et al.</i> (1993); Das & Dasgupta (1993); Yalamanchili <i>et al.</i> (1996); Gradi <i>et al.</i> (1998); Joachims <i>et al.</i> (1999); Gustin & Sarnow (2001); Belov <i>et al.</i> (2004)
Virus-encoded proteases cleave translation initiation factors eIF4-G and eIF4A	FMDV	Devaney <i>et al.</i> (1988); Kirchweiger <i>et al.</i> (1994); Chinsangaram <i>et al.</i> (1999); Belsham <i>et al.</i> (2000); Li <i>et al.</i> (2001a); de Los Santos <i>et al.</i> (2006)
M protein inhibits host-cell transcription and translation	VSV	Ferran & Lucas-Lenard (1997); Her <i>et al.</i> (1997); Ahmed & Lyles (1998); Yuan <i>et al.</i> (1998); Connor & Lyles (2002, 2005); Ahmed <i>et al.</i> (2003); Stojdl <i>et al.</i> (2003); Whitlow <i>et al.</i> (2006)
Disrupts nucleocytoplasmic transport and affects translation of capped mRNAs	EMCV	Svitkin <i>et al.</i> (2005); Porter <i>et al.</i> (2006)
L protein interacts with Ran-GTPase to interfere with nucleocytoplasmic trafficking	TMEV	van Pesch <i>et al.</i> (2001); Delhaye <i>et al.</i> (2004); Porter <i>et al.</i> (2006)
Virus-encoded NS1 protein inhibits processing and export of cellular mRNAs	Influenza A virus	Fortes <i>et al.</i> (1994); Chen <i>et al.</i> (1999); Li <i>et al.</i> (2001b); Kim <i>et al.</i> (2002); Noah <i>et al.</i> (2003); Satterly <i>et al.</i> (2007)
Virion host-cell shut off protein [vhs/U(L)41] together with ICP27 interfere with host-cell transcription and mRNA stability	HSV-1	Everly <i>et al.</i> (2002); Murphy <i>et al.</i> (2003); Chee & Roizman (2004); Doepker <i>et al.</i> (2004); Smiley (2004)
Virus-encoded NSs protein inhibits RNA polymerase II (RNAP II) to inhibit host-cell transcription	BUNV	Weber <i>et al.</i> (2002); Thomas <i>et al.</i> (2004)
Virus-encoded NSs protein inhibits host-cell transcription by interacting with the transcription factor TF _{II} H	RVFV	Bouloy <i>et al.</i> (2001); Billecocq <i>et al.</i> (2004); Le May <i>et al.</i> (2004)
Capsid protein, mechanism unknown	EEEV	Aguilar <i>et al.</i> (2007)
nsP2 inhibits cellular transcription (and translation)	SINV	Frolova <i>et al.</i> (2002); Gorchakov <i>et al.</i> (2005); Garmashova <i>et al.</i> (2006)

cellular protein synthesis is inhibited. Cells in which cellular protein synthesis is blocked might also be killed more rapidly by other innate immune responses, including the induction of apoptosis by cytokines such as TNF. It may be for these reasons that other viruses have evolved more subtle means to circumvent the IFN response.

(ii) Minimizing IFN production. As discussed above, different viruses, different virus strains and even different laboratory preparations of the same virus strain can vary considerably in their ability to induce IFN. The amount of IFN induced by a virus is dependent upon a number of factors that include (i) the amount and type of IFN inducer(s) produced by virus infection (this can be influenced heavily by the presence of DI particles), (ii) whether a virus produces a specific inhibitor of the IFN-induction pathways, its mode of action and its kinetics of synthesis, and (iii) the type of cell infected. Furthermore, the source of the virus may influence the induction of IFN. Thus, it has been reported that alphaviruses derived from mosquito cells induce less IFN than viruses derived from mammalian cells and that this might have an impact on the ability of the mosquito-borne virus to transfer to the vertebrate host (Shabman *et al.*, 2007).

Clearly, *in vivo*, it would be expected that viruses would keep the production of viral PAMPs to a minimum. This may be achieved by having a replication strategy that minimizes the production of PAMPs by: (i) tightly controlling virus transcription and replication to minimize the production of PAMPs such as dsRNA [e.g. poxviruses minimize the production of overlapping mRNAs by (a) transcribing genes in blocks in the same direction and (b) having multiple transcriptional terminators in intergenic regions where two early genes are transcribed convergently; reviewed by Smith *et al.*, 1998]; (ii) encapsidating both genomic and antigenomic RNA, in the case of negative-strand RNA viruses; (iii) protecting the 5' end of their mRNA from recognition by RIG-I, e.g. by capping their mRNAs (e.g. paramyxoviruses) or 'cap-snatching' (e.g. influenza viruses), coupling a protein to the 5' end of their RNA (e.g. picornaviruses) or not having 5' triphosphate ends of the genomic RNAs (e.g. hantaviruses); (iv) integrating their genomes into host chromosomes and thereafter using cellular machinery for virus transcription and replication (retroviruses); (v) 'hiding' any viral PAMPs from their cellular receptors (e.g. by positive-strand RNA viruses replicating within intracellular membrane vesicles).

However, it is likely that small amounts of dsRNA are produced as a by-product of the replication of most viruses and, thus, many viruses produce dsRNA-binding proteins that sequester dsRNA. This strategy has the additional advantage that it also minimizes the dsRNA-dependent activation of antiviral gene products such as PKR, OAS and ADAR, as well as blocking dsRNA-induced apoptosis. The reovirus major outer capsid protein sigma3 is a dsRNA-binding protein (Imani & Jacobs, 1988; reviewed by Jacobs & Langland, 1998; Olland *et al.*, 2001), as is the sigma A

protein of avian reovirus (Martinez-Costas *et al.*, 2000) and the NSP3 protein of porcine rotaviruses (Langland *et al.*, 1994). Reovirus strains vary significantly in their abilities to induce IFN- α/β (reviewed by Samuel, 1998), and strain differences in IFN sensitivity have been linked to differences in dsRNA affinity of the sigma3 protein (Bergeron *et al.*, 1998). The E3 protein of vaccinia virus (VACV) (Chang *et al.*, 1992) also binds dsRNA, as does the US11 protein of HSV-1 and the VP35 protein of Ebola virus (EBOV) (Cardenas *et al.*, 2006; Hartman *et al.*, 2006; Feng *et al.*, 2007). However, as with the NS1 protein of influenza A viruses (below), the importance of the dsRNA-binding activity of the VP35 protein of EBOV in blocking IFN induction is not clear (Cardenas *et al.*, 2006; Feng *et al.*, 2007). The nuclear protein BS-MLF1 (SM) of Epstein-Barr virus (EBV) (Poppers *et al.*, 2003) and TRS1 protein of HCMV (Cassady, 2005; Hakki & Geballe, 2005) bind dsRNA, whilst the m142 and m143 proteins of MCMV form a heterodimeric complex that binds dsRNA (Child *et al.*, 2006; Valchanova *et al.*, 2006). Furthermore, even though dsRNA is present in very low amounts in cells infected with negative-strand RNA viruses, including influenza virus (Weber *et al.*, 2006), the NS1 protein of influenza virus has a dsRNA-binding motif (Lu *et al.*, 1995). Whilst the primary function of this domain may be to mop up dsRNA to prevent activation of OAS (Min & Krug, 2006), depending on the virus strain (Hayman *et al.*, 2006; Kochs *et al.*, 2007), it may also play an important role in preventing dsRNA-mediated induction of IFN (Talon *et al.*, 2000a; Wang *et al.*, 2000; Li *et al.*, 2006b). Several other viral IFN antagonists, e.g. the HBV core antigen (Tsu & Schloemer, 1989; Whitten *et al.*, 1991), may block IFN production at the transcriptional level by sequestering dsRNA.

Intriguingly, there are examples of virus proteins that intervene specifically at most points in IFN-induction cascades (Fig. 7; Table 3). Bovine viral diarrhoea virus (BVDV) secretes a glycoprotein termed E^{tns} with dsRNA-binding and RNase activities that blocks the ability of extracellular dsRNA to activate TLR3 (Iqbal *et al.*, 2004). This may play an important role in preventing the induction of IFN by released PAMPs/apoptotic bodies in neighbouring or immune cells, such as pDCs. The NS3/4a protein of HCV cleaves the TLR3 adaptor protein TRIF, thereby blocking TLR3 signalling (Li *et al.*, 2005a). VACV proteins A52 and A46 target multiple TIR proteins, including TRIF, to block TLR3 and TLR4 induction of IFN (Harte *et al.*, 2003; Stack *et al.*, 2005). Other viruses can block TLR3 (and TLR7, 8 and 9, e.g. RSV and MeV; Schlender *et al.*, 2005) signalling by inhibiting downstream intracellular signalling components such as TBK and IRF-3 (below). The need to block TLR signalling within infected cells (including pDCs that may be infected productively or abortively) may reflect a requirement to prevent IFN induction through TLR activation following autophagy-dependent viral recognition (Lee *et al.*, 2007). Furthermore, the need to limit IFN induction by this




Fig. 7. Examples of viral IFN antagonists that block/limit the production of IFN- α/β from virus-infected cells. The signalling scheme presented is a composite of that shown in Figs 2 and 4 and shows IFN- β induction via both TLR3- and RNA helicase-dependent pathways. The sites of intervention by several antagonists are indicated. Note that some antagonists, such as N^{pro} of BVDV, are extremely effective in blocking IFN- α/β induction from a variety of PAMPs, because they target signalling molecules (IRF-3) that are far downstream in the IFN-induction cascades, whilst others, such as the V proteins of paramyxoviruses, act further upstream and may block only one arm of the induction pathway (mda-5). Also note that other IFN antagonists, such as HCV NS3/4a, have more than one cellular target. See text for details.

pathway may also partially explain why HSV-1 has evolved a mechanism to block autophagy specifically (Orvedahl *et al.*, 2007). In contrast, it has been suggested that some RNA viruses may activate components of the autophagic machinery both to promote their replication on cellular membranes (Prentice *et al.*, 2004; Jackson *et al.*, 2005) and to limit IFN production (Jounai *et al.*, 2007). This is because, during autophagy, the autophagy-related Atg family members Atg5 and Atg12 form conjugates that, amongst other functions, prevent the CARD domains of RIG-I/mda-5 from interacting with Cardif/VISA/MAVS/IPS-1, thereby preventing the induction of IFN through activation of RIG-I/mda-5 (Jounai *et al.*, 2007). Thus, given that autophagy appears to be an ancient pathway that plays an important role in innate and adaptive immune resistance to pathogens (Schmid *et al.*, 2006), we will probably soon learn more about how viruses specifically block or usurp this pathway.

Some viruses inhibit intracellular detectors of viral PAMPs. The V protein of most paramyxoviruses interacts with, and inhibits the activity of, mda-5 (Andrejeva *et al.*, 2004), but not RIG-I (Yoneyama *et al.*, 2005; Childs *et al.*, 2007; Komatsu *et al.*, 2007). It has also recently been reported that the C proteins of the paramyxovirus SeV can block RIG-I activation of the IFN-induction cascade (Strahle *et al.*, 2007). However, not all paramyxoviruses encode C proteins, suggesting that it may be more important for some paramyxoviruses, such as parainfluenza virus type 5 (PIV5; previously referred to as simian virus 5, SV5) and mumps virus (MuV), that do not encode C proteins to inhibit the function of mda-5 than RIG-I. However, this conclusion appears to contradict observations made on transgenic RIG-I or mda-5 knockout mice that suggest that the primary route for IFN induction by negative-strand viruses, including paramyxoviruses, is through RIG-I rather than mda-5 (Kato *et al.*, 2006; reviewed by Fujita

Table 3. Examples of viruses that block or limit IFN production specifically by infected cells

Mode of action	Virus	References
Degradation of dsRNA	BVDV (E ^{ms} has RNase activity)	Iqbal <i>et al.</i> (2004)
Sequestration of dsRNA	VACV (E3)	Chang <i>et al.</i> (1992)
	Reoviruses (sigma3/A)	Imani & Jacobs (1988); Martinez-Costas <i>et al.</i> (2000); Olland <i>et al.</i> (2001); reviewed by Jacobs & Langland (1998)
	Rotaviruses (NSP3)	Langland <i>et al.</i> (1994)
	Influenza A virus (NS1)	Lu <i>et al.</i> (1995); Talon <i>et al.</i> (2000a); Wang <i>et al.</i> (2000); Li <i>et al.</i> (2006b)
	HSV (US11)	Poppers <i>et al.</i> (2000); Mohr (2004)
	HCMV (TRS1)	Cassady (2005); Hakki & Geballe (2005)
	MCMV (m142/143)	Child <i>et al.</i> (2006); Valchanova <i>et al.</i> (2006)
	EBOV (VP35)	Cardenas <i>et al.</i> (2006); Hartman <i>et al.</i> (2006); Feng <i>et al.</i> (2007)
	EBV (SM)	Poppers <i>et al.</i> (2003)
	Inhibition of TLR signalling	HCV (NS3/4a)
VACV (A52, A46)		Harte <i>et al.</i> (2003); Stack <i>et al.</i> (2005)
Inhibition of mda-5/RIG-I	Paramyxoviruses (V)	Andrejeva <i>et al.</i> (2004); Yoneyama <i>et al.</i> (2005); Childs <i>et al.</i> (2006); Komatsu <i>et al.</i> (2007)
	Influenza A virus (NS1)	Hornung <i>et al.</i> (2006); Pichlmair <i>et al.</i> (2006); Mibayashi <i>et al.</i> (2007); Guo <i>et al.</i> (2006); Opitz <i>et al.</i> (2006)
Inactivation of Cardif/VISA/MAVS/IPS-1	HCV (NS3/4a)	Li <i>et al.</i> (2005c); Meylan <i>et al.</i> (2005); Cheng <i>et al.</i> (2006); Lin <i>et al.</i> (2006a); Loo <i>et al.</i> (2006)
Interferes with VAK	BDV (P)	Unterstab <i>et al.</i> (2005)
	VACV (N1)	DiPerna <i>et al.</i> (2004)
Inhibition or degradation of IRF-3	RABV (P)	Brzozka <i>et al.</i> (2005)
	HTNV (G1)	Alff <i>et al.</i> (2006)
	HHV-6 (E1)	Jaworska <i>et al.</i> (2007)
	Rotaviruses (NSP1)	Graff <i>et al.</i> (2007)
	BVDV (N ^{pro})	Hilton <i>et al.</i> (2006)
	CSFV (N ^{pro})	Bauhofer <i>et al.</i> (2005, 2007); La Rocca <i>et al.</i> (2005); Ruggli <i>et al.</i> (2005)
	HPV 16 (E6)	Ronco <i>et al.</i> (1998)
	HSV (ICP0)	Lin <i>et al.</i> (2004a); Melroe <i>et al.</i> (2004, 2007)
	BHV-1 (bICP0)	Saira <i>et al.</i> (2007)
	THOV (ML)	Hagmaier <i>et al.</i> (2004); Jennings <i>et al.</i> (2005)
	Bovine RSV	Bossert <i>et al.</i> (2003)
	Human RSV	Spann <i>et al.</i> (2005)
	SARS Co-V	Kopecky-Bromberg <i>et al.</i> (2007)
	Interferes with IKK complex	HPV (E7)
HCV (core)		Joo <i>et al.</i> (2005)
VACV		DiPerna <i>et al.</i> (2004)
AdV (E3)		Friedman & Horwitz (2002)
Inhibition of NF- κ B	ASFV (A238L)	Powell <i>et al.</i> (1996); Revilla <i>et al.</i> (1998); Tait <i>et al.</i> (2000)
	VACV (N1)	Shisler & Jin (2004)
	MYXV (MNF)	Camus-Bouclainville <i>et al.</i> (2004)
Interfere with IFN- β promoter activation	HHV-8 (IRF orthologues)	Offermann (2007)

et al., 2007). Whilst there are a number of potential explanations for these observations, including the fact that the paramyxoviruses used in these studies will have inhibited mda-5 activity (Childs *et al.*, 2007), an important consideration when interpreting such data is that different virus preparations used for experimentation, both in tissue culture and in animal studies, may not produce the same spectrum of IFN inducers as in nature. Indeed, viruses grown in tissue culture are not under the same set of

selective pressures as those encountered during replication *in vivo*, and this may lead to apparent inconsistencies. This is well illustrated by the ease with which DI particles, which are powerful inducers of IFN, are generated in cell culture (Johnston, 1981; Poole *et al.*, 2002; Strahle *et al.*, 2006).

In contrast to the V proteins of paramyxoviruses, the NS1 protein of influenza virus associates, either directly or indirectly, with RIG-I (but not mda-5) to inhibit its activity

(Guo *et al.*, 2006; Opitz *et al.*, 2006; Pichlmair *et al.*, 2006; Mibayashi *et al.*, 2007). As discussed previously, influenza viruses may need to inhibit RIG-I activity because uncapped ssRNAs with a 5' triphosphate group that may activate RIG-I may be produced at early times post-infection, before the virus can inhibit host gene expression.

Further downstream in the intracellular IFN-induction cascade, the HCV NS3/4a protein cleaves the C-terminal mitochondrial anchor of Cardif/VISA/MAVS/IPS-1, thereby disrupting its ability to signal to TBK1 and IKK ϵ (Li *et al.*, 2005c; Meylan *et al.*, 2005; Cheng *et al.*, 2006; Kaukinen *et al.*, 2006; Lin *et al.*, 2006a; Loo *et al.*, 2006). Hepatitis A virus (HAV) inhibits RIG-I activation of the IFN-induction cascade at some point upstream of TBK1/IKK ϵ (Fensterl *et al.*, 2005). Rabies virus (RABV) P protein (Brzozka *et al.*, 2005) and the G1 protein of hantavirus (HTNV) (Alff *et al.*, 2006) inhibit the activation of IRF-3 by TBK-1, as does the borna disease virus (BDV) P protein (Unterstab *et al.*, 2005) and the N1 protein of VACV (DiPerna *et al.*, 2004). N^{PRO} of both BVDV and classical swine fever virus (CSFV) (Bauhofer *et al.*, 2005, 2007; La Rocca *et al.*, 2005; Ruggli *et al.*, 2005) blocks IFN induction by targeting IRF-3 for proteasome-mediated degradation (Hilton *et al.*, 2006), whilst the E6 protein of human papillomavirus (HPV) 16 sequesters IRF-3 (Ronco *et al.*, 1998). Rotavirus non-structural protein 1 (NSP1) also targets IRF-3 for degradation (Graff *et al.*, 2007), as well as IRF-5 and IRF-7 (Barro & Patton, 2007). Similarly, ICP0 of HSV-1 inhibits IRF-3 and IRF-7 (Lin *et al.*, 2004a; Melroe *et al.*, 2004), reportedly by recruiting activated IRF-3 and CBP/p300 to nuclear structures away from host chromatin (Melroe *et al.*, 2007). In contrast, it appears that bovine herpesvirus 1 (BHV-1) bICP0 targets IRF-3 for degradation (Saira *et al.*, 2007). The ML protein of THOV acts as an IFN antagonist by blocking IRF-3 dimerization to inhibit its transcriptional activity (Hagmaier *et al.*, 2004; Jennings *et al.*, 2005), whilst severe acute respiratory syndrome coronavirus (SARS Co-V) appears to have multiple methods to inhibit IFN production (Spiegel *et al.*, 2005; Kopecky-Bromberg *et al.*, 2007). The immediate-early 1 protein of HHV-6 also inhibits IRF-3 function (Jaworska *et al.*, 2007), whilst that of HHV-8 targets IRF-7 for degradation (Yu *et al.*, 2005). HHV-8 also produces IRF orthologues, such as vIRF3, which interacts with IRF7 (Joo *et al.*, 2007), and these interfere with the activation of both the IFN- β promoter (Lin *et al.*, 2001) and IFN-responsive genes (reviewed by Rezaee *et al.*, 2006; Offermann, 2007). A number of other viral IFN antagonists, including the NS1 and NS2 proteins of bovine and human RSV (Bossert *et al.*, 2003; Spann *et al.*, 2005), also interfere with IRF-3 activation, by unknown mechanisms.

As discussed above, NF- κ B is also involved in the activation of the IFN- β promoter (as well as other cytokine genes) and its activity is manipulated by some viruses (reviewed by Hiscott *et al.*, 2006b). However, although most viruses do not normally activate NF- κ B (perhaps because they inhibit upstream targets in the IFN-induction

cascade), a few do suppress NF- κ B function directly. Thus, African swine fever virus (ASFV) encodes an IKK orthologue that inhibits the activity of NF- κ B (Powell *et al.*, 1996; Revilla *et al.*, 1998; Tait *et al.*, 2000). The HPV type 16 E7 protein (Spitkovsky *et al.*, 2002) and the HCV core protein (Joo *et al.*, 2005) target the IKK complex to inhibit NF- κ B signalling. The AdV E3 protein also inhibits the activation of NF- κ B (Friedman & Horwitz, 2002), as do the VACV K1 (Shisler & Jin, 2004) and N1 (DiPerna *et al.*, 2004) proteins, and the MNF protein of myxoma virus (MYXV) (Camus-Bouclainville *et al.*, 2004). However, as NF- κ B, as well as inducing proinflammatory cytokines, also induces anti-apoptotic genes (Li & Verma, 2002; Li *et al.*, 2005b; Karin, 2006), any virus that blocks NF- κ B activation specifically may leave itself susceptible to enhanced induction of apoptosis. Notably, the VACV N1 protein resembles Bcl-2 structurally and, as well as inhibiting NF- κ B activation, blocks apoptosis. Indeed, this latter function may be its more relevant biological function (Cooray *et al.*, 2007).

As well as inhibiting IFN induction, viruses may limit the release of IFN by inhibiting cellular protein secretion, as suggested for PV (Choe *et al.*, 2005). As PV also switches off host-cell protein expression (Table 2), this may reflect the need for PV (like other viruses) to inhibit several cellular processes to inhibit the IFN response efficiently.

(iii) Inhibition of IFN signalling. There are clear advantages to viruses in blocking IFN signalling and, as there are components in common between signalling pathways for type I, II and III IFNs, it is possible for a virus to block multiple pathways simultaneously. Using such strategies, not only would the induction of cellular antiviral enzymes, such as PKR, OAS, Mx, ISG15, PML and Sp100, be inhibited, but also there would be no upregulation of class I MHC molecules within infected cells, making them poorer targets for cytotoxic T lymphocytes (CTLs). Furthermore, virus-infected cells would be resistant to the actions of IFNs, regardless of whether the IFNs were produced by infected cells or by activated leukocytes. As with the IFN-induction cascade, there are examples of virus proteins blocking all aspects of the IFN-signalling cascade, from receptor signalling to the formation and activity of IFN-induced transcription factors (Fig. 8; Table 4). Furthermore, some viruses target multiple points of the IFN-signalling cascade. By blocking IFN signalling, some viruses also dismantle the IFN-induced antiviral state in cells. For example, although initially PIV5 cannot replicate efficiently in cells that have entered an antiviral state (Didcock *et al.*, 1999b; Carlos *et al.*, 2005), the degradation of STAT1 by the virion-associated V protein leads to an eventual decay (24–48 h) of the antiviral state (as cells cannot maintain their antiviral state indefinitely without continuous IFN signalling), thereby facilitating subsequent viral replication (Didcock *et al.*, 1999a). However, even if a virus can eventually dismantle the antiviral state of cells, the delay in virus replication induced by IFN should buy




Fig. 8. Examples of viral IFN antagonists that inhibit IFN- α/β signalling specifically within virus-infected cells. The signalling scheme presented is that shown in Fig. 5. Sites of intervention by several antagonists are indicated. Note that many, but not all, of these IFN antagonists will also inhibit IFN- γ signalling. For example, the V protein of PIV5 targets STAT1 for degradation and thus blocks IFN- α/β , IFN- γ and type III IFN signalling, whilst the V protein of hPIV2 targets STAT2 for degradation (in human cells) and thus does not block IFN- γ signalling. See text for details.

time for the host to mount an acquired immune response to help resolve the infection, and this may be why viruses that block IFN signalling also need to limit IFN production.

The diversity of molecular mechanisms that viruses have evolved to block IFN signalling is enormous. For example, viral IFN- γ -binding proteins, which can bind IFN- γ and thus neutralize it, are secreted by cells infected with MYXV, ectromelia virus (ECTV), cowpox virus (CPXV), camelpox virus (CMLV) and VACV (Upton *et al.*, 1992; Alcamí & Smith, 1995; Mossman *et al.*, 1995a, b). Similarly, VACV and most other orthopoxviruses also encode soluble IFN- α/β -binding proteins (Colamonici *et al.*, 1995; Symons *et al.*, 1995). Furthermore, the secreted vIFN- α/β -binding protein of VACV (B18 of VACV strain Western Reserve) also binds to the surface of infected and uninfected cells and prevents them from entering an antiviral state in response to IFN- α/β (Alcamí *et al.*, 2000). Yaba-like disease virus (YLDV) has a protein (YLDV 136) related to VACV B18, but this protein binds and inhibits both type I and type III IFNs (Huang *et al.*, 2007). The K3 and K5 proteins of HHV-8 downregulate IFNGR1 by targeting it for ubiquitination, endocytosis and degradation

(Li *et al.*, 2007). There are several viruses that interfere directly with IFN-receptor activity. The NS5 protein of Japanese encephalitis virus (JEV) interferes with Tyk2 activation, possibly by activating protein tyrosine phosphatases (Lin *et al.*, 2004b, 2006b). Other flaviviruses, including dengue virus (DV) (Ho *et al.*, 2005) and Langkat virus (LGTV) (Best *et al.*, 2005; Park *et al.*, 2007), also disrupt the IFN-receptor complex. HCMV has been reported to disrupt IFN signalling by (i) decreasing the levels of JAK1 (and IRF-9) by a mechanism involving the proteasome (Miller *et al.*, 1998, 1999), and (ii) by the HCMV immediate-early 1 (IE1) 72 kDa protein sequestering STAT1 and STAT2 (Paulus *et al.*, 2006). The T antigen of murine polyomavirus (MPyV) binds to JAK1, thereby blocking the activation of IFN- α/β - and IFN- γ -signalling pathways (Weihua *et al.*, 1998). HSV-1 gene products occlude IFN signalling at multiple points, including by rapidly reducing the levels of JAK1 and STAT2 (Chee & Roizman, 2004). Some paramyxoviruses, including MeV and MuV, as well as interfering with STAT function (below), have been reported to inhibit JAK kinase signalling by interacting with the scaffolding protein RACK1 (receptor for activated C kinase) (Kubota *et al.*, 2002; Yokota *et al.*, 2003).

Table 4. Examples of viruses that inhibit IFN signalling specifically

Mode of action	Virus	References	
Sequestration of IFN through soluble IFN receptor homologues	VACV, MYXV, ECTV, CPXV, CMLV, YLDV	Upton <i>et al.</i> (1992); Alcami & Smith (1995); Colamonici <i>et al.</i> (1995); Mossman <i>et al.</i> (1995a, b); Symons <i>et al.</i> (1995); Puehler <i>et al.</i> (1998); Alcami <i>et al.</i> (2000); Huang <i>et al.</i> (2007)	
Downregulation of IFN receptor	HHV-8	Li <i>et al.</i> (2007)	
Interference with JAK kinases	JEV	Lin <i>et al.</i> (2004b, 2006b)	
	DV	Ho <i>et al.</i> (2005)	
	LGTV	Best <i>et al.</i> (2005); Park <i>et al.</i> (2007)	
	HCMV	Miller <i>et al.</i> (1998, 1999)	
	MPyV	Weihua <i>et al.</i> (1998)	
	MeV	Yokota <i>et al.</i> (2003)	
	MuV	Croze <i>et al.</i> (2000); Usacheva <i>et al.</i> (2001); Kubota <i>et al.</i> (2002)	
	HSV	Chee & Roizman (2004)	
Interference with STAT activity by: <i>Inducing the degradation/loss of STATs</i>	HPV18	Li <i>et al.</i> (1999)	
	PIV2	Young <i>et al.</i> (2000); Nishio <i>et al.</i> (2001); Parisien <i>et al.</i> (2001); Ulane & Horvath (2002)	
	PIV5	Didcock <i>et al.</i> (1999a); Young <i>et al.</i> (2000); Andrejeva <i>et al.</i> (2002a, b); Parisien <i>et al.</i> (2002a, b); Ulane & Horvath (2002); Precious <i>et al.</i> (2005a, b); Ulane <i>et al.</i> (2005)	
	SV41	Kozuka <i>et al.</i> (2003)	
	MuV	Croze <i>et al.</i> (2000); Kubota <i>et al.</i> (2001); Nishio <i>et al.</i> (2002); Ulane <i>et al.</i> (2003); Kubota <i>et al.</i> (2005)	
	NDV	Huang <i>et al.</i> (2003)	
	RSV	Ramaswamy <i>et al.</i> (2004, 2006); Elliott <i>et al.</i> (2007)	
	SeV	Garcin <i>et al.</i> (2002)	
	HSV	Chee & Roizman (2004)	
	<i>Sequestration of STATs or altering their phosphorylation</i>	NiV	Rodriguez <i>et al.</i> (2002); Park <i>et al.</i> (2003); Butler (2004); Rodriguez & Horvath (2004); Shaw <i>et al.</i> (2004, 2005)
		HeV	Rodriguez <i>et al.</i> (2003)
		MeV	Palosaari <i>et al.</i> (2003); Shaffer <i>et al.</i> (2003); Takeuchi <i>et al.</i> (2003); Ohno <i>et al.</i> (2004)
		RPV	Nanda & Baron (2006)
HCV		Lin <i>et al.</i> (2006c)	
HCMV		Paulus <i>et al.</i> (2006)	
RABV		Brzozka <i>et al.</i> (2006); Vidy <i>et al.</i> (2007)	
SeV		Garcin <i>et al.</i> (1999, 2000, 2001); Kato <i>et al.</i> (2001); Takeuchi <i>et al.</i> (2001); Komatsu <i>et al.</i> (2002); Saito <i>et al.</i> (2002); Garcin <i>et al.</i> (2002, 2003); Gotoh <i>et al.</i> (2003a, b); Young <i>et al.</i> (2000); Marq <i>et al.</i> (2007)	
AdV		Look <i>et al.</i> (1998)	
HBV		Christen <i>et al.</i> (2007)	
VACV		Najarro <i>et al.</i> (2001)	
MPRV		Hagmaier <i>et al.</i> (2007)	
<i>Inhibiting STAT trafficking</i>		EBOV	Reid <i>et al.</i> (2006)
	SARS Co-V	Frieman <i>et al.</i> (2007)	
<i>Interfering with IRF-9</i>	HPV16	Barnard & McMillan (1999); Barnard <i>et al.</i> (2000)	
Induction of SOCS	HSV	Yokota <i>et al.</i> (2004)	
	HCV	Bode <i>et al.</i> (2003)	
Upregulation of protein phosphatase 2A	HBV	Christen <i>et al.</i> (2007)	
Interference with ISG promoter activity	HCV	Duong <i>et al.</i> (2006)	
	HHV-8	Reviewed by Lin <i>et al.</i> (2001); Rezaee <i>et al.</i> (2006); Offermann (2007)	
	HBV	Fernandez <i>et al.</i> (2003)	
Miscellaneous/unknown	AdV	Leonard & Sen (1996)	
	SARS Co-V	Kopecky-Bromberg <i>et al.</i> (2007)	
	DV, KUNV, WNV, YFV	Munoz-Jordan <i>et al.</i> (2003, 2005); Liu <i>et al.</i> (2004, 2005, 2006)	
	HCV	de Lucas <i>et al.</i> (2005); Ciccaglione <i>et al.</i> (2007)	
	HSV	Chee & Roizman (2004)	

STAT and IRF-9 proteins that form part of IFN-inducible ISGF3 and GAF transcription complexes are the targets for inhibition by several viruses. Most paramyxoviruses inhibit STAT function to block IFN signalling. However, the molecular mechanisms by which they achieve this differ. In human cells, the V protein of PIV5 blocks IFN signalling by targeting STAT1 for degradation; parainfluenza virus type 2 (HPIV2) targets STAT2 primarily, whilst MuV targets both STAT1 and STAT3. In contrast, the V and P proteins of Nipah (NiV) and Hendra (HeV) viruses sequester STAT1 and STAT2. SeV also blocks IFN signalling by sequestering STATs, increasing their turnover and altering the pattern of STAT1 phosphorylation, although it is another set of proteins encoded by the P/V/C gene, namely the C proteins, that are responsible for this aspect of IFN antagonism. The RSV NS1 protein also targets STAT2 for degradation (see Table 4 for references; for more extensive reviews of how paramyxoviruses block IFN signalling, see Horvath, 2004; Nagai & Kato, 2004; Conzelmann, 2005; Stock *et al.*, 2005).

The RABV P protein also interacts with STAT1 and STAT2, forming an inactive complex (Brzozka *et al.*, 2006; Vidy *et al.*, 2007). Amongst the reported ways that HCV blocks IFN signalling (see also below), the core protein interacts with STAT1 to inhibit its phosphorylation and interaction with STAT2 (Lin *et al.*, 2006c). However, whether this mechanism also operates *in vivo* is not clear, because treatment of HCV-infected or HCV-transfected cell cultures with IFN potently reduces HCV RNA replication (Frese *et al.*, 2001; Windisch *et al.*, 2005; Kim *et al.*, 2007). In contrast, DV clearly blocks IFN signalling (Munoz-Jordan *et al.*, 2003, 2005) and, indeed, in replicon-containing cells, DV RNA replication is resistant to IFN- α treatment (Jones *et al.*, 2005). The VH1 structural protein of VACV is a phosphatase that dephosphorylates STAT1 and has been reported to block IFN- γ signalling (Najarro *et al.*, 2001). Similarly, Kunjin virus (KUNV) and West Nile virus (WNV) (Liu *et al.*, 2004, 2005, 2006) also circumvent the IFN response at least partially by blocking IFN signalling. The HPV 16 E7 multifunctional protein interacts directly with IRF-9, preventing the formation of ISGF3 and thus the activation of IFN- α/β -inducible genes (Barnard & McMillan, 1999; Barnard *et al.*, 2000). In contrast to interfering directly with STAT1 or STAT2, VP24 of EBOV (Reid *et al.*, 2006) and the ORF6 protein of SARS-CoV (Frieman *et al.*, 2007) interact with karyopherin to inhibit the nuclear import of STATs. As part of its mechanism to circumvent the IFN response, HSV rapidly induces the expression of SOCS3, a cellular inhibitor of JAK/STAT pathways (Yokota *et al.*, 2004), as reportedly does the HCV core protein (Bode *et al.*, 2003). HCV and HBV have also been reported to upregulate the expression of protein phosphatase 2A (PP2A), which can interfere with STAT signalling (Duong *et al.*, 2004; Christen *et al.*, 2007). Furthermore, HBV core/precure proteins interact directly with the *MxA* gene promoter (Fernandez *et al.*, 2003). The AdV E1A protein disrupts transcriptional responses to

IFN- α/β and IFN- γ by decreasing the levels of STAT1 and IRF-9 (Leonard & Sen, 1996), by sequestering the transcriptional co-activator CBP/p300, which binds STAT1 and STAT2 and is involved in transcription responses mediated by these proteins (Bhattacharya *et al.*, 1996; Zhang *et al.*, 1996), and by interacting with STAT1 directly (Look *et al.*, 1998). Furthermore, viruses may limit long-term IFN signalling by targeting other transcription factors. For example, HCV core protein interferes with IRF-1, which, as discussed above, is a secondary transcription factor responsible for the prolonged induction of several IFN-stimulated antiviral and immunomodulatory genes (Ciccaglione *et al.*, 2007).

(iv) Inhibition of IFN-induced antiviral enzymes. Many viruses have evolved specific mechanisms to circumvent indirect IFN-induced antiviral responses, including stimulating cells to enter their cell cycle, inhibiting apoptosis and blocking the activity of CTLs and NK cells. However, here we will only briefly review the mechanisms that viruses have evolved to block IFN-induced enzymes with more direct effects on virus replication.

(a) PKR and OAS/RNase L. Several virus types, including pox-, herpes-, influenza and reoviruses, encode proteins that sequester dsRNA to minimize the activation of IFN-induced antiviral proteins PKR, OAS and ADAR by dsRNA. However, sequestration of dsRNA will not prevent the activation of PKR by PACT (PKR-associated activator), which is activated by different cellular-stress pathways that may be activated by virus infection, and it may be for this reason that some viruses, including influenza virus (Tan & Katze, 1998; Li *et al.*, 2006b), have evolved specific mechanisms to inhibit PKR activity or escape its action downstream. Furthermore, inhibition of PKR may also be important for viral evasion of autophagy (Taloczy *et al.*, 2006; Orvedahl *et al.*, 2007; reviewed by Munz, 2007). The mechanisms that viruses use to inhibit PKR have recently been reviewed extensively by Langland *et al.* (2006) and include the production of: (i) small, highly structured RNA molecules that prevent dimerization of PKR by dsRNA and hence its activation [VA-RNA of AdVs, possibly Epstein-Barr-encoded (early) RNAs, the internal ribosome entry sequence (IRES) of HCV]; (ii) proteins that bind directly to, and inhibit the activity of, PKR (HCV NS5A and E2, influenza virus NS1, HSV-1 US11, EBV SM, HHV-8 vIRF-2, baculovirus PK2, VACV E3); (iii) proteins that are pseudosubstrate competitive inhibitors of PKR (HIV Tat, VACV K3); (iv) proteins that degrade PKR (PV protease); (v) proteins that induce the expression of p58^{IPK} (influenza virus NS1), which is a cellular inhibitor of PKR; (vi) proteins that act indirectly on PKR activity, for example the γ_1 ICP34.5 protein of HSV-1 redirects the cellular protein phosphatase 1 α to dephosphorylate, and hence reactivate, eIF-2 α [as does the MyD116 protein of ASFV (Rivera *et al.*, 2007)]. For references, see Langland *et al.* (2006).

Unlike PKR, OAS lack canonical dsRNA-binding motifs and have a relatively low affinity for dsRNA (Hartmann *et al.*, 2003). Consequently, viral proteins that bind dsRNA (above) are particularly potent inhibitors of OAS, even if they also have relatively low affinity for dsRNA, such as influenza virus NS1 (Min & Krug, 2006). However, a few viruses may have evolved strategies that specifically counteract the antiviral activity of the OAS/RNase L system. For example, HIV-1 (Martinand *et al.*, 1999) and EMCV (Martinand *et al.*, 1998) apparently downregulate RNase L activity by inducing the expression of the cellular RNase L inhibitor (RLI), which antagonizes 2'5'-oligoadenylate binding to RNase L and hence prevents its activation. With regards to HIV, potentially the TAR sequence structural element could also bind and activate OAS and PKR, but this is prevented by the virus Tat protein binding the TAR element (Maitra *et al.*, 1994).

(b) ISG15. As described above, ISG15 is a ubiquitin-like molecule that is induced rapidly by IFN and has antiviral activity (Lenschow *et al.*, 2007). Indeed, overexpression of ISG15 in recombinant SINV attenuated the virus in in IFN- α/β receptor-knockout mice (Lenschow *et al.*, 2005). Although its mechanism(s) of action remain(s) unknown, its importance as an antiviral is suggested because several viruses interfere with its activity. It has been reported (Garcia-Sastre, 2007) that the N-terminal domain of the L protein of Crimean Congo hemorrhagic virus (CCHV) has de-ISGylating and de-ubiquitinating activity, which can restore the pathogenicity of the recombinant SINV that overexpresses ISG15 (above) in IFN- α/β receptor-knockout mice. The influenza B virus NS1 protein also binds to ISG15 and prevents it from interacting with its E1 ligase, thereby inhibiting conjugation of ISG15 to its target proteins (Yuan & Krug, 2001).

(c) PML and PML nuclear bodies. Much remains to be discovered of the role of PML and PML nuclear bodies both in virus replication and in antiviral defence (reviewed by Everett, 2001, 2006). Nevertheless, it is clear that many adenoviruses and herpesviruses encode regulatory proteins that localize to, and disrupt the functions of, PML nuclear bodies (reviewed by Everett & Chelbi-Alix, 2007). For example, to aid replication, the Adv E4orf3 protein reorganizes PML nuclear bodies into thread-like structures (Doucas *et al.*, 1996; Stracker *et al.*, 2005; Hoppe *et al.*, 2006; Ullman *et al.*, 2007); the ICP0 protein of HSV-1 has E3 ligase activity that induces the degradation of PML and SUMO-modified forms of Sp100 (reviewed by Hagglund & Roizman, 2004); EBV nuclear antigen 5 (EBNA5) interacts with Sp100A, but this may be to release PML-associated factors from PML nuclear bodies to aid viral gene expression (Ling *et al.*, 2005); HCMV IE72 induces the loss of SUMO-modified forms of PML to disrupt PML nuclear bodies (Ahn & Hayward, 2000; Kang *et al.*, 2006), and pp71 localizes to PML nuclear bodies to induce the

degradation of hDaxx, which inhibits HCMV IE gene expression (Cantrell & Bresnahan, 2006; Preston & Nicholl, 2006; Saffert & Kalejta, 2006). Even certain RNA viruses that replicate in the cytoplasm may block PML function. For example, PV induces the recruitment of p53, mdm2 and proteasome components to PML nuclear bodies, resulting in the degradation of p53 (Pampin *et al.*, 2006). Although it is not yet clear whether PML has a role in antiviral defence against LCMV and RABV, it may be significant that the LCMV Z protein and the RABV P protein, as well as smaller truncated products, interact with PML to cause a reorganization of PML nuclear bodies (Borden *et al.*, 1998a, b; Djavani *et al.*, 2001; Blondel *et al.*, 2002). HCV core protein can co-localize with PML and p53 in PML nuclear bodies (Herzer *et al.*, 2005), although the relevance of this is debated (Rouille *et al.*, 2006).

(d) APOBECs. As discussed earlier, the cytidine deaminases APOBEC3F and 3G, which are inducible by IFN in certain cell types, have antiviral activity against a broad range of retroviruses. However, HIV (and most other lentiviruses) ameliorate the effects of these enzymes at least partially by encoding a protein, termed vif, that forms a complex with components of cellular E3 ligases to target APOBEC3 proteins for ubiquitination and subsequent degradation by proteasomes (reviewed by Malim, 2006; Soros & Greene, 2007). Although it has been reported that these proteins also have activity against HBV, they are not primarily responsible for the anti-HBV activity of IFNs. Thus, HIV vif, even when used in combination with RNA interference to specific APOBEC3 proteins, does not abrogate the inhibitory effect of IFN on HBV (Jost *et al.*, 2007).

Consequences for the virus and host

(i) Consequences for the virus. It is increasingly clear that viruses have evolved an amazing variety of mechanisms to circumvent the IFN response. However, the knock-on consequences that a particular mechanism may have for both the virus and host are less clear. Intuitively, it would be expected that antagonists that block downstream in the IFN-induction cascade would be more effective than those that block upstream, because the multiple signalling pathways that can activate IFN production converge upon IRF-3 and NF- κ B. For example, although PIV5 blocks mda-5 activity (Andrejeva *et al.*, 2004), it does not block RIG-I-induced responses (Childs *et al.*, 2007) and is therefore much less effective in blocking IFN induction than N^{pro} of BVDV, which targets IRF-3 for degradation (Hilton *et al.*, 2006) and, consequently, blocks IFN induction from a number of convergent pathways. Presumably, PIV5 (and closely related paramyxoviruses) are less concerned with the RIG-I-induction pathway, as they limit the production of the inducer of RIG-I (Dillon & Parks, 2007). Nevertheless, it is of note that the ability of paramyxoviruses to circumvent the IFN response may not be absolute (Young *et al.*, 2003).

Although apparently less efficient, viruses may block upstream in the IFN-induction pathway because there may be negative consequences for viruses in blocking further downstream; for example, it may make the cells more susceptible to other host-defence responses. Indeed, given that the IFN response is a near-universal target for viruses, it would be surprising if, in the context of an infection, the immune system had not evolved ways to recognize and eliminate cells with defects in their IFN response.

Although the IFN response is an extremely powerful response that is critical in helping to control most virus infection, it is a programmed response in that, if stimulated in a given way, it will respond predictably. Consequently, as well as blocking the IFN response specifically, viruses may also have more subtle ways to take advantage of the IFN response by evolving particular lifestyles. For example, to survive in nature, certain simple RNA viruses, such as BVDV and HCV, establish persistent infections in a proportion of infected hosts. These relatively simple RNA viruses do not have the ability to establish latent infections in the way that herpesviruses or retroviruses do. Thus, it is possible that their ability to manipulate, and be manipulated by, the IFN response may be critical for the ability of small RNA viruses to establish persistent infections. Indeed, we have suggested such a scenario for PIV5 (Chatziandreu *et al.*, 2002; Carlos *et al.*, 2005). IFN may also help herpesviruses to establish and maintain latency. HSV-1 latency in neurons is characterized by low-level viral gene expression in some neurons, and HSV-1-specific CD8⁺ T cells in close proximity to the neurons prevent full reactivation by the production of IFN- γ and other factors (Khanna *et al.*, 2004; Decman *et al.*, 2005). As latency is central to the life cycle of HSV, if this model is correct, it could be argued that the virus has evolved a strategy to exploit the predictable CD8⁺ T-cell response and the resulting release of IFN- γ . It has also been reported that EBV latent membrane protein 1 (LMP-1) primes viral latency cells for production of IFN- α/β and this would help to preserve EBV latency in two ways; firstly, by preventing the cells from being superinfected by other viruses, and secondly, by inhibiting EBV replication itself (Zhang *et al.*, 2004; Xu *et al.*, 2006).

Viruses may also make use of IFN-induced antiviral proteins. For example, it is surprising that HIV has evolved such a complicated mechanism for controlling the activity of APOBEC3 proteins when it would seem easier to simply avoid packaging APOBEC3F and 3G into virions. An intriguing possibility is that HIV controls APOBEC activity, rather than eliminating it entirely, to increase virus sequence diversity in a way that is not lethal to the virus (Malim, 2006). Consequently, in the future, we are likely to learn more of the ways in which viruses have evolved lifestyles to take advantage of the programmed IFN response for their own benefit.

(ii) Consequences for the host. As long as a viral antagonist does not interfere with host-cell protein

expression in a gross way, persistently infected cells can survive with defects in their IFN response. However, their normal cellular functions may be affected adversely by the continuous expression of the viral IFN antagonist and this may lead to long-term sequelae. Indeed, certain cellular functions may be controlled by the actions of IFN in the absence of virus infection. For example, the activity of osteoclasts, which are involved in bone reabsorption, is regulated negatively by their release of small amounts of IFN (Takayanagi *et al.*, 2002, 2005). Paget's bone disease, in which there is thinning of the bone, has been linked both to increased osteoclast activity and to persistent paramyxovirus infections (Bender, 2003). As paramyxoviruses interfere with both IFN production and signalling, there is potentially a mechanistic link between these observations. Others have disputed this link (Nuovo *et al.*, 1992; Ooi *et al.*, 2000; Rima *et al.*, 2002; Ralston *et al.*, 2007), but the picture is complicated by the fact that most paramyxoviruses, and many other viruses, block the IFN-signalling pathways, raising the possibility that persistent infections by different viruses may cause the same manifestations of a chronic disease. Indeed, a number of different paramyxoviruses, including MeV, canine distemper virus (CDV), RSV, PIV3 and PIV5, have been reported to infect osteoclasts persistently (Mills *et al.*, 1984; Basle *et al.*, 1985, 1986, 1987; Gordon *et al.*, 1991; Mee *et al.*, 1998; Reddy *et al.*, 1999; Friedrichs *et al.*, 2002; Hoyland *et al.*, 2003). Whether or not paramyxoviruses are related to Paget's bone disease, the possibility that different viruses may cause similar chronic disease manifestations should be considered, especially as this scenario is well established for acute infections, e.g. respiratory and enteric viral diseases.

As well as possible adverse consequences of persistent/latent infections and their effects on the IFN response, perversely, the host may benefit from chronic/latent virus infections and the production of IFN. For example, mice infected latently with herpesviruses are more resistant to bacterial infections than non-infected mice (Barton *et al.*, 2007). The latency-induced protection was not antigen-specific, but rather correlated with the production of IFN- γ and systemic activation of macrophages. Furthermore, as IFNs also induce the production of IL-15, which stimulates memory CTLs (discussed above), chronic virus infections, in which there is continuous production of IFN, may help to maintain immune memory. Thus, it is important to be aware not only of surprising adverse consequences of virus-host relationships, but also of possible benefits.

(iii) IFN response and virus host range. Given that viruses must at least partially circumvent the IFN response, it is not surprising that an inability to do this can restrict host range. There are many examples where this is known to be the case. MYXV infects rabbits but will not replicate efficiently in murine or human cells unless their IFN response is disabled (Wang *et al.*, 2004; Johnston *et al.*, 2005; reviewed by McFadden, 2005). Neither MeV nor PV will replicate efficiently in mouse models unless the IFN

system has been compromised [and, for MeV, only if the appropriate virus receptors are expressed (Mrkic *et al.*, 1998; Ohka *et al.*, 2007; Ohno *et al.*, 2007)]. One restriction on the replication of human RSV in bovine cells, and bovine RSV in human cells, is their inability to circumvent the IFN response efficiently in cells of the other species (Schlender *et al.*, 2000; Bossert & Conzelmann, 2002; Young *et al.*, 2003). The IFN response can limit the replication of avian influenza viruses in human cells and thus contribute to host-range restriction (Hayman *et al.*, 2007). Similarly, PIV5 can not replicate in murine cells because it fails to block the IFN response in these cells (Didcock *et al.*, 1999b; Young *et al.*, 2001); furthermore, whilst PIV5 is non-pathogenic in SCID (severe combined immunodeficient) mice (which cannot make an adaptive immune response; Young *et al.*, 1990; Didcock *et al.*, 1999b), the virus kills STAT1^{0/0} mice (He *et al.*, 2001, 2002), because they cannot mount an effective IFN response. The failure of PIV5 to block IFN signalling in mice is because its V protein does not target STAT1 for proteasome-mediated degradation (Didcock *et al.*, 1999b; Young *et al.*, 2001; Chatziandreu *et al.*, 2002; Parisien *et al.*, 2002a), although only a single amino acid change is required for it to become functional in murine cells (Young *et al.*, 2001). Thus, the ability of a virus to adapt (mutate) so that its antagonist becomes functional in a new host species is unlikely to prove more or less of a barrier than any other virus function that requires a virus protein to interact with a cellular protein.

The failure of a virus to circumvent the IFN response in a particular species could be the result of (i) the viral IFN antagonist not functioning correctly in cells of its non-natural species, (ii) the virus replication cycle being too slow, thereby buying time for the cell to mount an effective antiviral response before the virus can produce enough of its antagonist to block the IFN response, or (iii) the production of more, or different, inducers of IFN, such as dsRNA, by the virus, due to a loss of control of virus transcription and replication. Thus, even if a particular viral IFN antagonist is fully functional, the IFN response may still be critical in controlling the infection, as there will be a race between the ability of a virus to block the IFN response and the host's ability to mount one. For example, one reason that the HTNV Andres strain is pathogenic in humans, whereas the Prospect Hill strain is not, may be because Prospect Hill induces more IFN at early times in infection, rather than the inability of the Prospect Hill strain to downregulate IFN signalling (Spiropoulou *et al.*, 2007).

Conversely, there are many examples where a viral IFN antagonist works in several species that are not the natural host for the virus. For example, soluble IFN-binding proteins released from orthopoxvirus-infected cells bind IFNs from a wide variety of species (Alcami & Smith, 1995; Symons *et al.*, 1995; Liptakova *et al.*, 1997); N^{PTO} of BVDV blocks IFN production in cells from many species that it does not infect naturally, including human, monkey and

canine cells (Hilton *et al.*, 2006; M. C. Galiano & R. E. Randall, unpublished observations); and the V protein of Mapuera virus (MPRV) blocks IFN signalling in human, monkey, equine, porcine and canine cells, amongst others, even though MPRV does not infect these species naturally (Hagmaier *et al.*, 2007). Thus, whilst the IFN response is undoubtedly a major barrier that all viruses must overcome to jump species, it is only one of many potential barriers that define species specificity for viruses.

Vaccines, antiviral drugs and oncolytic viruses

IFN-sensitive viruses as attenuated virus vaccines

As our molecular understanding of how viruses block the IFN response has increased, so new possibilities for the control of virus infections have opened up. Thus, attenuated virus vaccines may be developed by isolating viruses that are unable to circumvent the IFN response. This may be achieved either by using reverse genetics to target known genes that encode viral IFN antagonists or by selecting mutants that are sensitive to IFN. There are a number of examples where such an approach has been, or is being, tried, e.g. influenza virus (Talon *et al.*, 2000b; Ferko *et al.*, 2004; Fernandez-Sesma *et al.*, 2006; Richt *et al.*, 2006), paramyxoviruses (Teng *et al.*, 2000; Valarcher *et al.*, 2003; Bartlett *et al.*, 2006; Van Cleve *et al.*, 2006; Wright *et al.*, 2006), BUNV (Weber *et al.*, 2002), EBOV (Hartman *et al.*, 2006) and flaviruses (Liu *et al.*, 2006). However, there are a number of difficulties in using IFN-sensitive viruses as vaccines. The virus may not be as attenuated as expected or required or, alternatively, if the virus is completely sensitive to IFN, the vaccine candidate may be over-attenuated and thus not immunogenic enough. For example, IFN antagonists are often multifunctional proteins, and thus simple gene deletions may overattenuate the virus. Consequently, a better approach may be to introduce/select for point mutations that knock out the IFN-antagonist function of the protein without affecting its other functions. However, single point mutations raise the possibility that such attenuated viruses may revert relatively quickly to wild-type phenotypes. In addition, such IFN-sensitive viruses may be difficult to grow in culture, because most tissue-culture cells can produce and respond to IFN following infection. Vero cells can be used, as they do not produce IFN due to natural deletion of the IFN genes (Desmyter *et al.*, 1968; Mosca & Pitha, 1986). They have the additional advantage that they are already used in manufacture of virus vaccines. However, not all viruses grow in Vero cells, due to other host-cell restrictions, but in such cases it is possible to engineer cells to express different viral IFN antagonists constitutively, to prevent IFN synthesis or the response to IFN (Young *et al.*, 2003).

Antiviral drugs

Viral IFN antagonists might also be targets for novel antiviral drugs. Excitingly, although viruses have many ways to circumvent the IFN response, certain viruses target

the same cellular protein, raising the possibility of antiviral drugs with a fairly wide spectrum of activity. For example, the V proteins of most paramyxoviruses have a conserved cysteine-rich C-terminal domain that interacts with, and blocks the activity of, mda-5 (Andrejeva *et al.*, 2004; Childs *et al.*, 2007). Therefore, a drug or peptide that inhibits the ability of V proteins to interact with mda-5 may inhibit a wide spectrum of paramyxoviruses, including highly pathogenic zoonotic viruses, such as NiV and HeV. In addition, if antiviral drugs/peptides that bind to the cellular targets of viral IFN antagonists can be developed, it may be more difficult for the viruses to acquire drug resistance.

Although antiviral drugs that block viral circumvention of the IFN response may help to control acute infections, they may be particularly useful in chronic virus diseases. Indeed, one reason that IFN treatment of patients infected chronically with HCV or HBV is not always successful may be because these viruses interfere with IFN signalling (Heim *et al.*, 1999; Blindenbacher *et al.*, 2003; Bode *et al.*, 2003; Duong *et al.*, 2004; Christen *et al.*, 2007). If this is correct (see above), then drugs that prevent viral inhibition of IFN signalling may result in either the natural IFN response clearing the infection or a significant improvement in the success of IFN therapy. Similarly, the ability of the HPV E7 protein to block IFN signalling may be important/necessary for the development of papilloma-virus-derived cancers (Barnard & McMillan, 1999; Barnard *et al.*, 2000; Antonsson *et al.*, 2006). Thus, a drug preventing E7-mediated inhibition of IFN signalling might be effective in treating certain types of papillomavirus-related cancers.

IFN-sensitive oncolytic viruses

There is a great deal of interest in using oncolytic viruses for cancer therapy, particularly viruses unable to circumvent the IFN response in human cells, including NDV (Elankumaran *et al.*, 2006; Krishnamurthy *et al.*, 2006), VSV (Stojdl *et al.*, 2000, 2003; Gaddy & Lyles, 2007; Wollmann *et al.*, 2007), mutants of HSV (Mineta *et al.*, 1995; Hunter *et al.*, 1999) and poxviruses. Such viruses may be useful in treating cancers containing tumour cells deficient in their IFN response (Perou *et al.*, 1999; Pitha, 2000), because these viruses would replicate and lyse the tumour cells, but not normal cells with an intact IFN response.

Concluding remarks

In recent years, there has been a resurgence of interest in how viruses interface with the IFN response. This has been due to (i) an increased appreciation of the importance of innate immunity in controlling virus infections, (ii) the elucidation of the molecular pathways by which IFNs are induced and of the signalling response to IFN, and (iii) a realization that all mammalian viruses must have some

strategy for circumventing the IFN response. Ten years ago, it was not generally appreciated that even small RNA viruses with limited coding capacity could produce proteins whose primary function is to target some aspect of the IFN pathway. Whilst over the coming years we can expect an increase in our understanding of how viruses block the IFN response, it is likely that more attention will be placed on how IFN controls virus infections, the interplay between the IFN response and other host-defence mechanisms, and not only how viruses block the IFN response, but also how they have adapted the programmed IFN response to their own advantage. The application of the knowledge gained from such studies should enable better control of virus infections and other diseases.

Acknowledgements

We thank Kay Childs and John McCauley for their comments on the manuscript. Work in our laboratories is supported by the Wellcome Trust, Biotechnology and Biological Sciences Research Council and Medical Research Council (UK).

References

- Aguilar, P. V., Weaver, S. C. & Basler, C. F. (2007).** Capsid protein of eastern equine encephalitis virus inhibits host cell gene expression. *J Virol* **81**, 3866–3876.
- Ahmed, M. & Lyles, D. S. (1998).** Effect of vesicular stomatitis virus matrix protein on transcription directed by host RNA polymerases I, II, and III. *J Virol* **72**, 8413–8419.
- Ahmed, M., McKenzie, M. O., Puckett, S., Hojnacki, M., Poliquin, L. & Lyles, D. S. (2003).** Ability of the matrix protein of vesicular stomatitis virus to suppress beta interferon gene expression is genetically correlated with the inhibition of host RNA and protein synthesis. *J Virol* **77**, 4646–4657.
- Ahn, J. H. & Hayward, G. S. (2000).** Disruption of PML-associated nuclear bodies by IE1 correlates with efficient early stages of viral gene expression and DNA replication in human cytomegalovirus infection. *Virology* **274**, 39–55.
- Alcami, A. & Smith, G. L. (1995).** Vaccinia, cowpox, and camelpox viruses encode soluble gamma interferon receptors with novel broad species specificity. *J Virol* **69**, 4633–4639.
- Alcami, A., Symons, J. A. & Smith, G. L. (2000).** The vaccinia virus soluble alpha/beta interferon (IFN) receptor binds to the cell surface and protects cells from the antiviral effects of IFN. *J Virol* **74**, 11230–11239.
- Alexopoulou, L., Holt, A. C., Medzhitov, R. & Flavell, R. A. (2001).** Recognition of double-stranded RNA and activation of NF- κ B by Toll-like receptor 3. *Nature* **413**, 732–738.
- Alff, P. J., Gavrilovskaya, I. N., Gorbunova, E., Endriss, K., Chong, Y., Geimonen, E., Sen, N., Reich, N. C. & Mackow, E. R. (2006).** The pathogenic NY-1 hantavirus G1 cytoplasmic tail inhibits RIG-I- and TBK-1-directed interferon responses. *J Virol* **80**, 9676–9686.
- Andrejeva, J., Young, D. F., Goodbourn, S. & Randall, R. E. (2002a).** Degradation of STAT1 and STAT2 by the V proteins of simian virus 5 and human parainfluenza virus type 2, respectively: consequences for virus replication in the presence of alpha/beta and gamma interferons. *J Virol* **76**, 2159–2167.
- Andrejeva, J., Poole, E., Young, D. F., Goodbourn, S. & Randall, R. E. (2002b).** The p127 subunit (DDB1) of the UV-DNA damage repair

- binding protein is essential for the targeted degradation of STAT1 by the V protein of the paramyxovirus simian virus 5. *J Virol* **76**, 11379–11386.
- Andrejeva, J., Childs, K. S., Young, D. F., Carlos, T. S., Stock, N., Goodbourn, S. & Randall, R. E. (2004). The V proteins of paramyxoviruses bind the IFN-inducible RNA helicase, mda-5, and inhibit its activation of the IFN- β promoter. *Proc Natl Acad Sci U S A* **101**, 17264–17269.
- Ank, N., West, H. & Paludan, S. R. (2006). IFN- λ : novel antiviral cytokines. *J Interferon Cytokine Res* **26**, 373–379.
- Ankel, H., Westra, D. F., Welling-Wester, S. & Lebon, P. (1998). Induction of interferon- α by glycoprotein D of herpes simplex virus: a possible role of chemokine receptors. *Virology* **251**, 317–326.
- Antonsson, A., Payne, E., Hengst, K. & McMillan, N. A. (2006). The human papillomavirus type 16 E7 protein binds human interferon regulatory factor-9 via a novel PEST domain required for transformation. *J Interferon Cytokine Res* **26**, 455–461.
- Argyris, E. G., Acheampong, E., Wang, F., Huang, J., Chen, K., Mukhtar, M. & Zhang, H. (2007). The interferon-induced expression of APOBEC3G in human blood-brain barrier exerts a potent intrinsic immunity to block HIV-1 entry to central nervous system. *Virology* **367**, 440–451.
- Asaoka, K., Ikeda, K., Hishinuma, T., Horie-Inoue, K., Takeda, S. & Inoue, S. (2005). A retrovirus restriction factor TRIM5 α is transcriptionally regulated by interferons. *Biochem Biophys Res Commun* **338**, 1950–1956.
- Asefa, B., Klarmann, K. D., Copeland, N. G., Gilbert, D. J., Jenkins, N. A. & Keller, J. R. (2004). The interferon-inducible p200 family of proteins: a perspective on their roles in cell cycle regulation and differentiation. *Blood Cells Mol Dis* **32**, 155–167.
- Au, W. C., Moore, P. A., LaFleur, D. W., Tombal, B. & Pitha, P. M. (1998). Characterization of the interferon regulatory factor-7 and its potential role in the transcription activation of interferon A genes. *J Biol Chem* **273**, 29210–29217.
- Balachandran, S., Kim, C. N., Yeh, W. C., Mak, T. W., Bhalla, K. & Barber, G. N. (1998). Activation of the dsRNA-dependent protein kinase, PKR, induces apoptosis through FADD-mediated death signaling. *EMBO J* **17**, 6888–6902.
- Balachandran, S., Roberts, P. C., Brown, L. E., Truong, H., Pattnaik, A. K., Archer, D. R. & Barber, G. N. (2000a). Essential role for the dsRNA-dependent protein kinase PKR in innate immunity to viral infection. *Immunity* **13**, 129–141.
- Balachandran, S., Roberts, P. C., Kipperman, T., Bhalla, K. N., Compans, R. W., Archer, D. R. & Barber, G. N. (2000b). Alpha/beta interferons potentiate virus-induced apoptosis through activation of the FADD/caspase-8 death signaling pathway. *J Virol* **74**, 1513–1523.
- Balachandran, S., Venkataraman, T., Fisher, P. B. & Barber, G. N. (2007). Fas-associated death domain-containing protein-mediated antiviral innate immune signaling involves the regulation of *Irf7*. *J Immunol* **178**, 2429–2439.
- Banninger, G. & Reich, N. C. (2004). STAT2 nuclear trafficking. *J Biol Chem* **279**, 39199–39206.
- Barnard, P. & McMillan, N. A. (1999). The human papillomavirus E7 oncoprotein abrogates signaling mediated by interferon- α . *Virology* **259**, 305–313.
- Barnard, P., Payne, E. & McMillan, N. A. (2000). The human papillomavirus E7 protein is able to inhibit the antiviral and anti-growth functions of interferon- α . *Virology* **277**, 411–419.
- Barnes, B. J., Richards, J., Mancl, M., Hanash, S., Beretta, L. & Pitha, P. M. (2004). Global and distinct targets of IRF-5 and IRF-7 during innate response to viral infection. *J Biol Chem* **279**, 45194–45207.
- Barro, M. & Patton, J. T. (2007). Rotavirus NSP1 inhibits expression of type I interferon by antagonizing the function of interferon regulatory factors IRF3, IRF5, and IRF7. *J Virol* **81**, 4473–4481.
- Bartlett, E. J., Amaro-Carambot, E., Surman, S. R., Collins, P. L., Murphy, B. R. & Skiadopoulos, M. H. (2006). Introducing point and deletion mutations into the P/C gene of human parainfluenza virus type 1 (HPIV1) by reverse genetics generates attenuated and efficacious vaccine candidates. *Vaccine* **24**, 2674–2684.
- Barton, E. S., White, D. W., Cathelyn, J. S., Brett-McClellan, K. A., Engle, M., Diamond, M. S., Miller, V. L. & Virgin, H. W., IV (2007). Herpesvirus latency confers symbiotic protection from bacterial infection. *Nature* **447**, 326–329.
- Basle, M. F., Russell, W. C., Goswami, K. K., Rebel, A., Giraudon, P., Wild, F. & Filmon, R. (1985). Paramyxovirus antigens in osteoclasts from Paget's bone tissue detected by monoclonal antibodies. *J Gen Virol* **66**, 2103–2110.
- Basle, M. F., Fournier, J. G., Rozenblatt, S., Rebel, A. & Bouteille, M. (1986). Measles virus RNA detected in Paget's disease bone tissue by *in situ* hybridization. *J Gen Virol* **67**, 907–913.
- Basle, M. F., Rebel, A., Fournier, J. G., Russell, W. C. & Malkani, K. (1987). On the trail of paramyxoviruses in Paget's disease of bone. *Clin Orthop Relat Res* **217**, 9–15.
- Bauhofer, O., Summerfield, A., McCullough, K. C. & Ruggli, N. (2005). Role of double-stranded RNA and N^{pro} of classical swine fever virus in the activation of monocyte-derived dendritic cells. *Virology* **343**, 93–105.
- Bauhofer, O., Summerfield, A., Sakoda, Y., Tratschin, J. D., Hofmann, M. A. & Ruggli, N. (2007). N^{pro} of classical swine fever virus interacts with interferon regulatory factor 3 and induces its proteasomal degradation. *J Virol* **81**, 3087–3096.
- Baumert, T. F., Rosler, C., Malim, M. H. & von Weizsacker, F. (2007). Hepatitis B virus DNA is subject to extensive editing by the human deaminase APOBEC3C. *Hepatology* **46**, 682–689.
- Belov, G. A., Lidsky, P. V., Mikitas, O. V., Egger, D., Lukyanov, K. A., Bienz, K. & Agol, V. I. (2004). Bidirectional increase in permeability of nuclear envelope upon poliovirus infection and accompanying alterations of nuclear pores. *J Virol* **78**, 10166–10177.
- Belsham, G. J., McInerney, G. M. & Ross-Smith, N. (2000). Foot-and-mouth disease virus 3C protease induces cleavage of translation initiation factors eIF4A and eIF4G within infected cells. *J Virol* **74**, 272–280.
- Bender, I. B. (2003). Paget's disease. *J Endod* **29**, 720–723.
- Bergeron, J., Mabrouk, T., Garzon, S. & Lemay, G. (1998). Characterization of the thermosensitive ts453 reovirus mutant: increased dsRNA binding of $\sigma 3$ protein correlates with interferon resistance. *Virology* **246**, 199–210.
- Berghall, H., Siren, J., Sarkar, D., Julkunen, I., Fisher, P. B., Vainionpaa, R. & Matikainen, S. (2006). The interferon-inducible RNA helicase, mda-5, is involved in measles virus-induced expression of antiviral cytokines. *Microbes Infect* **8**, 2138–2144.
- Berkowitz, B., Huang, D. B., Chen-Park, F. E., Sigler, P. B. & Ghosh, G. (2002). The x-ray crystal structure of the NF- κ B p50.p65 heterodimer bound to the interferon β - κ B site. *J Biol Chem* **277**, 24694–24700.
- Best, S. M., Morris, K. L., Shannon, J. G., Robertson, S. J., Mitzel, D. N., Park, G. S., Boer, E., Wolfenbarger, J. B. & Bloom, M. E. (2005). Inhibition of interferon-stimulated JAK-STAT signaling by a tick-borne flavivirus and identification of NS5 as an interferon antagonist. *J Virol* **79**, 12828–12839.
- Bhattacharya, S., Eckner, R., Grossman, S., Oldread, E., Arany, Z., D'Andrea, A. & Livingston, D. M. (1996). Cooperation of Stat2 and p300/CBP in signalling induced by interferon- α . *Nature* **383**, 344–347.

- Billecocq, A., Spiegel, M., Vialat, P., Kohl, A., Weber, F., Bouloy, M. & Haller, O. (2004).** NSs protein of Rift Valley fever virus blocks interferon production by inhibiting host gene transcription. *J Virol* **78**, 9798–9806.
- Blakqori, G., Delhaye, S., Habjan, M., Blair, C. D., Sanchez-Vargas, I., Olson, K. E., Attarzadeh-Yazdi, G., Fragkoudis, R., Kohl, A. & other authors (2007).** La Crosse bunyavirus nonstructural protein NSs serves to suppress the type I interferon system of mammalian hosts. *J Virol* **81**, 4991–4999.
- Blindenbacher, A., Duong, F. H., Hunziker, L., Stutvoet, S. T., Wang, X., Terracciano, L., Moradpour, D., Blum, H. E., Alonzi, T. & other authors (2003).** Expression of hepatitis c virus proteins inhibits interferon α signaling in the liver of transgenic mice. *Gastroenterology* **124**, 1465–1475.
- Blondel, D., Regad, T., Poisson, N., Pavie, B., Harper, F., Pandolfi, P. P., De The, H. & Chelbi-Alix, M. K. (2002).** Rabies virus P and small P products interact directly with PML and reorganize PML nuclear bodies. *Oncogene* **21**, 7957–7970.
- Bluyssen, H. A. & Levy, D. E. (1997).** Stat2 is a transcriptional activator that requires sequence-specific contacts provided by stat1 and p48 for stable interaction with DNA. *J Biol Chem* **272**, 4600–4605.
- Bluyssen, H. A., Muzaffar, R., Vlieststra, R. J., van der Made, A. C., Leung, S., Stark, G. R., Kerr, I. M., Trapman, J. & Levy, D. E. (1995).** Combinatorial association and abundance of components of interferon-stimulated gene factor 3 dictate the selectivity of interferon responses. *Proc Natl Acad Sci U S A* **92**, 5645–5649.
- Bode, J. G., Ludwig, S., Ehrhardt, C., Albrecht, U., Erhardt, A., Schaper, F., Heinrich, P. C. & Haussinger, D. (2003).** IFN- α antagonistic activity of HCV core protein involves induction of suppressor of cytokine signaling-3. *FASEB J* **17**, 488–490.
- Boehme, K. W. & Compton, T. (2004).** Innate sensing of viruses by Toll-like receptors. *J Virol* **78**, 7867–7873.
- Bolitto, P., Voskoboinik, I., Trapani, J. A. & Smyth, M. J. (2007).** Apoptosis induced by the lymphocyte effector molecule perforin. *Curr Opin Immunol* **19**, 339–347.
- Bonvin, M., Achermann, F., Greeve, I., Stroka, D., Keogh, A., Inderbitzin, D., Candinas, D., Sommer, P., Wain-Hobson, S. & other authors (2006).** Interferon-inducible expression of APOBEC3 editing enzymes in human hepatocytes and inhibition of hepatitis B virus replication. *Hepatology* **43**, 1364–1374.
- Boone, D. L., Turer, E. E., Lee, E. G., Ahmad, R. C., Wheeler, M. T., Tsui, C., Hurley, P., Chien, M., Chai, S. & other authors (2004).** The ubiquitin-modifying enzyme A20 is required for termination of Toll-like receptor responses. *Nat Immunol* **5**, 1052–1060.
- Borden, K. L., Campbell Dwyer, E. J. & Salvato, M. S. (1998a).** An arenavirus RING (zinc-binding) protein binds the oncoprotein promyelocyte leukemia protein (PML) and relocates PML nuclear bodies to the cytoplasm. *J Virol* **72**, 758–766.
- Borden, K. L., Campbelldwyer, E. J., Carlile, G. W., Djavani, M. & Salvato, M. S. (1998b).** Two RING finger proteins, the oncoprotein PML and the arenavirus Z protein, colocalize with the nuclear fraction of the ribosomal P proteins. *J Virol* **72**, 3819–3826.
- Bossert, B. & Conzelmann, K. K. (2002).** Respiratory syncytial virus (RSV) nonstructural (NS) proteins as host range determinants: a chimeric bovine RSV with NS genes from human RSV is attenuated in interferon-competent bovine cells. *J Virol* **76**, 4287–4293.
- Bossert, B., Marozin, S. & Conzelmann, K. K. (2003).** Nonstructural proteins NS1 and NS2 of bovine respiratory syncytial virus block activation of interferon regulatory factor 3. *J Virol* **77**, 8661–8668.
- Bouloy, M., Janzen, C., Vialat, P., Khun, H., Pavlovic, J., Huerre, M. & Haller, O. (2001).** Genetic evidence for an interferon-antagonistic function of Rift Valley fever virus nonstructural protein NSs. *J Virol* **75**, 1371–1377.
- Boyman, O., Purton, J. F., Surh, C. D. & Sprent, J. (2007).** Cytokines and T-cell homeostasis. *Curr Opin Immunol* **19**, 320–326.
- Brideau-Andersen, A. D., Huang, X., Sun, S. C., Chen, T. T., Stark, D., Sas, I. J., Zadik, L., Dawes, G. N., Guptill, D. R. & other authors (2007).** Directed evolution of gene-shuffled IFN- α molecules with activity profiles tailored for treatment of chronic viral diseases. *Proc Natl Acad Sci U S A* **104**, 8269–8274.
- Brzozka, K., Finke, S. & Conzelmann, K. K. (2005).** Identification of the rabies virus alpha/beta interferon antagonist: phosphoprotein P interferes with phosphorylation of interferon regulatory factor 3. *J Virol* **79**, 7673–7681.
- Brzozka, K., Finke, S. & Conzelmann, K. K. (2006).** Inhibition of interferon signaling by rabies virus phosphoprotein P: activation-dependent binding of STAT1 and STAT2. *J Virol* **80**, 2675–2683.
- Butler, D. (2004).** Fatal fruit bat virus sparks epidemics in southern Asia. *Nature* **429**, 7.
- Caillaud, A., Hovanessian, A. G., Levy, D. E. & Marie, I. J. (2005).** Regulatory serine residues mediate phosphorylation-dependent and phosphorylation-independent activation of interferon regulatory factor 7. *J Biol Chem* **280**, 17671–17677.
- Camus-Bouclainville, C., Fiette, L., Bouchiha, S., Pignolet, B., Counor, D., Filipe, C., Gelfi, J. & Messud-Petit, F. (2004).** A virulence factor of myxoma virus colocalizes with NF- κ B in the nucleus and interferes with inflammation. *J Virol* **78**, 2510–2516.
- Cantrell, S. R. & Bresnahan, W. A. (2006).** Human cytomegalovirus (HCMV) UL82 gene product (pp71) relieves hDaxx-mediated repression of HCMV replication. *J Virol* **80**, 6188–6191.
- Cao, W. & Liu, Y. J. (2007).** Innate immune functions of plasmacytoid dendritic cells. *Curr Opin Immunol* **19**, 24–30.
- Capobianchi, M. R., Ankel, H., Ameglio, F., Paganelli, R., Pizzoli, P. M. & Dianzani, F. (1992).** Recombinant glycoprotein 120 of human immunodeficiency virus is a potent interferon inducer. *AIDS Res Hum Retroviruses* **8**, 575–579.
- Cardenas, W. B., Loo, Y. M., Gale, M., Jr, Hartman, A. L., Kimberlin, C. R., Martinez-Sobrido, L., Saphire, E. O. & Basler, C. F. (2006).** Ebola virus VP30 protein binds double-stranded RNA and inhibits alpha/beta interferon production induced by RIG-I signaling. *J Virol* **80**, 5168–5178.
- Carlos, T. S., Fearn, R. & Randall, R. E. (2005).** Interferon-induced alterations in the pattern of parainfluenza virus 5 transcription and protein synthesis and the induction of virus inclusion bodies. *J Virol* **79**, 14112–14121.
- Cassady, K. A. (2005).** Human cytomegalovirus TRS1 and IRS1 gene products block the double-stranded-RNA-activated host protein shutoff response induced by herpes simplex virus type 1 infection. *J Virol* **79**, 8707–8715.
- Castelli, J. C., Hassel, B. A., Maran, A., Paranjape, J., Hewitt, J. A., Li, X. L., Hsu, Y. T., Silverman, R. H. & Youle, R. J. (1998).** The role of 2'-5' oligoadenylate-activated ribonuclease L in apoptosis. *Cell Death Differ* **5**, 313–320.
- Cattaneo, R. (1994).** Biased (A→I) hypermutation of animal RNA virus genomes. *Curr Opin Genet Dev* **4**, 895–900.
- Chang, H. W., Watson, J. C. & Jacobs, B. L. (1992).** The E3L gene of vaccinia virus encodes an inhibitor of the interferon-induced, double-stranded RNA-dependent protein kinase. *Proc Natl Acad Sci U S A* **89**, 4825–4829.
- Chariot, A., Leonardi, A., Muller, J., Bonif, M., Brown, K. & Siebenlist, U. (2002).** Association of the adaptor TANK with the I κ B kinase (IKK)

- regulator NEMO connects IKK complexes with IKK ϵ and TBK1 kinases. *J Biol Chem* **277**, 37029–37036.
- Chatterjee-Kishore, M., van Den Akker, F. & Stark, G. R. (2000a).** Adenovirus E1A down-regulates LMP2 transcription by interfering with the binding of Stat1 to IRF1. *J Biol Chem* **275**, 20406–20411.
- Chatterjee-Kishore, M., Wright, K. L., Ting, J. P. & Stark, G. R. (2000b).** How Stat1 mediates constitutive gene expression: a complex of unphosphorylated Stat1 and IRF1 supports transcription of the LMP2 gene. *EMBO J* **19**, 4111–4122.
- Chatziandreou, N., Young, D., Andrejeva, J., Goodbourn, S. & Randall, R. E. (2002).** Differences in interferon sensitivity and biological properties of two related isolates of simian virus 5: a model for virus persistence. *Virology* **293**, 234–242.
- Chee, A. V. & Roizman, B. (2004).** Herpes simplex virus 1 gene products occlude the interferon signaling pathway at multiple sites. *J Virol* **78**, 4185–4196.
- Chen, Z. J. (2005).** Ubiquitin signalling in the NF- κ B pathway. *Nat Cell Biol* **7**, 758–765.
- Chen, Z., Li, Y. & Krug, R. M. (1999).** Influenza A virus NS1 protein targets poly(A)-binding protein II of the cellular 3'-end processing machinery. *EMBO J* **18**, 2273–2283.
- Chen, K., Huang, J., Zhang, C., Huang, S., Nunnari, G., Wang, F. X., Tong, X., Gao, L., Nikisher, K. & Zhang, H. (2006).** Alpha interferon potently enhances the anti-human immunodeficiency virus type 1 activity of APOBEC3G in resting primary CD4 T cells. *J Virol* **80**, 7645–7657.
- Cheng, G., Zhong, J. & Chisari, F. V. (2006).** Inhibition of dsRNA-induced signaling in hepatitis C virus-infected cells by NS3 protease-dependent and -independent mechanisms. *Proc Natl Acad Sci U S A* **103**, 8499–8504.
- Cheng, G., Zhong, J., Chung, J. & Chisari, F. V. (2007).** Double-stranded DNA and double-stranded RNA induce a common antiviral signaling pathway in human cells. *Proc Natl Acad Sci U S A* **104**, 9035–9040.
- Child, S. J., Hanson, L. K., Brown, C. E., Janzen, D. M. & Geballe, A. P. (2006).** Double-stranded RNA binding by a heterodimeric complex of murine cytomegalovirus m142 and m143 proteins. *J Virol* **80**, 10173–10180.
- Childs, K., Stock, N., Ross, C., Andrejeva, J., Hilton, L., Skinner, M., Randall, R. & Goodbourn, S. (2007).** mda-5, but not RIG-I, is a common target for paramyxovirus V proteins. *Virology* **359**, 190–200.
- Chin, K. C. & Cresswell, P. (2001).** Viperin (cig5), an IFN-inducible antiviral protein directly induced by human cytomegalovirus. *Proc Natl Acad Sci U S A* **98**, 15125–15130.
- Chin, Y. E., Kitagawa, M., Kuida, K., Flavell, R. A. & Fu, X. Y. (1997).** Activation of the STAT signaling pathway can cause expression of caspase 1 and apoptosis. *Mol Cell Biol* **17**, 5328–5337.
- Chinsangaram, J., Piccone, M. E. & Grubman, M. J. (1999).** Ability of foot-and-mouth disease virus to form plaques in cell culture is associated with suppression of alpha/beta interferon. *J Virol* **73**, 9891–9898.
- Choe, S. S., Dodd, D. A. & Kirkegaard, K. (2005).** Inhibition of cellular protein secretion by picornaviral 3A proteins. *Virology* **337**, 18–29.
- Christen, V., Duong, F., Bernsmeier, C., Sun, D., Nassal, M. & Heim, M. H. (2007).** Inhibition of alpha interferon signaling by hepatitis B virus. *J Virol* **81**, 159–165.
- Ciccaglione, A. R., Stellacci, E., Marcantonio, C., Muto, V., Equestre, M., Marsili, G., Rapicetta, M. & Battistini, A. (2007).** Repression of interferon regulatory factor 1 by hepatitis C virus core protein results in inhibition of antiviral and immunomodulatory genes. *J Virol* **81**, 202–214.
- Civas, A., Island, M. L., Genin, P., Morin, P. & Navarro, S. (2002).** Regulation of virus-induced interferon-A genes. *Biochimie* **84**, 643–654.
- Clark, M. E., Lieberman, P. M., Berk, A. J. & Dasgupta, A. (1993).** Direct cleavage of human TATA-binding protein by poliovirus protease 3C in vivo and in vitro. *Mol Cell Biol* **13**, 1232–1237.
- Clemens, M. J. (2003).** Interferons and apoptosis. *J Interferon Cytokine Res* **23**, 277–292.
- Clemens, M. J. (2005).** Translational control in virus-infected cells: models for cellular stress responses. *Semin Cell Dev Biol* **16**, 13–20.
- Colamonici, O. R., Domanski, P., Sweitzer, S. M., Larner, A. & Buller, R. M. L. (1995).** Vaccinia virus B18R gene encodes a type I interferon-binding protein that blocks interferon α transmembrane signaling. *J Biol Chem* **270**, 15974–15978.
- Connor, J. H. & Lyles, D. S. (2002).** Vesicular stomatitis virus infection alters the eIF4F translation initiation complex and causes dephosphorylation of the eIF4E binding protein 4E-BP1. *J Virol* **76**, 10177–10187.
- Connor, J. H. & Lyles, D. S. (2005).** Inhibition of host and viral translation during vesicular stomatitis virus infection. eIF2 is responsible for the inhibition of viral but not host translation. *J Biol Chem* **280**, 13512–13519.
- Conzelmann, K. K. (2005).** Transcriptional activation of alpha/beta interferon genes: interference by nonsegmented negative-strand RNA viruses. *J Virol* **79**, 5241–5248.
- Cooray, S., Bahar, M. W., Abrescia, N. G., McVey, C. E., Bartlett, N. W., Chen, R. A., Stuart, D. I., Grimes, J. M. & Smith, G. L. (2007).** Functional and structural studies of the vaccinia virus virulence factor N1 reveal a Bcl-2-like anti-apoptotic protein. *J Gen Virol* **88**, 1656–1666.
- Cröze, E., Usacheva, A., Asarnow, D., Minshall, R. D., Perez, H. D. & Colamonici, O. (2000).** Receptor for activated C-kinase (RACK-1), a WD motif-containing protein, specifically associates with the human type I IFN receptor. *J Immunol* **165**, 5127–5132.
- Cuddihy, A. R., Wong, A. H., Tam, N. W., Li, S. & Koromilas, A. E. (1999a).** The double-stranded RNA activated protein kinase PKR physically associates with the tumor suppressor p53 protein and phosphorylates human p53 on serine 392 in vitro. *Oncogene* **18**, 2690–2702.
- Cuddihy, A. R., Li, S., Tam, N. W., Wong, A. H., Taya, Y., Abraham, N., Bell, J. C. & Koromilas, A. E. (1999b).** Double-stranded-RNA-activated protein kinase PKR enhances transcriptional activation by tumor suppressor p53. *Mol Cell Biol* **19**, 2475–2484.
- Cusson-Hernance, N., Khurana, S., Lee, T. H., Fitzgerald, K. A. & Kelliher, M. A. (2005).** Rip1 mediates the Trif-dependent Toll-like receptor 3- and 4-induced NF- κ B activation but does not contribute to interferon regulatory factor 3 activation. *J Biol Chem* **280**, 36560–36566.
- Dai, C. & Krantz, S. B. (1999).** Interferon γ induces upregulation and activation of caspases 1, 3, and 8 to produce apoptosis in human erythroid progenitor cells. *Blood* **93**, 3309–3316.
- Das, S. & Dasgupta, A. (1993).** Identification of the cleavage site and determinants required for poliovirus 3CPro-catalyzed cleavage of human TATA-binding transcription factor TBP. *J Virol* **67**, 3326–3331.
- de Bouteiller, O., Merck, E., Hasan, U. A., Hubac, S., Benguigui, B., Trinchieri, G., Bates, E. E. & Caux, C. (2005).** Recognition of double-stranded RNA by human Toll-like receptor 3 and downstream receptor signaling requires multimerization and an acidic pH. *J Biol Chem* **280**, 38133–38145.
- De Clercq, E. (2006).** Interferon and its inducers – a never-ending story: “old” and “new” data in a new perspective. *J Infect Dis* **194**, S19–S26.

- de Los Santos, T., de Avila Botton, S., Weiblen, R. & Grubman, M. J. (2006). The leader proteinase of foot-and-mouth disease virus inhibits the induction of beta interferon mRNA and blocks the host innate immune response. *J Virol* **80**, 1906–1914.
- de Lucas, S., Bartolome, J. & Carreno, V. (2005). Hepatitis C virus core protein down-regulates transcription of interferon-induced antiviral genes. *J Infect Dis* **191**, 93–99.
- Deb, D. K., Sassano, A., Lekmine, F., Majchrzak, B., Verma, A., Kambhampati, S., Uddin, S., Rahman, A., Fish, E. N. & Platanias, L. C. (2003). Activation of protein kinase C δ by IFN- γ . *J Immunol* **171**, 267–273.
- Decker, T., Lew, D. J. & Darnell, J. E. J. (1991). Two distinct alpha-interferon-dependent signal transduction pathways may contribute to activation of transcription of the guanylate-binding protein gene. *Mol Cell Biol* **11**, 5147–5153.
- Decman, V., Freeman, M. L., Kinchington, P. R. & Hendricks, R. L. (2005). Immune control of HSV-1 latency. *Viral Immunol* **18**, 466–473.
- Delale, T., Paquin, A., Asselin-Paturel, C., Dalod, M., Brizard, G., Bates, E. E., Kastner, P., Chan, S., Akira, S. & other authors (2005). MyD88-dependent and -independent murine cytomegalovirus sensing for IFN- α release and initiation of immune responses in vivo. *J Immunol* **175**, 6723–6732.
- Delhay, S., van Pesch, V. & Michiels, T. (2004). The leader protein of Theiler's virus interferes with nucleocytoplasmic trafficking of cellular proteins. *J Virol* **78**, 4357–4362.
- Deng, L., Wang, C., Spencer, E., Yang, L., Braun, A., You, J., Slaughter, C., Pickart, C. & Chen, Z. J. (2000). Activation of the I κ B kinase complex by TRAF6 requires a dimeric ubiquitin-conjugating enzyme complex and a unique polyubiquitin chain. *Cell* **103**, 351–361.
- Desmyter, J., Melnick, J. L. & Rawls, W. E. (1968). Defectiveness of interferon production and of rubella virus interference in a line of African green monkey kidney cells (Vero). *J Virol* **2**, 955–961.
- Devaney, M. A., Vakharia, V. N., Lloyd, R. E., Ehrenfeld, E. & Grubman, M. J. (1988). Leader protein of foot-and-mouth disease virus is required for cleavage of the p220 component of the cap-binding protein complex. *J Virol* **62**, 4407–4409.
- Diao, F., Li, S., Tian, Y., Zhang, M., Xu, L. G., Zhang, Y., Wang, R. P., Chen, D., Zhai, Z. & other authors (2007). Negative regulation of MDA5- but not RIG-I-mediated innate antiviral signaling by the dihydroxyacetone kinase. *Proc Natl Acad Sci U S A* **104**, 11706–11711.
- Didcock, L., Young, D. F., Goodbourn, S. & Randall, R. E. (1999a). The V protein of simian virus 5 inhibits interferon signalling by targeting STAT1 for proteasome-mediated degradation. *J Virol* **73**, 9928–9933.
- Didcock, L., Young, D. F., Goodbourn, S. & Randall, R. E. (1999b). Sendai virus and simian virus 5 block activation of interferon-responsive genes: importance for virus pathogenesis. *J Virol* **73**, 3125–3133.
- Diebold, S. S., Kaisho, T., Hemmi, H., Akira, S. & Reis e Sousa, C. (2004). Innate antiviral responses by means of TLR7-mediated recognition of single-stranded RNA. *Science* **303**, 1529–1531.
- Diebold, S. S., Massacrier, C., Akira, S., Paturel, C., Morel, Y. & Reis e Sousa, C. (2006). Nucleic acid agonists for Toll-like receptor 7 are defined by the presence of uridine ribonucleotides. *Eur J Immunol* **36**, 3256–3267.
- Dillon, P. J. & Parks, G. D. (2007). Role for the phosphoprotein P subunit of the paramyxovirus polymerase in limiting induction of host cell antiviral responses. *J Virol* **81**, 11116–11127.
- DiPerna, G., Stack, J., Bowie, A. G., Boyd, A., Kotwal, G., Zhang, Z., Arvikar, S., Latz, E., Fitzgerald, K. A. & Marshall, W. L. (2004). Poxvirus protein NIL targets the I- κ B kinase complex, inhibits signaling to NF- κ B by the tumor necrosis factor superfamily of receptors, and inhibits NF- κ B and IRF3 signaling by Toll-like receptors. *J Biol Chem* **279**, 36570–36578.
- Djavani, M., Rodas, J., Lukashevich, I. S., Horejsh, D., Pandolfi, P. P., Borden, K. L. & Salvato, M. S. (2001). Role of the promyelocytic leukemia protein PML in the interferon sensitivity of lymphocytic choriomeningitis virus. *J Virol* **75**, 6204–6208.
- Doepker, R. C., Hsu, W. L., Saffran, H. A. & Smiley, J. R. (2004). Herpes simplex virus virion host shutoff protein is stimulated by translation initiation factors eIF4B and eIF4H. *J Virol* **78**, 4684–4699.
- Doucas, V., Ishov, A. M., Romo, A., Juguilon, H., Weitzman, M. D., Evans, R. M. & Maul, G. G. (1996). Adenovirus replication is coupled with the dynamic properties of the PML nuclear structure. *Genes Dev* **10**, 196–207.
- Dragan, A. I., Hargreaves, V. V., Makeyeva, E. N. & Privalov, P. L. (2007). Mechanisms of activation of interferon regulator factor 3: the role of C-terminal domain phosphorylation in IRF-3 dimerization and DNA binding. *Nucleic Acids Res* **35**, 3525–3534.
- Duong, F. H., Filipowicz, M., Tripodi, M., La Monica, N. & Heim, M. H. (2004). Hepatitis C virus inhibits interferon signaling through up-regulation of protein phosphatase 2A. *Gastroenterology* **126**, 263–277.
- Duong, F. H., Christen, V., Filipowicz, M. & Heim, M. H. (2006). S-Adenosylmethionine and betaine correct hepatitis C virus induced inhibition of interferon signaling in vitro. *Hepatology* **43**, 796–806.
- Ea, C. K., Deng, L., Xia, Z. P., Pineda, G. & Chen, Z. J. (2006). Activation of IKK by TNF α requires site-specific ubiquitination of RIP1 and polyubiquitin binding by NEMO. *Mol Cell* **22**, 245–257.
- Edelmann, K. H., Richardson-Burns, S., Alexopoulou, L., Tyler, K. L., Flavell, R. A. & Oldstone, M. B. (2004). Does Toll-like receptor 3 play a biological role in virus infections? *Virology* **322**, 231–238.
- Ehrhardt, C., Wolff, T., Pleschka, S., Planz, O., Beermann, W., Bode, J. G., Schmolke, M. & Ludwig, S. (2007). The influenza A virus NS1 protein activates the PI3K/Akt pathway to mediate antiapoptotic signaling responses. *J Virol* **81**, 3058–3067.
- Elankumaran, S., Rockemann, D. & Samal, S. K. (2006). Newcastle disease virus exerts oncolysis by both intrinsic and extrinsic caspase-dependent pathways of cell death. *J Virol* **80**, 7522–7534.
- Elliott, J., Lynch, O. T., Suessmuth, Y., Qian, P., Boyd, C. R., Burrows, J. F., Buick, R., Stevenson, N. J., Touzelet, O. & other authors (2007). Respiratory syncytial virus NS1 protein degrades STAT2 by using the Elongin-Cullin E3 ligase. *J Virol* **81**, 3428–3436.
- Ellis, M. J. & Goodbourn, S. (1994). NF- κ B-independent activation of β -interferon expression in mouse P9 embryonal carcinoma cells. *Nucleic Acids Res* **22**, 4489–4496.
- Epperson, D. E., Arnold, D., Spies, T., Cresswell, P., Pober, J. S. & Johnson, D. R. (1992). Cytokines increase transporter in antigen processing-1 expression more rapidly than HLA class I expression in endothelial cells. *J Immunol* **149**, 3297–3301.
- Erlandsson, L., Blumenthal, R., Eloranta, M. L., Engel, H., Alm, G., Weiss, S. & Leanderson, T. (1998). Interferon- β is required for interferon- α production in mouse fibroblasts. *Curr Biol* **8**, 223–226.
- Espert, L., Codogno, P. & Biard-Piechaczyk, M. (2007). Involvement of autophagy in viral infections: antiviral function and subversion by viruses. *J Mol Med* **85**, 811–823.
- Etchison, D., Milburn, S. C., Ederly, I., Sonenberg, N. & Hershey, J. W. (1982). Inhibition of HeLa cell protein synthesis following poliovirus infection correlates with the proteolysis of a 220,000-dalton polypeptide associated with eucaryotic initiation factor 3 and a cap binding protein complex. *J Biol Chem* **257**, 14806–14810.

- Everett, R. D. (2001). DNA viruses and viral proteins that interact with PML nuclear bodies. *Oncogene* **20**, 7266–7273.
- Everett, R. D. (2006). Interactions between DNA viruses, ND10 and the DNA damage response. *Cell Microbiol* **8**, 365–374.
- Everett, R. D. & Chelbi-Alix, M. K. (2007). PML and PML nuclear bodies: implications in antiviral defence. *Biochimie* **89**, 819–830.
- Everly, D. N., Jr, Feng, P., Mian, I. S. & Read, G. S. (2002). mRNA degradation by the virion host shutoff (Vhs) protein of herpes simplex virus: genetic and biochemical evidence that Vhs is a nuclease. *J Virol* **76**, 8560–8571.
- Feng, Z., Cerveny, M., Yan, Z. & He, B. (2007). The VP35 protein of Ebola virus inhibits the antiviral effect mediated by double-stranded RNA-dependent protein kinase PKR. *J Virol* **81**, 182–192.
- Fensterl, V., Grotheer, D., Berk, I., Schlemminger, S., Vallbracht, A. & Dotzauer, A. (2005). Hepatitis A virus suppresses RIG-I-mediated IRF-3 activation to block induction of β interferon. *J Virol* **79**, 10968–10977.
- Ferko, B., Stasakova, J., Romanova, J., Kittel, C., Sereinig, S., Katinger, H. & Egorov, A. (2004). Immunogenicity and protection efficacy of replication-deficient influenza A viruses with altered NS1 genes. *J Virol* **78**, 13037–13045.
- Fernandez, M., Quiroga, J. A. & Carreno, V. (2003). Hepatitis B virus downregulates the human interferon-inducible MxA promoter through direct interaction of precore/core proteins. *J Gen Virol* **84**, 2073–2082.
- Fernandez-Sesma, A., Marukian, S., Ebersole, B. J., Kaminski, D., Park, M. S., Yuen, T., Sealfon, S. C., Garcia-Sastre, A. & Moran, T. M. (2006). Influenza virus evades innate and adaptive immunity via the NS1 protein. *J Virol* **80**, 6295–6304.
- Ferran, M. C. & Lucas-Lenard, J. M. (1997). The vesicular stomatitis virus matrix protein inhibits transcription from the human beta interferon promoter. *J Virol* **71**, 371–377.
- Ferrantini, M., Capone, I. & Belardelli, F. (2007). Interferon- α and cancer: mechanisms of action and new perspectives of clinical use. *Biochimie* **89**, 884–893.
- Fitzgerald, K. A., Palsson-McDermott, E. M., Bowie, A. G., Jefferies, C. A., Mansell, A. S., Brady, G., Brint, E., Dunne, A., Gray, P. & other authors (2001). Mal (MyD88-adaptor-like) is required for Toll-like receptor-4 signal transduction. *Nature* **413**, 78–83.
- Fitzgerald, K. A., McWhirter, S. M., Faia, K. L., Rowe, D. C., Latz, E., Golenbock, D. T., Coyle, A. J., Liao, S. M. & Maniatis, T. (2003). IKK ϵ and TBK1 are essential components of the IRF3 signaling pathway. *Nat Immunol* **4**, 491–496.
- Fortes, P., Beloso, A. & Ortin, J. (1994). Influenza virus NS1 protein inhibits pre-mRNA splicing and blocks mRNA nucleocytoplasmic transport. *EMBO J* **13**, 704–712.
- Frahm, T., Hauser, H. & Koster, M. (2006). IFN-type-I-mediated signaling is regulated by modulation of STAT2 nuclear export. *J Cell Sci* **119**, 1092–1104.
- Frese, M., Pietschmann, T., Moradpour, D., Haller, O. & Bartenschlager, R. (2001). Interferon- α inhibits hepatitis C virus subgenomic RNA replication by an MxA-independent pathway. *J Gen Virol* **82**, 723–733.
- Friedman, J. M. & Horwitz, M. S. (2002). Inhibition of tumor necrosis factor alpha-induced NF- κ B activation by the adenovirus E3–10.4/14.5K complex. *J Virol* **76**, 5515–5521.
- Friedrichs, W. E., Reddy, S. V., Bruder, J. M., Cundy, T., Cornish, J., Singer, F. R. & Roodman, G. D. (2002). Sequence analysis of measles virus nucleocapsid transcripts in patients with Paget's disease. *J Bone Miner Res* **17**, 145–151.
- Frieman, M., Yount, B., Heise, M., Kopecky-Bromberg, S. A., Palese, P. & Baric, R. S. (2007). Severe acute respiratory syndrome coronavirus ORF6 antagonizes STAT1 function by sequestering nuclear import factors on the rough endoplasmic reticulum/Golgi membrane. *J Virol* **81**, 9812–9824.
- Frolova, E. I., Fayzuln, R. Z., Cook, S. H., Griffin, D. E., Rice, C. M. & Frolov, I. (2002). Roles of nonstructural protein nsP2 and alpha/beta interferons in determining the outcome of Sindbis virus infection. *J Virol* **76**, 11254–11264.
- Fujita, T., Onoguchi, K., Onomoto, K., Hirai, R. & Yoneyama, M. (2007). Triggering antiviral response by RIG-I-related RNA helicases. *Biochimie* **89**, 754–760.
- Gack, M. U., Shin, Y. C., Joo, C. H., Urano, T., Liang, C., Sun, L., Takeuchi, O., Akira, S., Chen, Z. & other authors (2007). TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated antiviral activity. *Nature* **446**, 916–920.
- Gaddy, D. F. & Lyles, D. S. (2007). Oncolytic vesicular stomatitis virus induces apoptosis via signaling through PKR, Fas, and Daxx. *J Virol* **81**, 2792–2804.
- Garcia, M. A., Gil, J., Ventoso, I., Guerra, S., Domingo, E., Rivas, C. & Esteban, M. (2006). Impact of protein kinase PKR in cell biology: from antiviral to antiproliferative action. *Microbiol Mol Biol Rev* **70**, 1032–1060.
- Garcia-Sastre, A. (2007). De-ISGylating and de-ubiquitinating activities of viral OUT-like proteins. In *Gordon Research Conference 2007: Viruses and Cells, 3–8 June 2007* (oral presentation). Tilton School, Tilton, NH, USA.
- Garcin, D., Latorre, P. & Kolakofsky, D. (1999). Sendai virus C proteins counteract the interferon-mediated induction of an antiviral state. *J Virol* **73**, 6559–6565.
- Garcin, D., Curran, J. & Kolakofsky, D. (2000). Sendai virus C proteins must interact directly with cellular components to interfere with interferon action. *J Virol* **74**, 8823–8830.
- Garcin, D., Curran, J., Itoh, M. & Kolakofsky, D. (2001). Longer and shorter forms of Sendai virus C proteins play different roles in modulating the cellular antiviral response. *J Virol* **75**, 6800–6807.
- Garcin, D., Marq, J. B., Strahle, L., le Mercier, P. & Kolakofsky, D. (2002). All four Sendai virus C proteins bind Stat1, but only the larger forms also induce its mono-ubiquitination and degradation. *Virology* **295**, 256–265.
- Garcin, D., Marq, J. B., Goodbourn, S. & Kolakofsky, D. (2003). The amino-terminal extensions of the longer Sendai virus C proteins modulate pY701-Stat1 and bulk Stat1 levels independently of interferon signaling. *J Virol* **77**, 2321–2329.
- Garmashova, N., Gorchakov, R., Frolova, E. & Frolov, I. (2006). Sindbis virus nonstructural protein nsP2 is cytotoxic and inhibits cellular transcription. *J Virol* **80**, 5686–5696.
- Georgel, P., Jiang, Z., Kunz, S., Janssen, E., Mols, J., Hoebe, K., Bahram, S., Oldstone, M. B. & Beutler, B. (2007). Vesicular stomatitis virus glycoprotein G activates a specific antiviral Toll-like receptor 4-dependent pathway. *Virology* **362**, 304–313.
- Giannakopoulos, N. V., Luo, J. K., Papov, V., Zou, W., Lenschow, D. J., Jacobs, B. S., Borden, E. C., Li, J., Virgin, H. W. & Zhang, D. E. (2005). Proteomic identification of proteins conjugated to ISG15 in mouse and human cells. *Biochem Biophys Res Commun* **336**, 496–506.
- Gil, J., Alcami, J. & Esteban, M. (1999). Induction of apoptosis by double-stranded-RNA-dependent protein kinase (PKR) involves the α subunit of eukaryotic translation initiation factor 2 and NF- κ B. *Mol Cell Biol* **19**, 4653–4663.
- Gitlin, L., Barchet, W., Gilfillan, S., Cella, M., Beutler, B., Flavell, R. A., Diamond, M. S. & Colonna, M. (2006). Essential role of mda-5 in type I IFN responses to polyriboinosinic:polyribocytidylic acid and encephalomyocarditis picornavirus. *Proc Natl Acad Sci U S A* **103**, 8459–8464.

- Goodbourn, S., Zinn, K. & Maniatis, T. (1985).** Human β -interferon gene expression is regulated by an inducible enhancer element. *Cell* **41**, 509–520.
- Gorchakov, R., Frolova, E. & Frolov, I. (2005).** Inhibition of transcription and translation in Sindbis virus-infected cells. *J Virol* **79**, 9397–9409.
- Gordon, M. T., Anderson, D. C. & Sharpe, P. T. (1991).** Canine distemper virus localised in bone cells of patients with Paget's disease. *Bone* **12**, 195–201.
- Gotoh, B., Takeuchi, K., Komatsu, T. & Yokoo, J. (2003a).** The STAT2 activation process is a crucial target of Sendai virus C protein for the blockade of alpha interferon signaling. *J Virol* **77**, 3360–3370.
- Gotoh, B., Komatsu, T., Takeuchi, K. & Yokoo, J. (2003b).** The C-terminal half-fragment of the Sendai virus C protein prevents the gamma-activated factor from binding to a gamma-activated sequence site. *Virology* **316**, 29–40.
- Gradi, A., Svitkin, Y. V., Imataka, H. & Sonenberg, N. (1998).** Proteolysis of human eukaryotic translation initiation factor eIF4GII, but not eIF4GI, coincides with the shutoff of host protein synthesis after poliovirus infection. *Proc Natl Acad Sci U S A* **95**, 11089–11094.
- Graff, J. W., Ewen, J., Ettayebi, K. & Hardy, M. E. (2007).** Zinc-binding domain of rotavirus NSP1 is required for proteasome-dependent degradation of IRF3 and autoregulatory NSP1 stability. *J Gen Virol* **88**, 613–620.
- Graham, R. R., Kyogoku, C., Sigurdsson, S., Vlasova, I. A., Davies, L. R., Baechler, E. C., Plenge, R. M., Koeth, T., Ortmann, W. A. & other authors (2007).** Three functional variants of IFN regulatory factor 5 (IRF5) define risk and protective haplotypes for human lupus. *Proc Natl Acad Sci U S A* **104**, 6758–6763.
- Guo, B. & Cheng, G. (2007).** Modulation of the interferon antiviral response by the TBK1/IKKi adaptor protein TANK. *J Biol Chem* **282**, 11817–11826.
- Guo, J., Hui, D. J., Merrick, W. C. & Sen, G. C. (2000).** A new pathway of translational regulation mediated by eukaryotic initiation factor 3. *EMBO J* **19**, 6891–6899.
- Guo, Z., Chen, L. M., Zeng, H., Gomez, J. A., Plowden, J., Fujita, T., Katz, J. M., Donis, R. O. & Sambhara, S. (2006).** NS1 protein of influenza A virus inhibits the function of intracytoplasmic pathogen sensor, RIG-I. *Am J Respir Cell Mol Biol* **36**, 263–269.
- Gustin, K. E. & Sarnow, P. (2001).** Effects of poliovirus infection on nucleocytoplasmic trafficking and nuclear pore complex composition. *EMBO J* **20**, 240–249.
- Hacker, H., Redecke, V., Blagoev, B., Kratchmarova, I., Hsu, L. C., Wang, G. G., Kamps, M. P., Raz, E., Wagner, H. & other authors (2006).** Specificity in Toll-like receptor signalling through distinct effector functions of TRAF3 and TRAF6. *Nature* **439**, 204–207.
- Hagglund, R. & Roizman, B. (2004).** Role of ICP0 in the strategy of conquest of the host cell by herpes simplex virus 1. *J Virol* **78**, 2169–2178.
- Hagmaier, K., Gelderblom, H. R. & Kochs, G. (2004).** Functional comparison of the two gene products of Thogoto virus segment 6. *J Gen Virol* **85**, 3699–3708.
- Hagmaier, K., Stock, N., Precious, B., Childs, K., Wang, L. F., Goodbourn, S. & Randall, R. E. (2007).** Mapuera virus, a rubulavirus that inhibits interferon signalling in a wide variety of mammalian cells without degrading STATs. *J Gen Virol* **88**, 956–966.
- Hakki, M. & Geballe, A. P. (2005).** Double-stranded RNA binding by human cytomegalovirus pTRS1. *J Virol* **79**, 7311–7318.
- Hale, B. G., Jackson, D., Chen, Y. H., Lamb, R. A. & Randall, R. E. (2006).** Influenza A virus NS1 protein binds p85 β and activates phosphatidylinositol-3-kinase signaling. *Proc Natl Acad Sci U S A* **103**, 14194–14199.
- Haller, O., Staeheli, P. & Kochs, G. (2007).** Interferon-induced Mx proteins in antiviral host defense. *Biochimie* **89**, 812–818.
- Harte, M. T., Haga, I. R., Maloney, G., Gray, P., Reading, P. C., Bartlett, N. W., Smith, G. L., Bowie, A. & O'Neill, L. A. (2003).** The poxvirus protein A52R targets Toll-like receptor signaling complexes to suppress host defense. *J Exp Med* **197**, 343–351.
- Hartman, A. L., Dover, J. E., Towner, J. S. & Nichol, S. T. (2006).** Reverse genetic generation of recombinant Zaire Ebola viruses containing disrupted IRF-3 inhibitory domains results in attenuated virus growth in vitro and higher levels of IRF-3 activation without inhibiting viral transcription or replication. *J Virol* **80**, 6430–6440.
- Hartmann, R., Justesen, J., Sarkar, S. N., Sen, G. C. & Yee, V. C. (2003).** Crystal structure of the 2'-specific and double-stranded RNA-activated interferon-induced antiviral protein 2'-5'-oligoadenylate synthetase. *Mol Cell* **12**, 1173–1185.
- Hartwig, D., Schutte, C., Warnecke, J., Dorn, I., Hennig, H., Kirchner, H. & Schlenke, P. (2006).** The large form of ADAR 1 is responsible for enhanced hepatitis delta virus RNA editing in interferon- α -stimulated host cells. *J Viral Hepat* **13**, 150–157.
- Hata, N., Sato, M., Takaoka, A., Asagiri, M., Tanaka, N. & Taniguchi, T. (2001).** Constitutive IFN- α/β signal for efficient IFN- α/β gene induction by virus. *Biochem Biophys Res Commun* **285**, 518–525.
- Hatzioannou, T., Perez-Caballero, D., Yang, A., Cowan, S. & Bieniasz, P. D. (2004).** Retrovirus resistance factors Ref1 and Lvl1 are species-specific variants of TRIM5 α . *Proc Natl Acad Sci U S A* **101**, 10774–10779.
- Hayden, M. S. & Ghosh, S. (2004).** Signaling to NF- κ B. *Genes Dev* **18**, 2195–2224.
- Hayman, A., Comely, S., Lackenby, A., Murphy, S., McCauley, J., Goodbourn, S. & Barclay, W. (2006).** Variation in the ability of human influenza A viruses to induce and inhibit 2'-IFN- β pathway. *Virology* **347**, 52–64.
- Hayman, A., Comely, S., Lackenby, A., Hartgroves, L. C., Goodbourn, S., McCauley, J. W. & Barclay, W. S. (2007).** NS1 proteins of avian influenza A viruses can act as antagonists of the human alpha/beta interferon response. *J Virol* **81**, 2318–2327.
- Haynes, L. M., Moore, D. D., Kurt-Jones, E. A., Finberg, R. W., Anderson, L. J. & Tripp, R. A. (2001).** Involvement of Toll-like receptor 4 in innate immunity to respiratory syncytial virus. *J Virol* **75**, 10730–10737.
- He, B., Lin, G. Y., Durbin, J. E., Durbin, R. K. & Lamb, R. A. (2001).** The SH integral membrane protein of the paramyxovirus simian virus 5 is required to block apoptosis in MDBK cells. *J Virol* **75**, 4068–4079.
- He, B., Paterson, R. G., Stock, N., Durbin, J. E., Durbin, R. K., Goodbourn, S., Randall, R. E. & Lamb, R. A. (2002).** Recovery of paramyxovirus simian virus 5 with a V protein lacking the conserved cysteine-rich domain: the multifunctional V protein blocks both interferon- β induction and interferon signaling. *Virology* **303**, 15–32.
- Heil, F., Ahmad-Nejad, P., Hemmi, H., Hochrein, H., Ampenberger, F., Gellert, T., Dietrich, H., Lipford, G., Takeda, K. & other authors (2003).** The Toll-like receptor 7 (TLR7)-specific stimulus loxoribine uncovers a strong relationship within the TLR7, 8 and 9 subfamily. *Eur J Immunol* **33**, 2987–2997.
- Heil, F., Hemmi, H., Hochrein, H., Ampenberger, F., Kirschning, C., Akira, S., Lipford, G., Wagner, H. & Bauer, S. (2004).** Species-specific recognition of single-stranded RNA via Toll-like receptor 7 and 8. *Science* **303**, 1526–1529.
- Heim, M. H., Moradpour, D. & Blum, H. E. (1999).** Expression of hepatitis C virus proteins inhibits signal transduction through the Jak-STAT pathway. *J Virol* **73**, 8469–8475.

- Helbig, K. J., Lau, D. T., Semendric, L., Harley, H. A. & Beard, M. R. (2005). Analysis of ISG expression in chronic hepatitis C identifies viperin as a potential antiviral effector. *Hepatology* **42**, 702–710.
- Hemmi, H., Kaisho, T., Takeuchi, O., Sato, S., Sanjo, H., Hoshino, K., Horiuchi, T., Tomizawa, H., Takeda, K. & Akira, S. (2002). Small antiviral compounds activate immune cells via the TLR7 MyD88-dependent signaling pathway. *Nat Immunol* **3**, 196–200.
- Her, L. S., Lund, E. & Dahlberg, J. E. (1997). Inhibition of Ran guanosine triphosphatase-dependent nuclear transport by the matrix protein of vesicular stomatitis virus. *Science* **276**, 1845–1848.
- Herzer, K., Weyer, S., Krammer, P. H., Galle, P. R. & Hofmann, T. G. (2005). Hepatitis C virus core protein inhibits tumor suppressor protein promyelocytic leukemia function in human hepatoma cells. *Cancer Res* **65**, 10830–10837.
- Hilton, L., Moganeradi, K., Zhang, G., Chen, Y. H., Randall, R. E., McCauley, J. W. & Goodbourn, S. (2006). The NPro product of bovine viral diarrhoea virus inhibits DNA binding by interferon regulatory factor-3 and targets it for proteasomal degradation. *J Virol* **80**, 11723–11732.
- Hiscott, J., Lin, R., Nakhaei, P. & Paz, S. (2006a). MasterCARD: a priceless link to innate immunity. *Trends Mol Med* **12**, 53–56.
- Hiscott, J., Nguyen, T. L., Arguello, M., Nakhaei, P. & Paz, S. (2006b). Manipulation of the nuclear factor- κ B pathway and the innate immune response by viruses. *Oncogene* **25**, 6844–6867.
- Ho, L. J., Hung, L. F., Weng, C. Y., Wu, W. L., Chou, P., Lin, Y. L., Chang, D. M., Tai, T. Y. & Lai, J. H. (2005). Dengue virus type 2 antagonizes IFN- α but not IFN- γ antiviral effect via down-regulating Tyk2-STAT signaling in the human dendritic cell. *J Immunol* **174**, 8163–8172.
- Hoebé, K., Du, X., Georgel, P., Janssen, E., Tabeta, K., Kim, S. O., Goode, J., Lin, P., Mann, N. & other authors (2003). Identification of Lps2 as a key transducer of MyD88-independent TIR signalling. *Nature* **424**, 743–748.
- Honda, K. & Taniguchi, T. (2006). IRFs: master regulators of signalling by Toll-like receptors and cytosolic pattern-recognition receptors. *Nat Rev Immunol* **6**, 644–658.
- Honda, K., Yanai, H., Mizutani, T., Negishi, H., Shimada, N., Suzuki, N., Ohba, Y., Takaoka, A., Yeh, W. C. & Taniguchi, T. (2004). Role of a transductional-transcriptional processor complex involving MyD88 and IRF-7 in Toll-like receptor signaling. *Proc Natl Acad Sci U S A* **101**, 15416–15421.
- Honda, K., Ohba, Y., Yanai, H., Negishi, H., Mizutani, T., Takaoka, A., Taya, C. & Taniguchi, T. (2005a). Spatiotemporal regulation of MyD88-IRF-7 signalling for robust type-I interferon induction. *Nature* **434**, 1035–1040.
- Honda, K., Yanai, H., Negishi, H., Asagiri, M., Sato, M., Mizutani, T., Shimada, N., Ohba, Y., Takaoka, A. & other authors (2005b). IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature* **434**, 772–777.
- Hoppe, A., Beech, S. J., Dimmock, J. & Leppard, K. N. (2006). Interaction of the adenovirus type 5 E4 Orf3 protein with promyelocytic leukemia protein isoform II is required for ND10 disruption. *J Virol* **80**, 3042–3049.
- Hornig, T., Barton, G. M. & Medzhitov, R. (2001). TIRAP: an adapter molecule in the Toll signaling pathway. *Nat Immunol* **2**, 835–841.
- Hornung, V., Ellegast, J., Kim, S., Brzozka, K., Jung, A., Kato, H., Poeck, H., Akira, S., Conzelmann, K. K. & other authors (2006). 5'-Triphosphate RNA is the ligand for RIG-I. *Science* **314**, 994–997.
- Horvath, C. M. (2004). Silencing STATs: lessons from paramyxovirus interferon evasion. *Cytokine Growth Factor Rev* **15**, 117–127.
- Hoshino, K., Sugiyama, T., Matsumoto, M., Tanaka, T., Saito, M., Hemmi, H., Ohara, O., Akira, S. & Kaisho, T. (2006). I κ B kinase- α is critical for interferon- α production induced by Toll-like receptors 7 and 9. *Nature* **440**, 949–953.
- Hoyland, J. A., Dixon, J. A., Berry, J. L., Davies, M., Selby, P. L. & Mee, A. P. (2003). A comparison of in situ hybridisation, reverse transcriptase-polymerase chain reaction (RT-PCR) and in situ-RT-PCR for the detection of canine distemper virus RNA in Paget's disease. *J Virol Methods* **109**, 253–259.
- Huang, Z., Krishnamurthy, S., Panda, A. & Samal, S. K. (2003). Newcastle disease virus V protein is associated with viral pathogenesis and functions as an alpha interferon antagonist. *J Virol* **77**, 8676–8685.
- Huang, J., Smirnov, S. V., Lewis-Antes, A., Balan, M., Li, W., Tang, S., Silke, G. V., Putz, M. M., Smith, G. L. & Kolenko, S. V. (2007). Inhibition of type I and type III interferons by a secreted glycoprotein from Yaba-like disease virus. *Proc Natl Acad Sci U S A* **104**, 9822–9827.
- Hui, D. J., Bhasker, C. R., Merrick, W. C. & Sen, G. C. (2003). Viral stress-inducible protein p56 inhibits translation by blocking the interaction of eIF3 with the ternary complex eIF2.GTP.Met-tRNAi. *J Biol Chem* **278**, 39477–39482.
- Hui, D. J., Terenzi, F., Merrick, W. C. & Sen, G. C. (2005). Mouse p56 blocks a distinct function of eukaryotic initiation factor 3 in translation initiation. *J Biol Chem* **280**, 3433–3440.
- Hunter, W. D., Martuza, R. L., Feigenbaum, F., Todo, T., Mineta, T., Yazaki, T., Toda, M., Newsome, J. T., Platenberg, R. C. & other authors (1999). Attenuated, replication-competent herpes simplex virus type 1 mutant G207: safety evaluation of intracerebral injection in nonhuman primates. *J Virol* **73**, 6319–6326.
- Huye, L. E., Ning, S., Kelliher, M. & Pagano, J. S. (2007). Interferon regulatory factor 7 is activated by a viral oncoprotein through RIP-dependent ubiquitination. *Mol Cell Biol* **27**, 2910–2918.
- Imani, F. & Jacobs, B. L. (1988). Inhibitory activity for the interferon-induced protein kinase is associated with the reovirus serotype 1 sigma 3 protein. *Proc Natl Acad Sci U S A* **85**, 7887–7891.
- Iordanov, M. S., Kirsch, J. D., Ryabinina, O. P., Wong, J., Spitz, P. N., Korcheva, V. B., Thorburn, A. & Magun, B. E. (2005). Recruitment of TRADD, FADD, and caspase 8 to double-stranded RNA-triggered death inducing signaling complexes (dsRNA-DISCs). *Apoptosis* **10**, 167–176.
- Iqbal, M., Poole, E., Goodbourn, S. & McCauley, J. W. (2004). Role for bovine viral diarrhoea virus Erns glycoprotein in the control of activation of beta interferon by double-stranded RNA. *J Virol* **78**, 136–145.
- Isaacs, A. & Lindenmann, J. (1957). Virus interference. I. The interferon. *Proc R Soc Lond B Biol Sci* **147**, 258–267.
- Ishii, K. J., Coban, C., Kato, H., Takahashi, K., Torii, Y., Takeshita, F., Ludwig, H., Sutter, G., Suzuki, K. & other authors (2006). A Toll-like receptor-independent antiviral response induced by double-stranded B-form DNA. *Nat Immunol* **7**, 40–48.
- Ito, T., Yang, M. & May, W. S. (1999). RAX, a cellular activator for double-stranded RNA-dependent protein kinase during stress signaling. *J Biol Chem* **274**, 15427–15432.
- Iwasaki, A. (2007). Role of autophagy in innate viral recognition. *Autophagy* **3**, 354–356.
- Jackson, W. T., Giddings, T. H., Jr, Taylor, M. P., Mulinyawe, S., Rabinovitch, M., Kopito, R. R. & Kirkegaard, K. (2005). Subversion of cellular autophagosomal machinery by RNA viruses. *PLoS Biol* **3**, e156.
- Jacobs, B. L. & Langland, J. O. (1998). Reovirus sigma 3 protein: dsRNA binding and inhibition of RNA-activated protein kinase. *Curr Top Microbiol Immunol* **233**, 185–196.
- Jaworska, J., Gravel, A., Fink, K., Grandvaux, N. & Flamand, L. (2007). Inhibition of transcription of the beta interferon gene by the human herpesvirus 6 immediate-early 1 protein. *J Virol* **81**, 5737–5748.

- Jennings, S., Martinez-Sobrido, L., Garcia-Sastre, A., Weber, F. & Kochs, G. (2005). Thogoto virus ML protein suppresses IRF3 function. *Virology* **331**, 63–72.
- Jiang, Z., Mak, T. W., Sen, G. & Li, X. (2004). Toll-like receptor 3-mediated activation of NF- κ B and IRF3 diverges at Toll-IL-1 receptor domain-containing adapter inducing IFN- β . *Proc Natl Acad Sci U S A* **101**, 3533–3538.
- Joachims, M., Van Breugel, P. C. & Lloyd, R. E. (1999). Cleavage of poly(A)-binding protein by enterovirus proteases concurrent with inhibition of translation in vitro. *J Virol* **73**, 718–727.
- Johnsen, I. B., Nguyen, T. T., Ringdal, M., Tryggestad, A. M., Bakke, O., Lien, E., Espevik, T. & Anthonson, M. W. (2006). Toll-like receptor 3 associates with c-Src tyrosine kinase on endosomes to initiate antiviral signaling. *EMBO J* **25**, 3335–3346.
- Johnson, C. L. & Gale, M., Jr (2006). CARD games between virus and host get a new player. *Trends Immunol* **27**, 1–4.
- Johnston, M. D. (1981). The characteristics required for a Sendai virus preparation to induce high levels of interferon in human lymphoblastoid cells. *J Gen Virol* **56**, 175–184.
- Johnston, J. B., Nazarian, S. H., Natale, R. & McFadden, G. (2005). Myxoma virus infection of primary human fibroblasts varies with cellular age and is regulated by host interferon responses. *Virology* **332**, 235–248.
- Jones, M., Davidson, A., Hibbert, L., Gruenwald, P., Schlaak, J., Ball, S., Foster, G. R. & Jacobs, M. (2005). Dengue virus inhibits alpha interferon signaling by reducing STAT2 expression. *J Virol* **79**, 5414–5420.
- Joo, M., Hahn, Y. S., Kwon, M., Sadikot, R. T., Blackwell, T. S. & Christman, J. W. (2005). Hepatitis C virus core protein suppresses NF- κ B activation and cyclooxygenase-2 expression by direct interaction with IkappaB kinase beta. *J Virol* **79**, 7648–7657.
- Joo, C. H., Shin, Y. C., Gack, M., Wu, L., Levy, D. & Jung, J. U. (2007). Inhibition of interferon regulatory factor 7 (IRF7)-mediated interferon signal transduction by the Kaposi's sarcoma-associated herpesvirus viral IRF homolog vIRF3. *J Virol* **81**, 8282–8292.
- Jost, S., Turelli, P., Mangeat, B., Protzer, U. & Trono, D. (2007). Induction of antiviral cytidine deaminases does not explain the inhibition of hepatitis B virus replication by interferons. *J Virol* **81**, 10588–10596.
- Jounai, N., Takeshita, F., Kobiyama, K., Sawano, A., Miyawaki, A., Xin, K. Q., Ishii, K. J., Kawai, T., Akira, S. & other authors (2007). The Atg5 Atg12 conjugate associates with innate antiviral immune responses. *Proc Natl Acad Sci U S A* **104**, 14050–14055.
- Jurk, M., Heil, F., Vollmer, J., Schetter, C., Krieg, A. M., Wagner, H., Lipford, G. & Bauer, S. (2002). Human TLR7 or TLR8 independently confer responsiveness to the antiviral compound R-848. *Nat Immunol* **3**, 499.
- Kanayama, A., Seth, R. B., Sun, L., Ea, C. K., Hong, M., Shaito, A., Chiu, Y. H., Deng, L. & Chen, Z. J. (2004). TAB2 and TAB3 activate the NF- κ B pathway through binding to polyubiquitin chains. *Mol Cell* **15**, 535–548.
- Kang, H., Kim, E. T., Lee, H. R., Park, J. J., Go, Y. Y., Choi, C. Y. & Ahn, J. H. (2006). Inhibition of SUMO-independent PML oligomerization by the human cytomegalovirus IE1 protein. *J Gen Virol* **87**, 2181–2190.
- Kariko, K., Buckstein, M., Ni, H. & Weissman, D. (2005). Suppression of RNA recognition by Toll-like receptors: the impact of nucleoside modification and the evolutionary origin of RNA. *Immunity* **23**, 165–175.
- Karin, M. (2006). Nuclear factor- κ B in cancer development and progression. *Nature* **441**, 431–436.
- Kato, A., Ohnishi, Y., Kohase, M., Saito, S., Tashiro, M. & Nagai, Y. (2001). Y2, the smallest of the Sendai virus C proteins, is fully capable of both counteracting the antiviral action of interferons and inhibiting viral RNA synthesis. *J Virol* **75**, 3802–3810.
- Kato, H., Sato, S., Yoneyama, M., Yamamoto, M., Uematsu, S., Matsui, K., Tsujimura, T., Takeda, K., Fujita, T. & other authors (2005). Cell type-specific involvement of RIG-I in antiviral response. *Immunity* **23**, 19–28.
- Kato, H., Takeuchi, O., Sato, S., Yoneyama, M., Yamamoto, M., Matsui, K., Uematsu, S., Jung, A., Kawai, T. & other authors (2006). Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. *Nature* **441**, 101–105.
- Katsoulidis, E., Li, Y., Mears, H. & Plataniias, L. C. (2005). The p38 mitogen-activated protein kinase pathway in interferon signal transduction. *J Interferon Cytokine Res* **25**, 749–756.
- Kaukinen, P., Sillanpaa, M., Kotenko, S., Lin, R., Hiscott, J., Melen, K. & Julkunen, I. (2006). Hepatitis C virus NS2 and NS3/4A proteins are potent inhibitors of host cell cytokine/chemokine gene expression. *Virol J* **3**, 66.
- Kaur, S., Uddin, S. & Plataniias, L. C. (2005). The PI3' kinase pathway in interferon signaling. *J Interferon Cytokine Res* **25**, 780–787.
- Kawai, T., Sato, S., Ishii, K. J., Coban, C., Hemmi, H., Yamamoto, M., Terai, K., Matsuda, M., Inoue, J. & other authors (2004). Interferon-alpha induction through Toll-like receptors involves a direct interaction of IRF7 with MyD88 and TRAF6. *Nat Immunol* **5**, 1061–1068.
- Kawai, T., Takahashi, K., Sato, S., Coban, C., Kumar, H., Kato, H., Ishii, K. J., Takeuchi, O. & Akira, S. (2005). IPS-1, an adaptor triggering RIG-I- and Mda5-mediated type I interferon induction. *Nat Immunol* **6**, 981–988.
- Kawakami, T., Matsumoto, M., Sato, M., Harada, H., Taniguchi, T. & Kitagawa, M. (1995). Possible involvement of the transcription factor ISGF3 γ in virus-induced expression of the IFN- β gene. *FEBS Lett* **358**, 225–229.
- Keckesova, Z., Ylilinen, L. M. & Towers, G. J. (2004). The human and African green monkey TRIM5 α genes encode Ref1 and Lv1 retroviral restriction factor activities. *Proc Natl Acad Sci U S A* **101**, 10780–10785.
- Kerscher, O., Felberbaum, R. & Hochstrasser, M. (2006). Modification of proteins by ubiquitin and ubiquitin-like proteins. *Annu Rev Cell Dev Biol* **22**, 159–180.
- Khanna, K. M., Lepisto, A. J., Decman, V. & Hendricks, R. L. (2004). Immune control of herpes simplex virus during latency. *Curr Opin Immunol* **16**, 463–469.
- Kim, M. J., Latham, A. G. & Krug, R. M. (2002). Human influenza viruses activate an interferon-independent transcription of cellular antiviral genes: outcome with influenza A virus is unique. *Proc Natl Acad Sci U S A* **99**, 10096–10101.
- Kim, C. S., Jung, J. H., Wakita, T., Yoon, S. K. & Jang, S. K. (2007). Monitoring the antiviral effect of alpha interferon on individual cells. *J Virol* **81**, 8814–8820.
- King, P. & Goodbourn, S. (1994). The β -interferon promoter responds to priming through multiple independent regulatory elements. *J Biol Chem* **269**, 30609–30615.
- Kirchweger, R., Ziegler, E., Lamphear, B. J., Waters, D., Liebig, H. D., Sommergruber, W., Sobrino, F., Hohenadl, C., Blaas, D. & other authors (1994). Foot-and-mouth disease virus leader proteinase: purification of the Lb form and determination of its cleavage site on eIF-4 γ . *J Virol* **68**, 5677–5684.
- Kochs, G., Trost, M., Janzen, C. & Haller, O. (1998). MxA GTPase: oligomerization and GTP-dependent interaction with viral RNP target structures. *Methods* **15**, 255–263.

- Kochs, G., Garcia-Sastre, A. & Martinez-Sobrido, L. (2007). Multiple anti-interferon actions of the influenza A virus NS1 protein. *J Virol* **81**, 7011–7021.
- Kolumam, G. A., Thomas, S., Thompson, L. J., Sprent, J. & Murali-Krishna, K. (2005). Type I interferons act directly on CD8 T cells to allow clonal expansion and memory formation in response to viral infection. *J Exp Med* **202**, 637–650.
- Komatsu, T., Takeuchi, K., Yokoo, J. & Gotoh, B. (2002). Sendai virus C protein impairs both phosphorylation and dephosphorylation processes of Stat1. *FEBS Lett* **511**, 139–144.
- Komatsu, T., Takeuchi, K. & Gotoh, B. (2007). Bovine parainfluenza virus type 3 accessory proteins that suppress beta interferon production. *Microbes Infect* **9**, 954–962.
- Kopecky-Bromberg, S. A., Martinez-Sobrido, L., Frieman, M., Baric, R. A. & Palese, P. (2007). Severe acute respiratory syndrome coronavirus open reading frame (ORF) 3b, ORF 6, and nucleocapsid proteins function as interferon antagonists. *J Virol* **81**, 548–557.
- Kozuka, Y., Yamashita, Y., Kawano, M., Tsurudome, M., Ito, M., Nishio, M., Komada, H. & Ito, Y. (2003). Identification of amino acids essential for the human parainfluenza type 2 virus V protein to lower the intracellular levels of the STAT2. *Virology* **317**, 208–219.
- Kozyrev, S. V., Lewen, S., Reddy, P. M., Pons-Estel, B., Witte, T., Junker, P., Lastrup, H., Gutierrez, C., Suarez, A. & other authors (2007). Structural insertion/deletion variation in IRF5 is associated with a risk haplotype and defines the precise IRF5 isoforms expressed in systemic lupus erythematosus. *Arthritis Rheum* **56**, 1234–1241.
- Krishnamurthy, S., Takimoto, T., Scroggs, R. A. & Portner, A. (2006). Differentially regulated interferon response determines the outcome of Newcastle disease virus infection in normal and tumor cell lines. *J Virol* **80**, 5145–5155.
- Krug, A., French, A. R., Barchet, W., Fischer, J. A., Dzionek, A., Pingel, J. T., Orihuela, M. M., Akira, S., Yokoyama, W. M. & Colonna, M. (2004). TLR9-dependent recognition of MCMV by IPC and DC generates coordinated cytokine responses that activate antiviral NK cell function. *Immunity* **21**, 107–119.
- Kubota, T., Yokosawa, N., Yokota, S. & Fujii, N. (2001). C terminal Cys-rich region of mumps virus structural V protein correlates with block of interferon α and γ signal transduction pathway through decrease of STAT 1- α . *Biochem Biophys Res Commun* **283**, 255–259.
- Kubota, T., Yokosawa, N., Yokota, S. & Fujii, N. (2002). Association of mumps virus V protein with RACK1 results in dissociation of STAT-1 from the alpha interferon receptor complex. *J Virol* **76**, 12676–12682.
- Kubota, T., Yokosawa, N., Yokota, S., Fujii, N., Tashiro, M. & Kato, A. (2005). Mumps virus V protein antagonizes interferon without the complete degradation of STAT1. *J Virol* **79**, 4451–4459.
- Kumar, K. P., McBride, K. M., Weaver, B. K., Dingwall, C. & Reich, N. C. (2000). Regulated nuclear-cytoplasmic localization of interferon regulatory factor 3, a subunit of double-stranded RNA-activated factor 1. *Mol Cell Biol* **20**, 4159–4168.
- Kumar, H., Kawai, T., Kato, H., Sato, S., Takahashi, K., Coban, C., Yamamoto, M., Uematsu, S., Ishii, K. J. & other authors (2006). Essential role of IPS-1 in innate immune responses against RNA viruses. *J Exp Med* **203**, 1795–1803.
- Kumar, P. P., Bischof, O., Purbey, P. K., Notani, D., Urlaub, H., Dejean, A. & Galande, S. (2007). Functional interaction between PML and SATB1 regulates chromatin-loop architecture and transcription of the MHC class I locus. *Nat Cell Biol* **9**, 45–56.
- Kurt-Jones, E. A., Popova, L., Kwinn, L., Haynes, L. M., Jones, L. P., Tripp, R. A., Walsh, E. E., Freeman, M. W., Golenbock, D. T. & other authors (2000). Pattern recognition receptors TLR4 and CD14 mediate response to respiratory syncytial virus. *Nat Immunol* **1**, 398–401.
- Kyogoku, C. & Tsuchiya, N. (2007). A compass that points to lupus: genetic studies on type I interferon pathway. *Genes Immun* **8**, 445–455.
- La Rocca, S. A., Herbert, R. J., Crooke, H., Drew, T. W., Wileman, T. E. & Powell, P. P. (2005). Loss of interferon regulatory factor 3 in cells infected with classical swine fever virus involves the N-terminal protease, Npro. *J Virol* **79**, 7239–7247.
- Langland, J. O., Pettiford, S., Jiang, B. & Jacobs, B. L. (1994). Products of the porcine group C rotavirus NSP3 gene bind specifically to double-stranded RNA and inhibit activation of the interferon-induced protein kinase PKR. *J Virol* **68**, 3821–3829.
- Langland, J. O., Cameron, J. M., Heck, M. C., Jancovich, J. K. & Jacobs, B. L. (2006). Inhibition of PKR by RNA and DNA viruses. *Virus Res* **119**, 100–110.
- Le Bon, A. & Tough, D. F. (2002). Links between innate and adaptive immunity via type I interferon. *Curr Opin Immunol* **14**, 432–436.
- Le Bon, A., Etchart, N., Rossmann, C., Ashton, M., Hou, S., Gewert, D., Borrow, P. & Tough, D. F. (2003). Cross-priming of CD8⁺ T cells stimulated by virus-induced type I interferon. *Nat Immunol* **4**, 1009–1015.
- Le Bon, A., Durand, V., Kamphuis, E., Thompson, C., Bulfone-Paus, S., Rossmann, C., Kalinke, U. & Tough, D. F. (2006). Direct stimulation of T cells by type I IFN enhances the CD8⁺ T cell response during cross-priming. *J Immunol* **176**, 4682–4689.
- Le May, N., Dubaele, S., Proietti De Santis, L., Billecocq, A., Bouloy, M. & Egly, J. M. (2004). TFIIF transcription factor, a target for the Rift Valley hemorrhagic fever virus. *Cell* **116**, 541–550.
- Lee, H. K., Dunzendorfer, S., Soldau, K. & Tobias, P. S. (2006). Double-stranded RNA-mediated TLR3 activation is enhanced by CD14. *Immunity* **24**, 153–163.
- Lee, H. K., Lund, J. M., Ramanathan, B., Mizushima, N. & Iwasaki, A. (2007). Autophagy-dependent viral recognition by plasmacytoid dendritic cells. *Science* **315**, 1398–1401.
- Leib, D. A., Machalek, M. A., Williams, B. R., Silverman, R. H. & Virgin, H. W. (2000). Specific phenotypic restoration of an attenuated virus by knockout of a host resistance gene. *Proc Natl Acad Sci U S A* **97**, 6097–6101.
- Lenschow, D. J., Giannakopoulos, N. V., Gunn, L. J., Johnston, C., O'Guin, A. K., Schmidt, R. E., Levine, B. & Virgin, H. W., IV (2005). Identification of interferon-stimulated gene 15 as an antiviral molecule during Sindbis virus infection in vivo. *J Virol* **79**, 13974–13983.
- Lenschow, D. J., Lai, C., Frias-Staheli, N., Giannakopoulos, N. V., Lutz, A., Wolff, T., Osiak, A., Levine, B., Schmidt, R. E. & other authors (2007). IFN-stimulated gene 15 functions as a critical antiviral molecule against influenza, herpes, and Sindbis viruses. *Proc Natl Acad Sci U S A* **104**, 1371–1376.
- Leonard, G. T. & Sen, G. C. (1996). Effects of adenovirus E1A protein on interferon-signaling. *Virology* **224**, 25–33.
- Li, Q. & Verma, I. M. (2002). NF- κ B regulation in the immune system. *Nat Rev Immunol* **2**, 725–734.
- Li, X., Leung, S., Qureshi, S., Darnell, J. E., Jr & Stark, G. R. (1996). Formation of STAT1–STAT2 heterodimers and their role in the activation of IRF-1 gene transcription by interferon- α . *J Biol Chem* **271**, 5790–5794.
- Li, S., Labrecque, S., Gauzzi, M. C., Cuddihy, A. R., Wong, A. H., Pellegrini, S., Matlashewski, G. J. & Koromilas, A. E. (1999). The human papillomavirus (HPV)-18 E6 oncoprotein physically associates with Tyk2 and impairs Jak-STAT activation by interferon- α . *Oncogene* **18**, 5727–5737.
- Li, W., Ross-Smith, N., Proud, C. G. & Belsham, G. J. (2001a). Cleavage of translation initiation factor 4AI (eIF4AI) but not eIF4AII

- by foot-and-mouth disease virus 3C protease: identification of the eIF4AI cleavage site. *FEBS Lett* **507**, 1–5.
- Li, Y., Chen, Z. Y., Wang, W., Baker, C. C. & Krug, R. M. (2001b). The 3'-end-processing factor CPSF is required for the splicing of single-intron pre-mRNAs in vivo. *RNA* **7**, 920–931.
- Li, C., Ni, C. Z., Havert, M. L., Cabezas, E., He, J., Kaiser, D., Reed, J. C., Satterthwait, A. C., Cheng, G. & Ely, K. R. (2002). Downstream regulator TANK binds to the CD40 recognition site on TRAF3. *Structure* **10**, 403–411.
- Li, K., Foy, E., Ferreon, J. C., Nakamura, M., Ferreon, A. C., Ikeda, M., Ray, S. C., Gale, M., Jr & Lemon, S. M. (2005a). Immune evasion by hepatitis C virus NS3/4A protease-mediated cleavage of the Toll-like receptor 3 adaptor protein TRIF. *Proc Natl Acad Sci U S A* **102**, 2992–2997.
- Li, Q., Withoff, S. & Verma, I. M. (2005b). Inflammation-associated cancer: NF- κ B is the lynchpin. *Trends Immunol* **26**, 318–325.
- Li, X. D., Sun, L., Seth, R. B., Pineda, G. & Chen, Z. J. (2005c). Hepatitis C virus protease NS3/4A cleaves mitochondrial antiviral signaling protein off the mitochondria to evade innate immunity. *Proc Natl Acad Sci U S A* **102**, 17717–17722.
- Li, H., Kobayashi, M., Blonska, M., You, Y. & Lin, X. (2006a). Ubiquitination of RIP is required for tumor necrosis factor α -induced NF- κ B activation. *J Biol Chem* **281**, 13636–13643.
- Li, S., Min, J. Y., Krug, R. M. & Sen, G. C. (2006b). Binding of the influenza A virus NS1 protein to PKR mediates the inhibition of its activation by either PACT or double-stranded RNA. *Virology* **349**, 13–21.
- Li, Q., Means, R., Lang, S. & Jung, J. U. (2007). Downregulation of gamma interferon receptor 1 by Kaposi's sarcoma-associated herpesvirus K3 and K5. *J Virol* **81**, 2117–2127.
- Lin, R., Heylbroeck, C., Pitha, P. M. & Hiscott, J. (1998). Virus-dependent phosphorylation of the IRF-3 transcription factor regulates nuclear translocation, transactivation potential, and proteasome-mediated degradation. *Mol Cell Biol* **18**, 2986–2996.
- Lin, R., Genin, P., Mamane, Y. & Hiscott, J. (2000). Selective DNA binding and association with the CREB binding protein coactivator contribute to differential activation of alpha/beta interferon genes by interferon regulatory factors 3 and 7. *Mol Cell Biol* **20**, 6342–6353.
- Lin, R., Genin, P., Mamane, Y., Sgarbanti, M., Battistini, A., Harrington, W. J., Jr, Barber, G. N. & Hiscott, J. (2001). HHV-8 encoded vIRF-1 represses the interferon antiviral response by blocking IRF-3 recruitment of the CBP/p300 coactivators. *Oncogene* **20**, 800–811.
- Lin, R., Noyce, R. S., Collins, S. E., Everett, R. D. & Mossman, K. L. (2004a). The herpes simplex virus ICP0 RING finger domain inhibits IRF3- and IRF7-mediated activation of interferon-stimulated genes. *J Virol* **78**, 1675–1684.
- Lin, R. J., Liao, C. L., Lin, E. & Lin, Y. L. (2004b). Blocking of the alpha interferon-induced Jak-Stat signaling pathway by Japanese encephalitis virus infection. *J Virol* **78**, 9285–9294.
- Lin, R., Lacoste, J., Nakhaei, P., Sun, Q., Yang, L., Paz, S., Wilkinson, P., Julkunen, I., Vitour, D. & other authors (2006a). Dissociation of a MAVS/IPS-1/VISA/Cardif- $IKK\epsilon$ molecular complex from the mitochondrial outer membrane by hepatitis C virus NS3-4A proteolytic cleavage. *J Virol* **80**, 6072–6083.
- Lin, R. J., Chang, B. L., Yu, H. P., Liao, C. L. & Lin, Y. L. (2006b). Blocking of interferon-induced Jak-Stat signaling by Japanese encephalitis virus NS5 through a protein tyrosine phosphatase-mediated mechanism. *J Virol* **80**, 5908–5918.
- Lin, W., Kim, S. S., Yeung, E., Kamegaya, Y., Blackard, J. T., Kim, K. A., Holtzman, M. J. & Chung, R. T. (2006c). Hepatitis C virus core protein blocks interferon signaling by interaction with the STAT1 SH2 domain. *J Virol* **80**, 9226–9235.
- Ling, P. D., Peng, R. S., Nakajima, A., Yu, J. H., Tan, J., Moses, S. M., Yang, W. H., Zhao, B., Kieff, E. & other authors (2005). Mediation of Epstein-Barr virus EBNA-LP transcriptional coactivation by Sp100. *EMBO J* **24**, 3565–3575.
- Liptakova, H., Kontsekova, E., Alcamí, A., Smith, G. L. & Kontsek, P. (1997). Analysis of an interaction between the soluble vaccinia virus-coded type I interferon (IFN)-receptor and human IFN- α 1 and IFN- α 2. *Virology* **232**, 86–90.
- Liu, W. J., Chen, H. B., Wang, X. J., Huang, H. & Khromykh, A. A. (2004). Analysis of adaptive mutations in Kunjin virus replicon RNA reveals a novel role for the flavivirus nonstructural protein NS2A in inhibition of beta interferon promoter-driven transcription. *J Virol* **78**, 12225–12235.
- Liu, W. J., Wang, X. J., Mokhonov, V. V., Shi, P. Y., Randall, R. & Khromykh, A. A. (2005). Inhibition of interferon signaling by the New York 99 strain and Kunjin subtype of West Nile virus involves blockage of STAT1 and STAT2 activation by nonstructural proteins. *J Virol* **79**, 1934–1942.
- Liu, W. J., Wang, X. J., Clark, D. C., Lobigs, M., Hall, R. A. & Khromykh, A. A. (2006). A single amino acid substitution in the West Nile virus nonstructural protein NS2A disables its ability to inhibit alpha/beta interferon induction and attenuates virus virulence in mice. *J Virol* **80**, 2396–2404.
- Loo, Y. M., Owen, D. M., Li, K., Erickson, A. K., Johnson, C. L., Fish, P. M., Carney, D. S., Wang, T., Ishida, H. & other authors (2006). Viral and therapeutic control of IFN- β promoter stimulator 1 during hepatitis C virus infection. *Proc Natl Acad Sci U S A* **103**, 6001–6006.
- Look, D. C., Roswit, W. T., Frick, A. G., Gris-Alevy, Y., Dickhaus, D. M., Walter, M. J. & Holtzman, M. J. (1998). Direct suppression of Stat1 function during adenoviral infection. *Immunity* **9**, 871–880.
- Lu, Y., Wambach, M., Katze, M. G. & Krug, R. M. (1995). Binding of the influenza virus NS1 protein to double-stranded RNA inhibits the activation of the protein kinase that phosphorylates the eIF-2 translation initiation factor. *Virology* **214**, 222–228.
- Lu, G., Reinert, J. T., Pitha-Rowe, I., Okumura, A., Kellum, M., Knobeloch, K. P., Hassel, B. & Pitha, P. M. (2006). ISG15 enhances the innate antiviral response by inhibition of IRF-3 degradation. *Cell Mol Biol (Noisy-le-grand)* **52**, 29–41.
- Lund, J. M., Alexopoulou, L., Sato, A., Karow, M., Adams, N. C., Gale, N. W., Iwasaki, A. & Flavell, R. A. (2004). Recognition of single-stranded RNA viruses by Toll-like receptor 7. *Proc Natl Acad Sci U S A* **101**, 5598–5603.
- Maher, S. G., Romero-Weaver, A. L., Scarzello, A. J. & Gamero, A. M. (2007). Interferon: cellular executioner or white knight? *Curr Med Chem* **14**, 1279–1289.
- Maitra, R. K., McMillan, N. A., Desai, S., McSwiggen, J., Hovanessian, A. G., Sen, G., Williams, B. R. & Silverman, R. H. (1994). HIV-1 TAR RNA has an intrinsic ability to activate interferon-inducible enzymes. *Virology* **204**, 823–827.
- Majumder, S., Zhou, L. Z., Chaturvedi, P., Babcock, G., Aras, S. & Ransohoff, R. M. (1998). p48/STAT-1 α -containing complexes play a predominant role in induction of IFN- γ -inducible protein, 10 kDa (IP-10) by IFN- γ alone or in synergy with TNF- α . *J Immunol* **161**, 4736–4744.
- Malakhov, M. P., Kim, K. I., Malakhova, O. A., Jacobs, B. S., Borden, E. C. & Zhang, D. E. (2003). High-throughput immunoblotting. Ubiquitin-like protein ISG15 modifies key regulators of signal transduction. *J Biol Chem* **278**, 16608–16613.
- Malakhova, O. A., Yan, M., Malakhov, M. P., Yuan, Y., Ritchie, K. J., Kim, K. I., Peterson, L. F., Shuai, K. & Zhang, D. E. (2003). Protein

- ISGylation modulates the JAK-STAT signaling pathway. *Genes Dev* 17, 455–460.
- Malathi, K., Dong, B., Gale, M., Jr & Silverman, R. H. (2007).** Small self-RNA generated by RNase L amplifies antiviral innate immunity. *Nature* 448, 816–819.
- Malim, M. H. (2006).** Natural resistance to HIV infection: the Vif-APOBEC interaction. *C R Biol* 329, 871–875.
- Marcus, P. I. (1983).** Interferon induction by viruses: one molecule of dsRNA as the threshold for interferon induction. *Interferon* 5, 115–180.
- Marcus, P. I. & Sekellick, M. J. (1977).** Defective interfering particles with covalently linked [+/-]RNA induce interferon. *Nature* 266, 815–819.
- Marie, I., Durbin, J. E. & Levy, D. E. (1998).** Differential viral induction of distinct interferon- α genes by positive feedback through interferon regulatory factor-7. *EMBO J* 17, 6660–6669.
- Marq, J. B., Brini, A., Kolakofsky, D. & Garcin, D. (2007).** Targeting of the Sendai virus C protein to the plasma membrane via a peptide-only membrane anchor. *J Virol* 81, 3187–3197.
- Marques, J. T., Rebouillat, D., Ramana, C. V., Murakami, J., Hill, J. E., Gudkov, A., Silverman, R. H., Stark, G. R. & Williams, B. R. (2005).** Down-regulation of p53 by double-stranded RNA modulates the antiviral response. *J Virol* 79, 11105–11114.
- Marrack, P., Kappler, J. & Mitchell, T. (1999).** Type I interferons keep activated T cells alive. *J Exp Med* 189, 521–530.
- Martens, S. & Howard, J. (2006).** The interferon-inducible GTPases. *Annu Rev Cell Dev Biol* 22, 559–589.
- Martinand, C., Salehzada, T., Silhol, M., Lebleu, B. & Bisbal, C. (1998).** RNase L inhibitor (RLI) antisense constructions block partially the down regulation of the 2-5A/RNase L pathway in encephalomyocarditis-virus-(EMCV)-infected cells. *Eur J Biochem* 254, 248–255.
- Martinand, C., Montavon, C., Salehzada, T., Silhol, M., Lebleu, B. & Bisbal, C. (1999).** RNase L inhibitor is induced during human immunodeficiency virus type 1 infection and down regulates the 2-5A/RNase L pathway in human T cells. *J Virol* 73, 290–296.
- Martinez-Costas, J., Gonzalez-Lopez, C., Vakharia, V. N. & Benavente, J. (2000).** Possible involvement of the double-stranded RNA-binding core protein sigmaA in the resistance of avian reovirus to interferon. *J Virol* 74, 1124–1131.
- Matsumoto, M., Kikkawa, S., Kohase, M., Miyake, K. & Seya, T. (2002).** Establishment of a monoclonal antibody against human Toll-like receptor 3 that blocks double-stranded RNA-mediated signaling. *Biochem Biophys Res Commun* 293, 1364–1369.
- Matsuyama, T., Kimura, T., Kitagawa, M., Pfeffer, K., Kawakami, T., Watanabe, N., Kundig, T., Amakawa, R., Kishihara, K. & other authors (1993).** Targeted disruption of IRF-1 or IRF-2 results in abnormal type I IFN gene induction and aberrant lymphocyte development. *Cell* 75, 83–97.
- McFadden, G. (2005).** Poxvirus tropism. *Nat Rev Microbiol* 3, 201–213.
- Meager, A., Visvalingam, K., Dilger, P., Bryan, D. & Wadhwa, M. (2005).** Biological activity of interleukins-28 and -29: comparison with type I interferons. *Cytokine* 31, 109–118.
- Mee, A. P., Dixon, J. A., Hoyland, J. A., Davies, M., Selby, P. L. & Mawer, E. B. (1998).** Detection of canine distemper virus in 100% of Paget's disease samples by in situ reverse transcriptase-polymerase chain reaction. *Bone* 23, 171–175.
- Melchjorsen, J., Jensen, S. B., Malmgaard, L., Rasmussen, S. B., Weber, F., Bowie, A. G., Matikainen, S. & Paludan, S. R. (2005).** Activation of innate defense against a paramyxovirus is mediated by RIG-I and TLR7 and TLR8 in a cell-type-specific manner. *J Virol* 79, 12944–12951.
- Melroe, G. T., DeLuca, N. A. & Knipe, D. M. (2004).** Herpes simplex virus 1 has multiple mechanisms for blocking virus-induced interferon production. *J Virol* 78, 8411–8420.
- Melroe, G. T., Silva, L., Schaffer, P. A. & Knipe, D. M. (2007).** Recruitment of activated IRF-3 and CBP/p300 to herpes simplex virus ICP0 nuclear foci: Potential role in blocking IFN- β induction. *Virology* 360, 305–321.
- Mennechet, F. J. & Uze, G. (2006).** Interferon- λ -treated dendritic cells specifically induce proliferation of FOXP3-expressing suppressor T cells. *Blood* 107, 4417–4423.
- Merika, M. & Thanos, D. (2001).** Enhanceosomes. *Curr Opin Genet Dev* 11, 205–208.
- Meylan, E., Burns, K., Hofmann, K., Blancheteau, V., Martinon, F., Kelliher, M. & Tschopp, J. (2004).** RIP1 is an essential mediator of Toll-like receptor 3-induced NF- κ B activation. *Nat Immunol* 5, 503–507.
- Meylan, E., Curran, J., Hofmann, K., Moradpour, D., Binder, M., Bartenschlager, R. & Tschopp, J. (2005).** Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted by hepatitis C virus. *Nature* 437, 1167–1172.
- Mibayashi, M., Martinez-Sobrido, L., Loo, Y. M., Cardenas, W. B., Gale, M., Jr & Garcia-Sastre, A. (2007).** Inhibition of retinoic acid-inducible gene I-mediated induction of beta interferon by the NS1 protein of influenza A virus. *J Virol* 81, 514–524.
- Miller, D. M., Rahill, B. M., Boss, J. M., Lairmore, M. D., Durbin, J. E., Waldman, J. W. & Sedmak, D. D. (1998).** Human cytomegalovirus inhibits major histocompatibility complex class II expression by disruption of the Jak/Stat pathway. *J Exp Med* 187, 675–683.
- Miller, D. M., Zhang, Y., Rahill, B. M., Waldman, W. J. & Sedmak, D. D. (1999).** Human cytomegalovirus inhibits IFN- α -stimulated antiviral and immunoregulatory responses by blocking multiple levels of IFN- α signal transduction. *J Immunol* 162, 6107–6113.
- Mills, B. G., Singer, F. R., Weiner, L. P., Suffin, S. C., Stabile, E. & Holst, P. (1984).** Evidence for both respiratory syncytial virus and measles virus antigens in the osteoclasts of patients with Paget's disease of bone. *Clin Orthop Relat Res* 183, 303–311.
- Min, J. Y. & Krug, R. M. (2006).** The primary function of RNA binding by the influenza A virus NS1 protein in infected cells: Inhibiting the 2'-5' oligo (A) synthetase/RNase L pathway. *Proc Natl Acad Sci U S A* 103, 7100–7105.
- Mineta, T., Rabkin, S. D., Yazaki, T., Hunter, W. D. & Martuza, R. L. (1995).** Attenuated multi-mutated herpes simplex virus-1 for the treatment of malignant gliomas. *Nat Med* 1, 938–943.
- Mohr, I. (2004).** Neutralizing innate host defenses to control viral translation in HSV-1 infected cells. *Int Rev Immunol* 23, 199–220.
- Morin, P., Braganca, J., Bandu, M. T., Lin, R., Hiscott, J., Doly, J. & Civas, A. (2002).** Preferential binding sites for interferon regulatory factors 3 and 7 involved in interferon-A gene transcription. *J Mol Biol* 316, 1009–1022.
- Mosca, J. D. & Pitha, P. M. (1986).** Transcriptional and posttranscriptional regulation of exogenous human beta interferon gene in simian cells defective in interferon synthesis. *Mol Cell Biol* 6, 2279–2283.
- Mossman, K., Upton, C. & McFadden, G. (1995a).** The myxoma virus-soluble interferon- γ receptor homolog, M-T7, inhibits interferon- γ in a species-specific manner. *J Biol Chem* 270, 3031–3038.
- Mossman, K., Upton, C., Buller, R. M. & McFadden, G. (1995b).** Species specificity of ectromelia virus and vaccinia virus interferon- γ binding proteins. *Virology* 208, 762–769.

- Mossman, K. L., Macgregor, P. F., Rozmus, J. J., Goryachev, A. B., Edwards, A. M. & Smiley, J. R. (2001). Herpes simplex virus triggers and then disarms a host antiviral response. *J Virol* **75**, 750–758.
- Mrkic, B., Pavlovic, J., Rulicke, T., Volpe, P., Buchholz, C. J., Hourcade, D., Atkinson, J. P., Aguzzi, A. & Cattaneo, R. (1998). Measles virus spread and pathogenesis in genetically modified mice. *J Virol* **72**, 7420–7427.
- Munoz-Fontela, C., Angel Garcia, M., Garcia-Cao, I., Collado, M., Arroyo, J., Esteban, M., Serrano, M. & Rivas, C. (2005). Resistance to viral infection of super p53 mice. *Oncogene* **24**, 3059–3062.
- Munoz-Jordan, J. L., Sanchez-Burgos, G. G., Laurent-Rolle, M. & Garcia-Sastre, A. (2003). Inhibition of interferon signaling by dengue virus. *Proc Natl Acad Sci U S A* **100**, 14333–14338.
- Munoz-Jordan, J. L., Laurent-Rolle, M., Ashour, J., Martinez-Sobrido, L., Ashok, M., Lipkin, W. I. & Garcia-Sastre, A. (2005). Inhibition of alpha/beta interferon signaling by the NS4B protein of flaviviruses. *J Virol* **79**, 8004–8013.
- Munz, C. (2007). Viral evasion of autophagy. *Cell Host Microbe* **1**, 9–11.
- Murphy, J. A., Duerst, R. J., Smith, T. J. & Morrison, L. A. (2003). Herpes simplex virus type 2 virion host shutoff protein regulates alpha/beta interferon but not adaptive immune responses during primary infection in vivo. *J Virol* **77**, 9337–9345.
- Murray, P. J. (2007). The JAK-STAT signaling pathway: input and output integration. *J Immunol* **178**, 2623–2629.
- Nagai, Y. & Kato, A. (2004). Accessory genes of the *Paramyxoviridae*, a large family of nonsegmented negative-strand RNA viruses, as a focus of active investigation by reverse genetics. *Curr Top Microbiol Immunol* **283**, 197–248.
- Najarro, P., Traktman, P. & Lewis, J. A. (2001). Vaccinia virus blocks gamma interferon signal transduction: viral VH1 phosphatase reverses Stat1 activation. *J Virol* **75**, 3185–3196.
- Nanda, S. K. & Baron, M. D. (2006). Rinderpest virus blocks type I and type II interferon action: role of structural and nonstructural proteins. *J Virol* **80**, 7555–7568.
- Navarro, L., Mowen, K., Rodems, S., Weaver, B., Reich, N., Spector, D. & David, M. (1998). Cytomegalovirus activates interferon immediate-early response gene expression and an interferon regulatory factor 3-containing interferon-stimulated response element-binding complex. *Mol Cell Biol* **18**, 3796–3802.
- Negishi, H., Fujita, Y., Yanai, H., Sakaguchi, S., Ouyang, X., Shinohara, M., Takayanagi, H., Ohba, Y., Taniguchi, T. & Honda, K. (2006). Evidence for licensing of IFN- γ -induced IFN regulatory factor 1 transcription factor by MyD88 in Toll-like receptor-dependent gene induction program. *Proc Natl Acad Sci U S A* **103**, 15136–15141.
- Netterwald, J. R., Jones, T. R., Britt, W. J., Yang, S. J., McCrone, I. P. & Zhu, H. (2004). Postattachment events associated with viral entry are necessary for induction of interferon-stimulated genes by human cytomegalovirus. *J Virol* **78**, 6688–6691.
- Nguyen, K. B., Salazar-Mather, T. P., Dalod, M. Y., Van Deusen, J. B., Wei, X. Q., Liew, F. Y., Caligiuri, M. A., Durbin, J. E. & Biron, C. A. (2002). Coordinated and distinct roles for IFN- $\alpha\beta$, IL-12, and IL-15 regulation of NK cell responses to viral infection. *J Immunol* **169**, 4279–4287.
- Nishio, M., Tsurudome, M., Ito, M., Kawano, M., Komada, H. & Ito, Y. (2001). High resistance of human parainfluenza type 2 virus protein-expressing cells to the antiviral and anti-cell proliferative activities of alpha/beta interferons: cysteine-rich V-specific domain is required for high resistance to the interferons. *J Virol* **75**, 9165–9176.
- Nishio, M., Garcin, D., Simonet, V. & Kolakofsky, D. (2002). The carboxyl segment of the mumps virus V protein associates with Stat proteins in vitro via a tryptophan-rich motif. *Virology* **300**, 92–99.
- Noah, D. L., Twu, K. Y. & Krug, R. M. (2003). Cellular antiviral responses against influenza A virus are countered at the posttranscriptional level by the viral NS1A protein via its binding to a cellular protein required for the 3' end processing of cellular pre-mRNAs. *Virology* **307**, 386–395.
- Nuovo, M. A., Nuovo, G. J., MacConnell, P., Forde, A. & Steiner, G. C. (1992). In situ analysis of Paget's disease of bone for measles-specific PCR-amplified cDNA. *Diagn Mol Pathol* **1**, 256–265.
- Offermann, M. K. (2007). Kaposi sarcoma herpesvirus-encoded interferon regulator factors. *Curr Top Microbiol Immunol* **312**, 185–209.
- Oganesyan, G., Saha, S. K., Guo, B., He, J. Q., Shahangian, A., Zarnegar, B., Perry, A. & Cheng, G. (2006). Critical role of TRAF3 in the Toll-like receptor-dependent and -independent antiviral response. *Nature* **439**, 208–211.
- Ohka, S., Igarashi, H., Nagata, N., Sakai, M., Koike, S., Nochi, T., Kiyono, H. & Nomoto, A. (2007). Establishment of a poliovirus oral infection system in human poliovirus receptor-expressing transgenic mice that are deficient in alpha/beta interferon receptor. *J Virol* **81**, 7902–7912.
- Ohno, S., Ono, N., Takeda, M., Takeuchi, K. & Yanagi, Y. (2004). Dissection of measles virus V protein in relation to its ability to block alpha/beta interferon signal transduction. *J Gen Virol* **85**, 2991–2999.
- Ohno, S., Ono, N., Seki, F., Takeda, M., Kura, S., Tsuzuki, T. & Yanagi, Y. (2007). Measles virus infection of SLAM (CD150) knockin mice reproduces tropism and immunosuppression in human infection. *J Virol* **81**, 1650–1659.
- Okumura, F., Zou, W. & Zhang, D. E. (2007). ISG15 modification of the eIF4E cognate 4EHP enhances cap structure-binding activity of 4EHP. *Genes Dev* **21**, 255–260.
- Olland, A. M., Jane-Valbuena, J., Schiff, L. A., Nibert, M. L. & Harrison, S. C. (2001). Structure of the reovirus outer capsid and dsRNA-binding protein $\sigma 3$ at 1.8 Å resolution. *EMBO J* **20**, 979–989.
- Onoguchi, K., Yoneyama, M., Takemura, A., Akira, S., Taniguchi, T., Namiki, H. & Fujita, T. (2007). Viral infections activate types I and III interferon genes through a common mechanism. *J Biol Chem* **282**, 7576–7581.
- Ooi, C. G., Walsh, C. A., Gallagher, J. A. & Fraser, W. D. (2000). Absence of measles virus and canine distemper virus transcripts in long-term bone marrow cultures from patients with Paget's disease of bone. *Bone* **27**, 417–421.
- Opitz, B., Rejaibi, A., Dauber, B., Eckhard, J., Vinzing, M., Schmeck, B., Hippenstiel, S., Suttrop, N. & Wolff, T. (2006). IFN β induction by influenza A virus is mediated by RIG-I which is regulated by the viral NS1 protein. *Cell Microbiol* **9**, 930–938.
- Orvedahl, A., Alexander, D., Tallozy, Z., Sun, Q., Wei, Y., Zhang, W., Burns, D., Leib, D. A. & Levine, B. (2007). HSV-1 ICP34.5 confers neurovirulence by targeting the Beclin 1 autophagy protein. *Cell Host Microbe* **1**, 23–35.
- Oshiumi, H., Sasai, M., Shida, K., Fujita, T., Matsumoto, M. & Seya, T. (2003). TIR-containing adapter molecule (TICAM)-2, a bridging adapter recruiting to Toll-like receptor 4 TICAM-1 that induces interferon- β . *J Biol Chem* **278**, 49751–49762.
- Paladino, P., Cummings, D. T., Noyce, R. S. & Mossman, K. L. (2006). The IFN-independent response to virus particle entry provides a first line of antiviral defense that is independent of TLRs and retinoic acid-inducible gene I. *J Immunol* **177**, 8008–8016.
- Palosaari, H., Parisien, J. P., Rodriguez, J. J., Ulane, C. M. & Horvath, C. M. (2003). STAT protein interference and suppression of cytokine

- signal transduction by measles virus V protein. *J Virol* **77**, 7635–7644.
- Pampin, M., Simonin, Y., Blondel, B., Percherancier, Y. & Chelbi-Alix, M. K. (2006).** Cross talk between PML and p53 during poliovirus infection: implications for antiviral defense. *J Virol* **80**, 8582–8592.
- Panne, D., Maniatis, T. & Harrison, S. C. (2004).** Crystal structure of ATF-2/c-Jun and IRF-3 bound to the interferon- β enhancer. *EMBO J* **23**, 4384–4393.
- Panne, D., Maniatis, T. & Harrison, S. C. (2007a).** An atomic model of the interferon- β enhanceosome. *Cell* **129**, 1111–1123.
- Panne, D., McWhirter, S. M., Maniatis, T. & Harrison, S. C. (2007b).** Interferon regulatory factor 3 is regulated by a dual phosphorylation-dependent switch. *J Biol Chem* **282**, 22816–22822.
- Parisien, J. P., Lau, J. F., Rodriguez, J. J., Sullivan, B. M., Moscona, A., Parks, G. D., Lamb, R. A. & Horvath, C. M. (2001).** The V protein of human parainfluenza virus 2 antagonizes type I interferon responses by destabilizing signal transducer and activator of transcription 2. *Virology* **283**, 230–239.
- Parisien, J. P., Lau, J. F. & Horvath, C. M. (2002a).** STAT2 acts as a host range determinant for species-specific paramyxovirus interferon antagonism and simian virus 5 replication. *J Virol* **76**, 6435–6441.
- Parisien, J. P., Lau, J. F., Rodriguez, J. J., Ulane, C. M. & Horvath, C. M. (2002b).** Selective STAT protein degradation induced by paramyxoviruses requires both STAT1 and STAT2 but is independent of alpha/beta interferon signal transduction. *J Virol* **76**, 4190–4198.
- Park, M. S., Shaw, M. L., Munoz-Jordan, J., Cros, J. F., Nakaya, T., Bouvier, N., Palese, P., Garcia-Sastre, A. & Basler, C. F. (2003).** Newcastle disease virus (NDV)-based assay demonstrates interferon-antagonist activity for the NDV V protein and the Nipah virus V, W, and C proteins. *J Virol* **77**, 1501–1511.
- Park, G. S., Morris, K. L., Hallett, R. G., Bloom, M. E. & Best, S. M. (2007).** Identification of residues critical for the interferon antagonist function of Langkat virus NS5 reveals a role for the RNA-dependent RNA polymerase domain. *J Virol* **81**, 6936–6946.
- Patel, C. V., Handy, I., Goldsmith, T. & Patel, R. C. (2000).** PACT, a stress-modulated cellular activator of interferon-induced double-stranded RNA-activated protein kinase, PKR. *J Biol Chem* **275**, 37993–37998.
- Paulus, C., Krauss, S. & Nevels, M. (2006).** A human cytomegalovirus antagonist of type I IFN-dependent signal transducer and activator of transcription signaling. *Proc Natl Acad Sci U S A* **103**, 3840–3845.
- Paun, A. & Pitha, P. M. (2007).** The IRF family, revisited. *Biochimie* **89**, 744–753.
- Pedersen, I. M., Cheng, G., Wieland, S., Volinia, S., Croce, C. M., Chisari, F. V. & David, M. (2007).** Interferon modulation of cellular microRNAs as an antiviral mechanism. *Nature* **449**, 919–922.
- Peng, G., Lei, K. J., Jin, W., Greenwell-Wild, T. & Wahl, S. M. (2006).** Induction of APOBEC3 family proteins, a defensive maneuver underlying interferon-induced anti-HIV-1 activity. *J Exp Med* **203**, 41–46.
- Perou, C. M., Jeffrey, S. S., van de Rijn, M., Rees, C. A., Eisen, M. B., Ross, D. T., Pergamenschikov, A., Williams, C. F., Zhu, S. X. & other authors (1999).** Distinctive gene expression patterns in human mammary epithelial cells and breast cancers. *Proc Natl Acad Sci U S A* **96**, 9212–9217.
- Perry, S. T. & Compton, T. (2006).** Kaposi's sarcoma-associated herpesvirus virions inhibit interferon responses induced by envelope glycoprotein gpK8.1. *J Virol* **80**, 11105–11114.
- Peters, K. L., Smith, H. L., Stark, G. R. & Sen, G. C. (2002).** IRF-3-dependent, NF κ B- and JNK-independent activation of the 561 and IFN- β genes in response to double-stranded RNA. *Proc Natl Acad Sci U S A* **99**, 6322–6327.
- Petricoin, E. F., III, Ito, S., Williams, B. L., Audet, S., Stancato, L. F., Gamero, A., Clouse, K., Grimley, P., Weiss, A. & other authors (1997).** Antiproliferative action of interferon- α requires components of T-cell-receptor signalling. *Nature* **390**, 629–632.
- Pichlmair, A., Schulz, O., Tan, C. P., Naslund, T. I., Liljestrom, P., Weber, F. & Reis e Sousa, C. (2006).** RIG-I-mediated antiviral responses to single-stranded RNA bearing 5'-phosphates. *Science* **314**, 997–1001.
- Pitha, P. M. (2000).** Introduction: interferon's connection to cancer. *Semin Cancer Biol* **10**, 69–72.
- Platanias, L. C. (2005).** Mechanisms of type-I- and type-II-interferon-mediated signalling. *Nat Rev Immunol* **5**, 375–386.
- Pomerantz, J. L. & Baltimore, D. (1999).** NF- κ B activation by a signaling complex containing TRAF2, TANK and TBK1, a novel IKK-related kinase. *EMBO J* **18**, 6694–6704.
- Poole, E., He, B., Lamb, R. A., Randall, R. E. & Goodbourn, S. (2002).** The V proteins of simian virus 5 and other paramyxoviruses inhibit induction of interferon- β . *Virology* **303**, 33–46.
- Poppers, J., Mulvey, M., Khoo, D. & Mohr, I. (2000).** Inhibition of PKR activation by the proline-rich RNA binding domain of the herpes simplex virus type 1 Us11 protein. *J Virol* **74**, 11215–11221.
- Poppers, J., Mulvey, M., Perez, C., Khoo, D. & Mohr, I. (2003).** Identification of a lytic-cycle Epstein-Barr virus gene product that can regulate PKR activation. *J Virol* **77**, 228–236.
- Porter, F. W., Bochkov, Y. A., Albee, A. J., Wiese, C. & Palmenberg, A. C. (2006).** A picornavirus protein interacts with Ran-GTPase and disrupts nucleocytoplasmic transport. *Proc Natl Acad Sci U S A* **103**, 12417–12422.
- Powell, P. P., Dixon, L. K. & Parkhouse, R. M. (1996).** An I κ B homolog encoded by African swine fever virus provides a novel mechanism for downregulation of proinflammatory cytokine responses in host macrophages. *J Virol* **70**, 8527–8533.
- Prakash, A., Smith, E., Lee, C. K. & Levy, D. E. (2005).** Tissue-specific positive feedback requirements for production of type I interferon following virus infection. *J Biol Chem* **280**, 18651–18657.
- Precious, B., Childs, K., Fitzpatrick-Swallow, V., Goodbourn, S. & Randall, R. E. (2005a).** Simian virus 5 V protein acts as an adaptor, linking DDB1 to STAT2, to facilitate the ubiquitination of STAT1. *J Virol* **79**, 13434–13441.
- Precious, B., Young, D. F., Andrejeva, L., Goodbourn, S. & Randall, R. E. (2005b).** In vitro and in vivo specificity of ubiquitination and degradation of STAT1 and STAT2 by the V proteins of the paramyxoviruses simian virus 5 and human parainfluenza virus type 2. *J Gen Virol* **86**, 151–158.
- Prentice, E., Jerome, W. G., Yoshimori, T., Mizushima, N. & Denison, M. R. (2004).** Coronavirus replication complex formation utilizes components of cellular autophagy. *J Biol Chem* **279**, 10136–10141.
- Preston, C. M. & Nicholl, M. J. (2006).** Role of the cellular protein hDaxx in human cytomegalovirus immediate-early gene expression. *J Gen Virol* **87**, 1113–1121.
- Preston, C. M., Harman, A. N. & Nicholl, M. J. (2001).** Activation of interferon response factor-3 in human cells infected with herpes simplex virus type 1 or human cytomegalovirus. *J Virol* **75**, 8909–8916.
- Puehler, F., Weining, K. C., Symons, J. A., Smith, G. L. & Staeheli, P. (1998).** Vaccinia virus-encoded cytokine receptor binds and neutralizes chicken interferon-gamma. *Virology* **248**, 231–240.
- Qing, J., Liu, C., Choy, L., Wu, R.-Y., Pagano, J. S. & Derynck, R. (2004).** Transforming growth factor β /Smad3 signaling regulates

- IRF-7 function and transcriptional activation of the beta interferon promoter. *Mol Cell Biol* **24**, 1411–1425.
- Ralston, S. H., Afzal, M. A., Helfrich, M. H., Fraser, W. D., Gallagher, J. A., Mee, A. & Rima, B. (2007). Multicentre blinded analysis of RT-PCR detection methods for paramyxoviruses in relation to Paget's disease of bone. *J Bone Miner Res* **22**, 569–577.
- Ramaswamy, M., Shi, L., Monick, M. M., Hunninghake, G. W. & Look, D. C. (2004). Specific inhibition of type I interferon signal transduction by respiratory syncytial virus. *Am J Respir Cell Mol Biol* **30**, 893–900.
- Ramaswamy, M., Shi, L., Varga, S. M., Barik, S., Behlke, M. A. & Look, D. C. (2006). Respiratory syncytial virus nonstructural protein 2 specifically inhibits type I interferon signal transduction. *Virology* **344**, 328–339.
- Randall, G., Chen, L., Panis, M., Fischer, A. K., Lindenbach, B. D., Sun, J., Heathcote, J., Rice, C. M., Edwards, A. M. & McGilvray, I. D. (2006). Silencing of USP18 potentiates the antiviral activity of interferon against hepatitis C virus infection. *Gastroenterology* **131**, 1584–1591.
- Rani, M. R. & Ransohoff, R. M. (2005). Alternative and accessory pathways in the regulation of IFN- β -mediated gene expression. *J Interferon Cytokine Res* **25**, 788–798.
- Reddy, S. V., Menaa, C., Singer, F. R., Cundy, T., Cornish, J., Whyte, M. P. & Roodman, G. D. (1999). Measles virus nucleocapsid transcript expression is not restricted to the osteoclast lineage in patients with Paget's disease of bone. *Exp Hematol* **27**, 1528–1532.
- Reich, N. C. & Liu, L. (2006). Tracking STAT nuclear traffic. *Nat Rev Immunol* **6**, 602–612.
- Reid, S. P., Leung, L. W., Hartman, A. L., Martinez, O., Shaw, M. L., Carbonnelle, C., Volchkov, V. E., Nichol, S. T. & Basler, C. F. (2006). Ebola virus VP24 binds karyopherin $\alpha 1$ and blocks STAT1 nuclear accumulation. *J Virol* **80**, 5156–5167.
- Revilla, Y., Callejo, M., Rodriguez, J. M., Culebras, E., Nogal, M. L., Salas, M. L., Vinuela, E. & Fresno, M. (1998). Inhibition of nuclear factor κB activation by a virus-encoded I κB -like protein. *J Biol Chem* **273**, 5405–5411.
- Rezaee, S. A., Cunningham, C., Davison, A. J. & Blackbourn, D. J. (2006). Kaposi's sarcoma-associated herpesvirus immune modulation: an overview. *J Gen Virol* **87**, 1781–1804.
- Richt, J. A., Lekcharoensuk, P., Lager, K. M., Vincent, A. L., Loiacono, C. M., Janke, B. H., Wu, W. H., Yoon, K. J., Webby, R. J. & other authors (2006). Vaccination of pigs against swine influenza viruses by using an NS1-truncated modified live-virus vaccine. *J Virol* **80**, 11009–11018.
- Rima, B. K., Gassen, U., Helfrich, M. H. & Ralston, S. H. (2002). The pro and con of measles virus in Paget's disease: con. *J Bone Miner Res* **17**, 2290–2292.
- Ritchie, K. J., Hahn, C. S., Kim, K. I., Yan, M., Rosario, D., Li, L., de la Torre, J. C. & Zhang, D. E. (2004). Role of ISG15 protease UBP43 (USP18) in innate immunity to viral infection. *Nat Med* **10**, 1374–1378.
- Rivera, J., Abrams, C., Hernaez, B., Alcazar, A., Escribano, J. M., Dixon, L. & Alonso, C. (2007). The MyD116 African swine fever virus homologue interacts with the catalytic subunit of protein phosphatase 1 and activates its phosphatase activity. *J Virol* **81**, 2923–2929.
- Rivieccio, M. A., Suh, H. S., Zhao, Y., Zhao, M. L., Chin, K. C., Lee, S. C. & Brosnan, C. F. (2006). TLR3 ligation activates an antiviral response in human fetal astrocytes: a role for viperin/cig5. *J Immunol* **177**, 4735–4741.
- Rodriguez, J. J. & Horvath, C. M. (2004). Host evasion by emerging paramyxoviruses: Hendra virus and Nipah virus v proteins inhibit interferon signaling. *Viral Immunol* **17**, 210–219.
- Rodriguez, J. J., Parisien, J. P. & Horvath, C. M. (2002). Nipah virus V protein evades alpha and gamma interferons by preventing STAT1 and STAT2 activation and nuclear accumulation. *J Virol* **76**, 11476–11483.
- Rodriguez, J. J., Wang, L. F. & Horvath, C. M. (2003). Hendra virus V protein inhibits interferon signaling by preventing STAT1 and STAT2 nuclear accumulation. *J Virol* **77**, 11842–11845.
- Ronco, L. V., Karpova, A. Y., Vidal, M. & Howley, P. M. (1998). Human papillomavirus 16 E6 oncoprotein binds to interferon regulatory factor-3 and inhibits its transcriptional activity. *Genes Dev* **12**, 2061–2072.
- Rothenfusser, S., Goutagny, N., DiPerna, G., Gong, M., Monks, B. G., Schoenemeyer, A., Yamamoto, M., Akira, S. & Fitzgerald, K. A. (2005). The RNA helicase Lgp2 inhibits TLR-independent sensing of viral replication by retinoic acid-inducible gene-I. *J Immunol* **175**, 5260–5268.
- Rouille, Y., Helle, F., Delgrange, D., Roingard, P., Voisset, C., Blanchard, E., Belouzard, S., McKeating, J., Patel, A. H. & other authors (2006). Subcellular localization of hepatitis C virus structural proteins in a cell culture system that efficiently replicates the virus. *J Virol* **80**, 2832–2841.
- Ruggli, N., Bird, B. H., Liu, L., Bauhofer, O., Tratschin, J. D. & Hofmann, M. A. (2005). N^{pro} of classical swine fever virus is an antagonist of double-stranded RNA-mediated apoptosis and IFN- α/β induction. *Virology* **340**, 265–276.
- Ryzhakov, G. & Randow, F. (2007). SINTBAD, a novel component of innate antiviral immunity, shares a TBK1-binding domain with NAPI and TANK. *EMBO J* **26**, 3180–3190.
- Saffert, R. T. & Kalejta, R. F. (2006). Inactivating a cellular intrinsic immune defense mediated by Daxx is the mechanism through which the human cytomegalovirus pp71 protein stimulates viral immediate-early gene expression. *J Virol* **80**, 3863–3871.
- Saira, K., Zhou, Y. & Jones, C. (2007). The infected cell protein 0 encoded by bovine herpesvirus 1 (bICP0) induces degradation of interferon response factor 3 and, consequently, inhibits beta interferon promoter activity. *J Virol* **81**, 3077–3086.
- Saito, S., Ogino, T., Miyajima, N., Kato, A. & Kohase, M. (2002). Dephosphorylation failure of tyrosine-phosphorylated STAT1 in IFN-stimulated Sendai virus C protein-expressing cells. *Virology* **293**, 205–209.
- Saito, T., Hirai, R., Loo, Y. M., Owen, D., Johnson, C. L., Sinha, S. C., Akira, S., Fujita, T. & Gale, M., Jr (2007). Regulation of innate antiviral defenses through a shared repressor domain in RIG-I and LGP2. *Proc Natl Acad Sci U S A* **104**, 582–587.
- Sakuma, R., Mael, A. A. & Ikeda, Y. (2007). Interferon alpha enhances the TRIM5 α -mediated antiviral activities in human and rhesus monkey cells. *J Virol* **81**, 10201–10206.
- Samuel, C. E. (1998). Reoviruses and the interferon system. *Curr Top Microbiol Immunol* **233**, 125–145.
- Sarkar, S. N., Peters, K. L., Elco, C. P., Sakamoto, S., Pal, S. & Sen, G. C. (2004). Novel roles of TLR3 tyrosine phosphorylation and PI3 kinase in double-stranded RNA signaling. *Nat Struct Mol Biol* **11**, 1060–1067.
- Sarkis, P. T., Ying, S., Xu, R. & Yu, X. F. (2006). STAT1-independent cell type-specific regulation of antiviral APOBEC3G by IFN- α . *J Immunol* **177**, 4530–4540.
- Sasai, M., Oshiumi, H., Matsumoto, M., Inoue, N., Fujita, F., Nakanishi, M. & Seya, T. (2005). NF- κB -activating kinase-associated protein 1 participates in TLR3/Toll-IL-1 homology domain-containing adapter molecule-1-mediated IFN regulatory factor 3 activation. *J Immunol* **174**, 27–30.

- Sasai, M., Shingai, M., Funami, K., Yoneyama, M., Fujita, T., Matsumoto, M. & Seya, T. (2006). NAK-associated protein 1 participates in both the TLR3 and the cytoplasmic pathways in type I IFN induction. *J Immunol* **177**, 8676–8683.
- Sato, M., Hata, N., Asagiri, M., Nakaya, T., Taniguchi, T. & Tanaka, N. (1998). Positive feedback regulation of type I IFN genes by the IFN-inducible transcription factor IRF-7. *FEBS Lett* **441**, 106–110.
- Sato, M., Suemori, H., Hata, N., Asagiri, M., Ogasawara, K., Nakao, K., Nakaya, T., Katsuki, M., Noguchi, S. & other authors (2000). Distinct and essential roles of transcription factors IRF-3 and IRF-7 in response to viruses for IFN- α/β gene induction. *Immunity* **13**, 539–548.
- Sato, S., Sugiyama, M., Yamamoto, M., Watanabe, Y., Kawai, T., Takeda, K. & Akira, S. (2003). Toll/IL-1 receptor domain-containing adaptor inducing IFN- β (TRIF) associates with TNF receptor-associated factor 6 and TANK-binding kinase 1, and activates two distinct transcription factors, NF- κ B and IFN-regulatory factor-3, in the Toll-like receptor signaling. *J Immunol* **171**, 4304–4310.
- Satterly, N., Tsai, P. L., van Deursen, J., Nussenzweig, D. R., Wang, Y., Faria, P. A., Levay, A., Levy, D. E. & Fontoura, B. M. (2007). Influenza virus targets the mRNA export machinery and the nuclear pore complex. *Proc Natl Acad Sci U S A* **104**, 1853–1858.
- Scheuner, D., Patel, R., Wang, F., Lee, K., Kumar, K., Wu, J., Nilsson, A., Karin, M. & Kaufman, R. J. (2006). Double-stranded RNA-dependent protein kinase phosphorylation of the α -subunit of eukaryotic translation initiation factor 2 mediates apoptosis. *J Biol Chem* **281**, 21458–21468.
- Schlender, J., Bossert, B., Buchholz, U. & Conzelmann, K. K. (2000). Bovine respiratory syncytial virus nonstructural proteins NS1 and NS2 cooperatively antagonize alpha/beta interferon-induced antiviral response. *J Virol* **74**, 8234–8242.
- Schlender, J., Hornung, V., Finke, S., Gunthner-Biller, M., Marozin, S., Brzozka, K., Moghim, S., Endres, S., Hartmann, G. & Conzelmann, K. K. (2005). Inhibition of Toll-like receptor 7- and 9-mediated alpha/beta interferon production in human plasmacytoid dendritic cells by respiratory syncytial virus and measles virus. *J Virol* **79**, 5507–5515.
- Schmid, D., Dengjel, J., Schoor, O., Stevanovic, S. & Munz, C. (2006). Autophagy in innate and adaptive immunity against intracellular pathogens. *J Mol Med* **84**, 194–202.
- Schmitz, F., Heit, A., Guggemoos, S., Krug, A., Mages, J., Schiemann, M., Adler, H., Drexler, I., Haas, T. & other authors (2007). Interferon-regulatory-factor 1 controls Toll-like receptor 9-mediated IFN- β production in myeloid dendritic cells. *Eur J Immunol* **37**, 315–327.
- Schoenemeyer, A., Barnes, B. J., Mancl, M. E., Latz, E., Goutagny, N., Pitha, P. M., Fitzgerald, K. A. & Golenbock, D. T. (2005). The interferon regulatory factor, IRF5, is a central mediator of Toll-like receptor 7 signaling. *J Biol Chem* **280**, 17005–17012.
- Schulz, O., Diebold, S. S., Chen, M., Naslund, T. I., Nolte, M. A., Alexopoulou, L., Azuma, Y. T., Flavell, R. A., Liljestrom, P. & Reis e Sousa, C. (2005). Toll-like receptor 3 promotes cross-priming to virus-infected cells. *Nature* **433**, 887–892.
- Seth, R. B., Sun, L., Ea, C. K. & Chen, Z. J. (2005). Identification and characterization of MAVS, a mitochondrial antiviral signaling protein that activates NF- κ B and IRF 3. *Cell* **122**, 669–682.
- Shabman, R. S., Morrison, T. E., Moore, C., White, L., Suthar, M. S., Hueston, L., Rulli, N., Lidbury, B., Ting, J. P. & other authors (2007). Differential induction of type I interferon responses in myeloid dendritic cells by mosquito and mammalian-cell-derived alphaviruses. *J Virol* **81**, 237–247.
- Shaffer, J. A., Bellini, W. J. & Rota, P. A. (2003). The C protein of measles virus inhibits the type I interferon response. *Virology* **315**, 389–397.
- Sharma, S., tenOever, B. R., Grandvaux, N., Zhou, G. P., Lin, R. & Hiscott, J. (2003). Triggering the interferon antiviral response through an IKK-related pathway. *Science* **300**, 1148–1151.
- Shaw, M. L., Garcia-Sastre, A., Palese, P. & Basler, C. F. (2004). Nipah virus V and W proteins have a common STAT1-binding domain yet inhibit STAT1 activation from the cytoplasmic and nuclear compartments, respectively. *J Virol* **78**, 5633–5641.
- Shaw, M. L., Cardenas, W. B., Zamarin, D., Palese, P. & Basler, C. F. (2005). Nuclear localization of the Nipah virus W protein allows for inhibition of both virus- and Toll-like receptor 3-triggered signaling pathways. *J Virol* **79**, 6078–6088.
- Shin, Y. K., Liu, Q., Tikoo, S. K., Babiuk, L. A. & Zhou, Y. (2007). Influenza A virus NS1 protein activates the phosphatidylinositol 3-kinase (PI3K)/Akt pathway by direct interaction with the p85 subunit of PI3K. *J Gen Virol* **88**, 13–18.
- Shinohara, M. L., Lu, L., Bu, J., Werneck, M. B., Kobayashi, K. S., Glimcher, L. H. & Cantor, H. (2006). Osteopontin expression is essential for interferon- α production by plasmacytoid dendritic cells. *Nat Immunol* **7**, 498–506.
- Shisler, J. L. & Jin, X. L. (2004). The vaccinia virus K1L gene product inhibits host NF- κ B activation by preventing I κ B α degradation. *J Virol* **78**, 3553–3560.
- Sigurdsson, S., Padyukov, L., Kurreeman, F. A., Liljedahl, U., Wiman, A. C., Alfredsson, L., Toes, R., Ronnelid, J., Klareskog, L. & other authors (2007). Association of a haplotype in the promoter region of the interferon regulatory factor 5 gene with rheumatoid arthritis. *Arthritis Rheum* **56**, 2202–2210.
- Silverman, R. H. (2007). Viral encounters with OAS and RNase L during the IFN antiviral response. *J Virol* (in press). doi:10.1128/JVI.01471-07
- Smiley, J. R. (2004). Herpes simplex virus virion host shutoff protein: immune evasion mediated by a viral RNase? *J Virol* **78**, 1063–1068.
- Smith, G. L., Symons, J. A. & Alcami, A. (1998). Poxviruses: interfering with interferon. *Semin Virol* **8**, 409–418.
- Song, B., Javanbakht, H., Perron, M., Park, D. H., Stremlau, M. & Sodroski, J. (2005). Retrovirus restriction by TRIM5 α variants from Old World and New World primates. *J Virol* **79**, 3930–3937.
- Soros, V. B. & Greene, W. C. (2007). APOBEC3G and HIV-1: strike and counterstrike. *Curr HIV/AIDS Rep* **4**, 3–9.
- Spann, K. M., Tran, K. C. & Collins, P. L. (2005). Effects of nonstructural proteins NS1 and NS2 of human respiratory syncytial virus on interferon regulatory factor 3, NF- κ B, and proinflammatory cytokines. *J Virol* **79**, 5353–5362.
- Spiegel, M., Pichlmair, A., Martinez-Sobrido, L., Cros, J., Garcia-Sastre, A., Haller, O. & Weber, F. (2005). Inhibition of beta interferon induction by severe acute respiratory syndrome coronavirus suggests a two-step model for activation of interferon regulatory factor 3. *J Virol* **79**, 2079–2086.
- Spiropoulou, C. F., Albarino, C. G., Ksiazek, T. G. & Rollin, P. E. (2007). Andes and Prospect Hill hantaviruses differ in early induction of interferon although both can downregulate interferon signaling. *J Virol* **81**, 2769–2776.
- Spitkovsky, D., Hehner, S. P., Hofmann, T. G., Moller, A. & Schmitz, M. L. (2002). The human papillomavirus oncoprotein E7 attenuates NF- κ B activation by targeting the I κ B kinase complex. *J Biol Chem* **277**, 25576–25582.
- Srivastava, S. P., Kumar, K. U. & Kaufman, R. J. (1998). Phosphorylation of eukaryotic translation initiation factor 2 mediates apoptosis in response to activation of the double-stranded RNA-dependent protein kinase. *J Biol Chem* **273**, 2416–2423.

- Stack, J., Haga, I. R., Schroder, M., Bartlett, N. W., Maloney, G., Reading, P. C., Fitzgerald, K. A., Smith, G. L. & Bowie, A. G. (2005). Vaccinia virus protein A46R targets multiple Toll-like-interleukin-1 receptor adaptors and contributes to virulence. *J Exp Med* **201**, 1007–1018.
- Stancato, L. F., David, M., Carter-Su, C., Lerner, A. C. & Pratt, W. B. (1996). Preassociation of STAT1 with STAT2 and STAT3 in separate signalling complexes prior to cytokine stimulation. *J Biol Chem* **271**, 4134–4137.
- Stetson, D. B. & Medzhitov, R. (2006a). Recognition of cytosolic DNA activates an IRF3-dependent innate immune response. *Immunity* **24**, 93–103.
- Stetson, D. B. & Medzhitov, R. (2006b). Type I interferons in host defense. *Immunity* **25**, 373–381.
- Stock, N., Goodbourn, S. & Randall, R. E. (2005). The anti-interferon mechanisms of paramyxoviruses. In *Modulation of Host Gene Expression and Innate Immunity by Viruses*, pp. 115–139. Edited by P. Palese. New York: Springer.
- Stojdl, D. F., Lichty, B., Knowles, S., Marius, R., Atkins, H., Sonenberg, N. & Bell, J. C. (2000). Exploiting tumor-specific defects in the interferon pathway with a previously unknown oncolytic virus. *Nat Med* **6**, 821–825.
- Stojdl, D. F., Lichty, B. D., tenOever, B. R., Paterson, J. M., Power, A. T., Knowles, S., Marius, R., Reynard, J., Poliquin, L. & other authors (2003). VSV strains with defects in their ability to shutdown innate immunity are potent systemic anti-cancer agents. *Cancer Cell* **4**, 263–275.
- Stracker, T. H., Lee, D. V., Carson, C. T., Araujo, F. D., Ornelles, D. A. & Weitzman, M. D. (2005). Serotype-specific reorganization of the Mre11 complex by adenoviral E4orf3 proteins. *J Virol* **79**, 6664–6673.
- Strahle, L., Garcin, D. & Kolakofsky, D. (2006). Sendai virus defective-interfering genomes and the activation of interferon- β . *Virology* **351**, 101–111.
- Strahle, L., Marq, J.-B., Brini, A., Hausmann, S., Kolakofsky, D. & Garcin, D. (2007). Activation of the IFN β promoter by unnatural Sendai virus infection requires RIG-I and is inhibited by the viral C proteins. *J Virol* (in press). doi:10.1128/JVI.01300-07
- Stremelau, M., Owens, C. M., Perron, M. J., Kiessling, M., Autissier, P. & Sodroski, J. (2004). The cytoplasmic body component TRIM5 α restricts HIV-1 infection in Old World monkeys. *Nature* **427**, 848–853.
- Sun, S., Zhang, X., Tough, D. F. & Sprent, J. (1998). Type I interferon-mediated stimulation of T cells by CpG DNA. *J Exp Med* **188**, 2335–2342.
- Sun, Q., Sun, L., Liu, H. H., Chen, X., Seth, R. B., Forman, J. & Chen, Z. J. (2006). The specific and essential role of MAVS in antiviral innate immune responses. *Immunity* **24**, 633–642.
- Surh, C. D., Boyman, O., Purton, J. F. & Sprent, J. (2006). Homeostasis of memory T cells. *Immunol Rev* **211**, 154–163.
- Svitkin, Y. V., Herdy, B., Costa-Mattioli, M., Gingras, A. C., Raught, B. & Sonenberg, N. (2005). Eukaryotic translation initiation factor 4E availability controls the switch between cap-dependent and internal ribosomal entry site-mediated translation. *Mol Cell Biol* **25**, 10556–10565.
- Symons, J. A., Alcami, A. & Smith, G. L. (1995). Vaccinia virus encodes a soluble type I interferon receptor of novel structure and broad species specificity. *Cell* **81**, 551–560.
- Tabeta, K., Georgel, P., Janssen, E., Du, X., Hoebe, K., Crozat, K., Mudd, S., Shamel, L., Sovath, S. & other authors (2004). Toll-like receptors 9 and 3 as essential components of innate immune defense against mouse cytomegalovirus infection. *Proc Natl Acad Sci U S A* **101**, 3516–3521.
- Taima, K., Imaizumi, T., Yamashita, K., Ishikawa, A., Fujita, T., Yoshida, H., Takanashi, S., Okumura, K. & Satoh, K. (2006). Expression of IP-10/CXCL10 is upregulated by double-stranded RNA in BEAS-2B bronchial epithelial cells. *Respiration* **73**, 360–364.
- Tait, S. W., Reid, E. B., Greaves, D. R., Wileman, T. E. & Powell, P. P. (2000). Mechanism of inactivation of NF- κ B by a viral homologue of I κ B α . Signal-induced release of I κ B α results in binding of the viral homologue to NF- κ B. *J Biol Chem* **275**, 34656–34664.
- Takahashi, K., Kawai, T., Kumar, H., Sato, S., Yonehara, S. & Akira, S. (2006). Roles of caspase-8 and caspase-10 in innate immune responses to double-stranded RNA. *J Immunol* **176**, 4520–4524.
- Takaoka, A., Hayakawa, S., Yanai, H., Stoiber, D., Negishi, H., Kikuchi, H., Sasaki, S., Imai, K., Shibue, T. & other authors (2003). Integration of interferon- α/β signalling to p53 responses in tumour suppression and antiviral defence. *Nature* **424**, 516–523.
- Takaoka, A., Yanai, H., Kondo, S., Duncan, G., Negishi, H., Mizutani, T., Kano, S., Honda, K., Ohba, Y. & other authors (2005). Integral role of IRF-5 in the gene induction programme activated by Toll-like receptors. *Nature* **434**, 243–249.
- Takaoka, A., Wang, Z., Choi, M. K., Yanai, H., Negishi, H., Ban, T., Lu, Y., Miyagishi, M., Kodama, T. & other authors (2007). DAI (DLM-1/ZBP1) is a cytosolic DNA sensor and an activator of innate immune response. *Nature* **448**, 501–505.
- Takayanagi, H., Kim, S. & Taniguchi, T. (2002). Signaling crosstalk between RANKL and interferons in osteoclast differentiation. *Arthritis Res* **4** (Suppl. 3), S227–S232.
- Takayanagi, H., Sato, K., Takaoka, A. & Taniguchi, T. (2005). Interplay between interferon and other cytokine systems in bone metabolism. *Immunol Rev* **208**, 181–193.
- Takeuchi, K., Komatsu, T., Yokoo, J., Kato, A., Shioda, T., Nagai, Y. & Gotoh, B. (2001). Sendai virus C protein physically associates with Stat1. *Genes Cells* **6**, 545–557.
- Takeuchi, K., Kadota, S. I., Takeda, M., Miyajima, N. & Nagata, K. (2003). Measles virus V protein blocks interferon (IFN)- α/β but not IFN- γ signaling by inhibiting STAT1 and STAT2 phosphorylation. *FEBS Lett* **545**, 177–182.
- Talloczy, Z., Jiang, W., Virgin, H. W., IV, Leib, D. A., Scheuner, D., Kaufman, R. J., Eskelinen, E. L. & Levine, B. (2002). Regulation of starvation- and virus-induced autophagy by the eIF2 α kinase signaling pathway. *Proc Natl Acad Sci U S A* **99**, 190–195.
- Talloczy, Z., Virgin, H. W., IV & Levine, B. (2006). PKR-dependent autophagic degradation of herpes simplex virus type 1. *Autophagy* **2**, 24–29.
- Talon, J., Horvath, C. M., Polley, R., Basler, C. F., Muster, T., Palese, P. & Garcia-Sastre, A. (2000a). Activation of interferon regulatory factor 3 is inhibited by the influenza A virus NS1 protein. *J Virol* **74**, 7989–7996.
- Talon, J., Salvatore, M., O'Neill, R. E., Nakaya, Y., Zheng, H., Muster, T., Garcia-Sastre, A. & Palese, P. (2000b). Influenza A and B viruses expressing altered NS1 proteins: a vaccine approach. *Proc Natl Acad Sci U S A* **97**, 4309–4314.
- Tamura, T., Ueda, S., Yoshida, M., Matsuzaki, M., Mohri, H. & Okubo, T. (1996). Interferon- γ induces *Ice* gene expression and enhances cellular susceptibility to apoptosis in the U937 leukemia cell line. *Biochem Biophys Res Commun* **229**, 21–26.
- Tan, S. L. & Katze, M. G. (1998). Biochemical and genetic evidence for complex formation between the influenza A virus NS1 protein and the interferon-induced PKR protein kinase. *J Interferon Cytokine Res* **18**, 757–766.
- Tang, X., Gao, J. S., Guan, Y. J., McLane, K. E., Yuan, Z. L., Ramratnam, B. & Chin, Y. E. (2007). Acetylation-dependent signal transduction for type I interferon receptor. *Cell* **131**, 93–105.

- Teng, M. N., Whitehead, S. S., Bermingham, A., St Claire, M., Elkins, W. R., Murphy, B. R. & Collins, P. L. (2000). Recombinant respiratory syncytial virus that does not express the NS1 or M2-2 protein is highly attenuated and immunogenic in chimpanzees. *J Virol* **74**, 9317–9321.
- tenOever, B. R., Servant, M. J., Grandvaux, N., Lin, R. & Hiscott, J. (2002). Recognition of the measles virus nucleocapsid as a mechanism of IRF-3 activation. *J Virol* **76**, 3659–3669.
- tenOever, B. R., Ng, S. L., Chua, M. A., McWhirter, S. M., Garcia-Sastre, A. & Maniatis, T. (2007). Multiple functions of the IKK-related kinase IKK ϵ in interferon-mediated antiviral immunity. *Science* **315**, 1274–1278.
- Terenzi, F., Pal, S. & Sen, G. C. (2005). Induction and mode of action of the viral stress-inducible murine proteins, P56 and P54. *Virology* **340**, 116–124.
- Thomas, D., Blakqori, G., Wagner, V., Banholzer, M., Kessler, N., Elliott, R. M., Haller, O. & Weber, F. (2004). Inhibition of RNA polymerase II phosphorylation by a viral interferon antagonist. *J Biol Chem* **279**, 31471–31477.
- Toth, A. M., Zhang, P., Das, S., George, C. X. & Samuel, C. E. (2006). Interferon action and the double-stranded RNA-dependent enzymes ADAR1 adenosine deaminase and PKR protein kinase. *Prog Nucleic Acid Res Mol Biol* **81**, 369–434.
- Tough, D. F., Borrow, P. & Sprent, J. (1996). Induction of bystander T cell proliferation by viruses and type I interferon in vivo. *Science* **272**, 1947–1950.
- Towers, G. J. (2007). The control of viral infection by tripartite motif proteins and cyclophilin A. *Retrovirology* **4**, 40.
- Twu, J. S. & Schloemer, R. H. (1989). Transcription of the human beta interferon gene is inhibited by hepatitis B virus. *J Virol* **63**, 3065–3071.
- Uddin, S., Sassano, A., Deb, D. K., Verma, A., Majchrzak, B., Rahman, A., Malik, A. B., Fish, E. N. & Platanius, L. C. (2002). Protein kinase C- δ (PKC- δ) is activated by type I interferons and mediates phosphorylation of Stat1 on serine 727. *J Biol Chem* **277**, 14408–14416.
- Uematsu, S., Sato, S., Yamamoto, M., Hirotani, T., Kato, H., Takeshita, F., Matsuda, M., Coban, C., Ishii, K. J. & other authors (2005). Interleukin-1 receptor-associated kinase-1 plays an essential role for Toll-like receptor (TLR)7- and TLR9-mediated interferon- α induction. *J Exp Med* **201**, 915–923.
- Ulane, C. M. & Horvath, C. M. (2002). Paramyxoviruses SV5 and HPIV2 assemble STAT protein ubiquitin ligase complexes from cellular components. *Virology* **304**, 160–166.
- Ulane, C. M., Rodriguez, J. J., Parisien, J. P. & Horvath, C. M. (2003). STAT3 ubiquitylation and degradation by mumps virus suppress cytokine and oncogene signaling. *J Virol* **77**, 6385–6393.
- Ulane, C. M., Kentsis, A., Cruz, C. D., Parisien, J. P., Schneider, K. L. & Horvath, C. M. (2005). Composition and assembly of STAT-targeting ubiquitin ligase complexes: paramyxovirus V protein carboxyl terminus is an oligomerization domain. *J Virol* **79**, 10180–10189.
- Ullman, A. J., Reich, N. C. & Hearing, P. (2007). Adenovirus E4 ORF3 protein inhibits the interferon-mediated antiviral response. *J Virol* **81**, 4744–4752.
- Unterstab, G., Ludwig, S., Anton, A., Planz, O., Dauber, B., Krappmann, D., Heins, G., Ehrhardt, C. & Wolff, T. (2005). Viral targeting of the interferon- β -inducing Traf family member-associated NF- κ B activator (TANK)-binding kinase-1. *Proc Natl Acad Sci U S A* **102**, 13640–13645.
- Upton, C., Mossman, K. & McFadden, G. (1992). Encoding of a homolog of the IFN-gamma receptor by myxoma virus. *Science* **258**, 1369–1372.
- Usacheva, A., Smith, R., Minshall, R., Baida, G., Seng, S., Croze, E. & Colamonic, O. (2001). The WD motif-containing protein receptor for activated protein kinase C (RACK1) is required for recruitment and activation of signal transducer and activator of transcription 1 through the type I interferon receptor. *J Biol Chem* **276**, 22948–22953.
- Uze, G. & Monneron, D. (2007). IL-28 and IL-29: newcomers to the interferon family. *Biochimie* **89**, 729–734.
- Valarcher, J. F., Furze, J., Wyld, S., Cook, R., Conzelmann, K. K. & Taylor, G. (2003). Role of alpha/beta interferons in the attenuation and immunogenicity of recombinant bovine respiratory syncytial viruses lacking NS proteins. *J Virol* **77**, 8426–8439.
- Valchanova, R. S., Picard-Maureau, M., Budt, M. & Brune, W. (2006). Murine cytomegalovirus m142 and m143 are both required to block protein kinase R-mediated shutdown of protein synthesis. *J Virol* **80**, 10181–10190.
- van Boxel-Dezaire, A. H., Rani, M. R. & Stark, G. R. (2006). Complex modulation of cell type-specific signaling in response to type I interferons. *Immunity* **25**, 361–372.
- Van Cleve, W., Amaro-Carambot, E., Surman, S. R., Bekisz, J., Collins, P. L., Zoon, K. C., Murphy, B. R., Skiadopoulos, M. H. & Bartlett, E. J. (2006). Attenuating mutations in the P/C gene of human parainfluenza virus type 1 (HPIV1) vaccine candidates abrogate the inhibition of both induction and signaling of type I interferon (IFN) by wild-type HPIV1. *Virology* **352**, 61–73.
- van Pesch, V., van Eyll, O. & Michiels, T. (2001). The leader protein of Theiler's virus inhibits immediate-early alpha/beta interferon production. *J Virol* **75**, 7811–7817.
- Venkataraman, T., Valdes, M., Elsby, R., Kakuta, S., Caceres, G., Saijo, S., Iwakura, Y. & Barber, G. N. (2007). Loss of DEXD/H box RNA helicase LGP2 manifests disparate antiviral responses. *J Immunol* **178**, 6444–6455.
- Vidy, A., El Bougrini, J., Chelbi-Alix, M. K. & Blondel, D. (2007). The nucleocytoplasmic rabies virus P protein counteracts interferon signaling by inhibiting both nuclear accumulation and DNA binding of STAT1. *J Virol* **81**, 4255–4263.
- Wang, X., Li, M., Zheng, H., Muster, T., Palese, P., Beg, A. A. & Garcia-Sastre, A. (2000). Influenza A virus NS1 protein prevents activation of NF- κ B and induction of alpha/beta interferon. *J Virol* **74**, 11566–11573.
- Wang, C., Deng, L., Hong, M., Akkaraju, G. R., Inoue, J. & Chen, Z. J. (2001). TAK1 is a ubiquitin-dependent kinase of MKK and IKK. *Nature* **412**, 346–351.
- Wang, F., Ma, Y., Barrett, J. W., Gao, X., Loh, J., Barton, E., Virgin, H. W. & McFadden, G. (2004). Disruption of Erk-dependent type I interferon induction breaks the myxoma virus species barrier. *Nat Immunol* **5**, 1266–1274.
- Wang, X., Hinson, E. R. & Cresswell, P. (2007). The interferon-inducible protein viperin inhibits influenza virus release by perturbing lipid rafts. *Cell Host Microbe* **2**, 96–105.
- Wathelet, M. G., Lin, C. H., Parekh, B. S., Ronco, L. V., Howley, P. M. & Maniatis, T. (1998). Virus infection induces the assembly of coordinately activated transcription factors on the IFN- β enhancer in vivo. *Mol Cell* **1**, 507–518.
- Weber, F., Haller, O. & Kochs, G. (2000). MxA GTPase blocks reporter gene expression of reconstituted Thogoto virus ribonucleo-protein complexes. *J Virol* **74**, 560–563.
- Weber, F., Bridgen, A., Fazakerley, J. K., Streitenfeld, H., Kessler, N., Randall, R. E. & Elliott, R. M. (2002). Bunyamwera bunyavirus nonstructural protein NSs counteracts the induction of alpha/beta interferon. *J Virol* **76**, 7949–7955.
- Weber, F., Wagner, V., Rasmussen, S. B., Hartmann, R. & Paludan, S. R. (2006). Double-stranded RNA is produced by positive-strand

- RNA viruses and DNA viruses but not in detectable amounts by negative-strand RNA viruses. *J Virol* **80**, 5059–5064.
- Weihua, X., Ramanujam, S., Lindner, D. J., Kudaravalli, R. D., Freund, R. & Kalvakolanu, D. V. (1998). The polyoma virus T antigen interferes with interferon-inducible gene expression. *Proc Natl Acad Sci U S A* **95**, 1085–1090.
- Whitlow, Z. W., Connor, J. H. & Lyles, D. S. (2006). Preferential translation of vesicular stomatitis virus mRNAs is conferred by transcription from the viral genome. *J Virol* **80**, 11733–11742.
- Whitten, T. M., Quets, A. T. & Schloemer, R. H. (1991). Identification of the hepatitis B virus factor that inhibits expression of the beta interferon gene. *J Virol* **65**, 4699–4704.
- Windisch, M. P., Frese, M., Kaul, A., Trippler, M., Lohmann, V. & Bartenschlager, R. (2005). Dissecting the interferon-induced inhibition of hepatitis C virus replication by using a novel host cell line. *J Virol* **79**, 13778–13793.
- Wollmann, G., Robek, M. D. & van den Pol, A. N. (2007). Variable deficiencies in the interferon response enhance susceptibility to vesicular stomatitis virus oncolytic actions in glioblastoma cells but not in normal human glial cells. *J Virol* **81**, 1479–1491.
- Wright, P. F., Karron, R. A., Madhi, S. A., Treanor, J. J., King, J. C., O'Shea, A., Ikizler, M. R., Zhu, Y., Collins, P. L. & other authors (2006). The interferon antagonist NS2 protein of respiratory syncytial virus is an important virulence determinant for humans. *J Infect Dis* **193**, 573–581.
- Wu, C. J., Conze, D. B., Li, T., Srinivasula, S. M. & Ashwell, J. D. (2006). Sensing of Lys 63-linked polyubiquitination by NEMO is a key event in NF- κ B activation. *Nat Cell Biol* **8**, 398–406.
- Wullaert, A., Heyninck, K., Janssens, S. & Beyaert, R. (2006). Ubiquitin: tool and target for intracellular NF- κ B inhibitors. *Trends Immunol* **27**, 533–540.
- Xu, L. G., Wang, Y. Y., Han, K. J., Li, L. Y., Zhai, Z. & Shu, H. B. (2005). VISA is an adapter protein required for virus-triggered IFN- β signaling. *Mol Cell* **19**, 727–740.
- Xu, D., Brumm, K. & Zhang, L. (2006). The latent membrane protein 1 of Epstein-Barr virus (EBV) primes EBV latency cells for type I interferon production. *J Biol Chem* **281**, 9163–9169.
- Yalamanchili, P., Harris, K., Wimmer, E. & Dasgupta, A. (1996). Inhibition of basal transcription by poliovirus: a virus-encoded protease (3C^{PRO}) inhibits formation of TBP-TATA box complex in vitro. *J Virol* **70**, 2922–2929.
- Yamamoto, M., Sato, S., Mori, K., Hoshino, K., Takeuchi, O., Takeda, K. & Akira, S. (2002). Cutting edge: a novel Toll/IL-1 receptor domain-containing adapter that preferentially activates the IFN- β promoter in the Toll-like receptor signaling. *J Immunol* **169**, 6668–6672.
- Yamamoto, M., Sato, S., Hemmi, H., Uematsu, S., Hoshino, K., Kaisho, T., Takeuchi, O., Takeda, K. & Akira, S. (2003a). TRAM is specifically involved in the Toll-like receptor 4-mediated MyD88-independent signaling pathway. *Nat Immunol* **4**, 1144–1150.
- Yamamoto, M., Sato, S., Hemmi, H., Hoshino, K., Kaisho, T., Sanjo, H., Takeuchi, O., Sugiyama, M., Okabe, M. & other authors (2003b). Role of adaptor TRIF in the MyD88-independent Toll-like receptor signaling pathway. *Science* **301**, 640–643.
- Yamashita, K., Imaizumi, T., Taima, K., Fujita, T., Ishikawa, A., Yoshida, H., Oyama, C. & Satoh, K. (2005). Polyinosinic-polycytidylic acid induces the expression of GRO- α in BEAS-2B cells. *Inflammation* **29**, 17–21.
- Yanai, H., Chen, H. M., Inuzuka, T., Kondo, S., Mak, T. W., Takaoka, A., Honda, K. & Taniguchi, T. (2007). Role of IFN regulatory factor 5 transcription factor in antiviral immunity and tumor suppression. *Proc Natl Acad Sci U S A* **104**, 3402–3407.
- Yang, H., Ma, G., Lin, C. H., Orr, M. & Wathelot, M. G. (2004). Mechanism for transcriptional synergy between interferon regulatory factor (IRF)-3 and IRF-7 in activation of the *interferon- β* gene promoter. *Eur J Biochem* **271**, 3693–3703.
- Yap, M. W., Nisole, S., Lynch, C. & Stoye, J. P. (2004). Trim5 α protein restricts both HIV-1 and murine leukemia virus. *Proc Natl Acad Sci U S A* **101**, 10786–10791.
- Yasuda, K., Richez, C., Maciaszek, J. W., Agrawal, N., Akira, S., Marshak-Rothstein, A. & Rifkin, I. R. (2007). Murine dendritic cell type I IFN production induced by human IgG-RNA immune complexes is IFN regulatory factor (IRF)5 and IRF7 dependent and is required for IL-6 production. *J Immunol* **178**, 6876–6885.
- Yeow, W. S., Au, W. C., Juang, Y. T., Fields, C. D., Dent, C. L., Gewert, D. R. & Pitha, P. M. (2000). Reconstitution of virus-mediated expression of interferon α genes in human fibroblast cells by ectopic interferon regulatory factor-7. *J Biol Chem* **275**, 6313–6320.
- Ying, S., Zhang, X., Sarkis, P. T., Xu, R. & Yu, X. (2007). Cell-specific regulation of APOBEC3F by interferons. *Acta Biochim Biophys Sin (Shanghai)* **39**, 297–304.
- Yokota, S., Saito, H., Kubota, T., Yokosawa, N., Amano, K. & Fujii, N. (2003). Measles virus suppresses interferon- α signaling pathway: suppression of Jak1 phosphorylation and association of viral accessory proteins, C and V, with interferon- α receptor complex. *Virology* **306**, 135–146.
- Yokota, S., Yokosawa, N., Okabayashi, T., Suzutani, T., Miura, S., Jimbow, K. & Fujii, N. (2004). Induction of suppressor of cytokine signaling-3 by herpes simplex virus type 1 contributes to inhibition of the interferon signaling pathway. *J Virol* **78**, 6282–6286.
- Yoneyama, M., Kikuchi, M., Natsukawa, T., Shinobu, N., Imaizumi, T., Miyagishi, M., Taira, K., Akira, S. & Fujita, T. (2004). The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. *Nat Immunol* **5**, 730–737.
- Yoneyama, M., Kikuchi, M., Matsumoto, K., Imaizumi, T., Miyagishi, M., Taira, K., Foy, E., Loo, Y. M., Gale, M., Jr & other authors (2005). Shared and unique functions of the DExD/H-box helicases RIG-I, MDA5, and LGP2 in antiviral innate immunity. *J Immunol* **175**, 2851–2858.
- Young, D. F., Randall, R. E., Hoyle, J. A. & Souberbielle, B. E. (1990). Clearance of a persistent paramyxovirus infection is mediated by cellular immune responses but not by serum-neutralizing antibody. *J Virol* **64**, 5403–5411.
- Young, D. F., Didcock, L., Goodbourn, S. & Randall, R. E. (2000). Paramyxoviridae use distinct virus-specific mechanisms to circumvent the interferon response. *Virology* **269**, 383–390.
- Young, D. F., Chatziandreou, N., He, B., Goodbourn, S., Lamb, R. A. & Randall, R. E. (2001). Single amino acid substitution in the V protein of simian virus 5 differentiates its ability to block interferon signaling in human and murine cells. *J Virol* **75**, 3363–3370.
- Young, D. F., Andrejeva, L., Livingstone, A., Goodbourn, S., Lamb, R. A., Collins, P. L., Elliott, R. M. & Randall, R. E. (2003). Virus replication in engineered human cells that do not respond to interferons. *J Virol* **77**, 2174–2181.
- Yu, Y., Wang, S. E. & Hayward, G. S. (2005). The KSHV immediate-early transcription factor RTA encodes ubiquitin E3 ligase activity that targets IRF7 for proteasome-mediated degradation. *Immunity* **22**, 59–70.
- Yuan, W. & Krug, R. M. (2001). Influenza B virus NS1 protein inhibits conjugation of the interferon (IFN)-induced ubiquitin-like ISG15 protein. *EMBO J* **20**, 362–371.
- Yuan, H., Yoza, B. K. & Lyles, D. S. (1998). Inhibition of host RNA polymerase II-dependent transcription by vesicular stomatitis virus results from inactivation of TFIID. *Virology* **251**, 383–392.

- Zahn, R. C., Schelp, I., Utermohlen, O. & von Laer, D. (2007). A-to-G hypermutation in the genome of lymphocytic choriomeningitis virus. *J Virol* **81**, 457–464.
- Zhang, P. & Samuel, C. E. (2007). Protein kinase PKR plays a stimulus- and virus-dependent role in apoptotic death and virus multiplication in human cells. *J Virol* **81**, 8192–8200.
- Zhang, J. J., Vinkemeier, U., Gu, W., Chakravarti, D., Horvath, C. M. & Darnell, J. E., Jr (1996). Two contact regions between Stat1 and CBP/p300 in interferon gamma signaling. *Proc Natl Acad Sci U S A* **93**, 15092–15096.
- Zhang, J., Das, S. C., Kotalik, C., Pattnaik, A. K. & Zhang, L. (2004). The latent membrane protein 1 of Epstein-Barr virus establishes an antiviral state via induction of interferon-stimulated genes. *J Biol Chem* **279**, 46335–46342.
- Zhang, Y., Burke, C. W., Ryman, K. D. & Klimstra, W. B. (2007). Identification and characterization of interferon-induced proteins that inhibit alphavirus replication. *J Virol* **81**, 11246–11255.
- Zhao, C., Denison, C., Huibregtse, J. M., Gygi, S. & Krug, R. M. (2005). Human ISG15 conjugation targets both IFN-induced and constitutively expressed proteins functioning in diverse cellular pathways. *Proc Natl Acad Sci U S A* **102**, 10200–10205.
- Zhao, T., Yang, L., Sun, Q., Arguello, M., Ballard, D. W., Hiscott, J. & Lin, R. (2007). The NEMO adaptor bridges the nuclear factor- κ B and interferon regulatory factor signaling pathways. *Nat Immunol* **8**, 592–600.
- Zhou, Z., Hamming, O. J., Ank, N., Paludan, S. R., Nielsen, A. L. & Hartmann, R. (2007). Type III interferon (IFN) induces a type I IFN-like response in a restricted subset of cells through signaling pathways involving both the Jak-STAT pathway and the mitogen-activated protein kinases. *J Virol* **81**, 7749–7758.
- Zou, W., Kim, J. H., Handidu, A., Li, X., Kim, K. I., Yan, M., Li, J. & Zhang, D. E. (2007). Microarray analysis reveals that type I interferon strongly increases the expression of immune-response related genes in Ubp43 (Usp18) deficient macrophages. *Biochem Biophys Res Commun* **356**, 193–199.