

Taxonomy and biogeography of the Nearctic *Raphia* Hübner (Lepidoptera, Noctuidae, Raphiinae)

B. Christian Schmidt¹, Gary G. Anweiler²

¹ Canadian Food Inspection Agency, Canadian National Collection of Insects, Arachnids and Nematodes, K.W. Neatby Bldg., 960 Carling Ave., Ottawa, ON, Canada K1A 0C6 ² E.H. Strickland Entomological Museum, 218 Earth Sciences Building, Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada, T6G 2E9

Corresponding author: B. Christian Schmidt (chris.schmidt@inspection.gc.ca)

Academic editor: D. Lafontaine | Received 14 March 2014 | Accepted 19 April 2014 | Published 27 June 2014

<http://zoobank.org/4DB3DA2D-21B1-4D26-9544-B4008028D304>

Citation: Schmidt BC, Anweiler GG (2014) Taxonomy and biogeography of the Nearctic *Raphia* Hübner (Lepidoptera, Noctuidae, Raphiinae). In: Schmidt BC, Lafontaine JD (Eds) Contributions to the systematics of New World macro-moths V. ZooKeys 421: 91–113. doi: 10.3897/zookeys.421.7517

Abstract

The taxonomic status and biogeography of the North American *Raphia* species is reviewed using adult morphology, larval host plants, geographic phenotypic variation, and variation of mtDNA COI barcode sequences. Lack of diagnostic morphological differences, combined with relatively low mtDNA barcode divergences and clinal phenotypic variation in key geographic regions indicate that the six previously recognized species of North American *Raphia* are best interpreted as parapatric subspecies. *Raphia frater abrupta* Grote, **stat. n.**, *R. f. coloradensis* Putnam-Cramer, **stat. r.**, *R. f. piazzzi* Hill, **stat. n.**, and *R. f. elbea* Smith, **stat. n.**, are accordingly revised to subspecies of *R. frater* Grote. Type locality restrictions are provided for *Raphia abrupta* and *Raphia frater* and a neotype is designated for *Raphia frater* var. *coloradensis*.

Keywords

DNA barcode, *Populus*, incipient species, subspecies, parapatry

Introduction

Raphia Hübner is a small genus of the Holarctic region, with a single African species (Poole 1989) questionably congeneric. As the sole genus currently comprising the Raphiinae, the phylogenetic placement of *Raphia* has an interesting history. Most early

works included the genus in older concepts of the Pantheinae. Smith (in Smith and Dyar 1898) excluded *Raphia* from the Pantheinae, but remained uncertain of its affinities within the Noctuidae. Hampson similarly excluded *Raphia* from the Pantheinae (Hampson 1913) and Acronictinae (Hampson 1909), and although never published, *Raphia* would presumably have been included in Hampson's volume covering "Ophi-derinae" (Kitching 1984), a catch-all subfamily for noctuoids with fully quadrifine hindwing venation and lacking other specialized features emphasized by Hampson. Forbes (1954) and Franclemont and Todd (1983) maintained *Raphia* in Pantheinae, and it was not until recent times that Beck (1996) proposed a separate subfamily to accommodate *Raphia*. Two additional genera were recently thought to be related to *Raphia*: *Diloba* Boisduval (Fibiger et al. 2009) and *Aon* Neumoegen (Fibiger and Lafontaine 2005), the former resulting in a family-level synonymy of Raphiinae under Dilobinae (Fibiger et al. 2009). Molecular study and re-assessment of morphological traits has since shown that Raphiinae and Dilobinae are best retained as valid subfamilies (Zahiri et al. 2013). Similarly, *Aon* has subsequently been reclassified as belonging to Erebidae: Hypocalinae (Lafontaine and Schmidt 2010).

Three *Raphia* species occur in temperate Asia, one in southern Europe (Poole 1989), and until now six species were recognized in North America (Lafontaine and Schmidt 2010). The current species concepts of Nearctic *Raphia* are essentially unchanged from those proposed by Smith (1908), although *R. coloradensis* was revised to the synonymy of *R. frater* by Schmidt and Anweiler (2010) and *R. piazzzi* (Hill 1927), described after Smith's work. Hence, over a century has elapsed since Smith's comprehensive synopsis of the Nearctic species. In this study, we revise the taxonomy of the six North American *Raphia* taxa based on geographic variation in adult phenotype, genitalia morphology, mtDNA barcode variation, and larval host plant use.

Methods and materials

Adult genitalia were prepared following the methods of Lafontaine (2004). Cleaned, stained genitalia were stored and examined in 30% ethanol, and slide-mounted in Euparal before being photographed using a Nikon D200 digital camera. Distribution maps were generated using SimpleMappr (<http://www.simplemappr.net/>). Tree distribution maps were adapted from USGS (2013).

We examined approximately 4000 specimens during the course of this study, primarily those of the CNC, EME, MEM, USNM, and UASM. Specimen repository abbreviations are as follows:

- AMNH** American Museum of Natural History, New York;
- ANSP** Academy of Natural Sciences, Philadelphia, PA;
- BIO** Biodiversity Institute of Ontario, Guelph, Ontario;
- BMNH** The Natural History Museum (statutorily British Museum [Natural History]), London;

| | |
|-------------|---|
| CNC | Canadian National Collection of Insects, Arachnids and Nematodes, Ottawa; |
| CSU | Colorado State University, Fort Collins; |
| EME | Essig Museum of Entomology, University of California, Berkeley, California; |
| MEM | Mississippi Entomological Museum, Mississippi State, MS; |
| UASM | University of Alberta Strickland Museum, Edmonton, Alberta; |
| USNM | National Museum of Natural History (formerly United States National Museum), Washington, D.C. |

DNA extraction, PCR amplification, and sequencing of the COI barcode region were performed at the Canadian Centre for DNA Barcoding (CCDB) and followed standard protocols (Hebert et al. 2013; <http://www.ccdb.ca/resources.php>). Only sequence records greater than 500bp (range 500bp – 658bp) are included.

Results and discussion

Morphology. Comparison of 20 genitalia dissections of each sex, representing all geographic entities, failed to reveal any diagnostic differences. The shape of the male valve apex and clasper varied slightly, but do so even within a single population. The shape of the inflated vesica, the most important diagnostic character in many noctuid species complexes, showed no discernible differences. Female genitalia were similarly conservative in variation. The European *R. hybris* Hübner, which is externally very similar to *R. frater* (and is in fact virtually indistinguishable from some *R. f. coloradensis* phenotypes), differs from *R. frater* in valve shape, vesica structure (including presence of spinules that are absent in *R. frater*) and shape and size of the corpus bursae. This indicates that *Raphia* genitalic morphology is not unusually homogeneous, where interspecific differences might be lacking.

The North American *Raphia* species have previously been delineated based on wing colour and pattern (Smith 1908), and geographic variation is considerable (Fig. 1). The most important forewing characters include ground colour, extent of medio-anal black shading, shape of the antemedial band, and the amount of fuscous shading of the hindwing. The colour of the prothoracic setae and overall size also vary.

Raphia has an extensive North American distribution, occupying virtually all biomes. Phenotypes are generally quite consistent regionally, but can appear drastically different in geographically disparate areas, which led early authors such as Smith (1908) to recognize multiple North American species. To assess phenotypic and mtDNA variation in these contact zones, we therefore attempted to locate and study specimens from key geographic regions where either two or more taxa would be expected to occur sympatrically or transition from one phenotype to another. The most comprehensive data were available for four such regions: **a**) the central Great Plains and **b**) the northeastern U.S., both where nominal ssp. *frater* interacts with ssp. *abrupta*; **b**) southern New Mexico where ssp. *elbea* meets ssp. *coloradensis*, and **c**) the Pacific Northwest / northern Rocky Mountains where ssp. *coloradensis*, *frater* and *elbea* meet (Figs 1, 3).




Figure 1. Geographic distribution and phenotypic variation of *Raphia frater* subspecies. Circles indicate specimens examined during this study: green – subsp. *piazzi*; pink – subsp. *abrupta*; black – subsp. *frater*; blue – subsp. *coloradensis*; yellow – subsp. *cinderella*. Multi-coloured circles indicate transitional populations and/or phenotypically intermediate specimens between respective subspecies. **a** *R. f. piazzi* (Zavalle Co., TX) **b** *R. f. abrupta* (Oktibeha Co., MS) **c** *R. f. abrupta* (Cottle Co., TX) **d** *R. f. abrupta* (Cottle Co., TX) **e** *R. f. abrupta* (Montgomery Co., MD) **f, g** *R. f. frater* (Edmunston, NB) **h** *R. f. abrupta* – *frater* intermediate (Anne Arundel Co., MD) **i** *R. f. abrupta* – *frater* – *coloradensis* intermediate from highly variable population in Cherry Co., NE **j** *R. f. coloradensis* (Alamosa Co., CO) **k** *R. f. coloradensis* (Milk River valley, AB) **l** *R. f. coloradensis* (Sanpete Co., UT) **m** *R. f. coloradensis* (Elko Co., NV) **n** *R. f. cinderella* (Ventura Co., CA) **o, p** *R. f. coloradensis* – *frater* intermediates (Chelan Co., WA) **q** *R. f. elbea* (Cochise Co., AZ) **r** *R. f. elbea* (San Juan Co., UT) **s** *R. f. elbea* (Santa Cruz Co., AZ). All specimens are males.

In the central Great Plains, a large series of over 60 specimens from northern Nebraska (Cherry Co.) is so variable that scarcely two individuals are alike, varying from the granular, dark grey forewing and white hindwing of ssp. *frater*, to the even, light grey forewing and slightly fuscous hindwing of *R. f. abrupta*; an intermediate specimen is shown in Fig. 1i. Some individuals show the blotchy black and grey pattern (with a contrasting black medio-anal shade) characteristic of *R. f. coloradensis*. A shorter series from Kansas (Riley Co.) falls within the variation of the Nebraska population. Interestingly, Smith (1908) remarked that Denver, Colorado specimens varied more towards ssp. *abrupta* than ssp. *coloradensis*. Single specimens from southeastern Montana and southwestern North Dakota have relatively pale forewings, like ssp. *coloradensis*, but some have fuscous hindwing shading and a dark prothoracic collar like *abrupta*. The zone of clinal variation may therefore be more extensive than the specimens from the few available sites in the northern Great Plains indicate, so further surveying in the region from southeastern Montana and southwestern North Dakota southward to western Nebraska / Kansas and eastern Colorado (particularly along major river corridors) would be helpful. Such a large transition zone appears to be the result of continuous, flat topography with a single, widespread *Populus* L. species (*P. deltoides* Bartr.) that is utilized by both ssp. *coloradensis* (to the northwest) and ssp. *abrupta* (to the southeast). This transition zone corresponds closely with the suture zone first proposed by Remington (1968) and recently verified by others (Swenson 2010 and references therein).

The nature of the ssp. *abrupta* – *frater* interface is somewhat different in the Northeast, and is seemingly more influenced by topography and host plant distribution (Fig. 2); at least three *Populus* species occur regionally among topography ranging from coastal floodplains to the Appalachian Mountains. Specimens from the Pocono Mtns. of Pennsylvania are *R. f. frater*, whereas nearby central Maryland (Ann Arundel Co.) specimens (Fig. 1h) show transitional features in having a forewing pattern much like ssp. *frater*, but with a fuscous hindwing and a darker prothoracic collar characteristic of ssp. *abrupta*. Coastal Maryland (Montgomery Co.) specimens are typical *R. f. abrupta* (Fig. 1e). As discussed below, the transition zone between ssp. *frater* and ssp. *abrupta* seems to be mediated by habitat and host plant differences, with *frater* largely associated with aspen (*P. tremuloides* Michx. and *P. grandidentata* Michx.; Fig. 2) and ssp. *abrupta* with cottonwood (*P. deltoides*; Fig. 2). Study of the populations on either side




Figure 2. Distribution of *Raphia frater* subspecies (circles) relative to range of *Populus* larval host plants (coloured shading) in eastern North America. Black circles - subsp. *frater*; white circles – subsp. *abrupta*; grey circles – subsp. *piazzii*; half-circles represent transitional populations and/or phenotypically intermediate specimens. Blue shading – combined ranges of *Populus tremuloides* and *P. grandidentata*; yellow shading – *P. deltooides*; range overlap depicted in green. *Populus* ranges adapted from USGS (2013).

of the Ohio River is needed because *frater* occurs throughout Ohio (Rings et al. 1992), whereas the few northeastern Kentucky specimens that were examined are mostly like ssp. *abrupta*, but show some ssp. *frater* traits, including a mostly white hindwing. The Ohio River valley is an important suture zone between other biota, but the relative limits of *R. f. frater* and ssp. *abrupta* from the northern Appalachians eastward appear to be further south than recognized suture zones (Swenson 2010).

In southern New Mexico where *elbea* and *coloradensis* meet, *elbea* is known from the Mimbres Mountains (Grant Co.) in the southwest, with the nearest documented *coloradensis* locations 180 km to the northeast in the Rio Grande valley, and 300 km to the east in Eddy County (Fig. 1). The phenotypic transition between the two taxa is more abrupt in southern New Mexico than it is in the Great Basin where *elbea* imperceptibly transitions to the very pale Great Basin forms of *coloradensis*. A series from

Twin Falls, Idaho, and some individuals from Leavenworth, Washington (Fig. 1p), are indistinguishable from Arizona *elbea* (Fig. 1q). *Raphia f. elbea* therefore grades into *R. f. coloradensis* in low-elevation habitats of the northern Great Basin. Specimens from the Eddy Co., New Mexico population are most like *coloradensis*, but some individuals are again indistinguishable from *elbea*. The mtDNA haplotypes associate this population with *coloradensis* (Fig. 3), and the available larval hosts are *Populus angustifolia* James and *P. deltoides* (National Park Service 2014), but not *P. fremonti* Wats. with which *elbea* is most often associated. Analysis of *Raphia* populations from the lower Rio Grande valley of New Mexico is desirable given the geographically intermediate position between *elbea* and *coloradensis*, and the presence of *Populus fremonti* (Fig. 3).

Populations at the edges of the Great Basin can be extremely variable, much like the situation between *abrupta* and *frater* in the north-central Great Plains. Series of specimens from Waterton Lakes, Alberta; Okanagan Valley, British Columbia; Baker County, Oregon; and Leavenworth, Washington range from the typical dark grey *frater* to pale yellowish *coloradensis* (Fig. 1). The Leavenworth population is remarkable in that it exhibits phenotypes ranging from *frater* to *coloradensis* (Fig 1o) and *elbea* (Fig 1p).

The geographic structure of California populations is not well documented; typical *cinderella* occurs from the San Francisco Bay area southward through the Central Valley to Los Angeles Co., but *R. frater* is apparently absent from southeastern California and the southern Sierra Nevada. Northern California (including the Sierra Nevada) specimens are most like Great Basin *coloradensis* but the transition from *cinderella* to this paler form is subtle, with Siskiyou Mountains material appearing intermediate. The Siskiyou Mountains are part of a northern California – southern Oregon suture zone also identified for other flora and fauna (see Swenson 2010 and references therein). Areas in central Texas where *abrupta* and *piazzi* meet, and eastern New Mexico/west Texas where the ranges of *abrupta*, *coloradensis* and *elbea* converge, remain unstudied.

Extreme phenotypic variation is therefore the modal geographic pattern at suture zones. In all cases where we examined suture zones between putative taxa, phenotype variation was moderate to extreme, and specimens could not consistently be assigned to existing taxonomic categories. Similarly, mtDNA variation patterns show no evidence of sympatric, reproductively isolated taxa, as discussed below.

Host plants. 72% of the 132 larval collections of *R. frater* from across Canada summarized by Prentice (1962) came from trembling aspen, 17% from other *Populus* species (*P. balsamifera* L., *P. trichocarpa* Torr. & Gray, *P. grandidentata*, *P. 'x canadensis'* (Alt.), *P. nigra* L. var. *italica* du Roi), and 3% from *Salix* spp. Three records from white birch and one from alder (both Betulaceae) reported by Prentice are exceptional and possibly accidental. Wagner et al. (2011) consider *Populus* to be the main hosts, and question the validity of records from birch and alder. The closely-related European species, *R. hybris*, is restricted to *Populus*. *Raphia* larvae possess an unusually large number of proleg crochets (Beck 1996), a trait also seen in the *Populus*-feeding genus *Ufeus* Grote (Noctuidae: Noctuinae); this trait is postulated to be an adaptation to maintaining a grip on the leaves of aspen species (Lafontaine and Walsh 2013), which tremble even in slight breezes. Based on these data, *Populus*, and to a lesser extent *Salix*




Figure 3. Distribution of *Raphia frater* subspecies (circles) relative to range of *Populus* species (coloured shading) in western North America. Half-circles represent transitional populations and/or phenotypically intermediate specimens. Ranges for *P. trichocarpa* + *P. balsamifera* and *P. angustifolia* + *P. deltoides* are combined, with both *P. balsamifera* and *P. deltoides* occurring in Alberta – Montana (upper right). *Populus* ranges adapted from USGS (2013).

L. (both Salicaceae), are certainly the primary and probably the only larval host plants. *Salix* may be used only incidentally or in certain regions/habitats; the parallel paucity of *Raphia* and *Populus* occurrence in the central Appalachian region is notable (Fig. 2). Eight species of *Populus* occur in North America, divided into four sections: *Leucooides* Spach (*P. heterophylla* L.), *Aigeiros* Duby (*P. deltoides*, *P. fremonti*), *Tacamahaca* Spach (*P. balsamifera*, *P. trichocarpa*, *P. angustifolia*) and *Populus* L. (*P. grandidentata*, *P. tremuloides*). One additional species, *P. mexicana* Wesmæl of central Mexico, is the sole constituent of section *Abaso* Eckenwalder (Eckenwalder 1996).

Although it is reasonably certain that *Raphia frater* larvae are Salicaceae specialists, the geographic variation in host use and extent of specialization is not well understood. Nonetheless, it is possible to extrapolate broader host use patterns based on larval collections, host plant distributions and habitat associations. Below, we outline some potential scenarios of host use among *R. frater* subspecies.

Host plant records for *R. f. frater* (Prentice 1962) indicate that *P. tremuloides* is probably the dominant, and certainly the most geographically widespread host (with the caveat that the high proportion of trembling aspen collections may simply reflect sampling bias). In eastern North America, the southern range limit of *R. f. frater* corresponds closely with the combined southern limits of the two aspen species (section *Populus*: *P. tremuloides* and *P. grandidentata*; Fig. 2). *Raphia frater* populations in riparian habitats of southern Alberta, where *P. tremuloides* is scarce or absent, are associated with other *Populus* species that form a complex zone of hybridization and overlap among four species (*P. deltoides*, *P. balsamifera*, *P. trichocarpa*, and *P. angustifolia*) along the major river valleys (Brayshaw 1965, Floate 2004). In this region *Raphia f. frater* phenotypes transition to *R. f. coloradensis* (Fig. 3). The northernmost extent of *P. deltoides* and *P. deltoides* × *balsamifera* hybrids in the Red Deer River valley at about 52° latitude (Floate 2004) also coincides with the northernmost extent of *R. f. coloradensis*-like phenotypes; north of there where *P. tremuloides* is the dominant species of the Aspen Parkland ecoregion, only pure *R. f. frater* phenotypes occur. Similarly, transitional *frater-coloradensis* populations occur in southwestern Alberta, southern British Columbia and central Washington at the range edges of *P. tremuloides*, where *P. trichocarpa* becomes the dominant *Populus* (Fig. 3). In northern Labrador, *R. f. frater* is at its northeastern range limit (not shown), occurring beyond the range of *Populus*; *Salix* species are the presumed hosts.

Throughout most of the range of *R. f. abrupta*, *P. deltoides* is the only *Populus* species present. Swamp cottonwood (*P. heterophylla*) has a small eastern North American range, occurring primarily along the Mississippi and Ohio River valleys and along the Atlantic seaboard (see e.g., Sibley 2009, USGS 2013), so this may serve as a host in some areas. The hosts for the southwest Texas taxon *R. f. piazzzi* are unknown, and may constitute willows rather than *Populus*, the latter being rare or absent where *R. f. piazzzi* occurs (Fig. 3).

In the Pacific Northwest, *R. f. frater* is associated with *P. tremuloides* in northern Washington and British Columbia, with *R. f. coloradensis* of dry, low-elevation habitats associated with *P. trichocarpa* (L. Crabo, pers. comm.). Crumb (1956) documented a larval collection from the latter species in south-central Washington. Throughout most of the Pacific Northwest, the only *Populus* species are *P. tremuloides* at upper elevations,




Figure 4. Neighbour-joining tree and associated sampling sites of mtDNA barcode haplotypes in *R. frater*. Haplogroup colour corresponds to that of sampling sites. Subspecies assignment based on morphology and sample size is indicated for each haplogroup. Width of triangles is proportional to number of haplotypes, height represents maximum divergence within haplogroup.

and *P. trichocarpa* in low-elevation riparian habitats and drier soils in moist regions (Fig. 3). A third species, *P. angustifolia*, occurs locally in the eastern parts of the Pacific Northwest, with habitats similar to *P. trichocarpa*. There are no specific host records for the Californian *R. f. cinderella*, with both *P. trichocarpa* and *P. fremonti* being the most likely hosts. Along the east slopes of the Oregon Cascade Ranges to the north, *R. f. coloradensis* associates with *P. trichocarpa* (L. Crabo, pers. comm.) Dyar (1894) cites “poplar” as a foodplant for larvae from Yosemite. Subspecies *coloradensis* in the Sierra Nevada, to the east of the central Californian range of *cinderella*, is associated with *P. tremuloides*.

Arizona populations of *R. f. elbea* feed on *P. fremonti* (Crumb 1956; D. Wagner, pers. comm.), and again this is the only available *Populus* in much of the range of ssp. *elbea*, excepting the higher elevations in the mountain ranges of central Arizona where *P. tremuloides* and *P. angustifolia* occur (Fig. 3). A population of *R. f. coloradensis* in the Rio Grande valley of central New Mexico is also associated with riparian *P. fremonti* (Fig. 3). Aspen-associated, *R. f. frater*-like populations may occur at high elevations in Arizona, similar to the situation in Colorado, but this has not been documented. This

raises the interesting possibility that *frater* occurs at higher elevations, with *elbea* occurring in the low-elevation floodplain. Surveying at the uppermost elevations of the Chiricahua and Santa Catalina Mountains of southeastern Arizona where aspen occurs have so far not yielded *Raphia* (BCS unpubl. data), but the more expansive range of *P. tremuloides* along, for example, the Mogollon Rim is poorly surveyed.

In summary, larval host plant associations of *R. frater* populations shows some broad congruencies between subspecies and *Populus* species distributions, but with limited evidence for high host fidelity: range edges of *R. frater* subspecies generally do not closely follow those of the various *Populus* hosts, suggesting that *Populus* availability rather than high host fidelity may be the limiting factor to *Raphia* distribution, and that climatic and topographic effects have a greater selective influence than host plant specialization. To what extent these congruencies reflect common co-evolutionary trajectories, and what factors drive intraspecific divergence, would be a fascinating and fruitful area of study.

Molecular variation. The deepest splits in mtDNA barcode variation (alluded to previously in Lafontaine and Schmidt 2010) segregate North American *Raphia* into four groups, but only one of these is private to a recognizable taxon (*R. f. elbea*). None of the remaining subspecies exhibited discrete haplotypic variation. Based on analysis of 192 specimens from localities across the range of *R. frater* representing all subspecies (Suppl. material 1), haplotypes segregated into five groups: **1**) a large group from across most of the eastern, northern and central range portions that includes *R. f. frater*, *R. f. coloradensis*, *R. f. abrupta* and *R. f. piazzzi*, varying by up to ~1.3% (Fig. 4); **2**) a discrete group closest to haplogroup 1, consisting of two *R. f. piazzzi* specimens (Fig. 4); **3**) a divergent group of geographically disparate samples of Eastern *R. f. frater* (Ontario, Manitoba) and Californian *R. f. cinderella* specimens, differing by a minimum of ~2.2% from all other groups (Fig. 4); **4**) a group private to *R. f. elbea* (Arizona, New Mexico, Utah) with a minimum ~1.0% divergence; **5**) A group of Californian *R. f. cinderella* with a minimum divergence of ~1.8% from all other haplotypes.

The combination of California and eastern Canada samples in haplogroup 3 to the exclusion of all others was quite unexpected, given the geographic structuring of other haplogroups. Two haplogroup-3 populations (Bird Hill, Manitoba; Bruce Peninsula, Ontario) also exhibited group 1 haplotypes (Fig. 4), the only sampled populations to yield more than one haplogroup. Representative specimens from these sites were of the same phenotype and from the same sampling event. This haplogroup could therefore be a retained ancestral mtDNA polymorphism, or indicative of *Wolbachia*-induced mtDNA lineage sorting similar to that documented by Kodandaramaiah et al. (2013). Determining the underlying cause of this interesting variation will require study using nuclear gene markers and *Wolbachia* assays.

Considering the general lack of taxonomic resolution of North American *Raphia* in the barcode sequence, and comparing divergences among Palearctic *Raphia* as a metric of mean species divergences within the genus, mtDNA variation is most parsimonious with a geographically structured, single-species interpretation. The contrast between often considerably different adult phenotypes and lack of significant mtDNA and morphological differentiation may reflect strong regional selection on bark-cryptic wing patterns, which in turn is dependent on dominant host trees that vary according to regional host preferences.

Systematics

Raphia Hübner, [1821]

Rhaphia Agassiz, 1847. An unjustified emendation of *Raphia* Hübner, [1821].

Anodonta Rambur, 1858. Type species: *Noctua hybris* Hübner, [1813], by monotypy.

A junior homonym of *Anodonta* Lamarck, 1799 [Mollusca].

Certila Walker, 1865. Type species: *Certila flexuosa* Walker, by monotypy. *Certila flexuosa* is a junior subjective synonym of *Raphia frater* Grote.

Saligena Walker, 1865. Type species: *Saligena personata* Walker, 1865, by monotypy.

Saligena personata is a junior subjective synonym of *Raphia frater* Grote.

Type species. *Noctua hybris* Hübner, [1813] by subsequent designation by Grote (1874).

Raphia frater Grote

Diagnosis. Despite variation in adult facies and lack of a particular diagnostic trait, *R. frater* is recognizable by the combination of a broad, rounded forewing, often conspicuous antemedial and postmedial band, obsolete medial band (rarely faint), an orbicular, reniform and usually also a claviform stigma that are clearly outlined, black shading in the anal angle of the hindwing, and the conspicuously bipectinate male antennae. *Pseudopanthea palata* (Grote) and *Colocasia* Ochseneheimer species share some superficial similarities with *Raphia frater*, but attention to the above-stated characters relative to those in *Pseudopanthea* McDunnough and *Colocasia* Ochseneheimer will provide an easy diagnosis of this unique species.

Description. **Head** – Male antennae bipectinate, anterior rami 3× longer than segment length, posterior rami 3.3× longer; female antennae simple; eyes round, with short, sparse interfacetal setae, visible only at high magnification; labial palpus with second segment clothed in long strap-like scales ventrally; third segment 0.6 × length of second segment (when denuded) and smoothly scaled; occiput and frons with mix of grey and black scales, frons with transverse line of black scales; frons rounded and moderately protuberant when denuded. **Thorax** – vestiture dark grey to yellowish grey, thoracic collar sometimes contrastingly darker than dorsum; tarsi smoothly scaled, with transverse bands of black and light to dark grey; tibia with similar scaling but with faint or indistinct banding; femur with long, shaggy hair-like scales. **Abdomen** – lacking specialized secondary sexual structures such as coremata; vestiture of smooth, short grey scales; small, rounded dorsal tufts on segments A3, A4 and A5, consisting of densely set spatulate scales. **Forewing** – ground colour varies from a dark charcoal grey to pale yellowish ochre; antemedial band a parallel-sided, double black line, varying from slightly irregular and rounded to nearly linear, acute, and angled at the cubital vein; medial band obsolete, usually reduced to a black bar or two diffuse lines at costa adjacent to reniform stigma, but band sometimes visible as

a faint, diffuse black line extending from bottom of reniform stigma to anal margin; postmedial band a single black line, sinuate and slightly sagittate at veins (often faint or absent in ssp. *coloradensis* and *elbea*), expanding to diffuse black patch at costa; orbicular stigma paler than ground colour, with black border and often with a diffuse dark pupil (orbicular often absent entirely in ssp. *coloradensis* and *elbea*); reniform stigma paler than ground colour, with a black border (border often lacking in *coloradensis* and *elbea*) and a diffuse black central crescent; subterminal band absent, faint, or diffusely sagittate with paler distal edging; terminal area often darker grey than subterminal area. Average size is greatest in subspecies *frater*, while *abrupta* and *piazzi* are smallest; forewing length varies from 16.2 mm and 18.5 mm in male and female *R. f. frater* to 13.7 mm and 15.2 mm in male and female *R. f. abrupta*, respectively. **Hindwing** – ground colour varying from white, white and dusted with fuscous grey (ssp. *frater*, *coloradensis*, *elbea*, *cinderella*, *piazzi*), or entirely pale fuscous grey (ssp. *abrupta*), females with more fuscous than males; crescentic discal spot diffuse or absent; postmedial band faint or absent, although nearly always with a contiguous diffuse black patch at anal angle. **Male genitalia** (Fig. 5) – uncus slightly compressed dorsoventrally, with slight medial bulge, apex blunt; valva tapering more or less evenly to a rounded point, sacculus poorly differentiated from remaining valva; ampulla long and flattened, $0.7 \times$ length of valva width, projecting mesially; aedeagus stout and sausage shaped, $2.1 \times$ longer than wide; vesica a simple kidney-shaped, unarmed chamber equal in length to aedeagus, tapering gradually into ductus. **Female genitalia** (Fig. 5) – bursa copulatrix membranous, lacking apparent differential sclerotization, including ostium, ante- and postvaginal plate; ductus bursae a simple rugose tube, $3.3 \times$ longer than diameter, connecting subbasally to corpus bursae; corpus bursae a simple kidney-shaped chamber, slighter wider over apical two-thirds; ductus seminalis arising from basal end of corpus bursae, caudad of ductus bursae; papillae anales short, bluntly rounded, with a broadly joined base, with two types of unusual, modified setae: 1) a dense band of thin, evenly curved setae arising from base of papillae and curving up to caudal margin, outer surface of lobe virtually encircled by a dense setal crown; and 2) highly modified thick, spatulate setae densely set along caudal margin of lobe. **Immature stages** – larva described by Thaxter (1883) based on eastern specimens, and by Dyar (1894) from Yosemite, California (possibly referable to *R. f. cinderella*). Illustrations in McCabe (1991) (head capsule and mandible), Wagner (2005) and Wagner et al. (2011). Mature larva stout, tapered only slightly anteriorly, bluish green to apple green with a slightly translucent quality, pinacula yellow, a dorsal transverse yellow band on A1, A5 and A8 extends to just above spiracle; T2 with short horn-like process middorsally, reddish with yellow base; these bands with whitish anterior border, those on A5 and A8 partially bordered with reddish orange; prolegs green, anal prolegs with yellow and reddish orange; head whitish green, usually retracted into T1, ocelli black, labrum whitish; total length 40 to 30 mm. Thaxter (1883) states that male larvae are more slender and smaller. Cocoon tough and firm, incorporating debris; pupa cylindrical with a rounded abdomen, cremaster short but broad and thick, lacking hooks. Eggs laid in small clusters or




Figure 5. Genitalia of *Raphia frater*. **a** female (Konza Prairie Biol. Stn., Riley Co., KS; leg. Metlevski), with inset showing apical spatulate setae and subapical dense setal crown of papillae anales **b** male vesica **c** male genital capsule (Pitchfork Ranch, Grant Co., NM; leg. C. Ferris); note magnification difference between sexes.

overlapping in linear groups; early instars much more elongate ‘semi-loopers’ with A3 and A4 prolegs reduced (Wagner et al. 2011). Larvae rest along midrib of leaf underside. Comparison among larvae of Californian *R. frater* (Dyar, 1894), *R. f. frater*, *R. f. abrupta* and *R. f. elbea* indicate no discernible differences among these subspecies.

Biology and distribution. *Raphia frater* occurs in virtually all wooded or shrubby habitats of the boreal region since the larval hosts form a dominant part of most non-coniferous forest types. It can be one of the most common late spring noctuids in aspen-dominated boreal forests of central Canada. In the West it becomes increasingly more restricted to riparian areas, particularly major river systems in drier regions of the Pacific Northwest and the desert of the Southwest. *Raphia frater* has a nearly trans-continental distribution, absent only from the arctic and most of the subarctic. The records from northern subarctic Labrador are surprising, but are based on three CNC specimens from two localities, so the data appear to be authentic. Handfield (2011) cites records from the northeast shore of the St. Lawrence, but the species is not known from Newfoundland. The range is essentially continuous south to northern Mexico, although very spotty throughout the Atlantic states, and spotty or absent in the central to southern Appalachians. Nominal *R. f. frater* occurs across the boreal region south to the northeastern States, southern Great Lakes region, and northern Rockies / Pacific Northwest; *R. f. abrupta* occurs from the Great Plains southward to eastern Texas and eastward to the Atlantic seaboard; *R. f. coloradensis* occurs from western California to the eastern slope of the Rockies; *R. f. elbea* occurs from at least southeastern Utah through Arizona, southwestern New Mexico and into Mexico; *R. f. cinderella* is restricted to western and central California; and *R. f. piazzzi* occurs from the Edwards Plateau into southern Texas. *Raphia frater* is univoltine across the boreal region and most of the west, with peak flight activity from late May to July. It is bivoltine in the eastern U.S., flying mostly in April to May, and July to August. In the Deep South, *R. f. abrupta* has three abundance peaks: March, May and a smaller flight (partial third brood?) in September (Brou 2014). Larvae are most common from late July to mid-August in Canada (Prentice 1962).

***Raphia frater frater* Grote**

Figs 1f, 1g, 2, 3

Raphia frater Grote, 1864

Saligena personata Walker, 1865

Type material. *Raphia frater* Grote, 1864 – # 7675 [ANSP]. Type locality: Middle States [eastern USA]; here restricted to Mount Pocono, Monroe Co., Pennsylvania. Grote (1864) simply stated the type locality as “Middle States,” and no additional information is available on the holotype label data. We interpret this as referring to the region south of the New England States, and north of the southern States. Given the complex variation of North American *R. frater*, it is advisable to restrict the type locality. As Grote’s material likely

originated from the eastern United States, we restrict the Type locality to Mount Pocono, Monroe Co., Pennsylvania, from which we examined typical *R. frater frater* specimens. *Raphia frater* and *R. abrupta* are the oldest available names for this species, and were published simultaneously. As first revisers, we designate *frater* as the senior name (ICZN, Article 24.2.2). Syn. *Saligena personata* Walker, 1865 - [BMNH]. Type locality: United States.

Diagnosis and description. The nominal subspecies of *R. frater* typically has an even, powdery, dark grey forewing ground colour with all of the markings complete, consisting of the antemedial and postmedial band, and the orbicular, reniform and usually the claviform stigmata. Average forewing length is 16.3 mm ($n = 9$) in males, 18.6 mm in females ($n = 9$). The male hindwing is white with little or no dusting of black scales in the subterminal area, and with a pronounced, diffusely-edged black patch in the anal angle, this often with an adjacent black line formed by the terminus of the postmedial band; females usually have some fuscous scales on the hindwing, especially on a slight postmedial band. This subspecies generally lacks the form with contrastingly darker medio-anal and costal black patches that is prevalent in *R. f. coloradensis*, but it does occur rarely even in Atlantic Canada (Fig. 1g). The yellowish-ochre forewing scales typical of *R. f. coloradensis* are absent. *Raphia f. abrupta* differs in having a more angulate and linear antemedial band, a paler grey and less powdery-appearing forewing, duskiest hindwing, and smaller size. As discussed in the section on *Raphia frater*, geographically intermediate populations are extremely variable with respect to these traits, and are considered to be transitional between subspecies *frater* and *coloradensis/abrupta*, the only two subspecies abutting the range of *R. f. frater*.

Biology and distribution. *Raphia f. frater* is primarily a boreal taxon, especially common in aspen (*Populus tremuloides* and *P. grandidentata*) dominated forests and the Aspen Parkland ecoregion of the Prairie Provinces. In the East, it extends south of the Great Lakes region into Pennsylvania, Ohio and Indiana, but apparently not southward into the southern Appalachians, which are essentially devoid of *Raphia* records. The transition zone between *R. f. frater* and *R. f. abrupta* extends from Maryland westward roughly along the Ohio River Valley to east-central Missouri, then northwestward through the northern Great Plains. The southeastern range edge of *R. f. frater* is virtually identical to that of both trembling and bigtooth aspens (Fig. 2). In the West, *R. f. frater* occurs south along mid-elevation mountain ranges of the Pacific Northwest into Washington, and southward along the Rocky Mountains. Specimens from high elevations in Colorado (Gilpin Co., 9500') and New Mexico (Sangre de Cristo Mtns., 7900') are of the typical *frater* phenotype, the *coloradensis* phenotypes occurring at lower elevations.

***Raphia frater abrupta* Grote, stat. n.**

Figs 1b–e, 2

Raphia abrupta Grote, 1864

Certila flexuosa Walker, 1865

Type material. *Raphia abrupta* - female holotype # 7675 [ANSP]. Type locality: not given; here restricted to Sycamore Landing, Seneca, Montgomery Co., Maryland. The female type bears no locality or collector label data, and since this is a widespread, geographically variable taxon, we restrict the type locality to Sycamore Landing, Seneca, Montgomery Co., Maryland; a series in USNM from this locality, collected by D. C. Ferguson, is phenotypically more similar to the female type than specimens from the Great Plains; it is also more likely that the holotype originated from the eastern US rather than the Great Plains, which were not well collected in the mid 1800's.

Certila flexuosa Walker - [BMNH; not examined]. Type locality: North America.

Diagnosis and description. *Raphia frater abrupta* replaces *R. f. frater* from the central Great Plains eastward to the mid-Atlantic seaboard, and southward to eastern Texas and Florida. It is on average smaller with a more evenly-coloured forewing, a more linear, angulate antemedial band and a fuscous hindwing. Average forewing length is 13.7 mm ($n = 9$) in males, 15.2 mm in females ($n = 9$). The thoracic collar is often darker than the dorsal thorax, not concolorous as in *R. f. frater*. The wing facies of subspecies *abrupta* is in many ways intermediate between *R. f. piazzzi* of central and southern Texas and *R. f. frater* to the north, but the exact nature of the interface between *abrupta* and *piazzi* in Texas remains unstudied.

Biology and distribution. Subspecies *abrupta* occurs south of the range of the aspen species favoured by *R. f. frater* larvae, and its riparian haunts suggest it feeds on eastern cottonwood (*Populus deltoides*), the only *Populus* species in much of its range. Swamp cottonwood (*P. heterophylla*) and willows (*Salix* spp.) may also be suitable hosts. This subspecies is apparently rare on the Atlantic seaboard and absent altogether in the Appalachians. We examined only a single historical specimen from New Jersey (Trenton), with records north of there assignable to *R. f. frater*. All Ohio records were attributed to *R. f. frater* by Rings et al. (1992), although specimens with a pale grey forewing and dusky hindwing, traits of the *abrupta* phenotype, rarely occur as far north as southernmost Ontario (Toronto) and southeastern Minnesota (Fillmore Co.).

***Raphia frater piazzzi* Hill**

Figs 1a, 2

Paphia [sic] *piazzi* Hill, 1927.

Type material. Holotype male [USNM]. Type locality: Brownsville, Texas [USA].

Diagnosis and description. *Raphia frater piazzzi* is the least-known member of the group with a restricted distribution in central and southern Texas. Most similar in size and facies to *R. f. abrupta*, it is distinguished from that subspecies by the paler, more evenly grey forewing with sharper transverse lines than in *R. f. abrupta*. The biology and biogeographic relationship to *R. f. abrupta*, which occurs to the northeast of *piazzi*'s range, is not known, and very few specimens of this taxon are present in collec-

tions. An additional enigma is whether or not Rio Grande *piazzi* populations interact with the vastly-different looking Sonoran *R. f. elbea*.

Biology and distribution. Described from southernmost Texas, this subspecies is otherwise known only from the Edwards Plateau region; a single specimen from Sinton County to the southeast is phenotypically intermediate between *abrupta* and *piazzi*, but clearly more field work is needed to establish the limits of both subspecies. mtDNA barcode data of three *piazzi* specimens (Sinton Co. and Zavalla Co.) are very similar to the haplotypes of *R. f. abrupta*, *R. f. frater*, and *R. f. coloradensis*.

Remarks. We were unable to obtain DNA sequence from topotypical specimens of *piazzi* from the lower Rio Grande near Brownsville, Texas. The unique haplotype of the Edwards Plateau specimens (Fig. 4) may represent nominal *piazzi*, but could equally represent a unique genetic lineage from the Edwards Plateau, with its unique fauna much of which is not shared with the Rio Grande fauna.

***Raphia frater coloradensis* Putnam-Cramer, stat. n.**

Figs 1j–m, 1o, 3

Raphia frater var. *coloradensis* Putnam-Cramer, 1886

Raphia pallula H. Edwards, 1886, **syn. nov.**

Type material. *Raphia frater* var. *coloradensis* - Neotype female, here designated [CNC]. Type locality: Deer Creek Cyn. Park, 39°33.18'N 105°08.49'W, 5950', SW Littleton, Jefferson Co., Colorado. None of the original types, three males and four females “taken in Colorado by D. Bruce,” could be located and are presumed lost. The primary type of *Xylena thoracica* Putnam-Cramer, the only other noctuid named by Putnam-Cramer, is housed at USNM. Prior to 1886, D. Bruce collected in the mountains and foothills near Denver (Brown 1966), and we accordingly select a specimen from the same region to designate as **neotype**: “Colorado: Jefferson Co. / 39°33.18'N 105°08.49'W / Deer Creek Cyn. Park / SW Littleton, w of hogback / 16–17 June 2008, 5950' elev / riparian area s. of road / leg: Chuck Harp uv trap”; “Neotype / *Raphia frater* var. / *coloradensis* Putnam-Cramer / Schmidt and Anweiler 2014.”

Raphia pallula - Holotype female [AMNH]. Type locality: Siskiyou Co., California [USA]. Published several months after *coloradensis* Putnam-Cramer, Edwards was apparently not aware of Putnam-Cramer’s name as it is not mentioned in his description.

Diagnosis and description. Within the range of *coloradensis*, specimens identical to the typical boreal *R. f. frater* are often present; in the most arid parts of the range of *coloradensis* in the southern Great Basin, *coloradensis* is more consistently pale ochre yellow with obsolete transverse lines and diffuse black costal/reniform blotches, overall very similar to *elbea*, but with less pronounced costal and reniform dark patches. Average forewing length is 14.9 mm ($n = 9$) in males, 16.8 mm in females ($n = 6$).

Biology and distribution. This subspecies occurs from southernmost British Columbia / Alberta to New Mexico, Utah, and California. It is most commonly associated

with riparian, low-elevation habitats. Northern populations fly from late May to July in a single generation. Flight dates spanning from May into August in the Great Basin and Southern Rocky mountain region indicate a second or partial second generation.

Remarks. *Raphia frater coloradensis* is the most weakly-differentiated subspecies, and may simply be an ecologically induced phenotype of *R. f. frater* that occurs in the warmer, drier regions of the West. Several populations, spanning a large geographical area, have been identified that exhibit a large range of phenotypic variation, as discussed above in the 'Morphology' section. Specimens from Siskiyou Co., California and the east slope of the northern Sierra Nevada (Sierra Co.) are phenotypically very similar to Great Basin *coloradensis*, and we therefore treat *pallula* as a junior subjective synonym. DNA barcodes of two specimens from the northern Sierra Nevada (Sierra Co.) belonged to the *frater-coloradensis-abrupta* haplogroup (Fig. 4).

***Raphia frater cinderella* Smith, stat. n.**

Figs 1n, 3

Raphia cinderella Smith, 1903.

Type material. A male lectotype was designated by Todd (1982) [AMNH]. Type locality: Los Angeles Co., Cal. [California, USA].

Diagnosis and description. *Raphia frater cinderella* is a Californian subspecies that is similar in size and colour to *R. f. coloradensis*, but with a more diffuse, poorly contrasting forewing pattern that usually lacks the pronouncedly darker reniform and costal dark patches. The forewing ground colour is also pale powdery grey, not pale ochre as it often is in *coloradensis*. The two taxa appear to intergrade in the Siskiyou and northern Sierra Nevada.

Biology and distribution. The range of this subspecies is restricted to central and southern California west of the Sierra Nevada. Fremont Cottonwood and willows are the most likely larval hosts, although records specific to this subspecies are lacking. Most collection dates are from June; Records from Stanislaus Co. for April - May and July may indicate a second generation.

***Raphia frater elbea* Smith, stat. n.**

Figs 1q-s, 3

Raphia elbea Smith, 1908

Type material. A male lectotype was designated by Todd (1982) [AMNH]. Type locality: Deming, [Luna Co.], New Mexico [USA].

Diagnosis and description. *Raphia frater elbea* is most similar to the pale yellowish-ochre forms of *R. f. coloradensis*, but differ from that subspecies in having both the costal and reniform dark patches more prominent; when present, the black medio-anal

patch is also darker and more elongate; additionally, *R. f. elbea* appears to exhibit a unique, divergent mtDNA haplotype group.

Biology and distribution. This subspecies occurs from southeastern Utah and western New Mexico southward through Arizona into northern Mexico. In southeastern Arizona it occurs in riparian areas in association with the larval host, *Populus fremonti*. Flight records are from February to October, with most being from March to May and August to September, indicating at least two generations annually.

Conclusions

The North American *Raphia* populations exhibit considerable geographic variation in phenotype, previously segregated into six species. Despite these geographically structured phenotypic differences, diagnostic morphological differences in genitalia and larvae are not evident. Scrutiny of geographic contact zones between putative taxa revealed populations with extensive phenotypic and conservative molecular variation, rather than bimodal phenotypic variation coupled with deep molecular divergences that would be expected for sympatric, reproductively isolated taxa. *Raphia frater* larvae are not highly restricted to a host species or genus, but do specialize on *Populus* and *Salix*, with a pattern of regional host availability and possibly also preference. Differences in host plant suitability among the various species of Salicaceae remain unstudied. Assessment of morphology, mtDNA variation, and biogeography therefore leads us to conclude that the geographic segregates of North American *Raphia* are best treated as subspecies of a single species. The regional adaptation to habitats representing nearly all North American biomes, combined with relatively discrete geographic ranges of unique adult phenotypes, suggest a pattern of young or incipient species in the *R. frater* group.

The taxonomy and biogeography of the North American *Raphia* populations is a complex interplay between topography, host plant use, phenotypic variation and evolutionary history. This study is only the first attempt at a better understanding of this interesting group. Many questions remain unanswered: what are the exact geospatial and host plant patterns of the contact zone between *R. f. abrupta* and *R. f. frater*? Is there geographic overlap with altitudinal segregation in the West between aspen-feeding *frater* and cottonwood feeding *elbea*? Does the mtDNA haplogroup 3 represent *Wolbachia* infection? Do the lower Rio Grande / Edwards Plateau *piazzi* populations grade into *abrupta*? *Raphia* would provide a fertile area of study in understanding large-scale patterns of host plant use and biogeography of a widely distributed continental Lepidopteran.

Acknowledgements

We thank James Adams, Charles Bordelon, Vernon A. Brou Jr., Richard Brown, Lars Crabo, Terhune Dickel, Cliff Ferris, Ed Knudson, Paul Opler, Jan Metlevski, Eric Metzler, Hugo Kons, Bruce Walsh and Chuck Harp for generously providing specimens

and/or data used in this study, and Jocelyn Gill for preparing the specimen images. Discussions with Lars Crabo and Don Lafontaine on *Raphia* biogeography and species limits provided insight and guidance during this project. Evgeny Zakharov, Paul Hebert and other members of the Barcode of Life Project at the University of Guelph, Ontario, Canada, provided DNA data. Molecular analyses were carried out through grants from the National Science and Engineering Research Council of Canada and Genome Canada through the Ontario Genomics Institute.

References

- Agassiz JLR (1847) Nomenclator Zoologici Index Universalis, continens nomina systematica classium, ordinem, familiarum et generum Animalium omnium, tam viventium quam fossilium, secundum ordinem alphabeticum unicum disposita, adjectis homonymiis plantarum, nec non variis adnotationibus et emendationibus. Fasciculus XII. In: Nomenclator Zoologicus continens nomina systematica generum animalium tam viventium quam fossilium, secundum ordinem alphabeticum disposita, adjectis auctoribus, libris, in quibus reperiuntur, anno editionis, etymologia et familiis, ad quas pertinent, in singulis classibus. Jent et Gassmann, Soloduri, viii + 393 pp.
- Beck H (1996) Systematische Liste der Noctuidae Europas (Lepidoptera, Noctuidae). *Neue Entomologische Nachrichten* 36: 1–122.
- Brayshaw TC (1965) Native poplars of southern Alberta and their hybrids. Canadian Department of Forestry, Ottawa. Publ. 1109.
- Brown FM (1966) David Bruce (1833–1903) and other entomological collectors in Colorado. *Journal of the New York Entomological Society* 74: 126–133.
- Brou VA Jr (2014) *Raphia abrupta* in Louisiana. *Southern Lepidopterists' News* 35: 173–174.
- Crumb SE (1956) The larvae of the Phalaenidae. United States Department of Agriculture Technical Bulletin 1135: 1–356.
- Dyar HG (1894) Some undescribed stages of Noctuidae larvae. *The Canadian Entomologist* 26: 17–21. doi: 10.4039/Ent2617-1
- Eckenwalder JE (1996) Systematics and evolution of *Populus*. In: Stettler RF, Bradshaw HD, Heilman PE, Hinckley TM (Eds) *Biology of Populus and its implications for management and conservation*. NRC Research Press, Ottawa, 7–32.
- Edwards H (1886) Apparently new forms of American Lepidoptera. *Entomologica Americana* 2: 165–171.
- Fibiger M, Lafontaine JD (2005) A review of the higher classification of the Noctuoidea (Lepidoptera) – with special reference to the Holarctic fauna. *Esperiana* 11: 7–92.
- Fibiger M, Ronkay L, Steiner A, Zilli A (2009) Noctuidae Europaeae. Vol. 11, Pantheinae, Dilobinae, Acronictinae, Eustrotiinae, Nolinae, Bagisarinae, Acontiinae, Metoponiinae, Heliothinae, and Bryophilinae. Entomological Press, Sorø, 504 pp.
- Floate KD (2004) Extent and patterns of hybridization among the three species of *Populus* that constitute the riparian forest of southern Alberta, Canada. *Canadian Journal of Botany*, 82: 253–264. doi: 10.1139/b03-135

- Forbes WTM (1954) Lepidoptera of New York and neighboring states. Part 3 Noctuidae. Cornell University Agriculture Experiment Station, Memoir 329: 1–433.
- Franclemont JG, Todd EL (1983) Noctuidae. In: Hodges RW, Dominick T, Davis DR, Ferguson DC, Franclemont JG, Munroe EG, Powell JA (1983) Check List of the Lepidoptera of America North of Mexico. E. W. Classey Ltd, London and The Wedge Entomological Research Foundation Washington, 284 pp.
- Grote AR (1864) Descriptions of North American Lepidoptera – no. 2. Proceedings of the Entomological Society of Philadelphia 2: 433–441.
- Grote AR (1874) List of the Noctuidae of North America. Bulletin of the Buffalo Society of Natural Sciences 2: 1–77.
- Hampson GF (1909) Catalogue of the Lepidoptera Phalaenae in the British Museum 8: 1–583.
- Hampson GF (1913) Catalogue of the Lepidoptera Phalaenae in the British Museum 13: 1–609.
- Handfield L (2011) Le guide des papillons du Québec. Version scientifique, Broquet, Ottawa, 1352 pp.
- Hebert PDN, deWaard JR, Zakharov EV, Prosser SWJ, Sones JE, McKeown JTA, Mantle B, La Salle J (2013) A DNA ‘Barcode Blitz’: Rapid digitization and sequencing of a natural history collection. PLoS ONE 8: e68535. doi: 10.1371/journal.pone.0068535
- Hill CA (1927) Three new moths from the southwest. Bulletin of the Southern California Academy of Sciences 26: 6–7.
- Hübner J (1813) Sammlung Europäischer Schmettlinge. Vol. 4. Eulen. Augsburg, 2–81.
- Hübner J (1821) Verzeichniss bekannter Schmettlinge. Augsburg, 431 pp.
- Kitching IJ (1984) An historical review of the higher classification of the Noctuidae (Lepidoptera). Bulletin of the British Museum (Natural History) (Entomology) 49: 153–234.
- Kodandaramaiah U, Simonsen TJ, Bromilow S, Wahlberg N, Sperling, F (2013) Deceptive single-locus taxonomy and phylogeography: *Wolbachia*-associated divergence in mitochondrial DNA is not reflected in morphology and nuclear markers in a butterfly species. Ecology and Evolution 3: 5167–5176. doi: 10.1002/ece3.886
- Lafontaine JD (2004) Noctuoidea: Noctuidae (part) – Agrotini. In: Hodges RW (Ed) The Moths of North America. Fascicle 27.1. The Wedge Entomological Research Foundation, Washington, 394 pp.
- Lafontaine JD, Schmidt BC (2010) Annotated check list of the Noctuoidea (Insecta, Lepidoptera) of North America north of Mexico. ZooKeys 40: 1–239. doi: 10.3897/zookeys.40.414
- Lafontaine JD, Walsh JB (2013) A revision of the genus *Ufeus* Grote with the description of a new species from Arizona (Lepidoptera, Noctuidae, Noctuinae, Xylenini, Ufeina). ZooKeys 264: 193–207. doi: 10.3897/zookeys.264.3526
- McCabe TL (1991) Atlas of Adirondack Caterpillars: With a Host List, Rearing Notes, and a Selected Bibliography of Works Depicting Caterpillars. New York State Museum Bulletin 470: 1–114.
- National Park Service (2014) Checklist of the Vascular Plants of Carlsbad Caverns National Park. http://www.nps.gov/cave/planyourvisit/upload/2007_CAVE_Plants.pdf [accessed 5 March 2014]
- Poole RW (1989) Lepidopterorum Catalogus (New Series). Fascicle 118 Noctuidae, Parts 1–3. E.J. Brill, New York, 1314 pp.

- Prentice RM (1962) Forest Lepidoptera of Canada reported by the Forest Insect Survey, Vol. 2: Nycteolidae, Notodontidae, Noctuidae, Liparidae. Canada Dept. of Forestry, Forest Entomology and Pathology Branch, Publication No. 1013, 77–281.
- Putnam-Cramer AW (1886) Two new varieties of noctuids. *Entomologica Americana* 2: 1–142.
- Rambur JP (1858) *Catalogue Systématique des Lépidoptères de l'Andalousie*. JB Bailliere, Paris, 422 pp. [22 plates]
- Remington CL (1968) Suture-zones of hybrid interaction between recently joined biotas. In: Dobzhansky T, Hecht MK, Steere WC (Eds) *Evolutionary Biolog.* Plenum, New York, 321–428. doi: 10.1007/978-1-4684-8094-8_8
- Rings RW, Metzler EH, Arnold FJ, Harris DH (1992) The owlet moths of Ohio Order Lepidoptera Family Noctuidae. *Ohio Biological Survey Bulletin, New Series* 9(2): 1–219. [pl. 1–16]
- Schmidt BC, Anweiler GG (2010) Taxonomic changes to Lepidoptera: Macro-moths. In: Pohl GR, Anweiler GG, Schmidt BC, Kondla NG (Eds) *An annotated list of the Lepidoptera of Alberta, Canada*. *ZooKeys* 38: 497–509. doi: 10.3897/zookeys.38.383
- Sibley DA (2009) *The Sibley Guide to Trees*. Alfred E. Knopf, New York, 426 pp.
- Smith JB (1903) New noctuids for 1903, no. 4, with notes on certain described species. *Transactions of the American Entomological Society* 29: 191–224.
- Smith JB (1908) New species of Noctuidae for 1908. I. With notes *Charadra*, *Raphia* and *Pseudanarta*. *Journal of the Entomological Society of New York* 16: 79–98.
- Smith JB, Dyar HG (1898) Contributions toward a monograph of the Lepidopterous family Noctuidae of boreal North America. A Revision of the species of *Acronycta* (Ochsenheimer) and of certain allied genera. *Proceedings of the United States National Museum* 21 (1140): 1–194. doi: 10.5479/si.00963801.21-1140.1
- Swenson NG (2010) Mapping the suturing of a continental biota. *Molecular Ecology* 19: 5324–5327. doi: 10.1111/j.1365-294X.2010.04880.x
- Thaxter R (1883) Descriptions of noctuid larvae found on Cutts' Island, Maine. *Papilio* 3: 10–19.
- USGS (2013) Digital Representations of Tree Species Range Maps from “Atlas of United States Trees” by Elbert L. Little, Jr. (and other publications). <http://esp.cr.usgs.gov/data/little/> [accessed 10 November 2010]
- Todd EL (1982) The noctuid type material of J. B. Smith (Lepidoptera). *United States Department of Agriculture Technical Bulletin* 1645: 1–228.
- Wagner DL (2005) *Caterpillars of eastern North America: a guide to identification and natural history*. Princeton University Press, Princeton, NJ, 512 pp.
- Wagner DL, Schweitzer DF, Sullivan JB, Reardon RC (2011) *Owlet Caterpillars of Eastern North America*. Princeton University Press, Princeton, NJ, 576 pp.
- Walker FD (1865) *List of the Specimens of Lepidopterous Insects in the Collection of the British Museum* 32. British Museum, London, 448 pp.
- Zahiri R, Lafontaine JD, Schmidt, BC, Holloway JD, Kitching IJ, Mutanen M, Wahlberg N (2013) Relationships of the basal lineages of Noctuidae (Lepidoptera, Noctuoidea) based on eight gene regions. *Zoologica Scripta* 42: 488–507. doi: 10.1111/zsc.12022

Supplementary material I

Table S1. Specimen data for mtDNA barcode vouchers.

Authors: B. Christian Schmidt, Gary G. Anweiler

Data type: data spreadsheet

Explanation note: Haplogroup numbers refer to those given in Fig. 4; Abbreviations for specimen depositories are as given in Methods and materials section.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: doi: 10.3897/zookeys.421.7517.app1