

Fixed Combinations of Glaucoma Medications

Nikola Babić^{1,2}

¹University of Novi Sad, Medical Faculty, Novi Sad, Serbia;

²Eye Clinic, Clinical Center of Vojvodina, Novi Sad, Serbia

SUMMARY

The first line treatment in the management of glaucoma is topical medical therapy. Many patients with glaucoma require multiple medications for adequate intraocular pressure control. For patients who need multi-dose regimens to control intraocular pressure, fixed combinations offer convenience, efficacy and safety. This review summarizes the role, efficacy, mechanism of action and indications for use of modern fixed combination of topical glaucoma medications. The review shows the advantages and disadvantages of a prescribing fixed combination in daily clinical practice.

Keywords: glaucoma; fixed combination therapy; drugs

INTRODUCTION

When initiating medical therapy for newly diagnosed glaucoma, it is typically desirable to begin with a single antiglaucoma medication [1]. Topical medical therapy remains the first-line treatment in the management of glaucoma, but over time monotherapy often fails to control intraocular pressure (IOP). The initial monotherapy fails to control IOP within the first two years of treatment in about 50% of glaucoma patients [2] and frequently more than one agent is required to achieve adequate IOP control. If the initial therapy seems to be ineffective or the first choice monotherapy is well tolerated and is effective in lowering IOP but does not succeed in reaching the target pressure, an additional drug is added to the therapeutic regimen [3, 4, 5]. Recent trials, such as the Normal Tension Glaucoma Treatment Study, the Collaborative Initial Glaucoma Treatment Study, and the Ocular Hypertension Treatment Study, demonstrate that in order to reach the target IOP, it is often necessary to use multiple drugs [6, 7]. Patients are often prescribed multiple medications from different classes of IOP-lowering drugs, including α -agonists and carbonic anhydrase inhibitors (CAI) to prostaglandin analogues and β -blockers, to help maintain IOP control. There have been a number of concerns about the use of multiple bottles of glaucoma medications, such as increased toxicity of ophthalmic preservatives in multiple drugs, compliance, costs and washout effects [7, 8]. Complex dosing regimens have been clearly associated with nonadherence to medical therapy. The washout effect depends not only on the increased tear fluid turnover, but also on addition of a second eye drop within a short period. When more than one bottle is used, patients do not always allow adequate time for ocular absorption of their first medication before the second drug is administered [9]. Furthermore, Robin and

Covert's study [10] showed that patients using two concomitant drugs often skip one drug in a few days. Low compliance with prescribed long-term glaucoma therapy is universal and impairs treatment outcome [11]. Inadequate compliance greatly diminishes drug efficacy, often leads to aggravation or undertreatment of health problems and inflates the cost of health care. Preservatives, especially benzalkonium chloride (BAK), are toxic to the eye in variety of ways, such as conjunctival inflammation, punctate keratopathy or dry eye syndrome [8, 12]. These concerns have spurred the search for glaucoma drugs that will work well when put together in one bottle.

RATIONALE FOR FIXED COMBINATION THERAPY

Combining two medications in one bottle may improve compliance by reducing the time required to administer drops, the frequency of use, and the total number of bottles [11, 13]. Additionally, the use of one bottle rather than two significantly reduces the inconvenience of filling prescriptions and reduces daily cost of therapy [14]. For patients who need multi-dose regimens to control IOP, fixed combinations offer convenience, efficacy and safety. Fixed combinations of glaucoma medications offer a reduction in the number of bottles, and reduction in the number of drops per day. They also offer reduced time for drop instillation and potentially greater efficacy by eliminating the washout effect. The cost and time saving can enhance compliance. Instilling two medications in a single drop reduces the amount of preservatives, which may improve the tolerability and eventual surgical outcomes in patients who require filtering procedures [13].

The major limitation of a fixed combination therapy is that dosing of the concomitant medi-

Correspondence to:

Nikola BABIĆ
Klinika za očne bolesti
Klinički centar Vojvodine
Hajduk Veljkova 1-7
21000 Novi Sad
Srbija
nikobab@sbb.rs

cation cannot be alerted within the concomitant product. For instance, a twice-daily regimen of a fixed combination product may contain too little or too much of one drug for a given patient. Morning dose may not be ideal for a prostaglandin analogue that seems to have more profound effects on IOP when dosed in the evening. Conversely, β -blockers show little effects on aqueous production during sleep and ideally should be given in the morning. Fixed combination medications may cause problems if a patient is allergic to any constituent of the fixed combination and make difficulties in finding which constituent causes allergy.

Fixed combination often includes timolol, a β -adrenergic antagonist that effectively decreases aqueous humor production and is generally well tolerated. For years these were dominant drugs used in the treatment of glaucoma [15, 16]. When we decide to initiate glaucoma therapy, compliance is of great importance. As the complexity of medical regimen grows, the likelihood of therapeutic adherence falls [7]. An important factor in the maintenance of adequate IOP control is patient's adherence to the prescribed treatment. Studies show that about 50% of glaucoma patients do not comply with dosing instructions and convenience is an important factor contributing to patients' adherence [17]. Simplification of the dosing regimen results in an increase in patients' adherence, but side effects may adversely affect it. A β -blocker, such as timolol, dosed twice daily, may cause cardiovascular, central nervous system, endocrine and pulmonary side effects [15]. Therefore, a once-daily prostaglandin/timolol fixed combination therapy may have important advantages over multi-dose regimens.

Prostaglandin analogues, another class of potent ocular hypotension substances, reduce IOP by increasing uveoscleral outflow of aqueous humor [18, 19]. Many studies showed that prostaglandin administered once daily efficiently lowers IOP in patients with primary open-angle glaucoma (POAG) and ocular hypertension (OHT) [20-23].

The complementary mechanisms of action of a prostaglandin analogue and a β -blocker are likely to show additive IOP-lowering effect in combination when compared with effects of either single agent [24-28].

THE AVAILABLE FIXED COMBINATIONS (TABLE 1)

Latanoprost/timolol

The first fixed combination of a prostaglandin analogue and β -blockers on the market was latanoprost 0.005% / timolol 0.5% fixed combination. Numerous studies have shown greater efficacy in decreasing of IOP compared to individual components. Mean diurnal IOP reduction from baseline was 33.5% and it was achieved in 73.5% of patients. Latanoprost/timolol fixed combination achieved a mean IOP reduction at baseline of -9.4 mmHg [29, 30]. Fixed combination of latanoprost plus timolol is effective in controlling IOP in patients with POAG and OHT who do not tolerate well multiple drops for reaching targeted IOP. The use of latanoprost/timolol fixed combination should reduce the exposure to ophthalmic preservatives

Table 1. Available fixed combinations of glaucoma medications

Antiglaucoma drug 1	Antiglaucoma drug 2	Commercial name
Bimatoprost 0.03%	Timolol 0.5%	Ganfort®
Latanoprost 0.005%	Timolol 0.5%	Xalacom®, Xalcom®
Travoprost 0.004%	Timolol 0.5%	DuoTrav®
Dorzolamide 2%	Timolol 0.5%	Cosopt®
Brinzolamide 1%	Timolol 0.5%	Azarga®
Brimonidine 0.2%	Timolol 0.5%	Combigan®
Brinzolamide 1%	Brimonidine 0.2%	Simbrinza®

compared (less BAK) with concomitant administration of multiple drops. Latanoprost/timolol is indicated in patients with POAG and OHT due to less adverse effects, better compliance and tolerability. Simplification of the regimen to once daily improves the side effect profile.

Travoprost/timolol

Travoprost 0.004% / timolol 0.5% is a second-line fixed combination therapy used if treatment with travoprost or timolol as monotherapy fails to reach target IOP, and can replace concomitant administration of non-fixed combination as adjunctive therapy. Clinical studies have found that fixed combination travoprost plus timolol achieved a mean IOP decrease at baseline of 32–38% [27, 31]. A recent study compared the efficacy and safety of the fixed combination of travoprost plus timolol with the fixed combination of dorzolamide plus timolol, and found that the mean reduction from baseline was -8.96 mmHg for the travoprost/timolol fixed combination and -8.07 mmHg for the dorzolamide/timolol fixed combination, with the highest difference of -1.07 mmHg in favor of the travoprost/timolol fixed combination. Mean percentage reduction in diurnal IOP was 36.28% in the travoprost/timolol fixed combination, and 35.66% in the dorzolamide/timolol fixed combination [32]. Travoprost/timolol is the only fixed combination without BAK (travoprost/timolol "BAK-free"). It contains Polyquad ophthalmic preservative, which has far less toxic effect on ocular surface. Fixed combination of travoprost plus timolol is effective in controlling IOP in patients with POAG and OHT who do not tolerate well multiple drops for reaching target IOP. The use of travoprost/timolol fixed combination should be ideal for patients who developed ocular surface disease, due to BAK toxicity. Travoprost/timolol is indicated in patients with POAG and OHT due to less adverse effects, better compliance and tolerability. Simplification of once-daily dosing improves the side effect profile. Travoprost/timolol fixed combination is among the antiglaucoma drugs with the highest efficacy in decreasing IOP, and in certain cases it can be used as initial monotherapy. Travoprost/timolol fixed combination is conveniently dosed once daily in the morning.

Bimatoprost/timolol

Bimatoprost 0.03% / timolol 0.5% is a new fixed combination which represents the second line therapy if treatment

with bimatoprost or timolol as monotherapy does not succeed in reaching the target pressure, and can replace concomitant administration of non-fixed combination as adjunctive therapy. A lot of clinical studies have shown that the average reduction in IOP from baseline was 35% [24, 33, 34]. Fixed combination of bimatoprost plus timolol is effective in controlling IOP in patients with POAG and OHT who do not tolerate well multiple drops for reaching target IOP. The use of bimatoprost/timolol should reduce the exposure to ophthalmic preservatives compared with concomitant administration of multiple drops. [35, 36, 37]. Bimatoprost/timolol is indicated in patients with POAG and OHT due to less adverse effects, better compliance and tolerability. Simplification of the regimen to once-daily dose improves the side effect profile. Bimatoprost/timolol is the most effective antiglaucoma drug with the highest efficacy in decreasing IOP, and in certain cases it can be used as initial monotherapy. Bimatoprost/timolol fixed combination is well tolerated and safe. It is used once daily, in the morning.

Dorzolamide/timolol

Dorzolamide is a topical carbonic anhydrase inhibitor which reduces aqueous humor production and it is frequently prescribed as adjunctive therapy to timolol for additional IOP reduction. In controlled clinical trials, dorzolamide showed additional IOP reduction when used as adjunctive therapy to timolol regardless of which of the two drugs was used as initial therapy [38]. Dorzolamide 2% / timolol 0.5% is a fixed combination product with a convenient dosing regimen for patients requiring multiple medications. The fixed combination of dorzolamide and timolol, dosed twice daily, similarly has been shown to lower IOP more than either of its individual components and has been found to be comparable in efficacy to the concomitant administration of the two medications [39]. Dorzolamide/timolol fixed combination is highly effective in lowering IOP by 34.4–34.6% as initial therapy in patients with POAG and OHT [40]. It is given twice daily.

Brinzolamide/timolol

A fixed combination of brinzolamide 1% plus timolol 0.5% is the latest fixed combination of β -blockers and CAI. Clinical trials have demonstrated that brinzolamide/timolol fixed combination provides clinically meaningful IOP reduction from baseline, non-inferior to those seen with dorzolamide/timolol fixed combination. IOP reduction from baseline was 28.4–34.9% in brinzolamide/timolol group vs. 29.2–33.9% in dorzolamide/timolol group [41]. The advantage of brinzolamide/timolol fixed combination is in a higher safety and tolerability profile. Patients with POAG and OHT prefer the brinzolamide/timolol fixed combination over the dorzolamide/timolol fixed combination. This is likely related to differences in the formulation, specifically differences in pH and buffering system between

the two products [42, 43]. Stronger patient's preference for greater comfort of brinzolamide/timolol may lead to better therapeutic compliance. The use of brinzolamide/timolol should reduce the exposure to ophthalmic preservatives compared with concomitant administration of multiple drops. Brinzolamide/timolol is indicated in patients with POAG and OHT due to less adverse effects, better compliance and tolerability. Due to high tolerability and comfort, brinzolamide/timolol fixed combination is a treatment of choice for patients on dorzolamide/timolol treatment who developed intolerability and adverse effects, and should be switched to a new drug. It is dosed twice daily.

Brimonidine/timolol

A fixed combination of brimonidine 2% / timolol 0.5% is a combination of α -agonist and β -blocker. The mechanism of decreasing IOP is complementary, even though both constituents cause IOP decrease by decreasing aqueous humor production by different ways of action. Brimonidine partially increases uveoscleral outflow of aqueous humor as well. The efficacy of IOP lowering ranged from 28% to 33.6% [44, 45]. Clinical studies found that brimonidine/timolol fixed combination is more effective in lowering IOP than any of its individual components and has been found to be comparable in efficacy to the concomitant use of the two medications [46, 47]. A simplified dosing regimen has a potential to improve the compliance. In comparison with dorzolamide/timolol fixed combination, clinical studies found a similar efficacy in IOP lowering, but brimonidine/timolol fixed combination showed better tolerability [47, 48, 49]. In patients on multiple-drug therapy, including prostaglandins, replacement of dorzolamide/timolol with brimonidine/timolol may help achieve a lower IOP of 2 mmHg, while replacement of brimonidine plus dorzolamide/timolol with brimonidine/timolol may help achieve low IOP with fewer medications [50]. It is used twice daily.

Brinzolamide/brimonidine

The newest fixed combination on the market is brinzolamide/brimonidine containing CAI – brinzolamide 1% and α -2 agonist – brimonidine 2% and it is the only fixed combination option that does not rely on a β -blocker to exert its IOP-lowering effect. Its mechanism of action differs from other fixed drug medications. Each component works differently to decrease the elevated IOP; brinzolamide decreases aqueous humor production and brimonidine decreases aqueous humor production and increases uveoscleral outflow. In recent studies, brinzolamide/brimonidine ophthalmic suspension resulted in greater IOP reduction (16.7–20.5 mmHg) than brinzolamide (19.8–21.4 mmHg) or brimonidine (18–22.5 mmHg). Brinzolamide/brimonidine suspension provided an average of 21.4–34.9% reduction in IOP, higher than either of its individual components (16.9–22.6% for brinzolamide; 14.3–25.8% for brimonidine). Patients receiving brinzolamide/brimonidine

achieved an average of 5.4–8.8 mmHg reduction of IOP from baseline [51, 52]. It is administered three times a day.

CONCLUSION

In patients with POAG and OHT, fixed combination therapy offers benefits over concomitant therapy, in terms of convenience, costs, and potentially compliance and efficacy. Combination therapy is not recommended as a first-

line treatment. However, in some cases, such as advanced glaucoma and/or very high IOP, the requested IOP reduction may exceed their efficacy range when used as a single agent. Therefore, a fixed combination can be used as a first-line therapy. If the combination therapy fails to reduce IOP sufficiently, one can either substitute the second drug or add a third medication to the fixed combination. A fixed combination with addition of a third medication is considered to be maximal medical therapy and at this stage laser or incisional surgery should be considered.

REFERENCES

- Denis P. Travoprost/timolol fixed combination in management of open-angle glaucoma: a clinical review. *Exp Opin Pharmacother*. 2011; 12(3):463-71. [DOI: 10.1517/14656566.2011.551007] [PMID: 21254951]
- Kobelt-Nguyen G, Gerdtham UG, Alm A. Costs of treating primary open-angle glaucoma and ocular hypertension: a retrospective observational two-year chart review of newly diagnosed patients in Sweden and the United States. *J Glaucoma*. 1998; 7:95-104. [PMID: 9559495]
- Kass M, Heuer DK, Higginbotham EJ, Johnson CA, Keltner JL, Miller JP, et al. The Ocular Hypertension Treatment Study: A randomized trial determines that topical ocular hypotensive medication delays or prevents the onset of primary open-angle glaucoma. *Arch Ophthalmol*. 2002; 120:701-13. [DOI: 10.1001/archophth.120.6.701] [PMID: 12049574]
- European Glaucoma Society. Terminology and Guidelines for Glaucoma. 4th ed. Savona, Italy: PubliComm; 2014.
- Babić N, Andreić V, Miljković A, Čanadanović V, Barišić S. Adjunctive therapy of brinzolamide in patients on travoprost treatment. *Med Pregl*. 2011; 64(5-6):310-4. [DOI: 10.2298/MPNS1106310B] [PMID: 21789924]
- Lichter PR, Musch DC, Gillespie BW, Guire KE, Janz NK, Wren PA, et al. Interim clinical outcomes in the Collaborative Initial Glaucoma Treatment Study comparing initial treatment randomized to medication or surgery. *Ophthalmology*. 2001; 108:1943-53. [DOI: 10.1016/S0161-6420(01)00873-9] [PMID: 11713061]
- Weinreb RN. Compliance with medical treatments of glaucoma. *J Glaucoma*. 1992; 1:134-6.
- Goto Y, Ibaraki N, Miyake K. Human lens epithelial cell damage and stimulation of their secretion of chemical mediators by benzalkonium chloride rather than latanoprost and timolol. *Arch Ophthalmol*. 2003; 121:835-9. [DOI: 10.1001/archophth.121.6.835] [PMID: 12796255]
- Katz JK. Modern alchemy: fixed combinations of glaucoma drugs. *Am J Ophthalmol*. 2005; 140(1):125-6. [DOI: 10.1016/j.ajo.2005.03.034] [PMID: 16038653]
- Robin AL, Covert D. Does adjunctive glaucoma therapy affect adherence to the initial primary therapy? *Ophthalmology*. 2005; 112:863-8. [DOI: 10.1016/j.ophtha.2004.12.026] [PMID: 15878067]
- Othoff CMG, Schouten JSAG, van den Borne BW, Webers CA. Noncompliance with ocular hypotensive treatment in patients with glaucoma or ocular hypertension. *Ophthalmology*. 2005; 112:953-61. [DOI: 10.1016/j.ophtha.2004.12.035] [PMID: 15885795]
- Noecker RJ, Herrygers LA, Anwaruddin R. Corneal and conjunctival changes caused by commonly used glaucoma medication. *Cornea*. 2004; 23:490-6. [DOI: 10.1097/01.ico.0000116526.57227.82] [PMID: 15220734]
- Fechner RD, Realini T. Fixed combination of topical glaucoma medications. *Curr Opin Ophthalmol*. 2004; 15:132-5. [DOI: 10.1097/00055735-200404000-00013] [PMID: 15021225]
- Hollo G, Thelen U, Teus MA, Quaranta L, Ferkova S, Babić N, et al. Long-term outcomes of prostaglandin analog versus timolol maleate in ocular hypertensive or primary open-angle glaucoma patients in Europe. *J Ocul Pharmacol Ther*. 2011; 27(5):493-8. [DOI: 10.1089/jop.2011.0051] [PMID: 21790326]
- Zimmermann TJ, Kaufman HE. Timolol. A beta-adrenergic blocking agent for the treatment of glaucoma. *Arch Ophthalmol*. 1977; 95:601-4. [DOI: 10.1001/archophth.1977.04450040067008] [PMID: 322648]
- Rafuse P. Adrenergic antagonist. In: Morrison JC, Pollack IP. *Glaucoma Science and Practice*. New York – Stuttgart: Thieme; 2003. p.376.
- Patel SC, Spaeth GL. Compliance in patients prescribed eye drops for glaucoma. *Ophthalmology Surg*. 1995; 26:233-6. [PMID: 7651690]
- Weinreb RN, Toris CB, Gabelt BA, Lindsey JD, Kaufman PL. Effects of prostaglandins on the aqueous humor outflow pathways. *Surv Ophthalmol*. 2002; 47(Suppl 1):S53-64. [DOI: 10.1016/S0039-6257(02)00306-5] [PMID: 12204701]
- Lawlor D, Toris CB, Camras CB. Prostaglandin analogs. In: Morrison JC, Pollack IP. *Glaucoma Science and Practice*. New York – Stuttgart, Thieme; 2003. p.391.
- Goldberg I, Cuhna-Vaz J, Jakobsen JE, Nordmann JF, Trost E, Sullivan KE; International Travoprost Study Group. Comparison of topical travoprost drops given once daily and timolol 0.5% given twice daily in patients with open-angle glaucoma or ocular hypertension. *J Glaucoma*. 2001; 10:414-22. [DOI: 10.1097/00061198-200110000-00010] [PMID: 11711841]
- Netland PA, Landry T, Sullivan EK, Andrew R, Silver L, Weiner A, et al; Travoprost Study Group. Travoprost compared with latanoprost and timolol in patients with open-angle glaucoma or ocular hypertension. *Am J Ophthalmol*. 2001; 4:472-84. [DOI: 10.1016/S0002-9394(01)01177-1] [PMID: 11589866]
- Fellman RL, Sullivan KE, Ratliff M, Silver LH, Whitson JT, Turner DF; Travoprost Study Group. Comparison of travoprost 0.0015% and 0.004% with timolol 0.5% in patients with elevated intraocular pressure. *Ophthalmology*. 2002; 109:998-1008. [DOI: 10.1016/S0161-6420(02)01010-2] [PMID: 11986110]
- Babić N, Čanadanović V, Žikić Z. Comparison of the efficacy and safety of travoprost and timolol throughout diurnal curve. *Asian Journal of Ophthalmology*. 2008; 10:215-20.
- Hommer A; Ganfort Investigators Group I. A double-masked, randomized, parallel comparison of a fixed combination of bimatoprost 0.03%/timolol 0.5% with non-fixed combination use in patients with glaucoma or ocular hypertension. *Eur J Ophthalmol*. 2007; 17:53-62. [PMID: 17294383]
- Diestelhorst M, Larsson LI; European Latanoprost Fixed Combination Study Group. A 12 week study comparing the fixed combination of latanoprost and timolol with the concomitant use of the individual components in patients with open angle glaucoma and ocular hypertension. *Br J Ophthalmol*. 2004; 88:199-203. [DOI: 10.1136/bjo.2003.018234] [PMID: 14736774]
- Hughes BA, Bacharach J, Craven ER, Kaback MB, Mallick S, Landry TA, et al. A three-month, multicenter, double-masked study of the safety and efficacy of travoprost 0.004%/timolol 0.5% ophthalmic solution compared to travoprost 0.004% ophthalmic solution and timolol 0.5% dosed concomitantly in subjects with open angle glaucoma or ocular hypertension. *J Glaucoma*. 2005; 14:392-9. [DOI: 10.1097/01.jig.0000176935.08392.14] [PMID: 16148589]
- Schuman JS, Katz GJ, Lewis RA, Mallick S, Wells DT, Sullivan EK, et al. Efficacy and safety of a fixed combination of travoprost 0.004%/timolol 0.5% ophthalmic solution once daily for open-angle glaucoma or ocular hypertension. *Am J Ophthalmol*. 2005; 140:242-50. [DOI: 10.1016/j.ajo.2005.02.058] [PMID: 16086946]
- Barnebey HS, Orengo-Nania S, Flowers BF, Samples J, Mallick S, Landry TA, et al. The safety and efficacy of travoprost 0.004%/timolol 0.5% fixed combination ophthalmic solution. *Am J Ophthalmol*. 2005; 140:1-7. [DOI: 10.1016/j.ajo.2005.02.043] [PMID: 15990081]

29. Higginbotham EJ, Olander KW, Kim EE, Grunden JW, Kwok KK, Tressler CS; United States Fixed-Combination Study Group. Fixed combination of latanoprost and timolol vs individual components for primary open-angle glaucoma and ocular hypertension: a randomized, double-masked study. *Arch Ophthalmol*. 2010; 128(2):165-72. [DOI: 10.1001/archophthalmol.2009.384] [PMID: 20142538]
30. Shin DH, Feldman RM, Sheu WP; Fixed Combination Latanoprost/Timolol Study Group. Fixed combination latanoprost/timolol study group. Efficacy and safety of the fixed combination latanoprost/timolol versus dorzolamide/timolol in patients with elevated intraocular pressure. *Ophthalmology*. 2004; 111(2):276-82. [DOI: 10.1016/j.ophtha.2003.05.019] [PMID: 20142538]
31. Teus MA, Miglior S, Laganovska G, Volkson L, Romanowska-Dixon B, Gos R, et al. Efficacy and safety of travoprost/timolol vs. dorzolamide/timolol in patients with open-angle glaucoma or ocular hypertension. *Clin Ophthalmol*. 2009; 3:629-36. [DOI: 10.2147/OPHT.S8011] [PMID: 19997566]
32. Babić N, Andreić V, Miljković A, Grković D, Jovanović P. Comparison of the efficacy and safety of the fixed combination travoprost/timolol and dorzolamide/timolol in patients with primary open-angle glaucoma and ocular hypertension. *Srp Arh Celok Lek*. 2013; 141(7-8):441-6. [DOI: 10.2298/SARH1308441B] [PMID: 24073548]
33. Brandt JD, Cantor LG, Batoosingh AL, Liu CC; for the Ganfort Investigators Group. A 3-month randomized study comparing bimatoprost/timolol fixed combination to monotherapy with bimatoprost or timolol in patients with glaucoma or ocular hypertension. *Proceeding of the 6th International Glaucoma Symposium*; 2007 Mar 28-31; Athens, Greece: Kenes; p.154.
34. Konstas AGP, Hollo, G, Mikropoulos D, Tsironi S, Heidich AB, Embeslidis T, et al. Twenty-four-hour intraocular pressure control with bimatoprost and bimatoprost/timolol fixed combination administered in the morning, or evening in exfoliative glaucoma. *Br J Ophthalmol*. 2010; 94:209-13. [DOI: 10.1136/bjo.2008.155317] [PMID: 19825835]
35. van der Valk R, Webers CA, Schouten JS, Zeegers MP, Hendrikse F, Prins MH. Intraocular pressure-lowering effects of all commonly used glaucoma drugs: a meta-analysis of randomized clinical trials. *Ophthalmology*. 2005; 112:1177-85. [DOI: 10.1016/j.ophtha.2005.01.042] [PMID: 15921747]
36. Stewart WC, Konstas AGP, Nelson LA, Kruf B. Meta-analysis of 24-hour intraocular pressure studies evaluating the efficacy of glaucoma medicines. *Ophthalmology*. 2008; 115(7):1117-1122.e1. [DOI: 10.1016/j.ophtha.2007.10.004] [PMID: 18082886]
37. Denis P, Lafuma A, Khoshnood B, Mimaud V, Berdeaux G. A meta-analysis of topical prostaglandin analogues intra-ocular pressure lowering in glaucoma therapy. *Curr Med Res Opin*. 2007; 23(3):601-8. [DOI: 10.1185/030079907X178720] [PMID: 17355741]
38. Boyle JE, Ghosh K, Gieser DK, Adamsons IA. A randomized trial comparing the dorzolamide-timolol combination given twice daily to monotherapy with timolol and dorzolamide. *Ophthalmology*. 1998; 105:1945-51. [PMID: 9787368]
39. Clineschmidt CM, Williams RD, Snyder E, Adamsons IA. A randomized trial in patients inadequately controlled with timolol alone comparing the dorzolamide-timolol combination to monotherapy with timolol or dorzolamide. *Dorzolamide-Timolol Combination Study Group*. *Ophthalmology*. 1998; 105:1952-9. [PMID: 9787369]
40. Radaković M, Čanadanović V, Babić N, Naumović N. Initial reduction of intraocular pressure with dorzolamide and timolol fixed combination in open-angle glaucoma and ocular hypertension. *Medicina danas*. 2011; 10(4-6):163-8.
41. Manni G, Denis P, Chew P, Sharpe ED, Orenge-Nania S, Coote MA, et al. The safety and efficacy of brinzolamide 1%/timolol 0.5% fixed combination versus dorzolamide 2%/timolol 0.5% in patients with open-angle glaucoma or ocular hypertension. *J Glaucoma*. 2009; 18(4):293-300. [DOI: 10.1097/IJG.0b013e31818fb434] [PMID: 19365194]
42. Vold SD, Evans RM, Stewart RH, Walters T, Mallick S. A one-week comfort study of bid-dosed brinzolamide 1%/timolol 0.5% ophthalmic suspension fixed combination compared to bid-dosed dorzolamide 2%/timolol 0.5% ophthalmic solution in patients with open-angle glaucoma or ocular hypertension. *J Ocul Pharmacol Ther*. 2008; 24(6):601-6. [DOI: 10.1089/jop.2008.0030] [PMID: 1904930]
43. Mundorf TK, Rauchman SH, Williams RD, Notivol R; Brinzolamide/Timolol Preference Study Group. A patient preference comparison of Azarga (brinzolamide/timolol fixed combination) vs. Cosopt (dorzolamide/timolol fixed combination) in patients with open-angle glaucoma or ocular hypertension. *Clin Ophthalmol*. 2008; 2(3):623-8. [DOI: 10.2147/OPHT.S4088] [PMID: 19668763]
44. Cheng JW, Cheng SW, Gao LD, Lu GC, Wei R. Intraocular pressure-lowering effects of commonly used fixed combination drugs with timolol: a systemic review and meta-analysis. *PLoS One*. 2012; 7(9):e45079. [DOI: 10.1371/journal.pone.0045079] [PMID: 23028770]
45. Hommer A, Sperl P, Resch H, Popa-Cherechenau A, Qiao C, Schmettler L, et al. A double-masked randomized crossover study comparing the effect of latanoprost/timolol and brimonidine/timolol fixed combination on intraocular pressure and ocular blood flow in patients with primary open-angle glaucoma or ocular hypertension. *J Ocul Pharmacol Ther*. 2012; 28(6):569-75. [DOI: 10.1089/jop.2011.0165] [PMID: 22775229]
46. Goñi FJ; Brimonidine/Timolol Fixed Combination Study Group. 12-week study comparing the fixed combination of brimonidine and timolol with concomitant use of the individual components in patients with glaucoma and ocular hypertension. *Eur J Ophthalmol*. 2005; 15(5):581-90. [PMID: 16167288]
47. Frampton JE. Topical brimonidine 0.2%/timolol 0.5% ophthalmic solution: in glaucoma and ocular hypertension. *Drugs Aging*. 2006; 23(9):753-61. [PMID: 17020399]
48. García-Feijó J, Sáenz-Francés F, Martínez-de-la-Casa JM, Méndez-Hernández C, Fernández-Vidal A, Calvo-González C, et al. Comparison of ocular hypotensive action of fixed combination of brimonidine/timolol and dorzolamide/timolol. *Curr Med Res Opin*. 2010; 26(7):1599-606. [DOI: 10.1185/03007995.2010.482017] [PMID: 20429818]
49. Gulkilik G, Oba E, Odabasi M. Comparison of fixed combination of brimonidine/timolol and dorzolamide/timolol in patients with primary open-angle glaucoma. *Int Ophthalmol*. 2011; 31(6):447-51. [DOI: 10.1007/s10792-011-9495-z] [PMID: 22207522]
50. Nguyen QH, Earl M. Fixed-combination brimonidine/timolol as adjunctive therapy to a prostaglandin analog: a 3-month, open-label, replacement study in glaucoma patients. *J Ocul Pharmacol Ther*. 2009; 25(6):541-4. [DOI: 10.1089/jop.2009.0045] [PMID: 20028261]
51. Nguyen QH, McMenemy MG, Realini T, Whitson JT, Goode SM. Phase 3 randomized 3-month trial with an ongoing 3 month safety extension of fixed-combination brinzolamide 1%/brimonidine 0.2%. *J Ocul Pharmacol Ther*. 2013; 29(3):290-7. [DOI: 10.1089/jop.2012.0235] [PMID: 23425430]
52. Whitson JT, Realini J, Nguyen QH, McMenemy MG, Goode SM. Six-month results from a Phase III randomized trial of fixed-combination brinzolamide 1% + brimonidine 0.2% versus brinzolamide or brimonidine monotherapy in glaucoma or ocular hypertension. *Clin Ophthalmol*. 2013; 7:1053-60. [DOI: 10.2147/OPHT.S46881] [PMID: 23766627]

Фиксне комбинације у терапији глаукома

Никола Бабић^{1,2}

¹Универзитет у Новом Саду, Медицински факултет, Нови Сад, Србија;

²Клиника за очне болести, Клинички центар Војводине, Нови Сад, Србија

КРАТАК САДРЖАЈ

Прва терапијска линија у лечењу од глаукома је медикаментна, у виду капи. Многи болесници који се лече од глаукома захтевају примену већег броја лекова за регулисање интраокуларног притиска. Код болесника којима је потребна употреба већег броја бочица за регулисање интраокуларног притиска, фиксне комбинације нуде једноставност,

ефикасност и безбедност у лечењу. Овај ревијски рад приказује улогу, ефикасност, механизам дејства и индикације за примену савремених фиксних комбинација. Рад указује на предности и недостатке код употребе фиксних комбинација у свакодневној клиничкој пракси.

Кључне речи: глауком; фиксне комбинације; лекови

Примљен • Received: 31/03/2015

Прихваћен • Accepted: 29/06/2015