

A review of Spinosyns, a derivative of biological acting substances as a class of insecticides with a broad range of action against many insect pests

L. Bacci,¹ D. Lupi,² S. Savoldelli,² B. Rossaro²

¹Dow Agrosiences Italia s.r.l., Bologna; ²Department of Food, Environmental and Nutritional Sciences, University of Milano, Milano, Italy

Abstract

Spinosyns are a class of insecticides with a broad range of action against many insect pests belonging to different orders, noxious to a wide variety of agricultural crops; spinosyns were also used against insects of sanitary interest. Spinosyns are derivative of biological active substances produced by soil Actinomycete *Saccharopolyspora spinosa*; being of biological origin, they are considered to have a low environmental impact and they are not much aggressive against non-target species. They act as allosteric activators of nicotinic acetylcholine receptors; thanks to their mode of action the resistance phenomena are uncommon, even few cases of resistance were recently reported.

For all these reasons at present they are one of the most interesting product to be used in fighting against agriculture pests.

Introduction

Spinosyns are a family of broad-spectrum insecticides including spinosad and spinetoram, all with a macrocyclic lactone structure, isolated from the actinomycete soil bacterium *Saccharopolyspora spinosa*,

Mertz and Yao, 1990 (Bacteria: Actinobacteridae) (Sparks *et al.*, 1998).

Spinosad is derived from the aerobic fermentation of *S. spinosa* in aqueous growth media (containing *e.g.*, corn solids, soya bean flour, and cottonseed flour) after extraction and recrystallization of technical spinosad. Spinosad is a mixture (85:15%) of spinosyn A and spinosyn D (Mertz & Yao, 1990; Kirst *et al.*, 1992; Sparks *et al.*, 1999) (Figure 1). Because of its microbial origin, spinosad can be considered a bio pesticide, and it is useful for the management of many insect pests, including caterpillars, leaf miners, thrips, flies, drywood termites, and some beetles, in various vegetables, field crops, and fruits.

A descendant of spinosad is spinetoram (Saglam *et al.*, 2013), a mixture of two synthetically modified spinosyns (spinosyn J and spinosyn L), which are metabolites of *S. spinosa*. Chemical modifications relating spinosyn A, J, L and spinetoram are summarized in Figure 1 (Dripps *et al.*, 2008).

As spinosad, spinetoram has a neurotoxic mode of action on insects, through contact or ingestion, it was introduced as a novel insecticide with greater potency and faster speed of action in comparison with the older spinosyn-based relative spinosad (Dripps *et al.*, 2008; Sparks *et al.*, 2008).

Recently combining genome shuffling with antibiotics resistance screening resulted in an effective approach to achieve rapid improvement of spinosad yield of *S. spinosa* (Wang *et al.*, 2015a, 2015b).

Spinosad has been evaluated and accepted in USA for listing by the World Health Organization Pesticide Evaluation Scheme (WHOPES) working group, the official WHO body in charge of the assessment of pesticides for their effectiveness and safety (WHO, 2007). Spinosad was authorized in 2007 in Europe with a EU Commission Directive 2007/6/EC (European Commission, 2007), with subsequent updates, *e.g.*, EU Regulations 556/2012 and 2015/603 (European Commission, 2012, 2015). It was registered to be used against a broad array of target pests as Lepidoptera (pests of cotton), Diptera (fruit flies and mosquitoes), Thysanoptera, Isoptera and Coleoptera (Thompson *et al.*, 2000; Legocki *et al.*, 2010; Tescari *et al.*, 2014).

Spinetoram was registered for the first time in New Zealand immediately followed by the United States in September 2007 under the United States Environmental Protection Agency (USEPA) reduced risk pesticide initiative. Submission for Annex I inclusion in Europe was completed in 2007 and accepted in 2014 with the EU implementing Regulation 140/2014 (European Commission, 2014).

Correspondence: Bruno Rossaro, Department of Food, Environmental and Nutritional Sciences (DeFENS), University of Milano, via Celoria 2, I 20133 Milano, Italy.

E-mail: bruno.rossaro@unimi.it

Key words: Spinosad; spinetoram; insecticides; target species; resistance.

Acknowledgements: research partly supported by Dow Agrosiences Italia S.r.l.

Received for publication: 29 November 2015.

Revision received: 19 February 2015.

Accepted for publication: 19 February 2015.

©Copyright L. Bacci *et al.*, 2016

Licencee PAGEPress, Italy

Journal of Entomological and Acarological Research 2016; 48:5653

doi:10.4081/jea.2016.5653

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (by-nc 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Toxicity on target organisms

Spinosyns were developed to control species detrimental to agriculture, but they were also used to control insects of sanitary interest. Spinosyns were tested on many pest species; the most investigated are summarized in Table 1.




Figure 1. Chemical modifications connecting spinosyn A, J, L and spinetoram.

The mode of action of spinosyns may be as direct contact through body surface *contact toxicity* or through ingestion *dietary toxicity* (Shimokawatoko *et al.*, 2012).

Spinosad acts through contact activity on all life stages of insects, including eggs, larvae, and adults. Eggs must be sprayed directly, but larvae and adults can be effectively dosed through contact with treated surfaces. Spinosad is most effective when ingested, generally showing a greater selectivity toward target insects and a lesser activity against many beneficial predators as well as mammals and other aquatic and avian animals (Cleveland *et al.*, 2001).

A useful library to analyse the results of toxicity tests is given in the R-package drc: <https://cran.r-project.org/web/packages/drc/drc.pdf>

A comparison with other insecticides can be also carried out in a series of field tests. For example the efficacy of a spinosyn, spinosad (DE-105), was compared with other insecticides on the tobacco budworm *Heliothis virescens* (Fabricius, 1777) (Lepidoptera: Noctuidae) a parasite of cotton in Louisiana during 1989-95 and on the corn earworms *Helicoverpa zea* Boddie, 1850 and *H. armigera* (Hübner, 1805) (Lepidoptera: Noctuidae). Spinosad provided control of Lepidoptera equal to or greater than that afforded by pyrethroid, organophosphate and carbamate standards. In larval topical tests, the range of spinosad lethal dose at 50% (LD₅₀) among several populations of *H. virescens* was from 0.4 to 8.5 µg/g (Leonard *et al.*, 1996).

A variety of bioassays can be planned to evaluate the activity of nine different molecules of spinosyns. As test species *H. virescens* was used. Spinosyn A resulted the most effective with an LC₅₀ of 0.3 ppm in the active range of many pyrethroids (Sparks *et al.*, 1998).

To explore the efficacy of spinosyn A compared with other insecticides, both drench (application to the larva in a large volume of solvent) and topical applications (direct application to the dorsum of the larva in a 1-µL drop of acetone) were used. For example topical bioassay performed on 3rd instar larvae of *H. virescens* gave an LD₅₀ between 1.28-2.25 µg/g after 48-72 h (Sparks *et al.*, 1998).

The oral efficacy of spinosyn A compared with cypermethrin was also evaluated using leaf-dip, diet/egg, leaf-spray bioassays and hot-probe bioassays: i) the leaf-dip bioassay involved dipping a cotton leaf disk (33 mm diameter) in a test solution and allowing it to dry; ii) the diet/egg bioassay consisted of spraying a 1-oz (30 mL) plastic cup containing artificial diet with a test solution; iii) the leaf-spray bioassay was carried out using a cotton leaf disk (33 mm diameter) placed in the bottom of a 1-oz (30-mL) plastic cup and sprayed using a track sprayer calibrated to evenly coat the leaves with 0.055 mL of test solution; iv) a hot-probe bioassay: a simple and rapid paralysis assay to detect and characterize knockdown resistance (Bloomquist & Miller, 1985).

Spinosad and spinetoram were also tested as cereal protector against several stored product pests. Hertlein *et al.* (2011) summarized the results of several laboratory and field tests on various grains to reach conclusion that the lesser grain borer, *Rhyzopertha dominica* (Fabricius, 1972) (Coleoptera: Bostrichidae); larger grain borer, *Prostephanus truncatus* (G. H. Horn, 1878) (Coleoptera: Bostrichidae); rusty grain beetle, *Cryptolestes ferrugineus* (Stephens, 1831) (Coleoptera: Laemophloeidae); flat grain beetle, *Cryptolestes pusillus* (Schönherr, 1817) (Coleoptera: Laemophloeidae); red flour beetle, *Tribolium castaneum* (Herbst, 1787) (Coleoptera: Tenebrionidae); confused flour beetle, *Tribolium confusum* Jacquelin du Val, 1868 (Coleoptera: Tenebrionidae); Indian meal moth, *Plodia interpunctella* (Hubner, 1813) (Lepidoptera: Pyralidae); rice moth, *Corcyra cephalonica* (Stainton, 1866) (Lepidoptera: Pyralidae); angoumois grain moth, *Sitotroga cerealella* (Olivier, 1789) (Lepidoptera: Gelechiidae); almond moth, *Cadra cautella* (Walker, 1863) (Lepidoptera: Pyralidae), and the psocid species *Lepinotus reticulatus* Enderlein, 1905 (Psocoptera: Trogiidae) and *Liposcelis entomophila* (Enderlein, 1907) (Psocoptera: Liposcelididae) are susceptible to spinosad and complete control is to be expected. Other pest species such as the maize weevil, *Sitophilus*

zeamais Motschulsky, 1855 (Coleoptera: Dryophthoridae); rice weevil, *Sitophilus oryzae* (Linnaeus, 1763) (Coleoptera: Dryophthoridae), and sawtoothed grain beetle, *Oryzaephilus surinamensis* (Linnaeus, 1758) (Coleoptera: Silvanidae) are susceptible to spinosad to varying degrees.

Likewise *P. truncatus* and *R. dominica* are very susceptible to spinetoram used to protect grain, while *T. confusum* and *O. surinamensis* showed a limited susceptibility to spinetoram (Vassilakos *et al.*, 2012). For *Sitophilus granarius* (Linnaeus, 1758) (Coleoptera: Dryophthoridae) and *S. oryzae*, spinetoram efficacy differed among the grain commodities (hard wheat, soft wheat, oats, rye, triticale, paddy rice and maize). In general, mortality was higher in hard wheat for both *Sitophilus* species in comparison with the other grains, while the lowest mortality levels were recorded in oats and soft wheat for *S. granarius* and in maize and soft wheat for *S. oryzae* (Vassilakos *et al.*, 2015).

Spinetoram was effective against *T. confusum* adults and young larvae. It revealed more effective than imidacloprid, thiamethoxam and chlorantraniliprole in controlling *T. castaneum*. On the other hand spinetoram was ineffective against *T. confusum* when applied directly on wheat, *Triticum aestivum* L. (Vassilakos & Athanassiou, 2013).

The toxicity of the bioinsecticide spinosad and the semi-synthetic insecticide spinetoram was compared with the toxicity of three insect growth regulator (IGR) compounds: lufenuron, chlorfluazuron and methoxyfenozide studying the response of 2nd instar larvae of the cotton leafworm *Spodoptera littoralis* Boisduval, 1833 (Lepidoptera: Noctuidae). Lufenuron, chlorfluazuron and methoxyfenozide were more toxic than spinosad by 9.8-, 9.4- and 9.2-fold, and more toxic than spinetoram by 2.0-, 1.9- and 1.8-fold, respectively, after 72 h of exposure. Spinetoram (LC₅₀=8.6 ppm) was more toxic than spinosad (LC₅₀=43.1 ppm) by approximately 5 folds, after 72 h of exposure. These results are summarized in Table 2.

Spinosad toxicity interacts with toxicity of IGR compounds. Spinosad/IGR compounds mixtures resulted in a potentiation effect more than the spinetoram/IGR compounds mixtures. The ratio spinosad or spinetoram at LC₂₅/IGR compounds at LC₂₅ revealed potentiating effects higher than the ratio spinosad or spinetoram at LC₂₅/IGR compounds at LC₁₀. The ratio spinosad or spinetoram at LC₁₀/IGR compounds at LC₂₅ resulted only in an additive effect.

Mixtures of spinosad-spinetoram/IGR were more toxic when spinosad-spinetoram were in higher proportion respect to IGR. Therefore, it was preferred to use high concentrations of spinosad or spinetoram with low concentrations of the IGR compounds and not the opposite. The toxicity of the three IGR compounds was enhanced with the time. Although the toxicity of tested IGR compounds appeared after 48 and 72 h of exposure, the potentiating effect of these compounds to spinosad and spinetoram appeared after 24 h (Rahman & Abou-Taleb, 2007).

Spinetoram toxicity was compared with the toxicity of Vertimec® (Syngenta AG, Basel, Switzerland). Vertimec® is a product containing abamectin, a mixture of avermectins, compounds derived from the soil bacterium *Streptomyces avermitilis*. Abamectin is a natural fermentation product of this bacterium. It is used to control insect and mite pests of a range of agronomic, fruit, vegetable and ornamental crops, and it is used by homeowners for control of fire ants on the moveable stages of *Tetranychus urticae* Koch, 1836 (Boisduval, 1867) (Acari: Tetranychidae). Spinetoram 12% was most effective than Vermectin®. The best reduction of pests was 100% after 13 and 19 days of egg treatment, 96.7% after 5 days for adult and 87.5 after 11 days for immature stages at dose 1 ml solution per liter of water (El-Kady *et al.*, 2007).

Spinetoram toxicity was compared with the toxicity of spinosad; at 61 g active ingredient (a.i.)/ha spinetoram was as effective as spinosad at 140 g a.i./ha against the western flower thrips and the other common thrips in Florida, *Frankliniella tritici* Fitch, 1855 and *F. bispinosa* (Morgan, 1913) (Thysanoptera: Thripidae).

Spinosad was used against the olive fly *Bactrocera oleae* (Gmelin,

Table 1. Target species investigated.

Higher taxon	Family	Scientific name	Common name	
Acari	Tetranychidae	<i>Tetranychus urticae</i>	Red spider mite	
Coleoptera	Bostrichidae	<i>Rhyzopertha dominica</i>	Lesser grain borer	
		<i>Prostephanus truncatus</i>	Larger grain borer	
	Bruchidae	<i>Callosobruchus maculatus</i>	Cowpea weevil	
		Dryophthoridae	<i>Sitophilus granarius</i>	Grain weevil
	<i>Sitophilus oryzae</i>		Rice weevil	
	<i>Sitophilus zeamais</i>		Maize weevil	
	Laemophloeidae		<i>Cryptolestes ferrugineus</i>	Rusty grain beetle
		<i>Cryptolestes pusillus</i>	Flat grain beetle	
	Silvanidae	<i>Oryzaephilus surinamensis</i>	Sawtoothed grain beetle	
	Tenebrionidae	<i>Tribolium confusum</i>	Confused flour beetle	
<i>Tribolium castaneum</i>		Red flour beetle		
Diptera	Agromyzidae	<i>Liriomyza sativae</i>	Tomato leafminer	
		<i>Liriomyza huidobrensis</i>	Pea leafminer	
		<i>Liriomyza trifolii</i>	American serpentine leafminer	
	Culicidae	<i>Culex quinquefasciatus</i>	Southern house mosquito	
	Drosophilidae	<i>Drosophila melanogaster</i>	Fruit fly	
	Muscidae	<i>Musca domestica</i>	Housefly	
	Tephritidae	<i>Rhagoletis cerasi</i>	Cherry fruit fly	
		<i>Ceratitis capitata</i>	Mediterranean fruit fly	
	<i>Bactrocera oleae</i>	Olive fruit fly		
	Hemiptera	Aleyrodidae	<i>Bemisia tabaci</i>	Sweetpotato whitefly
Aphididae		<i>Sitobion avenae</i>	English grain aphid	
Psyllidae		<i>Cacopsylla pyri</i>	European pear psylla	
Tessaratomidae		<i>Tessaratomia javanica</i>	Giant stink bugs	
Encyrtidae		<i>Tachardiaephagus tachardiae</i>	None	
Eulophidae		<i>Aprostocetus purpureus</i>	None	
Eupelmidae		<i>Eupelmus tachardiae</i>	None	
Lepidoptera	Blastobasidae	<i>Pseudohypatopa pulvereana</i>	Black caterpillar	
	Crambidae	<i>Chaphalocrocis medinalis</i>	Rice leafroller	
		Gelechiidae	<i>Sitotroga cerealella</i>	Angoumois grain moth
	<i>Tuta absoluta</i>		Tomato leafminer	
	Noctuidae		<i>Eublemma amabilis</i>	Lac insects moth
			<i>Helicoverpa armigera</i>	Cotton bollworm
			<i>Helicoverpa zea</i>	Corn earworm
			<i>Heliothis virescens</i>	Tobacco budworm
			<i>Spodoptera exigua</i>	Beet armyworm
			<i>Spodoptera littoralis</i>	Cotton leafworm
			<i>Spodoptera litura</i>	Common cutworm
	<i>Trichoplusia ni</i>		Cabbage looper	
	Pieridae	<i>Pieris rapae crucivora</i>	Common white	
	Plutellidae	<i>Plutella xylostella</i>	Diamondback moth	
	Pyralidae	<i>Plodia interpunctella</i>	Indian meal moth	
		<i>Cadra cautella</i>	Almond moth	
		<i>Corcyra cephalonica</i>	Rice moth	
		Tortricidae	<i>Adoxophyes honmai</i>	Smaller tea tortrix
			<i>Adoxophyes orana fasciata</i>	Summer fruit tortrix
			<i>Cydia molesta</i>	Oriental fruit moth
			<i>Cydia pomonella</i>	Codling moth
	<i>Eupoecilia ambiguella</i>		European grape berry moth	
	<i>Homona magnanima</i>		Oriental tea tortrix	
	<i>Lobesia botrana</i>	Gravelpine moth		
	<i>Pandemis cerasana</i>	Barred fruit-tree tortrix		
	Psocoptera	Liposcelididae	<i>Liposcelis entomophila</i>	Booklice
		Trogiidae	<i>Lepinotus reticulatus</i>	Reticulatedwinged booklouse
Thysanoptera	Thripidae	<i>Frankliniella occidentalis</i>	Western flower thrips	
		<i>Frankliniella bispinosa</i>	Florida flower thrips	
		<i>Frankliniella tritici</i>	Eastern flower thrips	
		<i>Scirtothrips dorsalis</i>	Yellow tea thrips	
		<i>Thrips palmi</i>	Melon thrips	

1790) (Diptera: Tephritidae) a dangerous parasite of olives in most of the countries around the Mediterranean Sea. However the study of the effect of the technological process on pesticide transfer in olive oil showed that spinosad tends to remain in the olive cake (Angioni *et al.*, 2011). The residue level found in olives after treatment was 0.30 mg/kg.

The spinosad GF-120 fruit fly bait (Dow AgroSciences, 2013) was tested in several experiments against *Rhagoletis cerasi* (Linnaeus, 1758) (Diptera: Tephritidae) together a treatment with entomopathogenic fungi (Daniel & Grunder, 2012).

Fruits treated with spinosad formulations with bait (Spintor®; Dow AgroSciences Italia s.r.l., Milano, Italy) suffered 8 times less damage than when treated with other attract-and-kill prototype devices against *Ceratitis capitata* (Wiedeman, 1824) (Navarro-Llopis *et al.*, 2013) (Diptera: Tephritidae).

In comparison with methoxyfenozide and chlorantraniliprole, spinetoram was effective against the Lepidoptera Tortricidae *Eupoecilia ambiguella* (Hübner, 1796) and even more on *Lobesia botrana* (Denis & Schiffermüller, 1775), the European grapevine moths, species causing serious damages to vineyards mainly with generations that feed on berries (Forte *et al.*, 2014). Spinetoram was also effective against *Cacopsylla pyri* (Linnaeus, 1761) (Hemiptera: Psyllidae) codling moth *Cydia pomonella* (Linnaeus, 1758), oriental fruit moth, *C. molesta* (Busck, 1916) and barred fruit-tree *Pandemis cerasana* Hübner, 1786 (Lepidoptera: Tortricidae) (Boselli & Scannavini, 2014; Tesconi *et al.*, 2014) with no permanent effects against beneficial arthropods, secondary insects or mites populations.

Spinetoram and *Bacillus thuringiensis* gradually replaced organophosphates (Bacci *et al.*, 2014) in the control of the grape berry moth, *Lobesia botrana* (Lepidoptera: Tortricidae) that mainly needs effective spray programmes against generations that feed on grape berries, in central and southern Italy vineyards.

Spinosad was used against litchi stinkbug *Tessaratomia javanica* (Thunberg, 1783) (Hemiptera: Tessaratomidae) a pest of subtropical fruit litchi. The relative toxicity against first instar nymphs of litchi stink bug revealed that chlorantraniliprole, thiacloprid, thiodicarb and spinosad were 36.83, 27.62, 22.10 and 10.04 times more toxic with reference to novaluron, respectively (Choudhary *et al.*, 2015).

A field experiment was conducted at Agronomy Instructional Farm, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar (Gujarat) during the season of monsoons, characterized by humid climate for crops, in 2012 and 2013. Based on number of larvae per plant, flower percentage and capsule infestation recorded after two sprays it was concluded that all the treatments showed significant difference in reducing larval population, and flower and capsule infestation over control. Spinosad 0.001% was found significantly most effective followed by profenofos 0.05%, dichlorvos 0.05%, acephate 0.075% and triazophos 0.04%, whereas imidacloprid 0.01%, neem oil 1% and the entomopathogen fungus *Beauveria bassiana* 2×10^8 cfu/gm were the least effective. Maximum yield was obtained with spinosad followed by profenofos and dichlorvos (Wazire & Patel, 2015).

Table 2. Relative toxicity of different insecticides and their lethal concentration at 50%.

	IGR/spinosad	IGR/spinetoram	LC ₅₀ 72 h
Lufenuron	9.8	2.0	4.4
Chlorfluazuron	9.4	1.9	4.6
Methoxyfenozide	9.2	1.8	4.7
Spinetoram	-	-	8.6
Spinosad	-	-	43.1

IGR, insect growth regulator; LC₅₀, lethal concentration at 50%.

Spinosad applied at doses varying between 48 to 120 g a.i./ha (one or 2 times at 14 days interval) was effective against tomato leafminer, *Tuta absoluta* (Meyrick, 1917) (Lepidoptera: Gelechiidae) and did not emphasize product phytotoxicity (Bratu *et al.*, 2015).

A storage experiment was conducted to study the effect of newer insecticide molecules in combating *Callosobruchus maculatus* (Fabricius, 1775) (Coleoptera: Bruchidae) on storability of chickpea variety JG-11 under ambient storage condition from August 2012 to May 2013. The treatment of spinosad 45 SC at 2 ppm followed by emamectin benzoate 5 SG at 2 ppm were found free from the seed damage (natural infestation) and presence of live *C. maculatus* adults were statistically at par with the rest of insecticidal treatments (Vidyashree *et al.*, 2015).

The results of toxicity tests on target species are summarized in Table 3.

Toxicity on insect predators or parasitoids of useful species

As *Kerria lacca* (Kerr, 1782) (Hemiptera: Tachardiidae) popularly known as *Laccifer lacca* is an insect used for lac production, the control of natural enemies of this insect is sometimes necessary. The response of insecticides on emergence of predators and parasitoids that can threaten brood-lac was assessed under laboratory conditions by dipping kusmi broodlac (kusmi is a strain of host plants) in insecticidal formulations and subsequent inoculation of treated broodlac on lac host plants, *Flemingia semialata* Roxburg, and on the Oken-Kusum tree *Schleichera oleosa* (Lour.). Insecticidal solution of indoxacarb (0.007, 0.014 and 0.021%), spinosad (0.005, 0.007 and 0.01%), fipronil (0.007, 0.014 and 0.02%) and ethofenprox (0.02, 0.03 and 0.04%) were applied for 15 min. No detrimental effect of insecticides on emergence and survival were noticed. Normal emergence and settlement on lac host *F. semialata* was seen clearly indicating the safety of insecticides. On the other side significant reduction in lepidopteran predators and hymenopteran parasitoids population was observed from the treated broodlac. Maximum reduction in the emergence of *Eublemma arabilis* Moore, 1884 (Lepidoptera: Noctuidae), a predator of lac insects, was observed with spinosad (100%) followed by indoxacarb (97.92 to 100%), ethofenprox (75 to 93.75%) and fipronil (72.92 to 91.67%). All the insecticides have shown very good response on the predator *Pseudohypatopa pulverea* (Meyrick, 1907) (Lepidoptera: Blastobasidae). Emergence of parasitoids of lac insects, *Tachardiaephagus tachardiae* Hovard, 1896 (Hymenoptera: Encyrtidae), *Aprostocetus purpureus* (Cameron, 1913) (Hymenoptera: Eulophidae) and *Eupelmus tachardiae* (Hovard, 1896) (Hymenoptera: Eupelmidae) was significantly low from treated broodlac. Reduction in population of *T. tachardiae* in different treatments varied from 47.06 to 89.71%, *A. purpureus* from 61.54 to 100%, *E. tachardiae* (male) from 38.46 to 100% and *E. tachardiae* (female) from 45.45 to 100%. This study clearly indicates that the treatment of broodlac prior to inoculation can be safely and effectively used as a tool in integrated pest management (IPM) programmes; selective insecticides namely indoxacarb, fipronil, spinosad and ethofenprox can be safely and effectively used (Singh & Jaiswal, 2015).

Toxicity on beneficial and non target organisms

The efficacy of an insecticide on target species is of primary importance to select it among other products, but its effects on beneficial and non target species and on human health must also be considered. The

toxicity to non-target taxa is summarized in Table 4. A summary of toxicity results is available at: http://www.sumitomo-chem.co.jp/english/red/report/theses/docs/2012E_1.pdf (accessed 9/4/2015).

A chronic toxicity test in mammals showed that spinosad has no carcinogenic, teratogenic, mutagenic, or neurotoxic effects, so it is not considered dangerous for human health. In fact there was no-observed-effect-level of spinosad in a 13-week studies on rats was 0.012% (24 mg/kg/day) and there was no evidence of a treatment-related increase in tumors in any rat tissue (Yano *et al.*, 2002).

A comparison of spinosyns with other insecticides was carried out to analyse the effect on non-target species. The results were that spinosyns were generally the least toxic of all insecticides for the non-target species analysed. Non-target species belong to different order of insects and to all the other living organisms, including mammals, birds, other vertebrates and plants.

Williams *et al.* (2003) in his review on the effect of spinosad on beneficials analysed a database of researches conducted on 228 observations and on 52 species of natural enemies (27 predators and 25 parasitoids), detailing laboratory, semi field and field tests. The results of this study evidenced that spinosad has little impact on predator populations, even if earwigs and ants can be vulnerable to spinosad (Malagnoux *et al.*, 2015). In the same article the Authors evidenced that spinosad can be more harmful to parasitoids. Other researches demonstrated that spinetoram might be toxic to some non-target parasitoids playing a prominent role in biological control of pests (Nasreen *et al.*, 2000; Tillman & Mulrooney, 2000; Consoli *et al.*, 2001). Spinosad, as other pesticides (abamectin, chlorfenapyr, emamectin benzoate, malathion, nitenpyram, permethrin) caused 100% mortality against the parasitoid *Aphidius gifuensis* Ashmead, 1906 (Hymenoptera: Braconidae). For this reason it could be categorized as seriously harmful based on the International Organization for Biological and Integrated Control/West Palaearctic Regional Section (IOBC/WPRS) guideline. However spinosad had no harmful effects on the emergence of *A. gifuensis* adults living in the mummified aphid *Sitobion avenae* (Fabricius, 1775) (Homoptera: Aphididae) while abamectin, malathion, thiamethoxam and tolfenpyrad determined mortalities >30% (Ohta & Takeda, 2015).

About predators Elzen *et al.* (1998) investigated the effect of 10 insecticides, including spinosad, in a spray chamber on the adults of the following species: *Ortus insidiosus* (Say, 1832) (Hemiptera: Anthocoridae), *Geocoris punctipes* (Say, 1832) (Hemiptera: Lygaeidae), *Hippodamia convergens* Guérin-Ménéville, 1842 (Coleoptera: Coccinellidae) and *Chrysoperla carnea* (Stephens, 1836) (Neuroptera: Chrysopidae).

They demonstrated a considerable variation in response among the species tested to the insecticides. In detail *C. carnea* was highly sensitive to most of the insecticides, and malathion was the most toxic to all species. Cyfluthrin, profenofos, endosulfan, spinosad and oxamyl caused no mortality in *G. punctipes*.

Spinosad is effective on the pest thrip *F. occidentalis* (Thysanoptera: Thripidae), but ineffective on its predator anthocorid *O. insidiosus* (Harrison & Rahman, 2014).

Cote *et al.* (2002) examining the effect of many acaricide including spinosad, on *Phytoseiulus persimilis* Athias-Henriot, 1957 (Acari: Phytoseiidae), in leaf-disk assays found no greater mortality of *P. persimilis* in spinosad-treated leaf disks than in water controls. Besides at rates recommended for control of pest insects, spinosad is harmless (<30% mortality) to most predatory mites (Williams *et al.*, 2003).

However a population resistant to insecticides (ETO6) including spinosad was selected (Yorulmaz Salman *et al.*, 2015), so spinosad can be used in conjunction with biological control.

Adult acetylcholinesterase and carboxylesterases activities were significantly reduced (28 to 67% of controls) in earwigs *Forficula auricularia* L. (Dermaptera: Forficulidae) after the application of spinosad and chlorpyrifos-ethyl, respectively. *F. auricularia* was considered as pest in stone-fruit orchards especially when fruit is close to maturity, but the insect revealed to be also an effective predator of aphids, leafrollers, and psyllids in apple orchards (Malagnoux *et al.*, 2015). It is an interesting case of an insect species that can be considered a target of treatment in some situations but also a useful species for biological control in other situations.

Stevens *et al.* (2005) also evidenced that spinosad is active against *Chironomus tepperi* Skuse, 1889 (Diptera: Chironomidae) with laboratory 24 h LC₅₀ and LC₉₀ estimated at 28.9 and 61.8 g L⁻¹, respectively. In outdoor containers treated with spinosad at 5 ppm (equivalent to 5 mg L⁻¹), chironomid larvae were not able to develop.

For *Chironomus circumdatus* Kieffer, 1916 (Diptera: Chironomidae) laboratory 48-h LC₉₀ of spinosad was estimated at 63 and 177 g L⁻¹ for the 1st and 4th larval instars, respectively (Kumar *et al.*, 2011). Duchet *et al.* (2015) found that following spinosad treatments, emergence of *Polypedilum nubifer* (Skuse, 1889) (Diptera: Chironomidae), dramatically decreased from day 4 up to the end of the observation period. Adult emergence never recovered in spite of the rapid degradation of the insecticide, with a half-life of 1-2 d for the sum of spinosyns A and D (Duchet *et al.*, 2008). Concerning *Tanytarsus curticornis* Kieffer, 1911 (Diptera: Chironomidae), adult emergence remained low in the enclo-

Table 3. Insecticidal activity of spinetoram on major pests.

	Name	Pests Scientific name	Growth stage	Crop	Methods	Days after treatment	LC ₅₀ (ppm)
Lepidoptera	Diamondback moth	<i>Plutella xylostella</i>	3 rd instar larva	Cabbage	Dipping (leaf)	4	0.01
	Common cutworm	<i>Spodoptera litura</i>	3 rd instar larva	Cabbage	Dipping (leaf)	4	1.17
	Common white	<i>Pieris rapae crucivora</i>	Mid-instar larva	Cabbage	Dipping (leaf)	4	0.02
	Cotton bollworm	<i>Helicoverpa armigera</i>	3 rd instar larva	Cabbage	Dipping (leaf)	4	0.08
	Cabbage looper	<i>Trichoplusia ni</i>	3 rd instar larva	Cabbage	Dipping (leaf)	4	0.01
	Smaller tea tortrix	<i>Adoxophyes honmai</i>	Mid-instar larva	Tea	Dipping (leaf)	10	0.94
	Oriental tea tortrix	<i>Homona magnanima</i>	3 rd instar larva	Tea	Dipping (leaf)	4	0.87
	Summer fruit tortrix	<i>Adoxophyes orana fasciata</i>	3 rd instar larva	Apple	Forliar spray	4	0.11
	Rice leafroller	<i>Cnaphalocrocis medinalis</i>	Late instar larva	Rice	Dipping (leaf)	4	0.06
	Thysanoptera	Melon thrips	<i>Thrips palmi</i>	Adult	Cucumber	Dipping (leaf)	3
Yellow tea thrips		<i>Scirtothrips dorsalis</i>	Adult	Tea	Dipping (leaf)	3	0.038
Diptera	Tomato leafminer	<i>Liriomyza sativae</i>	Early instar larva	Cucumber	Dipping (leaf)	3	23: 100%
	Pea leafminer	<i>Liriomyza huidobrensis</i>	Early instar larva	Cucumber	Dipping (leaf)	3	23: 100%
Hemiptera	Sweet potato whitefly (biotype Q)	<i>Bemisia tabaci</i>	First instar nymph	Cabbage	Dipping (leaf and insect)	4	47: 98%

LC₅₀, lethal concentration at 50%. Data from Shimokawatoko *et al.* (2012).

tures treated with spinosad at 17 and 33 g L⁻¹, though not statically different from control. The emergence was higher in the enclosures treated with spinosad at 8 g L⁻¹. The short half-life of spinosad resulted in the fast dissipation of such a low concentration of the insecticide from the water, so that *T. curticornis* could rapidly recolonize the enclosures from eggs laid by females. In addition, it cannot be excluded that even at low concentration, spinosad may have reduced the abundance of chironomid larvae competitors and/or predators. This may explain why adult emergence rate was higher in the enclosures treated with 8 g L⁻¹ spinosad as compared to the control enclosures and the enclosures treated with *Bti*, which is known to preserve non-target invertebrate communities in wetlands (Lagadic *et al.*, 2014).

The toxicity against non-target Crustacea was conducted in experiments with Conserve[®] 120SC (a product with spinosad as active ingredient by Dow AgroSciences Italia s.r.l.) in field enclosures implemented in Atlantic and Mediterranean wetlands (Duchet *et al.*, 2008; Duchet *et al.*, 2010b). Natural populations of *Daphnia pulex* Leydig, 1860 and *D. magna* Straus, 1820 (Crustacea: Cladocera) were dramatically reduced by the exposure to 8, 17 and 33 g L⁻¹ spinosad. Although recovery was observed in *D. pulex* in the Atlantic wetlands (Duchet *et al.*, 2008), population model forecasted quasi-extinction after 43.9 weeks (Duchet *et al.*, 2010a).

Special consideration has to be given also to bees (both social and solitary bees). Direct and indirect effects have to be considered; sublethal effects may have impacts on bees and pollination in addition to the more easily observable mortality, causing decreased pollination and affecting bee reproduction and development. Bee larvae, feeding on exogenous pollen containing residues of different active principles during development may result in lethal or sublethal effects during the adult stage. Scott-Dupree *et al.* (2009) in their work on the effect of dif-

ferent insecticides on *Bombus impatiens* (Hymenoptera: Apidae), *Megachile rotundata* (Hymenoptera: Megachilidae), and *Osmia lignaria* (Hymenoptera: Megachilidae) evidenced that spinosad was only moderately toxic for these bees. Adult stingless bee workers of *Melipona quadrifasciata* Lepeletier, 1836 (Hymenoptera: Apidae) exhibited high oral insecticide susceptibility, with LD₅₀ of 23.54 and 12.07 ng a.i./bee for imidacloprid and spinosad, respectively (Tomé *et al.*, 2015). Morandin *et al.* (2005) studied the effect on *Bombus impatiens* Cresson 1863 (Hymenoptera: Apidae) colony health evidencing a detrimental effect at the dose of spinosad at of 8.0 mg kg⁻¹ in pollen, and minimal effect at the dose of 0.8 mg kg⁻¹.

Miles (2003) studies demonstrated that spinosad was safe to *Apis mellifera* (Linnaeus, 1758) (Hymenoptera: Apidae) when applied to flowering crops during periods of low bee activity. Bayley *et al.* (2005) comparing direct and residual contact and oral toxicities to honey bees posed little to no risk.

Spinosad was initially considered safe for several non-target arthropods, and its use was allowed and expanded for crop protection (Miles, 2003; Sarfraz *et al.*, 2005), especially in organic production, but such perceived selectivity has been challenged (Biondi *et al.*, 2012). The LD₅₀ estimates obtained with the probit model were 23.54 (3.8-81) and 12.07 ng a.i./bee for imidacloprid and spinosad, respectively (Figure 2).

Action at cellular level

Spinosad has a unique mechanism of action involving the disruption of nicotinic acetylcholine receptors and γ -aminobutyric acid (GABA)-

Table 4. Ecotoxicological summary and environmental fate values of spinetoram on non-target organisms.

Test substance	Test	Species	Test type	Results
Spinetoram	Aquatic organisms	<i>Carp</i>	Acute (96 h)	LC ₅₀ =3.9 mg/L
		Rainbow trout	Acute	LC ₅₀ >3.46 mg/L (ppm)
		<i>Daphnia magna</i>	Acute (48 h)	EC ₅₀ >3.17 mg/L
	Honeybee	<i>Green algae</i>	Acute (72 h)	ErC ₅₀ =1.060 mg/L
		<i>Apis mellifera</i>	Acute contact (48 h)	LD ₅₀ =24.8 ng/bee
	Bird	<i>Bobwhite quail</i>	Acute oral	LD ₅₀ >2250 mg/kg
		Mallard duck	Acute	LD ₅₀ >2250 mg/kg
		Rat	Acute oral	LD ₅₀ >5000 mg/kg
		Rat	Acute dermal	LD ₅₀ >5000 mg/kg
		Earthworm	Acute	LC ₅₀ >1000 mg/kg soil
		Terrestrial dissipation half-life (field soil)		2 days minor factor 4 days major factor
	Aquatic dissipation half-life (natural surface water)		0.5 days minor factor 0.6 days major factor	
Spinetoram 25%WDG	Aquatic organisms	<i>Carp</i>	Acute (96 h)	LC ₅₀ =24 mg/L
		<i>Daphnia magna</i>	Acute (48 h)	EC ₅₀ >24 mg/L
		<i>Green algae</i>	Acute (72 h)	ErC ₅₀ =19 mg/L
	Honeybee	<i>Apis mellifera</i>	Residual toxicity test*2	≤3 days
	Silkworm	<i>Bombyx mori</i>	Residual toxicity test*3	≤31 days
	Spinetoram 11.7%SG	Aquatic organisms	<i>Carp</i>	Acute (96 h)
<i>Daphnia magna</i>			Acute (48 h)	EC ₅₀ >54 mg/L
<i>Green algae</i>			Acute (72 h)	ErC ₅₀ =530 mg/L
Honeybee		<i>Apis mellifera</i>	Residual toxicity test*4	? 7 days
		Natural enemy insects	<i>Paederus fuscipes (adult)</i>	Acute contact (48 h)
		<i>Harmonia axyridis (larvae)</i>	Acute contact (48 h)	Mortality 3.3% (at 50 mg a.i./L)
		<i>Chrysoperla carnea (larvae)</i>	Acute contact (96 h)	Mortality 10% (at 50 mg a.i./L)
Spinetoram 0.5%GR		Aquatic organisms	<i>Carp</i>	Acute contact (96 h)
	<i>Daphnia magna</i>		Acute contact (48 h)	EC ₅₀ >1000 mg/L
	<i>Green algae</i>		Acute contact (72 h)	ErC ₅₀ >1000 mg/L

LC₅₀, lethal concentration at 50%; EC₅₀, effective concentration at 50%; ErC₅₀, concentration related to growth at 50%; LD₅₀, lethal dose at 50%; a.i., active ingredient.

gated ion channels of insect nervous systems (Salgado & Sparks, 2005; Kirst, 2010) (Figure 3).

It is here remembered that there are two types of acetylcholine receptors (AChR) that bind acetylcholine and transmit its signal: muscarinic and nicotinic AChR, which are named after the agonists muscarine and nicotine, respectively. These receptors are functionally different, the muscarinic type being G-protein coupled receptors (Sadava *et al.*, 2014) that mediate a slow metabolic response via second messenger cascades, while the nicotinic type are ligand-gated ion channels that mediate a fast synaptic transmission of the neurotransmitter.

Spinosyns act as allosteric activators of nicotinic AChR receptors (Figure 4). Spinosyns act through a site in the nicotinic receptor that is distinct from other neo-nicotinoids or nicotinic actives.

In particular spinosyns act on a special type of nicotinic acetylcholine receptor (nAChR): Dm 6, which is known as D α 6 in *Drosophila melanogaster* Meigen, 1830 (Diptera: Drosophilidae) (Perry *et al.*, 2015), while spinosins do not respond to the similar nicotinic acetylcholine receptor D α 7 (Somers *et al.*, 2015).

nAChR is an excitatory receptor (cation-selective) for acetylcholine. It is the activation of this α 6-nAChR by the spinosyns that begins the cascade of events leading to insect death (Salgado & Saar, 2004; Dripps *et al.*, 2008).

In cockroach two nicotinic receptors were identified: non-desensitizing nACh (nAChN) and desensitizing nACh (nAChD) receptors; non-desensitizing receptors are not deactivated by repeated stimulations; nAChD is desensitized by neonicotinoids as imidacloprid, while nAChN is not desensitized but it is inhibitable with methylcaconitine; nAChN receptors are activated allosterically by spinosyn A with an effective concentration at 50% of 27 nM, that is at very low nanomolar concentrations, while are activated by other neonicotinoids only at much higher concentrations: that is micromolar instead of nanomolar. Spinosyn A




Figure 2. Mortality against active ingredient (a.i.) per bee using two different insecticides. LD₅₀, lethal dose at 50%.




Figure 3. Mechanism of transmission. GABA, γ -aminobutyric acid.

weakly antagonizes nAChD receptors at micromolar concentrations (10 μ M) (Salgado & Saar, 2004).

Spinosad is an agonist of nicotinic acetylcholine receptors and interferes also with receptors of GABA in the nervous system (Salgado, 1998; Sparks *et al.*, 2012) as an antagonistic. GABA receptors are responsible of inhibiting neural excitation, generating a hyperpolarization due to the opening of the chloride channels. As agonistic of nAChR and antagonistic of GABA receptors, spinosad acts exciting the insect nervous system. Initial investigations had concluded that the primary action of spinosyn A affected the insect nervous system and disrupted neuronal activity by exciting motor neurons and causing involuntary muscle contractions, eventually leading to paralysis and death.

Spinosyn probably ligates in a transmembrane site, in a manner sim-




Figure 4. Allosteric action of spinosad on nicotinic receptors; spinosad ligates to a different site respect to the acetylcholine site.

ilar to ivermectin, which ligates to a receptor situated in the glutamate-gated chloride channel (GluCl) (Puinean *et al.*, 2013). The mechanism of action of this family of transmembrane receptors was well studied in GluCl complex which interacts with the allosteric partial agonist ivermectin (Hibbs & Gouaux, 2011; Althoff *et al.*, 2014); this complex is assumed to have a structure similar to the one of nAChR receptors, in this case spinosyns instead of ivermectin act as allosteric effectors (Figure 5). The mechanism of action of spinosad on nAChR receptors is not well known as the one of ivermectin on GluCl receptors, but it is supposed that the mechanism of action are similar (Figure 5).

Spinosad manner of action differs deeply from the pyrethroids action operating on sodium channels. The toxic effects of pyrethroids are mediated through preventing the closure of the voltage-gated sodium channels in the axonal membranes, while spinosad acts with an allosteric mechanism on nAChR receptors.

Resistance and cross-resistance

The emergence of resistant insects is a common situation when an insecticide is spread for long time (Roush & Tabashnik, 1991; Lawrence & Sarjeet, 2010), thus the potential development of a resistance in an insect should be evaluated.

Resistance to spinosad was forced in larvae of *H. virescens* in laboratory after 14 generation using a specific protocol (Bailey *et al.*, 1999). Similar results were obtained on *Musca domestica* Linnaeus, 1758 (Diptera: Muscidae) in laboratory (Shono & Scott, 2003). In addition to the above-mentioned studies, several other studies have demonstrated the development of spinosad resistance in the laboratory and the field in such diverse species as *D. melanogaster*, *Liriomyza trifolii* (Burgess 1880) (Diptera: Agromyzidae), *Helicoverpa armigera* (Hübner, 1809) (Lepidoptera: Noctuidae), and *F. occidentalis* (Sparks *et al.*, 2012).

At present it is clear that in *D. melanogaster* the resistance to spinosad is bound to a mutation of the D 6 receptors (Perry *et al.*, 2007; Somers *et al.*, 2015). The Da6P146S mutation was recreated using the clustered regularly interspaced short palindromic repeats/Cas9 system, this is the first use of this technology to introduce a resistant mutation into a controlled genetic background (Somers *et al.*, 2015). Shono & Scott (2003) demonstrated that spinosad resistance is linked to a non-sex linked single recessive gene. It was observed that sublethal-spinosad-treated (Spin-Sub) strains of *F. occidentalis* developed physiological and biochemical adaptations after a long-term treatment with spinosad (You-Hui *et al.*, 2015).

Cloning the nicotinic acetylcholine receptor $\alpha 6$ subunit from *F. occi-*

dentalis (Foc6), it was demonstrated (Puinean *et al.*, 2013) that in nAChR a glycine replacement with glutamic acid is responsible of insect resistance to spinosad. As stated above GluCl found in different insects has a binding site for ivermectin, a lactone with similar insecticidal effect and it is supposed that a similar binding site for spinosad is present in nAChR; all evidences are in favour of the hypothesis that spinosad has an allosteric mechanism of action on nAChR as ivermectin has on GluCl (Figure 5). It is interesting to note that spinosad has an excitatory effect on insects' receptor nAChR $\alpha 6$, but an inhibitory effect of human nAChR 7 receptor, which is similar but not identical with the insects' nAChR 6 receptor.

A problem connected with resistance is also the cross-resistance, which is observed when the same mechanism of resistance allows the insect to resist to different insecticides. To date, in nearly 90 studies that examined cross-resistance to the spinosyns in strains resistant to a wide spectrum of other classes of insecticides, including dichlorodiphenyltrichloroethane, pyrethroids, avermectins, oxadiazines (*i.e.*, indoxacarb), carbamates, methoxyfenozide, and neonicotinoids (*i.e.*, imidacloprid), the level of cross-resistance observed to spinosad or spinetoram was none to very low (Sparks *et al.*, 2012).

The allosteric mechanism of action of spinosyns is different from the one of many other insecticides, so it reduces the risk of appearance of resistance phenomena.

Mutants resistant to spinosyns and spinetoram were recently reported (Puinean *et al.*, 2013; Su & Cheng, 2014), but spinosyns generally do not show cross-resistance with other existing chemicals, that is spinosyns resistant insects are not resistant to other insecticides (Lawrence & Sarjeet, 2010), demonstrating its utility against resistant strain. For example it was observed absence of cross-resistance to Bti and to a combination of Bti with other pesticides in spinosad-resistant population of *Culex quinquefasciatus* Say, 1823 (Diptera: Culicidae) (Su & Cheng, 2014). The reasons of lack of cross-resistance must be better investigated, but are probably bound to the different mechanism of action of spinosyns respect to other biocides including *B. thuringiensis* and *B. sphaericus*. Bti, a microbial pesticide, produces toxins causing destruction of the larval midgut leading to larval mortality, while spinosad acts as a neurotoxin.

A rare case of cross-resistance was observed in spinosad resistant populations of *C. quinquefasciatus*, which showed lower susceptibility respect to reference colony to a combination of *B. sphaericus*, spinetoram, abamectin and fipronil, emphasizing the existence of cross-resistance; it remains unknown why this spinosad resistant population resulted in cross-resistance to *B. sphaericus* (Su & Cheng, 2014). In any case at present there are no known important episodes of cross-resistance in insect species of interest in agriculture. In the same man-




Figure 5. Hypothetic structure of acetylcholine receptor (AChR) with detail of the allosteric site of attachment of spinosyns.

ner insects of sanitary interest as *M. domestica* when treated with the insecticide profenofos emphasized a similar response; selected strains of *M. domestica* resistant to profenofos did not show cross-resistance to spinosad (Khan *et al.*, 2015).

Recently, cases of resistance in field have been reported. A population strain of *S. exigua*, manifesting a reduced susceptibility to spinosad, has been detected in Thailand (Moulton *et al.*, 2000). An effective resistance to spinosins was reported on *Plutella xylostella* (Linnaeus, 1758) (Lepidoptera: Plutellidae) in Hawaii, where spinosad seemed to be the only opportunity to control this pest. After two years of intensive use the moth developed resistance to this insecticide (Zhao *et al.*, 2002).

Conclusions

Spinosyns are an important group of insecticides effective in the control of economically important pests in field and in post harvest control. The wide application of these actives ingredients is attributed to their selective and unique mode of action that sometimes makes their use as the only alternative to other insecticides (Zhao *et al.*, 2002).

The versatility in application is determined by both contact and ingestion that makes it efficient against numerous pests in the orders of Lepidoptera, Diptera, Thysanoptera, Coleoptera, Orthoptera, Hymenoptera, and others (Sparks *et al.*, 1995; Bret *et al.*, 1997).

Foliar applications of spinosad are not highly systemic in plants although some trans-laminar movement in leaf tissue has been demonstrated. Besides the addition of an appropriate surfactant increases trans-laminar movement and activity on pests that mine leaves (Larson, 1997).

Relating to plants no case of phytotoxicity has been signaled so far (Bratu *et al.*, 2015).

Considering that they are insecticides with precise target sites of action, the possibility that insects can develop a resistance exists. So far few cases of resistance were obtained in laboratory and also very few cases were reported from field. Thanks to its peculiar mode of action at cellular level no cross-resistance with other group of insecticides are signaled (Salgado & Sparks, 2005).

The results obtained in the attract-and-kill techniques, in which spinosyns are baited with different attractants, allow the use of small doses of the insecticides with a good efficacy.

Spinosyn are characterized by a broad spectrum of action, but they are also characterized by a low toxicity for natural enemies, especially predators. Trials on bees demonstrated a low toxicity also for these insects; nevertheless, there is still a need for testing field-realistic concentrations at relevant exposure and durations and, especially for honeybees, to continue side-effect evaluation over winter and the next year in spring.

Spinosad exhibits favorable safety profile with low mammalian toxicity, low toxicity to most non-target organisms and rapid degradation in several environmental matrices (Cleveland *et al.*, 2001). It is considered a natural product and thus approved for use in organic agriculture by numerous national and international certification bodies (Cleveland, 2007; Racke, 2007).

References

ALTHOFF T., HIBBS R.E., BANERJEE S., GOUAUX E., 2014 - X-ray structures of GluCl in apo states reveal a gating mechanism of Cys-loop receptors. - *Nature* 512: 333-339.
 ANGIONI A., PORCU L., PIRISI F., 2011 - LC/DAD/ESI/MS Method for the Determination of Imidacloprid, Thiacloprid, and Spinosad in olives

and olive oil after field treatment. - *J. Agric. Food Chem.* 59: 11359-11366.

- BACCI L., DESTEFANI G., CRIVELLI L., SITTI A., D'ASCENZO D., 2014 - Strategie con Methoxyfenozide e Spinetoram per il contenimento di *Lobesia botrana* su vite da vino. - *Atti Giorn. Fitopatol.* 1: 157-166.
 BAILEY J., SCOTT-DUPREE C., HARRIS R., TOLMAN J., HARRIS B., 2005 - Contact and oral toxicity to honey bees (*Apis mellifera*) of agents registered for use for sweet corn insect control in Ontario, Canada. - *Apidologie* 36: 623-633.
 BAILEY W.D., YOUNG H.P., ROE R.M., 1999 - Laboratory selection of a Tracer® resistant strain of tobacco budworm and comparisons with field selected strains from the southeastern. - *US. Proc. 1999 Beltwide Cotton Production Conf.*: 1221-1224.
 BIONDI A., MOMMAERTS V., SMAGGHE G., VIÑUELA E., ZAPPALÀ L., DESNEUX N., 2012 - The non-target impact of spinosyns on beneficial arthropods. - *Pest Manage. Sci.* 68: 1523-1536.
 BLOOMQUIST J.R., MILLER T.A., 1985 - A simple bioassay for detecting and characterising insecticide resistance. - *Pest. Sci.* 16: 611-614.
 BOSELLI M., SCANNAVINI M., 2014 - Efficacia di Spinetoram (Delegate Wg) nei confronti dei principali insetti dannosi al pero. - *Atti Giorn. Fitopatol.* 1: 27-36.
 BRATU E., PETUCI A.M., SOVAREL G., 2015 - Efficacy of the product Spinosad an insecticide used in the control of tomato leafminer (*Tuta absoluta* - Meyrick, 1917). *Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca. - Horticulture* 72: 209-210.
 BRET B.L., LARSON L.L., SCHOONOVER J.R., SPARKS T.C., THOMPSON G.D., 1997 - Biological properties of spinosad. - *Down Earth* 52: 6-13.
 CHOUDHARY J., MOANARO S., NAAZ N., IDRIS M.D., 2015 - Determination bio-efficacy of insecticides against litchi stink bug, *Tessaratoma javanica* (Thunberg) (Hemiptera: Tessaratomidae): an emerging major pest of litchi, *Litchi chinensis* Sonn. - *Bioscan* 10: 217-222.
 CLEVELAND C.B., 2007 - Environmental and health assessments for spinosad against the backdrop of organic certification. In: FELSOT A.J., RACKE K.D. (Eds.), *Certified organic and biologically-derived pesticides: environmental, health, and efficacy assessment. Symposium series.* - American Chemical Society, Washington, DC: 109-130.
 CLEVELAND C.B., MAYES M.A., CRYER S.A., 2001 - An ecological risk assessment for spinosad use on cotton. - *Pest Manage. Sci.* 58: 70-84.
 CONSOLI F.L., BOTELHO P.S.M., PARRA J.R.P., 2001 - Selectivity of insecticides to the egg parasitoid *Trichogramma galloi* Zucchi, 1988 (Hym., Trichogrammatidae). - *J. App. Entomol.* 125: 37-43.
 COTE K.W., EDWIN E.L., SCHULTZ P.B., 2002 - Compatibility of acaricide residues with *Phytoseiulus persimilis* and their effects on *Tetranychus urticae*. - *HortSci.* 37: 906-909.
 DANIEL C., GRUNDER J., 2012 - Review. Integrated management of European cherry fruit fly *Rhagoletis cerasi* (L.): situation in Switzerland and Europe. - *Insects* 3: 956-988.
 DOW AGROSCIENCES, 2013 - SPINETORAM: Insecticide for fruit, vegetables, grapes, and row crops. Technical Bulletin. - Dow AgroSciences Italia s.r.l., Milano, available from: http://msdssearch.dow.com/PublishedLiteratureDOWCOM/dh_091b/0901b8038091bc16.pdf?filepath=productsafety/pdfs/noreg/233-00382.pdf&fromPage=GetDoc
 DRIPPS J., OLSON B., SPARKS T., CROUSE G., 2008 - Spinetoram: how artificial intelligence combined natural fermentation with synthetic chemistry to produce a new spinosyn insecticide. - *Plant Health Progress August PHP-2008-0822-01-PS.*
 DUCHET C., CAQUET T., FRANQUET E., LAGNEAU C., LAGADIC L., 2010a - Influence of environmental factors on the response of a natural population of *Daphnia magna* (Crustacea: Cladocera) to

- spinosad and *Bacillus thuringiensis israelensis* in Mediterranean coastal wetlands. - Environ. Pollut. 158: 1825-1833.
- DUCHET C., COUTELLEC M.A., FRANQUET E., LAGNEAU C., LAGADIC L., 2010b - Population-level effects of spinosad and *Bacillus thuringiensis israelensis* in *Daphnia pulex* and *Daphnia magna*: comparison of laboratory and field microcosm exposure conditions. - Ecotoxicol. 19: 1224-1237.
- DUCHET C., FRANQUET E., LAGADIC L., LAGNEAU C., 2015 - Effects of *Bacillus thuringiensis israelensis* and spinosad on adult emergence of the non-biting midges *Polypedilum nubifer* (Skuse) and *Tanytarsus curticornis* Kieffer (Diptera: Chironomidae) in coastal wetlands. - Ecotoxicol. Environ. Saf. 115: 272-278.
- DUCHET C., LARROQUE M., CAQUET TH., FRANQUET E., LAGNEAU C., LAGADIC L., 2008 - Effects of spinosad and *Bacillus thuringiensis israelensis* on a natural population of *Daphnia pulex* in field microcosms. - Chemosphere 74: 70-77.
- EL-KADY G.A., EL-SHARABASY H.M., MAHMOUD M.F., BAHGAT I.M., 2007 - Toxicity of two potential bio-insecticides against moveable stages of *Tetranychus urticae* Koch. - J. Appl. Sci. Res. Nov. 1315-1319.
- ELZEN G.W., ELZEN P.J., KING E.G., DUGGER P., RICHTER D., 1998 - Laboratory toxicity of insecticide residues to *Orius insidiosus*, *Geocoris punctipes*, *Hippodamia convergens*, and *Chrysoperla carnea*. - Proc. Beltwide Cotton Conf., 5-9 January 1998, San Diego, CA, 2: 1235-1238.
- EUROPEAN COMMISSION, 2007 - Commission Directive 2007/6/EC of 14 February 2007 amending Council Directive 91/414/EEC to include metrafenone, *Bacillus subtilis*, spinosad and thiamethoxam as active substances. - Official Journal L 43, 15/2/2007, pp 13-18.
- EUROPEAN COMMISSION, 2012 - Commission Regulation (EU) No 556/2012 of 26 June 2012 amending Annex III to Regulation (EC) No 396/2005 of the European Parliament and of the Council as regards maximum residue levels for spinosad in or on raspberries. - Official Journal L 166, 27/6/2012, pp 67-80.
- EUROPEAN COMMISSION, 2014 - Commission Implementing Regulation (EU) No 140/2014 of 13 February 2014 approving the active substance spinetoram, in accordance with Regulation (EC) No 1107/2009 of the European Parliament and of the Council concerning the placing of plant protection products on the market, and amending the Annex to Commission Implementing Regulation (EU) No 540/2011. - Official Journal L 44, 14/2/2014, pp 35-39.
- EUROPEAN COMMISSION, 2015 - Commission Regulation (EU) 2015/603 of 13 April 2015 amending Annexes II, III and V to Regulation (EC) No 396/2005 of the European Parliament and of the Council as regards maximum residue levels for 2-naphthoxyacetic acid, acetochlor, chloropicrin, diflufenican, flurprimidol, flutolanil and spinosad in or on certain products. - Official Journal L 100, 17/4/2015, pp 10-59.
- FORTE V., ANGELINI E., CRESTANI D., PATRIARCA E., GANZINI L., PAVAN F., 2014 - Attività di metoxyfenozide e spinetoram contro le tignole della vite nell'Italia Nord-Orientale. - Atti Giorn. Fitopatol. 1: 147-156.
- HARRISON J., RAHMAN T., 2014 - Effect of new and old pesticides on *Orius armatus* (Gross) - an Australian predator of western flower thrips, *Frankliniella occidentalis* (Pergande) Sonya Broughton. - Pest Manage. Sci. 70: 389-397.
- HERTLEIN M.B., THOMPSON G.D., SUBRAMANYAM Bh., ATHANASIOU C.G., 2011 - Spinosad: a new natural product for stored grain protection. - J. Stored Prod. Res. 47: 131-146.
- HIBBS R.E., GOUAUX E., 2011 - Principles of activation and permeation in an anion-selective Cys-loop receptor. - Nature 474: 54-62.
- KHAN H.A.A., AKRAM W., IQBAL N., 2015 - Selection and preliminary mechanism of resistance to profenofos in a field strain of *Musca domestica* (Diptera: Muscidae) from Pakistan. - J. Med. Entomol. 52: 1013-1017.
- KIRST H.A., 2010 - The spinosyn family of insecticides: realizing the potential of natural products research. - J. Antibiotics 63: 101-111.
- KIRST H.A., MICHEL K.H., MYNDERSE J.S., CHAO E.H., YAO R.C., NAKATSUKASA W.M., BOECK L.D., OCCLOWITZ J., PASCHEL J.W., DEETER J.B., THOMPSON G.D., 1992 - Discovery, isolation and structure elucidation of a family of structurally unique fermentation-derived tetracyclic macrolides. In: BAKER D.R., FENYES J.G., STEFFENS J.J. (Eds.), Synthesis and chemistry of Agrochemicals, Vol. 3. - American Chemical Society, Washington, DC: 214-225.
- KUMAR A.N., MURUGAN K., MADHIYAZHAGAN P., PRABHU K., 2011 - Spinosad and neem seed kernel extract as bio-controlling agents for malarial vector, *Anopheles stephensi* and non-biting midge, *Chironomus circumdatus*. - Asian Pacif. J. Trop. Med. 4: 614-618.
- LAGADIC L., ROUCAUTE M., CAQUET T., 2014 - *Bti* sprays do not adversely affect non-target aquatic invertebrates in French Atlantic coastal wetlands. - J. Appl. Ecol. 51: 102-113.
- LARSON L.L., 1997 - Effects of adjuvants on the activity of Tracer® 480SC on cotton in the laboratory, 1996. - Arthropod Manage. Tests 22: 415-416.
- LAWRENCE I.G., SARJEET S.G., 2010 - Insect control: biological and synthetic agents. - Elsevier, London: 490.
- LEGOCKI J., POLEC I., ZELECHOWSKI K., 2010 - Contemporary trends in development of active substances possessing the pesticidal properties: spinosyn insecticides. - Pestycydy 1: 59-71.
- LEONARD B.R., GRAVES J.B., BURRIS E., MICINSKI S., MASCARENHAS V., MARTIN S.H., 1996 - Evaluation of selected commercial and experimental insecticides against lepidopteran cotton pests in Louisiana. - Proc. Beltwide Cotton Conf., January 9-12, 1996, Nashville, TN: 285-830.
- MALAGNOUX L., CAPOWIEZ Y., RAULT M., 2015 - Impact of insecticide exposure on the predation activity of the European earwig *Forficula auricularia*. - Environ. Sci. Pollut. Res. 22: 14116-14126.
- MERTZ F.P., YAO R.C., 1990 - *Saccharopolyspora spinosa* sp. nov. isolated from soil collected in a sugar rum still. - Int. J. Syst. Bacteriol. 40: 34-39.
- MILES M., 2003 - The effects of spinosad, a naturally derived insect control agent to the honeybee. - Bull. Insectol. 56: 119-124.
- MORANDIN L.A., WINSTON M.L., FRANKLIN M.T., ABBOTT V.A., 2005 - Lethal and sublethal effects of spinosad on bumblebees (*Bombus impatiens* Cresson). - Pest Manage. Sci. 61: 619-626.
- MOULTON J.K., PEPPER D.A., DENNEHY T.J., 2000 - Beet armyworm (*Spodoptera exigua*) resistance to spinosad. - Pest Manage. Sci. 56: 842-848.
- NASREEN A., ASHFAQ M., MUSTAFA G., 2000 - Intrinsic toxicity of some insecticides to egg parasitoid *Trichogramma chilonis* (Hym. Trichogrammatidae). - Bull. Inst. Trop. Agric. 23: 41-44.
- NAVARRO-LLOPIS V., PRIMO J., VACAS S., 2013 - Efficacy of attract-and-kill devices for the control of *Ceratitidis capitata*. - Pest Manage. Sci. 69: 478-482.
- OHTA I., TAKEDA M., 2015 - Acute toxicities of 42 pesticides used for green peppers to an aphid parasitoid, *Aphidius gifuensis* (Hymenoptera: Braconidae), in adult and mummy stages. - Appl. Entomol. Zool. 50: 207-212.
- PERRY T., MCKENZIE J.A., BATTERHAM P., 2007 - A D 6 knockout strain of *Drosophila melanogaster* confers a high level of resistance to spinosad. - Insect Biochem. Mol. Biol. 37: 184-188.
- PERRY T., SOMERS J., TING YANG Y., BATTERHAM P., 2015 - Expression of insect 6-like nicotinic acetylcholine receptors in *Drosophila melanogaster* highlights a high level of conservation of the receptor: spinosyn interaction. - Insect Biochem. Mol. Biol. 64: 106-115.
- PUINEAN A.M., LANSDELL S.J., COLLINS T., BIELZA P., MILLAR NS., 2013 - A nicotinic acetylcholine receptor transmembrane point mutation (G275E) associated with resistance to spinosad in *Frankliniella occidentalis*. - J. Neurochem. 124: 590-601.

- RACKE K.D., 2007 - A reduced risk insecticide for organic agriculture. In: FELSOT A.J., RACKE K.D. (Eds.), Certified organic and biologically-derived pesticides: environmental, health, and efficacy assessment. Symposium Series. - American Chemical Society, Washington, DC: 92-108.
- RAHMAN S.M.A., ABOU-TALEB H.K., 2007 - Joint toxic action of spinosad and spinetoram with certain IGR compounds against cotton leafworm. - Alexandria J. Agric. Res. 52: 45-51.
- ROUSH R.T., TABASHNIK B.E. (Eds.), 1991 - Pesticide resistance in Arthropods. - Springer, New York, NY: 303.
- SADAVA D., HILLIS D.M.H., CRAIG H., HELLER M., BEERENBAUM R., 2014 - Biologia 1. La cellula. Life: The science of biology. - Freeman & Company, New York; Transl. Zanichelli ed., Bologna: 208.
- SAGLAM O., ATHANASSIOU C.G., VASSILAKOS T.N., 2013 - Comparison of spinetoram, imidacloprid, thiamethoxam and chlorantranilprole against life stages of *Tribolium confusum* Jacquelin du Val (Coleoptera: Tenebrionidae) on concrete. - Crop Prot. 53: 85-95.
- SALGADO V.L., 1998 - Studies on the mode of action of spinosad: insect symptoms and physiological correlates. - Pest. Biochem. Physiol. 60: 91-102.
- SALGADO V.L., SAAR R., 2004 - Desensitizing and non-desensitizing subtypes of alpha-bungarotoxin sensitive nicotinic acetylcholine receptors in cockroach neurons. - J. Insect Physiol. 50: 867-879.
- SALGADO V.L., SPARKS T.C., 2005 - The spinosyns: chemistry, biochemistry, mode of action and resistance. - In: GILBERT L.I., IATROU K., GILL S. (Eds.), Comprehensive insect molecular science. Vol. 6. Elsevier, London: 137-173.
- SARFRAZ M., DOSDALL L. M., KEDDIE B.A., 2005 - Spinosad: a promising tool for integrated pest management. - Outlooks Pest Manage. 16: 78-84.
- SCOTT-DUPREE C.D., CONROY L., HARRIS C.R., 2009 - Impact of currently used or potentially useful insecticides for canola agroecosystems on *Bombus impatiens* (Hymenoptera: Apidae), *Megachile rotundata* (Hymenoptera: Megachilidae), and *Osmia lignaria* (Hymenoptera: Megachilidae). - J. Econ. Entomol. 102: 177-182.
- SHIMOKAWATOKO Y., SATO N., YAMAGUCHI Y., TANAKA H., 2012 - Development of the novel insecticide spinetoram (Diana®). - Sumitomo Chemical Co., Ltd., Tokyo.
- SHONO T., SCOTT J.G., 2003 - Spinosad resistance in the housefly, *Musca domestica*, is due to a recessive factor on autosome 1. - Pestic. Biochem. Physiol. 75: 1-7.
- SINGH J. P., JAISWAL A. K., 2015 - Response of broodlac treatment with insecticides on predators and parasitoids of lac insect, *Kerria lacca* (Kerr) harbouring broodlac. - Indian J. Entomol. 77: 21-26.
- SOMERS J., NGUYEN J., LUMB C., BATTERHAM P., PERRY T., 2015 - In vivo functional analysis of the *Drosophila melanogaster* nicotinic acetylcholine receptor Da6 using the insecticide spinosad. - Insect Biochem. Mol. Biol. 64: 116-127.
- SPARKS T.C., CROUSE G.D., DRIPPS J.E., ANZEVENO P., MARTYNOW J., DEAMICIS C.V., GIFFORD J., 2008 - Neural network-based QSAR and insecticide discovery: spinetoram. - J. Comput. Mol. Design 22: 393-401.
- SPARKS T.C., DRIPPS J.E., WATSON G.B., PAROONAGIAN D., 2012 - Resistance and cross-resistance to the spinosyns. A review and analysis. - Pest. Biochem. Physiol. 102: 1-10.
- SPARKS T.C., THOMPSON G.D., KIRST H.A., HERTLEIN M.B., LARSON L.L., WORDEN T.V., THIBAUT S.T., 1998 - Biological activity of the spinosyns, new fermentation derived insect control agents, on tobacco budworm (Lepidoptera: Noctuidae) larvae. - J. Econ. Entomol. 91: 1277-1283.
- SPARKS T.C., THOMPSON G.D., KIRST H.A., HERTLEIN M.B., MYNDERSE J.S., TURNER J.R., WORDEN T.V., 1999 - Fermentation-derived insect control agents. In: HALL F.R., MENN J.J. (Eds.), Biopesticides: use and delivery. - Humana Press, Totowa: 171-188.
- SPARKS T.C., THOMPSON G.D., LARSON L.L., KIRST H.A., JANTZ O.K., WORDEN T.V., HERTLEIN M.B., BUSACCA J.D., 1995 - Biological characteristics of the spinosyns: new naturally derived insect control agents. In: Proceedings of the Beltwide Cotton Conference, San Antonio, Texas, 4-7 January, 1995. - National Cotton Council of America, Memphis, TN: 903-907.
- SRIVASTAVA M., BOSCO L., FUNDERBURK J., OLSON S., WEISS A., 2008 - Spinetoram is compatible with the key natural enemy of *Frankliniella* species thrips in pepper. - Online. Plant Health Progress doi: 10.1094/PHP-2008-0118-02-RS.
- STEVENS M.M., HELLIWELL S., HUGHES P.A., 2005 - Toxicity of *Bacillus thuringiensis* var. *israelensis* formulations, spinosad, and selected synthetic insecticides to *Chironomus tepperi* larvae. - J. Am. Mosquito Control Assoc. 21: 446-450.
- SU T., CHENG M.L., 2014 - Cross resistances in spinosad-resistant *Culex quinquefasciatus* (Diptera: Culicidae). - J. Med. Entomol. 51: 428-35.
- TESCARI E., CHLORIDIS A., BACCI L., BRADASCIO R., GIBERTI A., 2014 - Spinetoram (Delegate®, Radiant®), Nuovo insetticida ad ampio spettro d'azione. - Atti Giorn. Fitopatol. 1: 11-20.
- THOMPSON G.D., DUTTON R., SPARKS T.C., 2000 - Spinosad - a case study: an example from a natural products discovery programme. - Pest Manage. Sci. 56: 696-702.
- TILLMAN P.G., MULROONEY J.E., 2000 - Effects of selected insecticides on the natural enemies *Coleomegilla maculata* and *Hippodamia convergens* (Coleoptera: Coccinellidae), *Geocoris punctipes* (Hemiptera: Lygaeidae), and *Bracon mellitor*, *Cardiochiles nigriceps*, and *Cotesia marginiventris* (Hymenoptera: Braconidae) in cotton. - J. Econ. Entomol. 93: 1638-1643.
- TOMÉ H.V.V., BARBOSA W.F., MARTINS G.F., NARCISO R., GUEDES C., 2015 - Spinosad in the native stingless bee *Melipona quadrifasciata*: regrettable non-target toxicity of a bioinsecticide. - Chemosphere 124: 103-109.
- VASSILAKOS T.N., ATHANASSIOU C.G., SAGLAM O., CHLORIDIS A.S., DRIPPS J.E., 2012 - Insecticidal effect of spinetoram against six major stored grain insect species. - J. Stored Prod. Res. 51: 69-73.
- VASSILAKOS T.N., ATHANASSIOU C.G., TSIROPOULOS N.G., 2015 - Influence of grain type on the efficacy of spinetoram for the control of *Rhyzopertha dominica*, *Sitophilus granarius* and *Sitophilus oryzae*. - J. Stored Prod. Res. 64: 1-7.
- VASSILAKOS T.N., ATHANASSIOU G.C., 2013 - Effect of temperature and relative humidity on the efficacy of spinetoram for the control of three stored product beetle species. - J. Stored Prod. Res. 55: 73-77.
- VIDYASHREE A.S., THIRUMALARAJU G.T., JYOTHI B.L., 2015 - Evaluation of newer molecule insecticides for the management of *Callosobruchus maculatus* Fab. on chickpea. - Environ. Ecol. 33: 877-879.
- WANG H., JUN Y., WEI X., YONG MEI H., RI-HE P., QUAN HONG Y., 2015a - Improving spinosad yield in *Saccharopolyspora spinosa* through antibiotic resistance screening technology and genome shuffling. - Acta Agric. Shanghai 31: 5-10.
- WANG H., WEI X., YONG MEI H., RI-HE P., QUAN-HONG Y., 2015b - Improvement of the ability to produce spinosad in *Saccharopolyspora spinosa* through the acquisition of drug resistance and genome shuffling. - Ann. Microbiol. 65: 771-777.
- WAZIRE N.S., PATEL J.I., 2015 - Efficacy of insecticides against sesamum leaf webber and capsule borer, *Antigastra catalaunalis* (Duponchel). - Indian J. Entomol. 77: 11-17.
- WHO (WORLD HEALTH ORGANIZATION), 2007 - Review of: Spinosad 0.5% GR and 12% SC, Lambda-Cyhalothrin 10% CS, K-O Table 1-2-3(r), Interceptor(r). - Report of the Tenth WHOPEs working group meeting; WHO/ CDS/ NTD/ WHOPEs/ 20 07.1.
- WILLIAMS T., VALLE J., VINUELA E., 2003 - Is the naturally derived

- insecticide spinosad compatible with insect natural enemies? - *Biocontr. Sci. Technol.* 13: 459-475.
- YANO B.L., BOND D.M., NOVILLA M.N., MCFADDEN L.G., REASOR M.J., 2002 - Spinosad insecticide: subchronic and chronic toxicity and lack of carcinogenicity in fischer 344 rats. - *Toxicol. Sci.* 65: 288-298.
- YORULMAZ SALMAN S., AYDINLI F., RECEP A.Y., 2015 - Etoxazole resistance in predatory mite *Phytoseiulus persimilis* A.H. (Acari: Phytoseiidae): Cross-resistance, inheritance and biochemical resistance mechanisms. - *Pest. Biochem. Physiol.* 122: 96-102.
- YOU-HUI G., BAO-YUN X., YOU-JUN Z., XI-WU G., QING-JUN W., 2015 - Demonstration of an adaptive response to preconditioning *Frankliniella occidentalis* (Pergande) to sublethal doses of spinosad: a hormetic-dose response. - *Ecotoxicology* 24: 1141-1151.
- ZHAO J.Z., LI Y.X., COLLINS H.L., GUSUKUMA-MINUTO L., MAU R.F.L., THOMPSON G.D., SHELTON A.M., 2002 - Monitoring and characterization of diamondback moth (Lepidoptera: Plutellidae) resistance to spinosad. - *J. Econ. Entomol.* 9: 430-436.

Non-commercial use only