

The subtribes and genera of the tribe Listroderini (Coleoptera, Curculionidae, Cyclominae): Phylogenetic analysis with systematic and biogeographical accounts

Juan J. Morrone¹

¹ Museo de Zoología “Alfonso L. Herrera”, Departamento de Biología Evolutiva, Facultad de Ciencias, Universidad Nacional Autónoma de México (UNAM), Apartado Postal 70-399, 04510 Mexico D.F., Mexico

Corresponding author: Juan J. Morrone (juanmorrone2001@yahoo.com.mx)

Academic editor: M. Alonso-Zarazaga | Received 12 November 2012 | Accepted 31 January 2013 | Published 28 February 2013

Citation: Morrone JJ (2013) The subtribes and genera of the tribe Listroderini (Coleoptera, Curculionidae, Cyclominae): Phylogenetic analysis with systematic and biogeographical accounts. ZooKeys 273: 15–71. doi: 10.3897/zookeys.273.4116

Abstract

The phylogenetic relationships of the genera of Listroderini LeConte, 1876 are analyzed based on 58 morphological characters. The genera are grouped in four clades, which are given subtribal status: Macrostyphlina new subtribe (*Adioristidius*, *Amathynetoides*, *Andesianellus*, *Macrostyphlus*, *Nacodius* and *Puranius*), Palaechthina Brinck, 1948 (*Anorthorhinus*, *Gunodes*, *Haversiella*, *Inaccodes*, *Listronotus*, *Neopachytychius*, *Palaechthus*, *Palaechtodes*, *Steriphus* and *Tristanodes*), Falklandiina new subtribe (*Falklandiellus*, *Falklandiopsis*, *Falklandius*, *Gromilus*, *Lanteriella*, *Liparogetus*, *Nestrius* and *Telurus*), and Listroderina (*Acroliellus*, *Acrorius*, *Acrostomus*, *Antarctobius*, *Germainiellus*, *Hyperoides*, *Lamiarhinus*, *Listroderes*, *Methypora*, *Philippius*, *Rupanius* and *Trachodema*). The subtribes are characterized and keys to identify them and their genera are provided. Listroderini have four main biogeographical patterns: Andean (Macrostyphlina), Andean-New Zealand (Falklandiina), Andean-Neotropical-Australian (Listroderina) and Andean-Neotropical-Australian-New Zealand-Nearctic-Tristan da Cunha-Gough islands (Palaechthina). Geographical paralogy, particularly evident in the Subantarctic subregion of the Andean region, suggests that Listroderini are an ancient Gondwanic group, in which several extinction events might have obscured relationships among the areas.

Keywords

Cyclominae, weevils, Americas, Australia, New Zealand, Tristan da Cunha-Gough islands

Introduction

Listroderini LeConte, 1876 are one of the largest tribes of Cyclominae (Oberprieler 2010, in press). They are widely distributed in the Southern Hemisphere, with the genus *Listronotus* also occurring in North America (Morrone 2011) and fossils known from Antarctica (Ashworth and Kuschel 2003). The tribe was originally proposed by LeConte (1876) for the New World genera *Listroderes*, *Listronotus* and *Macrops*. In the following decades additional new taxa were described from Chile (Germain 1895–1896; Kuschel 1949, 1950, 1952), Argentina (Enderlein 1907, 1912; Hustache 1926), North and Central America (e.g., Henderson 1940; O'Brien 1977, 1981) and Peru (Voss 1954). Kuschel (1950, 1952, 1955) transferred some listroderine species to genera of Entiminae. Additionally, the circumscription of the tribe was expanded, because several genera that have been originally assigned to other tribes (and even subfamilies) from Australia (Erichson 1842; Pascoe 1865, 1870; Blackburn 1890; Lea 1928), New Zealand (Broun 1893a, b, 1909, 1913, 1915) and the Tristan da Cunha-Gough islands (Brinck 1948) were transferred to Listroderini (Kuschel 1962, 1964, 1971, 1986; May 1994; Zimmerman 1994; Morrone 1997a). Recently, Oberprieler (2010) transferred *Rhigopsidius* from Rhythirrinini to Listroderini and reassigned the listroderine genus *Telurus* to the tribe Cylydrorhinini (Entiminae). According to the last checklist (Morrone 2011), a total of 407 species classified into 36 genera are assigned to Listroderini. Due to all these changes the taxa currently assigned to Listroderini constitute an assemblage that is difficult to characterize, and there is no complete treatment of all the genera.

Listroderini were originally assigned to the subfamily Cylydrorhininae (e.g., Enderlein 1907, 1912; Hustache 1926; Schenckling and Marshall 1931; Voss 1954; Kuschel 1955, 1958; O'Brien and Wibmer 1982). Kuschel (1964) transferred Listroderini to Rhyparosominae, which Kuschel (1971) later treated as a synonym of Rhythirrininae, and was followed by several authors (e.g., Wibmer and O'Brien 1986; Morrone et al. 1992; Morrone 1997a). Later, Rhythirrininae were demoted to a tribe of Cyclominae (Morrone 1997b), and thus Listroderini were considered as a subtribe (Anderson and Morrone 1996; Morrone 1997a, 2002a; Anderson 2002). More recently, Oberprieler (2010), while analysing the circumscription of Cyclominae and their tribes, reassigned tribal status to the listroderines.

Morrone (1997a) undertook a cladistic analysis of the South American genera of the tribe, considering that they represented a paraphyletic group, because the genera from Australia, New Zealand and the Tristan da Cunha-Gough islands are probably closely related to some of the American genera. The phylogenetic placement of these genera is not known, and the inclusion of *Rhigopsidius* and the exclusion of *Telurus* from the tribe, proposed by Oberprieler (2010), need to be tested.

My objective is to analyse the cladistic relationships of the genera of Listroderini, especially to determine the phylogenetic placement of the genera from Australia, New Zealand and the Tristan da Cunha-Gough islands. I intend to provide a phylogenetic framework for future studies and to summarize the systematics and biogeography of the genera to date.

Material and methods

The studied specimens were provided by the following collections:


AMNH	American Museum of Natural History, New York, USA.
AMPC	Amyan MacFadyen, private collection, Coleraine, Northern Ireland.
ARPC	Alexander Riedel, private collection, Friedberg, Germany.
BMNH	The Natural History Museum, London, England.
BPBM	Bernice P. Bishop Museum, Honolulu, USA.
CADIC	Centro Austral de Investigaciones Científicas, Ushuaia, Argentina.
CBPC	Carlos Bordón, private collection, Maracay, Venezuela.
CMNC	Canadian Museum of Nature, Ottawa, Canada.
CNCI	Canadian National Collection of Insects, Arachnids and Nematodes, Agriculture and Agri-Food Canada, Ottawa, Canada.
CWOB	Charles W. O'Brien private collection, Arizona, USA.
DEI	Deutsches Entomologisches Institut, Eberswalde-Finow, Germany.
DZUP	Departamento de Zoología, Universidade Federal do Paraná, Curitiba, Brazil.
FIML	Fundación e Instituto Miguel Lillo, San Miguel de Tucumán, Argentina.
FMNH	Field Museum of Natural History, Illinois, USA.
GJWC	Guillermo J. Wibmer, private collection, Tallahassee, USA.
IADIZA	Instituto Argentino de Investigaciones de las Zonas Áridas, Mendoza, Argentina.
ICNB	Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Santafé de Bogotá, Colombia.
IPUM	Instituto de la Patagonia, Universidad de Magallanes, Punta Arenas, Chile.
MACN	Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Buenos Aires, Argentina.
MCZ	Museum of Comparative Zoology, Harvard University, Massachusetts, USA.
MHNS	Museo Nacional de Historia Natural, Santiago, Chile.
MLP	Museo de La Plata, La Plata, Argentina.
MNHN	Museum National d'Histoire Naturelle, Paris, France.
MZFC	Museo de Zoología "Alfonso L. Herrera", Facultad de Ciencias, UNAM, Mexico City, Mexico.
NZAC	New Zealand Arthropod Collection, Auckland, New Zealand.
SMTD	Staatliches Museum für Tierkunde, Dresden, Germany.
USNM	National Museum of Natural History, Washington D.C., USA.
ZMC	Zoologisk Museum, Copenhagen, Denmark.
ZMHU	Zoologische Museum der Humboldt Universität, Berlin, Germany.

Habitus drawing were made with a camera lucida attached to a stereoscopic microscope. Photographs were taken using a Scanning Electron Microscope at the Facultad de Ciencias, UNAM.

For the present study I examined species of the genera previously recognized for the tribe (Morrone 2011). The outgroup taxa included the genera *Hyomora* (Hipporhinini), *Aphela* (Notiomimetini), *Rhythirrinus* (Rhythirrinini) and *Telurus* (Cylindrornithini). *Epichthonius* (Cyclomini) was used to root the cladograms.

The 58 morphological characters used in the analysis were taken from external structures (53) and male and female genitalia (5). The distribution of character states is shown in the data matrix (Table I). The characters and their corresponding character states are as follows:

- 1 Body: length. (0) large to very large (> 15.0 mm); (1) medium-sized (7.1–14.9 mm); (2) small to very small (< 7.0 mm) [additive].
- 2 Vestiture: scales. (0) present; (1) absent.
- 3 Vestiture: scale shape. (0) seta-like (Fig. 1); (1) subcircular (Fig. 2); (2) lanceolate (Fig. 3); (3) with finger-like processes (Fig. 4) [non-additive].
- 4 Vestiture: setae. (0) present; (1) absent.
- 5 Rostrum: shape. (0) stout, very short (Fig. 5); (1) relatively stout, medium-sized, shorter than pronotum (Fig. 6); (2) slender, as long as or longer than pronotum [additive].
- 6 Rostrum: dorsal carinae. (0) present (Fig. 6); (1) absent (Figs 5, 8).
- 7 Scrobes: shape. (0) long, deep, sharply bordered, reaching eyes; (1) short, ill-defined, broad.
- 8 Epistome. (0) poorly demarcated; (1) raised.
- 9 Scrobes: position. (0) dorsolateral to dorsal; (1) lateral.
- 10 Suprascrobal keels. (0) absent; (1) present.
- 11 Scrobes: ventral tooth. (0) absent; (1) present (Fig. 7).
- 12 Pterygia. (0) simple, not exposed (Fig. 6); (1) auriculate, exposed (Fig. 5).
- 13 Mandibles. (0) with one apical cusp; (1) with two apical cusps.
- 14 Mandible and pharyngeal processes. (0); short and strong; (1) long and narrow.
- 15 Mandibles. (0) plurisetose (more than 4 setae); (1) paucisetose (1–4 setae).
- 16 Maxillary malae: teeth. (0) present; (1) absent.
- 17 Eyes: shape. (0) subcircular (Fig. 5); (1) transverse (Fig. 7).
- 18 Eyes: size. (0) large to medium (more than 30 facets); (1) small (10–25 facets); (2) very small (8 or fewer facets) [additive].
- 19 Eyes: position. (0) lateral (Fig. 6); (1) dorsal (Fig. 5).
- 20 Eyes: convexity. (0) strong; (1) slight; (2) flat [additive].
- 21 Antennal insertions. (0) distal; (1) at the middle of the rostrum.
- 22 Scapes: length. (0) long (surpassing posterior margin of eyes when resting in scrobe); (1) medium-sized (reaching eyes when resting in scrobe); (2) short (not reaching anterior margin of eyes when resting in scrobe) [additive].
- 23 Funicles: segment 1. (0) elongate; (1) globose.
- 24 Funicles: segments 2. (0) elongate; (1) globose.
- 25 Funicles: relative lengths of segments 1 and 2. (0) 1 longer than 2 (Fig. 8); (1) 1 subequal to or shorter than 2.



Figures 1–12. Some of the characters analysed. **1** Seta-like scales **2** subcircular scales **3** lanceolate scales **4** scales with finger-like processes **5, 6, 8** face and rostrum, dorsal view **7** face and rostrum, lateral view **8, 9** elytra, dorsal view **11, 12** tarsomere 3, ventral view. **1, 5** *Falklandius antarcticus*; **2, 8, 11** *Falklandiellus suffodens*; **3** *Hyperoides subcinctus*; **4, 12** *Philippius superbus*; **6, 7, 10** *Listroderes costirostris*; **9** *Lamiarhinus aelficus*.

- 26 Funicles: segments 3–6. (0) elongate; (1) globose (Fig. 8).
- 27 Clubs: shape. (0) fusiform; (1) inflated.
- 28 Pronotum: shape. (0) subcircular; (1) transverse; (2) subtrapezoidal; (3) subquadrate; (4) subcylindrical [non-additive].
- 29 Pronotum: width. (0) larger than that of elytra; (1) smaller than that of elytra.
- 30 Pronotum: disc. (0) rugose; (1) smooth, polished.
- 31 Pronotum: tubercles. (0) absent; (1) present.
- 32 Postocular lobes. (0) present, well-developed; (1) present, slightly developed; (2) absent [additive].
- 33 Prosternum. (0) non-excavate; (1) excavate.
- 34 Metanepisternal sutures. (0) posteriorly fused or obliterated; (1) present, complete.
- 35 Scutellum. (0) not visible; (1) visible.
- 36 Elytra: shape. (0) oblong-oval (Fig. 10); (1) subrectangular (Fig. 9); (2) elongate-oval [non-additive].
- 37 Elytra. (0) not fused; (1) fused along interelytral suture.
- 38 Elytral disc. (0) convex; (1) slightly convex; (2) flat [additive].
- 39 Elytral intervals. (0) convex; (1) flat.
- 40 Elytral basal margin. (0) not raised; (1) raised, subcarinate.
- 41 Elytral humeri. (0) rounded; (1) subquadrate.
- 42 Elytral humeral tubercles. (0) absent; (1) present.
- 43 Several tubercles on elytral disc. (0) present, small, rounded; (1) absent; (2) present, strong (Fig. 9) [non-additive].
- 44 Series of three tubercles restricted to elytral interval 3. (0) absent; (1) present.
- 45 Series of declivital tubercles on elytra. (0) absent; (1) present.
- 46 Carina on elytral apical declivity. (0) absent; (1) present.
- 47 Anteapical elytral tubercle. (0) absent; (1) present.
- 48 Elytral apex, female. (0) not produced; (1) produced.
- 49 Femora: shape. (0) subcylindrical, clavate; (1) dorsoventrally compressed, clavate; (2) subcylindrical, markedly clavate [non-additive].
- 50 Tibiae: shape. (0) subcylindrical, laterally not expanded; (1) apically expanded.
- 51 Tibial spurs. (0) present; (1) absent.
- 52 Tarsomeres 3. (0) bilobed (Fig. 11); (1) subcylindrical (Fig. 12).
- 53 Ventrites 3 and 4, female. (0) combined shorter than 5; (1) combined longer than 5.
- 54 Aedeagus, lateral view. (0) robust; (1) slender.
- 55 Distal gonocoxites. (0) strongly sclerotized; (1) membranous.
- 56 Styli. (0) well-developed, claw-like; (1) well-developed, finger-like; (2) reduced to a few vibrissae [non-additive].
- 57 Apodeme of female sternum 8. (0) short (< 3 times longer than plate); (1) long (> 4 times longer than plate).
- 58 Plate of female sternum 8. (0) developed; (1) reduced.

Table 1. Data matrix analysed. Character states of polymorphic taxa are indicated between square brackets.

<i>Methypora</i>	2010100010001010100102000104000101110200000000110000001100
<i>Nacodius</i>	200010001100101010010000010001020110001000000000011100
<i>Neopachytichius</i>	201020001000111010010200010000010110001000100000000001100
<i>Nestrius</i>	2000101000011010010200001140002000000000100000000001100
<i>Palaeochthys</i>	1000121001000101010010200111200010112001000100000000001100
<i>Palaeochthodes</i>	100020001000101010010200111400010112001000100000000001100
<i>Philippius</i>	003010101000101011020100010010110101200101010100011001100
<i>Puranus</i>	201010001100101010010100011100010110001000000000000011110
<i>Rhigopsidius</i>	1030000110001010100201000101001011100010201010000000000
<i>Rupanius</i>	2000100010001010100101000101001011100010201010000000000
<i>Steriphus</i>	101020001000101010010000010000010112000010001000000001100
<i>Trachodema</i>	2030101010001010100100000101001101100100102010100000001100
<i>Tristanodes</i>	200020001000101010010200111400010112001000100000000001100

The cladograms were constructed using software TNT (Goloboff et al. 2008). A first analysis was conducted treating all characters under equal weights. Then, the effect of homoplasy on the results was explored by conducting different implied weights analyses (Goloboff 1993), with constants of concavity (k) set to a different integer value of 1–12, where 1 is weighted most severely against homoplastic characters. Implied weights analyses were conducted using the heuristic “traditional search” algorithm of TNT, with 1000 replications and tree-bisection-reconnection branch-swapping (TBR), holding 1000 trees during each replication.


Results

Phylogenetic Analysis

The analysis of the data matrix (Table I) under equal weights and with different concavity constants led to different cladograms: 100 cladograms under equal weights (Fig. 13); three cladograms with $k=3$ (Fig. 14); six cladograms with $k=6$ (Fig. 15); and two cladograms with $k=12$ (Fig. 16). In all the analyses the tribe Listroderini is recovered as a monophyletic taxon. *Rhigopsidius*, previously placed by Oberprieler (2010) in Listroderini, resulted to be the sister taxon to *Rhythirrinus* (Rhythirrinini). *Telurus*, excluded from Listroderini by Oberprieler (2010), was placed within Listroderini. In the analyses with $k=3$ and 6, *Aphela* (Notiomimetini) is the sister taxon to Listroderini. In spite of the different results, there are some larger clades that were fairly constant.

I consider that the results of the analysis with $k=6$ are not as extreme as the others and show more clearly the four main clades, which are treated herein as subtribes (Fig. 17):

- 1 Macrostyphlina new subtribe: genera *Adioristidius*, *Amathynetoides*, *Andesianellus*, *Macrostyphlus*, *Nacodius* and *Puranius*.
- 2 Palaechthina Brinck, 1948: genera *Anorthorhinus*, *Gunodes*, *Haversiella*, *Inaccodes*, *Listronotus*, *Neopachytychius*, *Palaechthus*, *Palaechtodes*, *Steriphus* and *Tristanodes*.
- 3 Falklandiina new subtribe: genera *Falklandiellus*, *Falklandiopsis*, *Falklandius*, *Gromilus*, *Lanteriella*, *Liparogetus*, *Nestrius* and *Telurus*.
- 4 Listroderina LeConte, 1876: genera *Acroriellus*, *Acrorius*, *Acrostomus*, *Antarctobiuss*, *Germainiellus*, *Hyperoides*, *Lamiarhinus*, *Listroderes*, *Methypora*, *Philippius*, *Rupanius* and *Trachodema*.



Figures 13–16. Consensus cladograms of the different analyses. **13** equal weights **14** $k=3$ **15** $k=6$ **16** $k=12$.




Figure 17. Consensus cladogram of the cladograms obtained with $k=6$ with character state changes indicated.

Systematic Account

Tribe Listroderini LeConte, 1876

Figs 18–35

Listroderi LeConte, 1876: 124.
Listroderitos Germain, 1895: 287.
Listroderina Champion, 1902: 120.
Listroderini Hustache, 1926: 175.
Listroderinae Thompson, 1992: 876.

Type genus. *Listroderes* Schönherr, 1826.

Diagnosis. Very small to very large (1.0–22.8 mm); integument reddish brown (black in *Acrostomus*); vestiture consisting mostly of dense scales and setae (rarely only scales or setae), setae on rostrum and pronotum directed anteriad or mesad, on elytra posteriad; rostrum stout and very short to slender, as long as or longer than pronotum; scrobes usually lateral; epistome poorly demarcated, rarely raised (*Acrostomus*); eyes usually large, flat, transverse or subcircular; mandibles with two apical cusps and paucisetose (1–4 setae); antennae with funicle 7-segmented, segments 1 and usually 2 elongate, clubs fusiform or inflated; prothorax with or without postocular lobes; prosternum long, non-excavate; elytra oblong-oval, elongate-oval or subrectangular; tibiae mucronate, generally with spurs (when present pro- and mesotibiae with 1 spur and metatibiae with 1–2 spurs); claws divaricate, simple or with slight basal swelling; aedeagus with tegmen lacking parameres (reduced in *Methypora*); distal gonocoxites membranous, generally simple, with large, apical or subapical stylus carrying a tuft of setae, but occasionally without stylus and apex of gonocoxite flattened and bent outwards.

Comparative notes. Listroderini were formerly considered as related to Rhytidinini (Kuschel 1971; Anderson and Morrone 1996; Morrone 1997a, b, 2002a; Anderson 2002). Oberprieler (2010, in press) considered Notiomimetini to be close relatives of Listroderini, although he suggested that more detailed studies would be required to decide whether they should be merged into a single tribe or not. Based on the results of this analysis, Listroderini and Notiomimetini (*Aphela*) are hypothesized to be sister tribes.

Biology. Larvae of Listroderini are generally oligophagous ectophytic root-feeders (Oberprieler in press). Adults feed on the leaves of a variety of angiosperms (Morrone 2011).

Key to the subtribes of Listroderini

- 1 Rostrum slender, as long as or longer than pronotum (except shorter than pronotum in some species of *Listronotus*); scrobes long, sharply bordered, reaching eyes; funicular segment 1 usually subequal to or shorter than 2; commonly associated with aquatic or semiaquatic plants **Palaechthrina**

- Rostrum stout or relatively stout, shorter than pronotum; scrobes usually short, ill-defined, broad; funicular segment 1 longer than 2; associated to terrestrial plants 2
- 2 Rostral dorsal carinae usually absent; pterygia auriculate, exposed (Fig. 5) **Falklandiina**
- Rostral dorsal carinae present; pterygia simple, not exposed (Fig. 6) 3
- 3 Scrobes short, ill-defined, broad, lacking suprascrobal keel; elytra with intervals convex, with antepical tubercle (except for *Rupanius*) **Listroderina**
- Scrobes long, deep, sharply bordered, reaching eyes, with suprascrobal keel; elytra with intervals usually flat, lacking antepical tubercle **Macrostyphlina**

Macrostyphlina, subtr. n.


Type genus. *Macrostyphlus* Kirsch, 1889.

Diagnosis. Scrobes long, deep, sharply bordered, reaching eyes, with suprascrobal keel; elytra oblong-oval, with intervals usually flat, lacking antepical tubercle.

Included taxa. This new subtribe, which basically corresponds to the *Macrostyphlus* generic group of Morrone (1994c, 1997a), includes the genera *Adioristidius*, *Amathynetoides*, *Andesianellus*, *Macrostyphlus*, *Nacodius* and *Puranius*. All these genera are distributed in South America, in the Andean region and the South American Transition Zone (*sensu* Morrone 2006).

Key to the genera of Macrostyphlina

- 1 Postocular lobes present 2
- Postocular lobes absent 4
- 2 Pronotum transverse to strongly transverse **Puranius** (Fig. 19)
- Pronotum subcircular or subcylindrical 3
- 3 Pronotum subcircular with subparallel flanks, disc smooth, polished; metanepisternal sutures present, complete; elytra with intervals flat... **Amathynetoides**
- Pronotum subcylindrical, disc rugose; metanepisternal sutures posteriorly fused or obliterated; elytra with intervals convex **Adioristidius** (Fig. 18)
- 4 Vestiture consisting of subcircular scales and setae; elytra with small, rounded tubercles **Macrostyphlus**
- Vestiture consisting of seta-like scales and setae or only setae; elytra lacking tubercles 5
- 5 Vestiture consisting of seta-like scales and setae; eyes large, slightly convex; pronotum disc smooth, polished; basal elytral margin not raised ... **Nacodius**
- Vestiture consisting of setae only; eyes very small, microphthalmic (8 or fewer facets), flat; pronotum disc rugose; basal elytral margin raised, subcarinate ... **Andesianellus**



Figures 18–26. Habitus of representative Listroderini. **18** *Adioristidius hirsutus* **19** *Puranius nigrinus* **20** *Haversiella albolimbata* **21** *Listronotus bosqi* **22** *Neopachytychius squamosus* **23** *Falklandiellus suffodens* **24** *Falklandiopsis magellanica* **25** *Falklandius antarcticus* **26** *Gromilus veneris*.

***Adioristidius* Morrone, 1994**

<http://species-id.net/wiki/Adioristidius>

Fig. 18

Adioristidius Voss, 1954: 242 (not available, type species not designated).

Anchadioristus Voss, 1954: 242 (not available, type species not designated).

Adioristidius Edwards & Hopwood, 1966: 5 (lapsus).

Adioristidius Morrone, 1994c: 13.

Type species. *Adioristus similaris* Voss, 1954.

Diagnosis. Small to very small (1.5–4.1 mm); vestiture consisting of seta-like scales and setae; antennal clubs fusiform; pronotum subcylindrical, disc rugose; metanepisternal sutures posteriorly fused or obliterated; elytral intervals convex.

Relationships. *Adioristidius* is the sister genus of *Macrostyphlus-Andesianellus*.

Species included. *Adioristidius anchonoideus* (Hustache, 1938); *A. carinicollis* (Voss, 1954); *A. chilensis* Morrone, 1994; *A. costulatus* (Hustache, 1938); *A. crassirostris* (Hustache, 1938); *A. cuprisquamatus* (Voss, 1954); *A. granulatus* (Hustache, 1938); *A. hirsutus* Morrone, 1994; *A. hydanius* Morrone, 1994; *A. jorgei* Morrone, 1994; *A. lidiae* Morrone, 1994; *A. manu* Morrone, 1994; *A. morio* (Voss, 1954); *A. nivalis* (Kuschel, 1949); *A. pampaensis* (Voss, 1954); *A. peruvianus* (Voss, 1954); *A. puncticollis* (Hustache, 1938); *A. scrobicollis* (Voss, 1954); *A. similaris* (Voss, 1954); *A. subimpresus* (Voss, 1954); *A. subtuberculatus* (Voss, 1954); *A. sulcicollis* (Hustache, 1938); *A. tuberculatus* (Voss, 1954); *A. variegatus* (Voss, 1954).

Host plants. *Adioristidius chilensis*: *Mulinum* spp. (Apiaceae); *A. tuberculatus*: *Solanum tuberosum* L. (Solanaceae) (Morrone 1994c).

Geographical distribution. South American Transition Zone (Puna biogeographical province) and Central Chilean and Subantarctic subregions (Andean region), from Peru to Central Chile (Morrone 1994c).

Material examined. *Adioristidius anchonoideus* (CMNC, DEI, MLP, MZFC), *A. chilensis* (MHNS), *A. costulatus* (DEI), *A. crassirostris* (DEI), *A. granulatus* (DEI), *A. hirsutus* (MHNS, MLP, MZFC), *A. hydanius* (DEI), *A. jorgei* (MHNS, MLP, MZFC), *A. lidiae* (CMNC), *A. manu* (CMNC, FMNH), *A. morio* (CWOB, MLP, MZFC), *A. nivalis* (MHNS, NZAC), *A. puncticollis* (DEI, MZFC), *A. similaris* (DEI), *A. sulcicollis* (DEI), *A. tuberculatus* (CWOB, MZFC, USNM), *A. variegatus* (DEI).

***Amathynetoides* Morrone, 1994**

<http://species-id.net/wiki/Amathynetoides>

Amathynetes Kuschel, 1949: 43 (non Olliff, 1891; misidentification, in part).

Amathynetoides Morrone, 1994c: 28.

Type species. *Amathynetes appendiculatus* Kuschel, 1949.

Diagnosis. Small to very small (3.0–6.6 mm); vestiture consisting of seta-like scales and setae; pronotum subcircular with subparallel flanks, disc smooth, polished; metanepisternal sutures present, complete; elytral intervals flat.

Relationships. *Amathynetoides* is the sister genus of *Nacodius*.

Species included. *Amathynetoides appendiculatus* (Kuschel, 1949); *A. ebeninus* (Hustache, 1938); *A. intemperatus* Morrone, 1994; *A. longulus* (Kuschel, 1949); *A. morbeamus* Morrone, 1994; *A. nitidiventris* (Hustache, 1938); *A. normae* Morrone, 1994; *A. palustris* (Kuschel, 1949); *A. sparsesetosus* (Hustache, 1938); *A. sundrianus* Morrone, 1994.

Host plants. *Amathynetoides nitidiventris*: *Ullucus tuberosus* Caldas (Basellaceae) (López and Hermann 2004).

Geographical distribution. South American Transition Zone (Puna and Coastal Peruvian Desert biogeographical provinces), from Peru to northern Chile (Morrone 1994c).

Material examined. *Amathynetoides appendiculatus* (CWOB, CMNC, MHNS, MZFC, NZAC, USNM), *A. ebeninus* (BPBM, CWOB, DEI, MZFC), *A. intemperatus* (AMNH, CWOB, MLP, MZFC), *A. longulus* (CWOB, MHNS, NZAC, MZFC, USNM), *A. morbeamus* (FIML), *A. nitidiventris* (DEI), *A. normae* (CMNC, MLP, MZFC), *A. palustris* (CWOB, FIML, MHNS, MZFC, NZAC, USNM), *A. sparsesetosus* (CWOB, DEI, CMNC, MZFC), *A. sundrianus* (BMNH, CWOB, FIML, MLP, MZFC).

Andesianellus Anderson & Morrone, 1996

<http://species-id.net/wiki/Andesianellus>

Andesianellus Anderson & Morrone, 1996: 260.

Type species. *Andesianellus microphthalmicus* Anderson & Morrone, 1996.

Diagnosis. Very small (1.9–3.3 mm); vestiture consisting of setae only; eyes very small, (8 or fewer facets), flat; postocular lobes absent; basal elytral margin raised, subcarinate.

Relationships. *Andesianellus* is the sister genus of *Macrostyphlus*, as hypothesized in previous analyses (Anderson and Morrone 1996; Morrone 1997a).

Biology. Species of this genus have been reported as leaf-litter inhabitants (Anderson and Morrone 1996).

Species included. *Andesianellus carltoni* Anderson & Morrone, 1996; *A. cotopaxi* Anderson & Morrone, 1996; *A. fulgidus* Anderson & Morrone, 1996; *A. hermani* Anderson & Morrone, 1996; *A. masneri* Anderson & Morrone, 1996; *A. microphthalmicus* Anderson & Morrone, 1996; *A. minutus* Anderson & Morrone, 1996; *A. planirostris* Anderson & Morrone, 1996; *A. tricarinatus* Anderson & Morrone, 1996.

Geographical distribution. South American Transition Zone (North Andean Paramo biogeographical province), in Colombia, Ecuador and Peru (Anderson and Morrone 1996).

Material examined. *Andesianellus carltoni* (CMNC), *A. cotopaxi* (AMNH), *A. fulgidus* (CMNC), *A. hermani* (AMNH), *A. masneri* (CMNC), *A. microphthalamicus* (CMNC, MLP), *A. minutus* (CMNC, FMNH), *A. planirostris* (AMNH, BMNH, CMNC, CWOB, FMNH, MLP, USNM), *A. tricarinatus* (CMNC, FMNH).

***Macrostyphlus* Kirsch, 1889**

<http://species-id.net/wiki/Macrostyphlus>

Macrostyphlus Kirsch, 1889: 25.

Type species. *Macrostyphlus gualcalae* Kirsch, 1889 (by indication, monotypy).

Diagnosis. Very small (1.9–3.5 mm); vestiture consisting of subcircular scales and setae; pronotum subcylindrical; metanepisternal sutures posteriorly fused or obliterated; elytra with intervals convex.

Relationships. *Macrostyphlus* is the sister genus of *Andesianellus*, as hypothesized in a previous analysis (Morrone 1997a).

Species included. *Macrostyphlus bilbo* Morrone, 1994; *M. coelorum* (Olliff, 1891); *M. frodo* Morrone, 1994; *M. gandalf* Morrone, 1994; *M. gualcalae* Kirsch, 1889; *M. howdenorum* Morrone, 1994; *M. peruvianus* Morrone, 1994; *M. sturmi* Morrone, 1994; *M. transatlanticus* (Kirsch, 1889); *M. venezolanus* Morrone, 1994.

Geographical distribution. South American Transition Zone (North Andean Paramo and Puna biogeographical provinces), from eastern Venezuela to southern Peru (Morrone 1994c).

Material examined. *Macrostyphlus bilbo* (CNCI), *M. coelorum* (CWOB), *M. frodo* (ICNB, USNM), *M. gandalf* (CMNC, CNCI, MLP, MZFC), *M. gualcalae* (SMTD), *M. howdenorum* (CMNC), *M. peruvianus* (FMNH), *M. sturmi* (ICNB), *M. transatlanticus* (SMTD), *M. venezolanus* (MZFC).

***Nacodius* Morrone, 1994**

<http://species-id.net/wiki/Nacodius>

Nacodius Morrone, 1994e: 3.

Type species. *Nacodius martitiae* Morrone, 1994.

Diagnosis. Small (4.6–6.9 mm); vestiture of seta-like scales and setae; eyes large, slightly convex; pronotum lacking postocular lobes, with disc smooth, polished; elytra with intervals flat.

Relationships. *Nacodius* is the sister genus to *Amathynetoides*, and both are placed in Macrostyphlini. In a previous analysis (Morrone 1997a) *Nacodius* was placed in the *Antarctobius* generic group (= Listroderina).

Species included. *Nacodius alecrus* Morrone, 1994; *N. brevirostris* (Voss, 1954); *N. martitiae* Morrone, 1994; *N. omissus* (Kuschel, 1952).

Geographical distribution. South American Transition Zone (North Andean Paramo and Puna biogeographical provinces), in Ecuador and Peru (Morrone 1994e).

Material examined. *Nacodius alecrus* (CWOB), *N. brevirostris* (SMTD), *N. martitiae* (AMNH, CWOB, MLP, MZFC) and *N. omissus* (BMNH).

Puranus Germain, 1895

<http://species-id.net/wiki/Puranus>

Fig. 19

Puranus Germain, 1895: 313.

Puranus Germain, 1911: 205 (lapsus).

Reichertia Enderlein, 1912: 31 (type species: *Listroderes sculpticollis* Enderlein, 1907, by original designation).

Type species. *Puranus inaequalis* Germain, 1896 (subsequent designation by Morrone, 1994c).

Relationships. *Puranus* is the sister genus to *Amathynetoides-Nacodius*.

Diagnosis. Small to very small (1.9–6.5 mm); vestiture of subcircular scales and setae; pronotum transverse to strongly transverse; metanepisternal suture present, complete; elytra oblong-oval, with small, rounded tubercles.

Species included. *Puranus argentinensis* Morrone, 1994; *P. australis* Germain, 1896; *P. championi* (Kuschel, 1952); *P. dubius* (Germain, 1896); *P. elguetai* Morrone, 1994; *P. exsculpticollis* (Enderlein, 1907); *P. fasciculiger* (Blanchard, 1851); *P. hispidus* (Germain, 1896); *P. inaequalis* Germain, 1896; *P. midas* Morrone, 1994; *P. nigrinus* (Fairmaire, 1884); *P. obrienorum* Morrone, 1994; *P. pusillus* Morrone, 1994; *P. scaber* (Enderlein, 1907); *P. sylvanicus* Morrone, 1994; *P. torosus* Morrone, 1994; *P. tothus* Morrone, 1994; *P. tuberosus* Germain, 1896; *P. verrucosus* (Germain, 1896); *P. vulgaris* Morrone, 1994.

Host plants. *Puranus argentinensis*: *Mulinum* sp. (Apiaceae); *P. championi*: *Poa flabellata* (Lam.) Raspail (Poaceae); *P. fasciculiger*: *Senecio smithii* DC (Asteraceae); *P. nigrinus*: *Taraxacum officinale* Weber ex F. H. Wigg. (Asteraceae) and *Nothofagus* sp. (Nothofagaceae); *P. vulgaris*: *Mulinum* sp. (Apiaceae); *P. scaber*: *Baccharis* sp. (Asteraceae) and *Ephedra* sp. (Ephedraceae) (Morrone, 1994c).

Geographical distribution. Andean region (Subantarctic and Central Chilean subregions) and South American Transition Zone, from southern Argentina, including the Falkland Islands (Islas Malvinas), to Peru (Morrone 1994c; Posadas 2008, 2012).

Material examined. *Puranus argentinensis* (AMNH, BMNH, MLP, MZFC), *P. australis* (AMNH, CWOB, MHNS, NZAC), *P. championi* (BMNH, CWOB, NZAC), *P. dubius* (CWOB, MHNS, NZAC), *P. elguetai* (AMNH, MHNS, MLP, MZFC), *P. exsculpticollis* (BMNH), *P. fasciculiger* (CWOB, MHNS, NZAC, USNM), *P. his-*

pidus (CWOB, MHNS, NZAC), *P. inaequalis* (CMNC, CWOB, MHNS, MZFC, NZAC), *P. midas* (AMNH), *P. nigrinus* (ARPC, BMNH, CADIC, CBPC, CMNC, CNCI, CWOB, DEI, FIML, IPUM, MCZ, MHNS, MZFC, NZAC, USNM), *P. obrienorum* (AMNH, CMNC, CWOB, MLP, MZFC), *P. pusillus* (MHNS, MLP, MZFC), *P. scaber* (AMPC, BMNH, CWOB, NZAC), *P. sylvanius* (AMNH, BMNH, CMNC, MLP, MZFC), *P. torosus* (MHNS, MLP, MZFC), *P. tothus* (MHNS), *P. tuberosus* (CWOB, MHNS, NZAC), *P. verrucosus* (CMNC, CWOB, MHNS, MZFC, NZAC) and *P. vulgaris* (AMNH, BMNH, CMNC, MHNS, MLP, MZFC).

Subtribe Palaechthina Brinck, 1948, stat. n.

Palaechtini Brinck, 1948: 43; Bouchard et al. 2011: 603 (incorrect original stem formation, not in prevailing usage).

Type genus. *Palaechthus* C. O. Waterhouse, 1884 (by original designation, as *Palaechtus*, incorrect subsequent spelling).

Diagnosis. Rostrum slender, as long as or longer than pronotum (except for some species of *Listronotus* where the rostrum is shorter than pronotum); scrobes long, deep, sharply bordered, reaching eyes; scape usually short (not reaching anterior margin of eye when resting in scrobe); pronotum usually subcylindrical or subcircular; elytra oblong-oval to elongate-oval.

Biology. Most of the species of Palaechthina are associated to aquatic or semi-aquatic plants, being found in wet or damp conditions (May 1970; O'Brien 1977, 1981; Marvaldi 1994; Morrone and O'Brien 2000). In contrast with the remaining Listroderini, larvae usually lead a more endophytic way of life inside the stems of several aquatic plants (Oberprieler, in press).

Included taxa. This subtribe includes the genera *Anorthorhinus*, *Gunodes*, *Haversiella*, *Inaccodes*, *Listronotus*, *Neopachytychius*, *Palaechthus*, *Palaechtodes*, *Steriphus* and *Tristanodes*. *Anorthorhinus* and *Steriphus* are Australian; *Gunodes*, *Inaccodes*, *Palaechthus*, *Palaechtodes* and *Tristanodes* are distributed in the Tristan da Cunha-Gough islands; and the remaining three genera are found in the Americas: *Haversiella* and *Neopachytychius* in South America and *Listronotus* has a disjunct distribution in South and North America.

Key to the genera of Palaechthina

1	Funicular segment 1 subequal to or shorter than 2.....	2
-	Funicular segment 1 longer than 2.....	6
2	Elytra with intervals convex; North and South America ...	<i>Listronotus</i> (Fig. 21)
-	Elytra with intervals flat; Tristan da Cunha-Gough islands.....	3
3	Small to very small (3.7–6.5 mm).....	<i>Tristanodes</i>
-	Medium-sized to large (7.0–12.0 mm)	4

- 4 Vestiture of subcircular scales and setae; pronotum subcircular; elytra oblong-oval..... ***Gunodes***
- Vestiture of seta-like scales and setae; pronotum subtrapezoidal or subcylindrical; elytra elongate-oval..... 5
- 5 Large (11.0–12.0 mm); rostral dorsal carinae absent; pronotum subtrapezoidal ***Palaechthus***
- Medium-sized (7.0–7.5 mm); rostral dorsal carinae present; pronotum subcylindrical ***Palaechtodes***
- 6 Scape long (surpassing posterior margin of eye when resting in scrobe); elytra with anteapical tubercle ***Steriphus***
- Scape short (not reaching anterior margin of eye when resting in scrobe); elytra lacking anteapical tubercle..... 7
- 7 Vestiture of seta-like scales and setae; Australia and Tristan da Cunha-Gough islands..... 8
- Vestiture of subcircular scales ans setae; South America 9
- 8 Funicular segments 3-6 elongate; club fusiform; pronotum subcylindrical; elytra with intervals convex; Australia ***Anorthorhinus***
- Funicular segments 3-6 globose; club inflated; pronotum subcircular; elytra with intervals flat; Tristan da Cunha-Gough islands ***Inaccodes***
- 9 Vestiture of subcircular scales and setae; rostral dorsal carinae present; mandibles long and narrow; antennal insertion distal; postocular lobes slightly developed; elytra oblong-oval; tibiae with spurs .. ***Neopachytychius*** (Fig. 22)
- Vestiture of subcircular scales only; rostral dorsal carinae absent; mandibles robust; antennal insertion at the middle of the rostrum; postocular lobes absent; elytra elongate-oval; tibiae lacking spurs ***Haversiella*** (Fig. 20)

***Anorthorhinus* Blackburn, 1890**

<http://species-id.net/wiki/Anorthorhinus>

Anorthorhinus Blackburn, 1890: 327.

Anorthorrhinus Sharp, 1892: 148 (lapsus).

Type species. *Anorthorhinus pictipes* Blackburn, 1890 (by indication, monotypy).

Diagnosis. Small to very small (2.5–6.0 mm); vestiture of seta-like scales and setae; funicular segments 3-6 elongate; club fusiform; pronotum subcylindrical; elytra with intervals convex.

Relationships. *Anorthorhinus* is the sister genus to the clade comprising *Haversiella*, *Neopachytychius* and the five genera from the Tristan da Cunha-Gough islands.

Species included. *Anorthorhinus apicalis* Lea, 1899; *A. brevicornis* Lea, 1899; *A. pictipes* Blackburn, 1890.

Geographical distribution. Australia (Oberprieler 2010).

Material examined. *Anorthorhinus apicalis* (MZFC) and *A. pictipes* (MZFC).

***Gunodes* Brinck, 1948**

<http://species-id.net/wiki/Gunodes>

Gunodes Brinck, 1948: 55.

Type species. *Gunodes major* Brinck, 1948.

Diagnosis. Medium-sized (7.5 mm); vestiture of subcircular scales and setae; pronotum subcircular; elytra oblong-oval.

Relationships. *Gunodes* is the sister genus to *Palaechthus-Paleachthodes-Tristanodes*. Oberprieler (1992) considered that the distinction between *Gunodes* and *Tristanodes* is not without doubt.

Species included. *Gunodes major* Brinck, 1948.

Geographical distribution. Tristan da Cunha-Gough islands (Brinck 1948).

***Haversiella* Schweiger, 1959**

<http://species-id.net/wiki/Haversiella>

Fig. 20

Haversia Champion, 1918a: 185 (*non* Röwer, 1913).

Haversiella Schweiger, 1959: 42 (replacement name for *Haversia*).

Type species. *Haversiella albolimbata* Champion, 1918 (by original designation).

Relationships. *Haversiella* is the sister genus to *Neopachytychius*, and both constitute the sister group to the five genera from the Tristan da Cunha-Gough islands.

Diagnosis. Very small (3.0–3.9 mm); vestiture of subcircular scales only; maxillary mala lacking teeth; antennal insertion at the middle of the rostrum; pronotum subcircular; elytra elongate-oval; tibiae lacking spurs; plate of female sternum 8 reduced.

Species included. *Haversiella albolimbata* (Champion, 1918).

Host plants. Bryophytes (Morrone 1994d).

Geographical distribution. Southern Argentina, including the Falkland Islands (Islas Malvinas), and southern Chile (Morrone 1994d; Posadas 2008, 2012).

Material examined. *Haversiella albolimbata* (BMNH, MHNS, MZFC, USNM).

***Inaccodes* Brinck, 1948**

<http://species-id.net/wiki/Inaccodes>

Inaccodes Brinck, 1948: 52.

Type species. *Inaccodes oblongus* Brinck, 1948.

Diagnosis. Small (4.5 mm); vestiture of seta-like scales and setae; funicular segments 3–6 globose; club inflated; pronotum subcircular; elytra with intervals flat.

Relationships. *Inaccodes* is the sister genus to the clade comprising the four remaining genera from the Tristan da Cunha-Gough islands. Oberprieler (1992) considered that the distinction between *Inaccodes* and *Tristanodes* is not without doubt.

Species included. *Inaccodes oblongus* Brinck, 1948.

Geographical distribution. Tristan da Cunha-Gough islands (Brinck 1948).

Listronotus Jekel, 1865

<http://species-id.net/wiki/Listronotus>

Fig. 21

Macrops Kirby, 1837: 199 (non Wagler 1830, nec Burmeister 1835) (type species: not designated).

Hyperodes Jekel, 1865: 566 (type species: *Listroderes humilis* Gyllenhal, 1834, by original designation).

Listronotus Jekel, 1865: 566.

Anchodemus LeConte, 1876: 181 (type species: *A. hubbardi* LeConte, 1876, subsequent designation by Kuschel 1950: 14).

Lixellus LeConte, 1876: 182 (type species: *L. filiformis* LeConte, 1876, by indication, monotypy).

Mascarauxia Desbrochers des Loges, 1898: 52 (type species: *M. cyrtica* Desbrochers des Loges 1898, by indication, monotypy).

Relistrodes Brèthes, 1910: 209 (type species: *R. breyeri* Brèthes, 1910, by indication, monotypy).

Aulametopiellus Brèthes, 1926: 415 (type species: *A. dauci* Brèthes, 1926, by indication, monotypy).

Mascaranxia Bosq, 1935: 330 (lapsus).

Pseudhyperodes Hustache, 1939a: 49 (type species: *P. elongatus* Hustache, 1939).

Type species. *Rhynchaenus caudatus* Say, 1824 (subsequent designation by Henderson 1940).

Diagnosis. Very small to medium-sized (1.0–14.0 mm); vestiture of subcircular scales and setae; antennal insertion distal; funicular segment 1 subequal to or shorter than 2; postocular lobes present, well-developed; elytra oblong-oval to elongate-oval, with intervals convex.

Relationships. *Listronotus* is the sister genus to *Steriphus* (Australia). In a previous analysis based only on American taxa (Morrone 1997a), *Listronotus* was considered to be the sister genus to *Neopachytychius*.

Species included. *Listronotus alternatus* (Dietz, 1889); *L. americanus* LeConte, 1876; *L. angustatus* (Champion, 1902); *L. annulipes* (Blatchley, 1925); *L. anthracinus* (Dietz, 1889); *L. apicalis* (Hustache, 1926); *L. appendiculatus* (Boheman, 1842); *L. argentinensis* (Hustache, 1926); *L. arizonicus* O'Brien, 1981; *L. blandus* Henderson, 1940; *L. blatchleyi* Henderson, 1940; *L. bonariensis* (Kuschel, 1955); *L. borrichiae* O'Brien,

1981; *L. bosqi* (Hustache, 1926); *L. breyeri* (Brèthes, 1910); *L. burkei* O'Brien, 1981; *L. californicus* (Dietz, 1889); *L. callosus* LeConte, 1876; *L. carinatus* (Blatchley, 1928); *L. carinicollis* (Hustache, 1926); *L. caudatus* (Say, 1824); *L. cinnamoneus* (Hustache, 1926); *L. conabilis* O'Brien, 1981; *L. crypticus* O'Brien, 1981; *L. cryptops* (Dietz, 1889); *L. cyrticus* (Desbrochers des Loges, 1898); *L. dauci* (Brèthes, 1926); *L. debilis* Blatchley, 1916; *L. deceptus* (Blatchley, 1916); *L. delumbis* (Gyllenhal, 1834); *L. dietrichi* (Stockton, 1963); *L. dietzi* O'Brien, 1979; *L. distinctus* Henderson, 1940; *L. dorsalis* (Dietz, 1889); *L. dorytomoides* (Hustache, 1926); *L. durangoensis* O'Brien, 1977; *L. echinatus* (Dietz, 1889); *L. echinodori* O'Brien, 1977; *L. elegans* Van Dyke, 1929; *L. elegantulus* O'Brien, 1981; *L. elongatus* (Hustache, 1939); *L. fasciatus* O'Brien, 1981; *L. filiformis* (LeConte, 1876); *L. frontalis* LeConte, 1876; *L. geminatus* (Hustache, 1926); *L. griseus* (Hustache, 1926); *L. grypidioides* (Dietz, 1889); *L. haldemani* (Burke, 1963); *L. hirtellus* (Dietz, 1889); *L. hoodi* (Stockton, 1963); *L. hornii* (Dietz, 1889); *L. hubbardi* (LeConte, 1876); *L. humilis* (Gyllenhal, 1834); *L. hyperodes* (Dietz, 1889); *L. incompletus* (Hatch, 1971); *L. ingens* Henderson, 1940; *L. insignis* Henderson, 1940; *L. laevis* (Hustache, 1926); *L. laramiensis* (Angell, 1893); *L. latinasus* (Blatchley, 1922); *L. lineolaticollis* (Blanchard, 1851); *L. lodingi* (Blatchley, 1920); *L. lucens* (Hustache, 1926); *L. lutulentus* (Bohemian, 1843); *L. maculatus* (Hatch, 1971); *L. maculicollis* (Kirby, 1837); *L. manifestus* Henderson, 1940; *L. marginalis* O'Brien, 1977; *L. marginicollis* (Hustache, 1926); *L. marshalli* O'Brien, 1981; *L. meridionalis* O'Brien, 1977; *L. minutus* (Blanchard, 1851); *L. montanus* (Dietz, 1889); *L. nebulosus* LeConte, 1876; *L. neocallosus* O'Brien, 1981; *L. nevadicus* LeConte, 1876; *L. nigropunctatus* (Suffrian, 1871); *L. novellus* (Blatchley, 1916); *L. obscurellus* (Dietz, 1889); *L. obtectus* (Dietz, 1889); *L. oregonensis* (LeConte, 1876); *L. ornatipennis* (Blanchard, 1851); *L. pallidus* O'Brien, 1981; *L. palustris* Blatchley, 1916; *L. pampaensis* (Voss, 1954); *L. peninsularis* (Blatchley, 1916); *L. plumosiventris* O'Brien, 1977; *L. porcellus* (Say, 1831); *L. poseyensis* (Blatchley, 1916); *L. pseudosetosus* O'Brien, 1981; *L. puncticollis* (Hustache, 1926); *L. punctiger* LeConte, 1876; *L. pusillus* (Hustache, 1926); *L. rotundicollis* LeConte, 1876; *L. rubtzoffi* O'Brien, 1981; *L. rufomarginatus* (Hustache, 1939); *L. salicorniae* O'Brien, 1981; *L. scapularis* Casey, 1895; *L. setosipennis* (Hustache, 1926); *L. setosus* LeConte, 1876; *L. similis* Henderson, 1940; *L. sondondoanus* (Voss, 1954); *L. sordidus* (Gyllenhal, 1834); *L. sparsus* (Say, 1831); *L. squamiger* (Say, 1831); *L. sulcipennis* (Bohemian, 1834); *L. suturalis* O'Brien, 1981; *L. teretirostris* (LeConte, 1857); *L. testaceipes* (Champion, 1902); *L. texanus* (Stockton, 1963); *L. truncatus* (Hatch, 1971); *L. tuberosus* LeConte, 1876; *L. turbatus* O'Brien, 1981; *L. vitticollis* (Kirby, 1837); *L. vulgaris* (Hustache, 1926); *L. wallacei* (Stockton, 1963); *L. weiseri* (Hustache, 1926).

Host plants. *Listronotus appendiculatus*: *Sagittaria latifolia* Willdenow (Alismataceae); *L. argentinensis*: *Triticum aestivum* L. (Poaceae); *L. blandus*: *Polygonum hydropiperoides* Michx. (Polygonaceae); *L. bonariensis*: *Dactylis glomerata* L., *Festuca arundinacea* Schreber, *Hordeum vulgare* L., *Lolium multiflorum* L., *L. perenne* L., *Poa annua* L., *Triticum aestivum* L., *Zea mays* L. (Poaceae) and *Trifolium repens* L. (Fabaceae); *L. borrichiae*: *Borrichia frutescens* (L.) DC (Asteraceae) and *Salvinia* sp. (Salviniaceae); *L. caudatus*: *Polygonum bincorne* Raf. (Polygonaceae); *L. cinnamoneus*: *Limnobium stoloniferum* (G.

F. W. Meyer) Griseb. (Hydrocharitaceae); *L. cryptops*: *Sagittaria lancifolia* L. (Alismataceae); *L. dauci*: *Daucus carota* L. (Apiaceae); *L. dietrichii*: *Dahlia* sp. (Asteraceae), *Gossypium* sp. (Malvaceae), *Persus* sp. (Lauraceae), *Phaseolus* sp. (Fabaceae), *Cenchrus* sp., *Chloris* sp., *Cynodon* sp., *Eleusine* sp., *Zea* sp. (Poaceae), *Coffea* sp. (Rubiaceae), *Lycopersicum* sp. (Solanaceae) and *Menta* sp. (Lamiaceae); *L. echinodori*: *Echinodorus cordifolius* (L.) Griseb. and *Sagittaria latifolia* Willdenow (Alismataceae); *L. elongatus*: *Hydrocotyle ranunculoides* L. f. (Apiaceae); *L. haldemani*: *Juncus nodatus* Coville in N. L. Britton and A. Brown (Juncaceae); *L. maculicollis*: *Agrostis palustris* Huds. and *Poa annua* L. (Poaceae); *L. manifestus*: *Sagittaria longiloba* Engelm. ex J. G. Sm. (Alismataceae); *L. marginicollis*: *Myriophyllum aquaticum* (Velloso) Verde (Haloragaceae); *L. montanus*: *Triticum aestivum* L. (Poaceae); *L. neocallosus*: *Sagittaria engelmanniana* J. G. Smith, S. graminea Michaux and S. stagnorum Small (Alismataceae); *Daucus carota* L. and *Petroselinum crispum* (Miller) A. W. Hill. (Apiaceae) (*L. oregonensis*); *L. plumosiventris*: *Sagittaria latifolia* Willdenow (Alismataceae); *L. rotundicollis*: *Crinum* sp. (Amaryllidaceae); *L. rubtzoffii*: *Sagittaria cuneata* Sheldon (Alismataceae); *L. salicorniae*: *Salicornia virginica* L. (Amaranthaceae); *L. scapularis*: *Sagittaria longiloba* Engelm. ex J. G. Sm. and *Sagittaria* sp. (Alismataceae); *L. setosipennis*: *Parthenium hysterophorus* L. (Asteraceae); *L. similis*: *Paspalum distichum* L. (Poaceae) and *Polygonum bincorne* Raf. (Polygonaceae); *L. teretirostris*: *Eleocharis macrostachya* Britton (Cyperaceae); *L. texanus*: *Daucus carota* L. (Apiaceae); *L. turbatus*: *Sagittaria* sp. (Alismataceae) (Brèthes 1926; Burke 1963; Martel et al. 1976; May 1977; O'Brien 1977, 1981; Cordo and DeLoach 1982; Cordo et al. 1982; Edelson 1985; Barker 1989; Boivin et al. 1990; Kuschel 1990; Maes and O'Brien 1990; Anderson 1992; Wild et al. 1992; May 1993; Cagnolini 1994; Blodgett et al. 1997; Lanteri et al. 2002; Torres and Casey 2002; Rothwell 2003).

Immature stages. *Listronotus bonariensis* (May, 1977, 1993, 1994; Marvaldi, 1998).

Geographical distribution. Widespread in the Americas, from Canada to Argentina and Chile (O'Brien 1977, 1981; O'Brien and Wibmer 1982; Wibmer and O'Brien 1986; Maes and O'Brien 1990; Anderson 1992). This distribution corresponds to the Nearctic, Neotropical and Andean regions, as well as the South American and Mexican Transition Zones.

Material examined. *Listronotus americanus* (BMNH), *L. apicalis* (MLP), *L. appendiculatus* (AMNH, BMNH), *L. argentinensis* (AMNH, MACN, MLP9), *L. blan-*
dus (AMNH, BMNH), *L. bonariensis* (BMNH, MHNS), *L. bosqi* (BMNH, MLP, MZFC), *L. breyeri* (MACN, MZFC), *L. californicus* (AMNH), *L. callosus* (BMNH, AMNH), *L. caudatus* (BMNH, AMNH), *L. cinnamomeus* (MLP), *L. cryptops* (BMNH, AMNH), *L. cyrticus* (AMNH, MACN, MLP), *L. dauci* (MACN), *L. debilis* (AMNH), *L. delumbis* (BMNH, AMNH), *L. dietzi* (AMNH), *L. distinctus* (BMNH), *L. durangoensis* (AMNH, BMNH), *L. echinatus* (AMNH), *L. echinodori* (AMNH, BMNH), *L. elongatus* (MLP, MZFC), *L. filiformis* (BMNH, AMNH), *L. frontalis* (AMNH, BMNH), *L. geminatus* (MACN, MLP), *L. griseus* (AMNH, MACN, MLP), *L. grypidioides* (AMNH), *L. haldemani* (BMNH), *L. hornii* (AMNH), *L. hubbardi* (BMNH), *L. humilis* (AMNH), *L. hyperodes* (AMNH), *L. incompletus* (AMNH), *L. ingens* (AMNH), *L. lineolaticollis* (MLP), *L. lutulentus* (BMNH), *L. maculicollis* (AMNH), *L. manifestus*

(AMNH, BMNH), *L. marginalis* (BMNH), *L. marginicollis* (MACN, MLP), *L. meridionalis* (BMNH), *L. minutus* (AMNH), *L. nebulosus* (AMNH), *L. novellus* (AMNH), *L. oregonensis* (AMNH, BMNH, MZFC), *L. ornatipennis* (MHNS), *L. palustris* (AMNH, BMNH), *L. plumosiventris* (BMNH), *L. porcellus* (AMNH), *L. puncticollis* (MLP), *L. punctiger* (AMNH, BMNH), *L. pusillus* (MLP, MZFC), *L. rotundicollis* (AMNH, BMNH), *L. rubtzoffi* (AMNH), *L. rufomarginatus* (MLP), *L. scapularis* (AMNH), *L. setosipennis* (MLP), *L. setosus* (AMNH), *L. similis* (AMNH, BMNH), *L. sordidus* (AMNH, BMNH), *L. sparsus* (AMNH, BMNH), *L. squamiger* (AMNH, BMNH), *L. teretirostris* (AMNH, BMNH), *L. texanus* (AMNH), *L. truncatus* (AMNH), *L. tuberosus* (AMNH, BMNH), *L. vitticollis* (AMNH) and *L. vulgaris* (MLP).

***Neopachytychius* Hustache, 1939**

<http://species-id.net/wiki/Neopachytychius>

Fig. 22

Neopachytychius Hustache, 1939b: 55.

Pernotaris Voss, 1943: 232 (type species: *P. squamiger* Voss, 1943 [= *Neopachytychius squamosus* Hustache, 1939]).

Type species. *Neopachytychius squamosus* Hustache, 1939.

Diagnosis. Small (3.8–6.5 mm); vestiture of subcircular scales and setae; mandible and pharyngeal process long and narrow; rostral dorsal carinae present; antennal insertion distal; postocular lobes slightly developed; elytra oblong-oval.

Relationships. *Neopachytychius* is the sister genus to *Haversiella*, and both constitute the sister group to the five genera from the Tristan da Cunha-Gough islands. In a previous analysis based only on American genera (Morrone 1997a), *Neopachytychius* was considered to be the sister genus to *Listronotus*.

Species included. *Neopachytychius squamosus* Hustache, 1939.

Geographical distribution. Neotropical region, in Argentina, Bolivia, Chile and Uruguay (Marvaldi 1994).

Material examined. *Neopachytychius squamosus* (FIML, IADIZA, MACN, MHNS, MLP, MZFC).

***Palaechthus* C. O. Waterhouse, 1884**

<http://species-id.net/wiki/Palaechthus>

Palaechthus C. O. Waterhouse, 1884: 277.

Palaechthus Brinck, 1948: 47 (lapsus).

Type species. *Palaechthus glabratus* Waterhouse, 1884 (subsequent designation by Brinck 1948).

Diagnosis. Medium-sized (11.0–12.0 mm); vestiture of seta-like scales and setae; rostral dorsal carinae absent; pronotum subtrapezoidal.

Relationships. *Palaechthus* is the sister genus to both *Paleachthodes* and *Tristanodes*. Oberprieler (1992) considered that the distinction between *Palaechthus* and *Palaechthodes* needs to be reevaluated.

Species included. *Palaechthus glabratus* C. O. Waterhouse, 1884.

Geographical distribution. Tristan da Cunha-Gough islands (Brinck 1948).

Material examined. *Palaechthus glabratus* (BMNH).

***Palaechthodes* Brinck, 1948**

<http://species-id.net/wiki/Palaechthodes>

Palaechthodes Brinck, 1948: 50.

Type species. *Palaechthus cossoides* C. O. Waterhouse, 1884 (by original designation).

Diagnosis. Medium-sized (7.0–7.5 mm); vestiture of seta-like scales and setae; rostral dorsal carinae present; pronotum subcylindrical.

Relationships. *Palaechthodes* is the sister genus to both *Paleachthodes* and *Tristanodes*.

Oberprieler (1992) considered that the distinction between *Palaechthodes* and *Palaechthus* needs to be reevaluated.

Species included. *Palaechthodes cossoides* (C. O. Waterhouse, 1884).

Geographical distribution. Tristan da Cunha-Gough islands (Brinck 1948).

Material examined. *Palaechthodes cossoides* (BMNH).

***Steriphus* Erichson, 1842**

<http://species-id.net/wiki/Steriphus>

Steriphus Erichson, 1842: 190.

Desiantha Pascoe, 1870: 193 (type species: *D. caudata* Pascoe, 1870, subsequent designation by Zimmerman 1994: 697).

Brexius Pascoe, 1870: 201 (type species: *B. murinus* Pascoe, 1870, subsequent designation by Zimmerman 1994: 697).

Dryopais Broun, 1885: 387 (type species: *D. variabilis* Broun, 1885, by indication, monotypy).

Xerostyggnus Broun, 1903: 79 (type species: *X. binodus* Broun, 1903, by indication, monotypy).

Type species. *Steriphus solidus* Erichson, 1842 (by indication, monotypy).

Diagnosis. Small to very small (3.0–6.5 mm); vestiture of subcircular scales and setae; scape long (surpassing posterior margin of eye when resting in scrobe); elytra with anteapical tubercle.

Relationships. *Steriphus* is the sister genus to the American genus *Listronotus*.

Species included. *Steriphus albidoparsus* (Lea, 1928); *S. alpinus* (Lea, 1928); *S. angusticollis* (Pascoe, 1870); *S. ascitus* (Pascoe, 1876); *S. binodulus* Broun, 1903; *S. caudatus* (Pascoe, 1870); *S. curvisetosus* (Lea, 1928); *S. diversipes* (Pascoe, 1870); *S. humeralis* (Lea, 1928); *S. incotaminatus* (Lea, 1899); *S. inermis* (Lea, 1928); *S. irrasus* (Lea, 1899); *S. longus* (Lea, 1928); *S. major* (Blackburn, 1890); *S. mecapispis* (Lea, 1899); *S. metallicus* (Lea, 1928); *S. mucronatus* (Lea, 1928); *S. murinus* (Pascoe, 1870); *S. parvicornis* (Lea, 1928); *S. parvonigrus* (Lea, 1928); *S. parvus* (Blackburn, 1890); *S. pullus* (Broun, 1910); *S. sericeus* (Blackburn, 1890); *S. solidus* Erichson, 1842; *S. stenoderes* (Lea, 1928); *S. variabilis* (Broun, 1885); *S. vittatus* (Blackburn, 1893).

Host plants. *Steriphus ascitus*: *Baumea articulata* (R. Br.) Blake, *B. rubiginosa* (Spreng.) Boeck., *Scirpus fluviatilis* (Torr.) Sojak (Cyperaceae) and *Typha orientalis* C. B. Presl. (Typhaceae); *S. diversipes*: *Medicago sativa* L. (Fabaceae) and *Rumex acetosella* L. (Polygonaceae); *S. variabilis*: *Cotula* spp. (Asteraceae), *Dichondra* sp. (Convolvulaceae) and *Myriophyllum* sp. (Haloragaceae) (May, 1977; Kuschel, 1990).

Immature stages. *Steriphus ascitus*, *S. caudatus*, *S. diversipes* and *S. variabilis* (May, 1970, 1977, 1993, 1994).

Geographical distribution. Australia and New Zealand (Schenkling and Marshall 1931; Kuschel 1972, 1990; May 1977; Zimmerman 1994).

Material examined. *Steriphus ascitus* (MZFC) and *S. variabilis* (MZFC).

***Tristanodes* Brinck, 1948**

<http://species-id.net/wiki/Tristanodes>

Tristanodes Brinck, 1948: 58.

Type species. *Tristanodes craterophilus* Brinck, 1948.

Diagnosis. Small to very small (3.7–6.5 mm); vestiture of seta-like scales and setae; pronotum subcylindrical.

Relationships. *Tristanodes* is the sister genus to both *Palaechthus* and *Palaechtodes*. Oberprieler (1992) considered that the distinction between *Tristanodes*, *Gunodes* and *Inaccodes* is not without doubt.

Species included. *Tristanodes attai* Brinck, 1948; *T. conicus* Brinck, 1948; *T. craterophilus* Brinck, 1948; *T. echinatus* Brinck, 1948; *T. insolitus* Brinck, 1948; *T. integer* Brinck, 1948; *T. medioides* Brinck, 1948; *T. minor* Brinck, 1948; *T. reppetonis* Brinck, 1948; *T. scirpophilus* Brinck, 1948; *T. sivertseni* Brinck, 1948.

Immature stages. *Tristanodes scirpophilus* (Kuschel, 1962).

Geographical distribution. Tristan da Cunha-Gough islands (Brinck 1948; Kuschel 1962).

Material examined. *Tristanodes attai* (BMNH) and *Tristanodes* spp. (BMNH).

Falklandiina subtr. n.

Type genus. *Falklandius* Enderlein, 1907.

Diagnosis. Small to very small (except *Liparogetus* and some species of *Gromilus*, which are medium-sized); rostrum stout, shorter than pronotum (except *Gromilus* and *Nestrius*, with relatively stout, medium-sized rostrum); pterygiae auriculate, exposed (Fig. 5); scrobes short, ill-defined, broad; eyes usually flat; postocular lobes usually absent (except *Gromilus* and *Falklandiopsis*); pronotum usually subcircular or subcylindrical; metanepisternal suture usually posteriorly fused or obliterated; elytra oblong-oval.

Included taxa. This new subtribe, which basically corresponds to the *Falklandius* generic group of Morrone (1997a), includes the genera *Falklandiellus*, *Falklandiopsis*, *Falklandius*, *Gromilus*, *Lanteriella*, *Liparogetus*, *Nestrius* and *Telurus*. The genera *Gromilus*, *Liparogetus* and *Nestrius* are distributed in New Zealand, whereas the five remaining genera are South American, distributed in the Subantarctic subregion of the Andean region (*sensu* Morrone, 2006).

Key to the genera of Falklandiina

- | | | |
|---|---|---------------------------------|
| 1 | Scrobes lateral | 2 |
| — | Scrobes dorsolateral to dorsal (Fig. 5) | 6 |
| 2 | Eyes transverse (Fig. 8); female elytral apex not produced; female ventrites 3 and 4 combined shorter than 5 | 3 |
| — | Eyes subcircular (Fig. 5); female elytral apex produced; female ventrites 3 and 4 combined longer than 5 | <i>Telurus</i> (Fig. 24) |
| 3 | Eyes slightly convex; postocular lobes absent..... | 4 |
| — | Eyes flat; postocular lobes slightly developed..... | 5 |
| 4 | Vestiture of setae only; rostrum very short, stout; rostral dorsal carinae absent; scrobes short, ill-defined; eyes dorsal; scape medium-sized (reaching eye when resting in scrobe); funicular segments 3-6 globose; pronotum subcircular; elytra with humeral tubercles; femora subcylindrical, markedly clavate; southern South America | <i>Falklandiopsis</i> (Fig. 24) |
| — | Vestiture of seta-like scales and setae; rostrum medium-sized, relatively stout; rostral dorsal carinae present; scrobes long, deep, sharply bordered, reaching eyes; eyes lateral; scape long (surpassing posterior margin of eye when resting in scrobe); funicular segments 3-6 elongate; pronotum subcylindrical; elytra lacking humeral tubercles; femora subcylindrical, clavate; New Zealand | <i>Gromilus</i> (Fig. 26) |
| 5 | Very small (2.6–3.5 mm); vestiture of subcircular scales and setae; rostrum lacking dorsal carinae; antennal insertion distal; club fusiform; pronotum transverse; metanepisternal suture posteriorly fused or obliterated; elytra with | |

- series of declivital tubercles; tibiae with spurs; southern South America
..... *Falklandiellus* (Fig. 23)
- Medium-sized (6.0–10.0 mm); vestiture of seta-like scales and setae; rostrum with dorsal carinae; antennal insertion at the middle of the rostrum; club inflated; pronotum subquadrate; metanepisternal suture complete; elytra lacking series of declivital tubercles; tibiae lacking spurs; New Zealand *Liparogetus*
- 6 Rostrum relatively stout, medium-sized, with dorsal carinae; eyes lateral; funicular segment 2 elongate; pronotum subcylindrical; scutellum not visible; New Zealand *Nestrius*
- Rostrum very short, stout, lacking dorsal carinae; eyes dorsal; funicular segment 2 globose; pronotum subcircular; scutellum visible; southern South America 7
- 7 Vestiture of seta-like scales and setae; eyes small; club inflated; pronotum with disc rugose; elytra with intervals convex; femora subcylindrical; tibiae subcylindrical..... *Falklandius* (Fig. 25)
- Vestiture of setae only; eyes very small, microphthalmic; club fusiform; pronotum with disc smooth, polished; elytra with intervals flat; femora dorsoventrally compressed; tibiae apically expanded *Lanteriella* (Fig. 27)

***Falklandiellus* Kuschel, 1950**

<http://species-id.net/wiki/Falklandiellus>

Fig. 23

Falklandiellus Kuschel, 1950: 14.

Type species. *Falklandius suffodens* Enderlein, 1907 (by original designation).

Diagnosis. Very small (2.6–3.5 mm); vestiture of subcircular scales and setae; rostrum lacking dorsal carinae; antennal insertion distal; club fusiform; pronotum transverse; metanepisternal suture posteriorly fused or obliterated; elytra with series of declivital tubercles; tibiae with spurs.

Relationships. *Falklandiellus* is the sister genus to *Telurus-Nestrius-Falklandius-Lanteriella*.

Species included. *Falklandiellus suffodens* (Enderlein, 1907).

Host plants. Bryophytes (Morrone 1995a).

Geographical distribution. Andean region (Subantarctic subregion), in southern Argentina, including the Falkland Islands (Islas Malvinas), and southern Chile (Morrone 1995a; Posadas 2008, 2012).

Material examined. *Falklandiellus suffodens* (BMNH, CADIC, MACN, MLP, MZFC, USNM, ZMHU).

***Falklandiopsis* Morrone and Anderson, 1995**

<http://species-id.net/wiki/Falklandiopsis>

Fig. 24

Falklandiopsis Morrone and Anderson, 1995: 5.

Type species. *Falklandius magellanicus* Morrone, 1992.

Diagnosis. Very small (3.5–4.0 mm); vestiture of setae only; rostrum very short, stout; rostral dorsal carinae absent; scrobes short, ill-defined; eyes dorsal; scape medium-sized (reaching eye when resting in scrobe); funicular segments 3–6 globose; pronotum subcircular; elytra with humeral tubercles; femora subcylindrical, markedly clavate.

Relationships. *Falklandiopsis* is the sister genus to both *Liparogetus* and the clade *Falklandiellus-Telurus-Nestrius-Falklandius-Lanteriella*.

Species included. *Falklandiopsis magellanica* (Morrone, 1992).

Host plants. *Nothofagus betuloides* (Mirb.) Oerst. (Nothofagaceae) (Morrone 1992b).

Geographical distribution. Andean region (Subantarctic subregion), in southern Chile (Morrone 1992b; Posadas 2012).

Material examined. *Falklandiopsis magellanica* (MLP, MZFC, NZAC, ZMHU).

***Falklandius* Enderlein, 1907**

<http://species-id.net/wiki/Falklandius>

Fig. 25

Falklandius Enderlein, 1907: 65.

Type species. *Falklandius brachyomma* Enderlein, 1907 (= *Otiorhynchus antarcticus* Stierlin, 1903) (by original designation).

Diagnosis. Small to very small (1.9–6.1 mm); vestiture of seta-like scales and setae; eyes small; club inflated; pronotal disc rugose; elytra with intervals convex.

Relationships. *Falklandius* is the sister genus to *Lanteriella*, as found in a previous analysis (Morrone, 1997a).

Species included. *Falklandius antarcticus* (Stierlin, 1903); *F. chilensis* Morrone and Anderson, 1995; *F. goliath* Morrone, 1992; *F. kuscheli* Morrone, 1992; *F. peckorum* Morrone and Anderson, 1995; *F. turbificatus* Enderlein, 1907.

Host plants. *Falklandius antarcticus*: *Callitricha* sp. (Callitrichaceae), *Myrteola nummularia* (Poir.) O. Berg (Myrtaceae), *Nothofagus antarctica* (G. Forster) Oerst. (Nothofagaceae) and *Poa flabellata* (Lam.) Raspail (Poaceae) (Morrone, 1992b); *F. turbificatus*: *Myrteola nummularia* (Poir.) O. Berg (Myrtaceae) (Morrone, 1992b).

Geographical distribution. Andean region (Subantarctic subregion), in southern Argentina, including the Falkland Islands (Islas Malvinas) and southern Chile (Morrone 1992b; Morrone and Anderson 1995; Posadas 2008, 2012).

Material examined. *Falklandius antarcticus* (AMPC, BMNH, CADIC, CMNC, CWOB, MACN, MHNS, MZFC, USNM), *F. chilensis* (AMNH, BMNH, CMNC, CWOB, FMNH, MLP, MZFC, USNM), *F. goliath* (BMNH), *F. kuscheli* (BMNH), *F. peckorum* (AMNH, BMNH, CMNC, CWOB, FMNH, MLP, MZFC, USNM), *F. turbificatus* (BMNH) and *Falklandius* spp. (MZFC).

Gromilus Blanchard, 1853

<http://species-id.net/wiki/Gromilus>

Fig. 26

Gromilus Blanchard, 1853: 208.

Clypeorrhynchus Sharp, 1883: 26 (type species: *C. gracilipes* Sharp, 1883, by indication, monotypy).

Clypeorrhynchus Kirby, 1885: 100 (unjustified emendation).

Dacnophylla Broun, 1893a: 1471 (type species: *D. setosa* Broun, 1893a, by indication, monotypy).

Hycanus Broun, 1905: 545 (type species: *H. cockaynei* Broun, 1905, by indication, monotypy).

Stilbodiscus Broun, 1909: 117 (type species: *S. setarius* Broun, 1909, by indication, monotypy).

Phygothalpus Broun, 1913: 117 (type species: *P. sulcicollis* Broun, 1913, by indication, monotypy).

Heteromias Broun, 1913: 120 (*non* Faust, 1897) (type species: *H. foveirostris* Broun, 1913, by indication, monotypy).

Pseudohycanus Brookes, 1951: 57 (type species: *P. fallai* Brookes, 1951).

Type species. *Gromilus insularis* Blanchard, 1853 (by indication, monotypy).

Diagnosis. Small to medium-sized (3.5–7.5 mm); vestiture of seta-like scales and setae; rostrum medium-sized, relatively stout; rostral dorsal carinae present; scrobes long, deeply bordered, reaching eyes; eyes lateral; scape long (surpassing posterior margin of eye when resting in scrobe); funicular segments 3–6 elongate; pronotum subcylindrical; elytra lacking humeral tubercles.

Relationships. *Gromilus* is the sister genus to the remaining genera of Falklandiina. Kuschel (1964) already noted the close relationship of *Gromilus* with *Nestrius*, *Liparogetus* and *Falklandius*.

Species included. *Gromilus anthracinus* (Broun, 1921); *G. aucklandicus* Kuschel, 1971; *G. bicarinatus* (Broun, 1921); *G. bifoveatus* (Broun, 1923); *G. brevicornis* (Broun, 1893); *G. brounii* Morrone, 2011; *G. calvulus* (Broun, 1913); *G. caudatus* (Broun, 1913); *G. clarulus* (Broun, 1917); *G. cockaynei* (Broun, 1905); *G. cordipennis* (Broun, 1893); *G. cristatus* (Broun, 1893); *G. dorsalis* (Broun, 1921); *G. exiguus* (Brookes, 1951); *G. fallai* (Brookes, 1951); *G. foveirostris* (Broun, 1913); *G. furvus* (Broun, 1921); *G. gracilipes* (Sharp, 1883); *G. granissimus* (Broun, 1917); *G. halli* (Broun, 1917); *G. im-*

pressus (Broun, 1893); *G. inophloeoides* (Broun, 1904); *G. insularis* Blanchard, 1853; *G. kuschelii* Morrone, 2011; *G. laqueorum* Kuschel, 1964; *G. majusculus* (Broun, 1915); *G. merus* (Broun, 1917); *G. narinosus* Kuschel, 1971; *G. nitidellus* (Broun, 1917); *G. nitidulus* (Broun, 1915); *G. nodiceps* (Broun, 1914); *G. philpotti* (Broun, 1917); *G. setosus* (Broun, 1893); *G. sparsus* (Broun, 1921); *G. striatus* (Broun, 1915); *G. sulcicollis* (Broun, 1913); *G. sulcipennis* (Broun, 1917); *G. tenuiculus* (Broun, 1921); *G. thoracicus* (Broun, 1893); *G. variegatus* (Broun, 1893); *G. veneris* (Kirsch, 1877).

Host plants. *Gromilus fallai*: *Blechnum capense* Burm. f. (Blechnaceae); *G. insularis*: *Colobanthus* sp. (Caryophyllaceae), *Pleurophyllum* sp. (Asteraceae), *Poa litorosa* Cheeseman (Poaceae), *Polystichum vestitum* (G. Forst.) C. Presl. (Dryopteridaceae), *Pleurophyllum criniferum* Hook. f. (Asteraceae), *Stilbocarpa polaris* (Homb. and Jacq.) Gray (Araliaceae) and *Tillaea moschata* DC (Crassulaceae); *G. setosus*: *Blechnum* sp. (Blechnaceae) and *Gahnia* sp. (Cyperaceae); *G. veneris*: *Blechnum capense* (L.) Schlecht. (Blechnaceae), *Polystichum* sp. (Dryopteridaceae) and *Pteris* sp. (Pteridaceae); *G. thoracicus*: *Anisotome latifolia* Hook. f. (Apiaceae), *Bulbinella* sp. (Liliaceae), *Cotula pluma* Hook. f. and *Pleurophyllum criniferum* Hook. f. (Asteraceae) and *Poa litorosa* Cheeseman (Poaceae) (May 1977, 1993; Kuschel 1964, 1971, 1990).

Immature stages. *Gromilus exiguis*, *G. insularis*, *G. thoracicus* and *G. veneris* (May, 1971).

Geographical distribution. New Zealand (Schenkling and Marshall 1929; Kuschel 1964, 1971, 1990).

Material examined. *Gromilus gracilipes* (MZFC), *G. insularis* (MZFC), *G. laqueorum* (MZFC), *G. merus* (MZFC), *G. nitidellus* (MZFC) and *G. veneris* (MZFC).

Lanteriella Morrone, 1992

<http://species-id.net/wiki/Lanteriella>

Fig. 27

Lanteriella Morrone, 1992b: 167.

Type species. *Lanteriella microptalma* Morrone, 1992.

Diagnosis. Very small (3.4–3.8 mm); vestiture of setae only; eyes very small, microphthalmic; pronotal disc smooth, polished; femora dorsoventrally compressed; tibiae apically expanded.


Relationships. *Lanteriella* is the sister genus to *Falklandius*, as found in a previous analysis (Morrone, 1997a).

Biology. The only species of this genus was hypothesized to live in litter or soil (Morrone 1992b).

Species included. *Lanteriella microptalma* Morrone, 1992.

Geographical distribution. Andean region (Subantarctic subregion), in the Falkland Islands (Islas Malvinas) (Morrone 1992b; Posadas 2008).

Material examined. *Lanteriella microptalma* (BMNH).



Figures 27–35. Habitus of representative Listroderini. **27** *Lanteriella microphthalmia* **28** *Telurus caudiculatus* **29** *Acrorius papallacta* **30** *Acrostomus bruchi* **31** *Antarctobius lacunosus* **32** *Germainiellus dentipennis* **33** *Lamiarhinus aelficus* **34** *Listroderes annulipes* **35** *Philippius superbus*.

***Liparogetus* Broun, 1915**

<http://species-id.net/wiki/Liparogetus>

Liparogetus Broun, 1915: 331.

Type species. *Liparogetus sulcatissimus* Broun, 1915 (by indication, monotypy).

Diagnosis. Small to medium-sized (6.0–10.0 mm); vestiture of seta-like scales and setae; rostrum with dorsal carinae; antennal insertion at the middle of the rostrum; club inflated; pronotum subquadrate; metanepisternal suture complete; tibiae lacking spurs.

Relationships. *Liparogetus* is the sister genus to both *Falklandiopsis* and the clade *Falklandiellus-Telurus-Nestrius-Falklandius-Lanteriella*.

Species included. *Liparogetus sulcatissimus* Broun, 1915.

Geographical distribution. New Zealand (Alonso-Zarazaga and Lyal 1999).

Material examined. *Liparogetus sulcatissimus* (MZFC).

***Nestrius* Broun, 1893**

<http://species-id.net/wiki/Nestrius>

Nestrius Broun, 1893a: 1480.

Phyllodytes Broun, 1893a: 1479 (*non* Wagler 1830, *nec* Gistel 1848, *nec* Finsch 1873) (type species: *P. foveatus* Broun, 1893a, by indication, monotypy).

Plotnus Broun, 1893a: 1481 (type species: *P. ovithorax* Broun, 1893a, by indication, monotypy).

Proboscoelus Broun, 1909: 55 (type species: *P. sculpturatus* Broun, 1909, by indication, monotypy).

Drymaria Broun, 1909: 56 (type species: *D. cilipes* Broun, 1909, by indication, monotypy).

Phyllodytesius Schenkling & Marshall, 1929: 57 (replacement name for *Phyllodytes* Broun).

Type species. *Nestrius serripes* Broun, 1893a (by indication, monotypy).

Diagnosis. Small to very small (2.8–5.0 mm); vestiture of seta-like scales and setae; rostrum relatively stout, medium-sized, with dorsal carinae; eyes lateral; funicular segment 2 elongate; pronotum subcylindrical; scutellum not visible.

Relationships. *Nestrius* is the sister genus to *Falklandius-Lanteriella*, confirming Kuschel's (1964) suggestion that it was intermediate between *Gromilus* and *Falklandius*.

Species included. *Nestrius bifurcus* Kuschel, 1964; *N. cilipes* Broun, 1909; *N. crassicornis* Broun, 1915; *N. foveatus* (Broun, 1893); *N. hudsoni* Marshall, 1953; *N. irregularis* (Broun, 1910); *N. laqueorum* Kuschel, 1964; *N. ovithorax* (Broun, 1893); *N. prolixus* Broun, 1917; *N. serripes* Broun, 1893; *N. sculpturatus* (Broun, 1909); *N. simmondsi* Broun, 1921; *N. sulcirostris* Broun, 1917; *N. zenoscelis* Broun, 1921.

Geographical distribution. New Zealand (Schenkling and Marshall 1929; Kuschel 1964, 1971).

Material examined. *Nestrius foveatus* (MZFC) and *N. sculpturatus* (MZFC).

***Telurus* Kuschel, 1955, rev. placement**

<http://species-id.net/wiki/Telurus>

Fig. 28

Telurus Kuschel, 1955: 288.

Type species. *Antarctobius laticauda* Champion, 1918 (= *T. dissimilis* [Fairmaire, 1885]) (by original designation).

Diagnosis. Small (3.9–6.5 mm); vestiture of setae only; eyes subcircular, slightly convex; female elytral apex produced; female ventrites 3 and 4 combined longer than 5.

Relationships. *Telurus* is closely related to *Falklandius-Lanteriella*, as found in a previous analysis (Morrone 1997a), and to *Nestrius*, from New Zealand. Based on the presence of small scars (due to deciduous cusps) on the mandibles of *T. caudiculatus*, Oberprieler (2010) excluded the genus from Listroderini and transferred it to Cylydrorhininae (Entiminae). Future molecular analyses are required to corroborate its precise placement.

Species included. *Telurus caudiculatus* Morrone and Anderson, 1995; *T. dissimilis* (Fairmaire, 1885).

Geographical distribution. Andean region (Subantarctic subregion), in southern Chile (Morrone and Anderson 1995; Posadas 2012).

Material examined. *Telurus caudiculatus* (AMNH, BMNH, CMNC, CNCI, CWOB, MCZ, MHNS, MLP, MZFC, USNM, ZMC) and *T. dissimilis* (BMNH, IPUM, MHNS, MZFC, NZAC).

Subtribe Listroderina LeConte, 1876

Type genus. *Listroderes* Schönherr, 1826.

Diagnosis. Rostrum relatively stout, medium-sized, shorter than pronotum; scrobes short, ill-defined, broad; funicular segmen 1 longer than 2; elytra usually oblong-oval (subrectangular in *Lamiarhinus* and *Philippius*), with intervals convex and with anteapical tubercle (except for *Rupanius*).

Included taxa. This subtribe, representing the listroderines in the strictest sense, includes the genera *Acroriellus*, *Acrorius*, *Acrostomus*, *Antarctobius*, *Germainiellus*, *Hyperoides*, *Lamiarhinus*, *Listroderes*, *Methypora*, *Philippius*, *Rupanius* and *Trachodema*. In a previous analysis restricted to American taxa (Morrone 1997a) most of these genera were placed as a grade basal to other listroderines. *Methypora* is distributed in Australia; and the remaining genera are South American: *Listroderes* is widely ranged in the Andean and Neotropical regions, and the other genera are found in the Andean region and the South American Transition Zone (*sensu* Morrone 2006).

Key to the genera of Listroderina

- 1 Elytral disc slightly convex to flat..... **2**
- Elytral disc convex **6**
- 2 Elytra oblong-oval (Fig. 10) **3**
- Elytra subrectangular (Fig. 9) **4**
- 3 Vestiture of seta-like scales and setae; scape medium-sized (reaching eye when resting in scrobe)..... *Acrorius* (Fig. 29)
- Vestiture of scales with finger-like processes and setae; scape long (surpassing posterior margin of eye when resting in scrobe) *Trachodema*
- 4 Pronotum transverse; elytra with carina on apical declivity, disc slightly convex, lacking anteapical tubercle *Rupanius*
- Pronotum subcircular or subcylindrical; elytra lacking carina on apical declivity, disc flat, with anteapical tubercle **5**
- 5 Large (17.5–22.8 mm); mandibles with 3-4 setae; pronotum subcircular, wider than elytra, with tubercles; scutellum not visible; elytra fused along interelytral suture, with series of declivital tubercles; female elytral apex not produced; tibiae lacking spurs; tarsomeres 3 subcylindrical (Fig. 12); southern South America *Philippius* (Fig. 35)
- Small (4.0–7.0 mm); mandibles with 2 setae; pronotum subcylindrical, narrower than elytra, lacking tubercles; scutellum visible; elytra not fused along interelytral suture, lacking series of declivital tubercles; female elytral apex produced; tibiae with spurs; tarsomeres 3 bilobed (Fig. 11); Australia *Methypora*
- 6 Funicular segments 3-6 globose; pronotum lacking tubercles; elytra oblong-oval, not fused along interelytral suture..... **7**
- Funicular segments 3-6 elongate; pronotum with tubercles; elytra subrectangular, fused along interelytral suture *Lamiarbinus* (Fig. 33)
- 7 Pronotum transverse or subquadrate; postocular lobes present **8**
- Pronotum subcircular; postocular lobes absent *Antarctobius* (Fig. 31)
- 8 Integument reddish brown; epistome not raised; pronotum transverse..... **9**
- Integument black; epistome raised; pronotum subquadrate..... *Acrostomus* (Fig. 30)
- 9 Vestiture of seta-like or lanceolate scales and setae; scrobal ventral tooth absent..... **10**
- Vestiture of subcircular scales and setae; scrobal ventral tooth usually present..... *Listroderes* (Fig. 34)
- 10 Vestiture of seta-like scales and setae **11**
- Vestiture of lanceolate scales and setae *Hyperoides*
- 11 Elytral interval 3 lacking series of three declivital tubercles..... *Germaniellus* (Fig. 32)
- Elytral interval 3 with series of three declivital tubercles..... *Acroriellus*

***Acroriellus* Morrone and Ocampo, 1995**

<http://species-id.net/wiki/Acroriellus>

Acroriellus Morrone & Ocampo, 1995: 257.

Type species. *Acroriellus viridisquamulosus* Morrone and Ocampo, 1995.

Diagnosis. Very small (2.5–3.8 mm); vestiture of seta-like scales and setae; elytra with small, rounded tubercles and series of three tubercles on interval 3.

Relationships. *Acroriellus* is the sister genus to *Acrostomus-Hyperoides*. Originally, it was suggested that it was close to *Acrorius* (Morrone and Ocampo 1995).

Species included. *Acroriellus bobi* Morrone and Ocampo, 1995; *A. carinatus* Morrone and Ocampo, 1995; *A. similaris* Morrone and Ocampo, 1995; *A. tuberculatus* Morrone and Ocampo, 1995; *A. viridisquamulosus* Morrone and Ocampo, 1995; *A. vittatae* Morrone and Ocampo, 1995.

Geographical distribution. South American Transition Zone (North Andean Paramo and Puna biogeographical provinces), in Colombia, Ecuador and Peru (Morrone and Ocampo 1995).

Material examined. *Acroriellus bobi* (AMNH, CMNC), *A. carinatus* (CMNC), *A. similaris* (CMNC), *A. tuberculatus* (CMNC), *A. viridisquamulosus* (CMNC, FMNH) and *A. vittatae* (AMNH, USNM).

***Acrorius* Kirsch, 1889**

<http://species-id.net/wiki/Acrorius>

Fig. 29

Acrorius Kirsch, 1889: 25.

Ocromis Sharp, 1890: 152 (lapsus).

Type species. *Acrorius puncticollis* Kirsch, 1889 (by indication, monotypy).

Diagnosis. Small (4.0–6.8 mm); vestiture of seta-like scales and setae; scape medium-sized (reaching eye when resting in scrobe); elytra with small, rounded tubercles.

Relationships. *Acrorius* is the sister genus to *Trachodema-Lamiarhinus-Philippius*, taxa that in a previous analysis (Morrone 1997a) constituted a paraphyletic group.

Species included. *Acrorius andersoni* Morrone, 1994; *A. bolivianus* Ocampo and Morrone, 1996; *A. cuprinus* Morrone, 1994; *A. nymphalis* Morrone, 1994; *A. otramas* Ocampo and Morrone, 1996; *A. papallacta* Morrone, 1994; *A. pillahuata* Morrone, 1994; *A. plicatifrons* Morrone, 1994; *A. puncticollis* Kirsch, 1889; *A. sisyphus* Morrone, 1994.

Geographical distribution. Bolivia, Ecuador and Peru (Morrone 1994a; Ocampo and Morrone 1996).

Material examined. *Acrorius andersoni* (CMNC), *A. bolivianus* (CMNC, MZFC), *A. cuprinus* (CMNC), *A. nymphalis* (CMNC), *A. otramas* (CMNC), *A. papallacta* (CMNC, MZFC), *A. pillahuata* (CMNC, FMNH), *A. plicatifrons* (FMNH) and *A. sisyphus* (CNCI, CMNC).

***Acrostomus* Kuschel, 1955**

<http://species-id.net/wiki/Acrostomus>

Fig. 30

Acrostomus Kuschel, 1955: 287.

Type species. *Adioristus bruchi* Hustache, 1926 (by original designation).

Diagnosis. Medium-sized (7.3–13.8 mm); integument black; vestiture of seta-like scales and setae; epistome raised; scrobal ventral tooth usually present; pronotum subquadrate.

Relationships. *Acrostomus* is the sister genus to *Hyperoides*.

Species included. *Acrostomus bruchi* (Hustache, 1926); *A. cruralis* Kuschel, 1958; *A. foveicollis* Kuschel, 1958; *A. griseus* (Guérin-Ménéville, 1839); *A. magellanicus* Kuschel, 1958; *A. mordor* Morrone, 1994; *Acrostomus vianai* Kuschel, 1958.

Host plants. *Acrostomus magellanicus* and *A. vianai*: *Azorella trifurcata* (Gaertner) Pers., *Bolax gummifera* (Lam.) Spreng. and *Mulinum spinosum* (Cav.) Pers. (Apiaceae) (Morrone, 1994b).

Geographical distribution. Andean region (Patagonian subregion), in southern Argentina and southern Chile (Morrone 1994b).

Material examined. *Acrostomus bruchi* (CWOB, IPCN, MACN, MLP, MZFC), *A. cruralis* (MACN, USNM), *A. foveicollis* (CBPC, CWOB, MHNS, MZFC), *A. griseus* (CWOB, FIML, IPUM, MHNS, MLP, MZFC), *A. magellanicus* (BMNH, MHNS, USNM), *A. mordor* (AMNH, MACN, MLP, MZFC) and *A. vianai* (BMNH, MHNS).

***Antarctobius* Fairmaire, 1885**

<http://species-id.net/wiki/Antarctobius>

Fig. 31

Antarctobius Fairmaire, 1885: 58.

Type species. *Antarctobius lacunosus* Fairmaire, 1885 (subsequent designation by Morrone, 1992a).

Diagnosis. Small to medium-sized (3.7–9.5 mm); vestiture of seta-like or subcircular scales and setae; pronotum subcircular; postocular lobes absent.

Relationships. *Antarctobius* is closely related to *Germainiellus*, *Listroderes* and the clade *Methylpora-Rupanius-Acrorius-Trachodema-Lamiarhinus-Philippius*. The distinction between *Antarctobius*, *Germainiellus* and *Listroderes* is not without doubt (see Morrone and Marvaldi, 1998), and future analyses may determine if they are merged into a single genus.

Species included. *Antarctobius abditus* (Enderlein, 1907); *A. bidentatus* (Champion, 1918); *A. falklandicus* (Enderlein, 1907); *A. germaini* (Kolbe, 1907); *A. hyadesii* Fairmaire, 1885; *A. lacunosus* Fairmaire, 1885; *A. malvinensis* Posadas and Morrone, 2004; *A. rugirostris* Champion, 1918; *A. vulsus* (Enderlein, 1907); *A. yefacel* Morrone, 1992.

Host plants. *Antarctobius abditus*: *Senecio candidans* DC (Asteraceae); *A. hyadesii*: *Senecio alloeophyllus* O. Hoffm. and *S. candidans* DC (Asteraceae) (Morrone, 1992a; Marvaldi, 1998).

Immature stages. *Antarctobius abditus* and *A. falklandicus* (Marvaldi, 1998).

Geographical distribution. Andean region (Subantarctic subregion), in southern Chile and southern Argentina, including the Falkland Islands (Islas Malvinas) (Morrone 1992a; Posadas and Morrone 2004; Posadas 2008, 2012).

Material examined. *Antarctobius abditus* (BMNH), *A. bidentatus* (BMNH), *A. falklandicus* (AMPC, BMNH, MZFC), *A. germaini* (AMNH, BMNH, CADIC, CMNC, CWOB, IPUM, MHNS, MLP, MZFC), *A. hyadesii* (BPBM, CMNC, MHNS, MZFC), *A. lacunosus* (BMNH, MCZ, MHNS), *A. rugirostris* (BMNH), *A. vulsus* (BMNH, USNM) and *A. yefacel* (AMNH).

***Germainiellus* Morrone, 1993**

<http://species-id.net/wiki/Germainiellus>

Fig. 32

Germainiellus Morrone, 1993a: 125.

Type species. *Listroderes dentipennis* Germain, 1895 (by original designation).

Diagnosis. Small to medium-sized (6.0-8.4 mm); vestiture of seta-like scales and setae; pronotum transverse; postocular lobes present.

Relationships. *Germainiellus* is closely related to *Antarctobius*, *Listroderes* and the clade *Methypora-Rupanius-Acrorius-Trachodema-Lamiarhinus-Philippius*. It was originally described as intermediate between *Antarctobius* and *Listroderes* (Morrone, 1993a). The distinction between *Antarctobius*, *Germainiellus* and *Listroderes* is not without doubt (see Morrone and Marvaldi, 1998), and future analyses may determine if they are merged into a single genus.

Species included. *Germainiellus angulipennis* (Germain, 1895); *G. attenuatus* (Germain, 1895); *G. dentipennis* (Germain, 1895); *G. fulvicornis* (Germain, 1895); *G. laevirostris* (Germain, 1895); *G. lugens* (Germain, 1895); *G. ovatus* (Boheman, 1842); *G. philippii* (Germain, 1896); *G. planipennis* (Blanchard, 1851); *G. punctiventris* (Germain, 1895); *G. rugipennis* (Blanchard, 1851); *G. salebrosus* (Enderlein, 1907); *Germainiellus* spp. (MZFC).

Host plants. *Germainiellus dentipennis* and *G. fulvicornis*: *Nothofagus* sp. (Nothofagaceae); *G. laevirostris*: *Senecio smithii* DC (Asteraceae); *G. planipennis*: *Nothofagus dombeyi* (Mirb.) Oerst. (Nothofagaceae) and *Peumus boldus* Mol. (Monimiaceae); *G. salebrosus*: *Empetrum rubrum* Vahl ex Willd. (Empetraceae) (Morrone, 1993a).

Geographical distribution. Andean region (Subantarctic subregion), in southern Chile and southern Argentina, including the Falkland Islands (Islas Malvinas) (Morrone 1993a, 1994e; Posadas 2008, 2012).

Material examined. *Germainiellus angulipennis* (MHNS), *G. attenuatus* (ARPC, MHNS), *G. dentipennis* (CBPC, CMNC, CWOB, MHNS, USNM), *G. fulvicornis* (AMNH, BPBM, CBCP, CMNC, CWOB, FIML, MCZ, MHNS, MLP, MZFC, USNM), *G. laevirostris* (BPBM, IPUM, MCZ, MHNS, MLP, USNM), *G. lugens* (CMNC, CWOB, IPUM, MACN, MCZ, MHNS, MZFC), *G. ovatus* (BMNH, MHNS, USNM), *G. philippii* (CMNC, DEI, MHNS, MZFC), *G. planipennis* (BMNH, CWOB, MHNS), *G. punctiventris* (MHNS), *G. rugipennis* (AMNH, BPBM, CADIC, CBCP, CMNC, MHNS, MLP, MZFC, USNM) and *G. salebrosus* (BMNH).

Hyperoides Marshall, 1914

<http://species-id.net/wiki/Hyperoides>

Hyperoides Marshall, 1914: 236.

Type species. *Hyperoides fragariae* Marshall, 1914 (by indication, monotypy).

Diagnosis. Small to medium-sized (5.1–7.5 mm); vestiture of lanceolate scales and setae; postocular lobes present; elytra lacking anteapical tubercle.

Relationships. *Hyperoides* is the sister genus to *Acrostomus*, contrasting with its more isolated position in a previous analysis (Morrone, 1997a).

Species included. *Hyperoides balfourbrowniae* (Kuschel, 1952); *H. fragariae* Marshall, 1914; *H. murinus* (Germain, 1896); *H. subcinctus* (Boheman, 1842); *H. victus* (Germain, 1896).

Host plants. *Hyperoides fragariae*: *Fragaria vesca* L. (Rosaceae); *H. subcinctus*: *Senecio* sp. (Asteraceae); *H. murinus*: *Citrulus vulgaris* Schrad. (Cucurbitaceae), *Phaseolus* sp. (Fabaceae) and *Solanum tuberosum* L. (Solanaceae); *H. victus*: *Senecio bahioides* Hook. et Arn. (Asteraceae) (Morrone 1993b; Lanteri et al. 2002).

Geographical distribution. Neotropical region and Andean region (Central Chilean subregion), in Argentina, Chile and Uruguay, and introduced into South Africa (Morrone 1993b).

Material examined. *Hyperoides balfourbrowniae* (MLP, MZFC), *H. fragariae* (BMNH, CBPC, MNHN), *H. murinus* (BMNH, CWOB, MHNS, MZFC), *H. subcinctus* (AMNH, BMNH, CBPC, CMNC, CWOB, IADIZA, MACN, MHNS, MNHN, MZFC) and *H. victus* (BMNH, CMNC, CWOB, MHNS).

Lamiarhinus Morrone, 1992

<http://species-id.net/wiki/Lamiarhinus>

Fig. 33

Lamiarhinus Morrone, 1992c: 419.

Type species. *Lamiarhinus aelficus* Morrone, 1992.

Diagnosis. Small to medium-sized (5.7–6.8 mm); vestiture of seta-like scales and setae; funicular segments 3–6 elongate; pronotum with tubercles; elytra subrectangular, fused along interelytral suture.

Relationships. *Lamiarhinus* is the sister genus to *Philippius*. In a previous analysis (Morrone 1997a), it was considered to be related to *Trachodema*.

Species included. *Lamiarhinus aelficus* Morrone, 1992; *L. horridus* (Germain, 1896).

Host plants. *Lamiarhinus aelficus*: *Podanthus ovatifolius* Lag. (Asteraceae) (Morrone 1992c).

Geographical distribution. Andean region (Central Chilean subregion) (Morrone 1992c).

Material examined. *Lamiarhinus aelficus* (CMNC, CWOB, MLP, MZFC) and *L. horridus* (MHNS).

Listroderes Schönherr, 1826

<http://species-id.net/wiki/Listroderes>

Fig. 34

Listroderes Schönherr, 1823: col. 1142 (*nom. nud.*).

Listroderes Schönherr, 1826: 158.

Listroderus Erichson, 1847: 129 (lapsus).

Listoderes Kuschel, 1990: 71 (lapsus).

Type species. *Listroderes costirostris* Schönherr, 1826 (by original designation, combined description).

Diagnosis. Small to medium-sized (3.9–12.5 mm); vestiture of subcircular scales and setae; scrobal ventral tooth usually present.

Relationships. *Listroderes* is closely related to *Antarctobius*, *Germainiellus* and the clade *Methypora-Rupanius-Acrorius-Trachodema-Lamiarhinus-Philippius*. The distinction between *Antarctobius*, *Germainiellus* and *Listroderes* is not without doubt (see Morrone and Marvaldi 1998), and future analyses may determine if they are merged into a single genus.

Species included. *Listroderes affinis* Hustache, 1926; *L. angusticeps* Blanchard, 1851; *L. annulipes* Blanchard, 1851; *L. apicalis* Waterhouse, 1841; *L. bimaculatus* Boheman, 1842; *L. brevirostris* Germain, 1895; *L. brevisetis* Hustache, 1926; *L. bruchi* Hustache, 1926; *L. charybdis* Morrone, 1993; *L. cinerarius* Blanchard, 1851; *L. confusus* Hustache, 1926; *L. costirostris* Schönherr, 1826; *L. curvipes* Germain, 1895; *L. delaiguei* Germain, 1895; *Listroderes desertorum* Germain, 1895; *L. difficilis* Germain, 1895; *L. elegans* Hustache, 1926; *L. erinaceus* Germain, 1895; *Listroderes fallax* Germain, 1895; *L. foveatus* (Lea, 1928); *L. hoffmanni* Germain, 1895; *L. howdenae* Morrone, 1993; *L. leviculus* Kuschel, 1952; *L. montanus* Germain, 1895; *L. nodifer* Boheman, 1842; *L. obliquus* Klug, 1829; *L. obrieni* Morrone, 1993; *L. paranensis* Hustache, 1926; *L. punicola* Kuschel, 1949; *L. pusillus* Hustache, 1926; *L. robustior* Schenkling and Marshall, 1931; *L. robustus* Waterhouse, 1841;

L. scylla Morrone, 1993; *L. trivialis* Germain, 1895; *L. tuberculifer* Blanchard, 1851; *L. uruguayensis* Kuschel, 1952; *L. wagneri* Hustache, 1926; *Listroderes wittei* Hustache, 1926.

Host plants. *Listroderes apicalis*: *Beta vulgaris* L. (Chenopodiaceae), *Helianthus annus* L. (Asteraceae) and *Triticum aestivum* L. (Poaceae); *L. bimaculatus*: *Baccharis linearis* (Ruiz and Pav.) Pers. (Asteraceae) and *Puya chilensis* Molina (Bromeliaceae); *L. bruchi*: *Baccharis salicifolia* (Ruiz and Pavón) Pers. and *Senecio subulatus* Don Hooker et Arnott (Asteraceae); *L. cinerarius*: *Atriplex* sp. (Chenopodiaceae); *L. costirostris*, *L. difficilis* and *L. obliquus*: *Apium graveolens* L. and *Daucus carota* L. (Apiaceae), *Brassica rapa* L., *B. oleracea* L. and *Coronopus didymus* (L.) Smith (Brassicaceae), *Rumex altissimus* Wood (Polygonaceae), *Nicotiana tabacum* L. and *Solanum tuberosum* L. (Solanaceae) and *Stellaria* spp. (Caryophyllaceae); *L. robustus*: *Atriplex semibaccata* R. Br. (Chenopodiaceae); *L. uruguayensis*: *Hydrocotyle bonariensis* Lam. (Apiaceae) (Morrone 1993d, 1995b; Marvaldi 1998; Lanteri et al. 2002).

Immature stages. *Listroderes bruchi*, *L. delaiguei* and *L. difficilis* (May 1977, 1993, 1994; Marvaldi 1998).

Geographical distribution. Andean region (Subantarctic, Central Chilean and Patagonian subregions), South American Transition Zone and Neotropical region, in Argentina, Brazil, Chile, Paraguay, Peru and Uruguay, and introduced into Australia, Easter Island, Israel, Japan, New Zealand, South Africa, Spain and USA (Wibmer and O'Brien 1986; Kuschel 1990; Morrone 1993c-e, 1995b; Morrone 2002b; Friedman 2009; Posadas 2012).

Material examined. *Listroderes affinis* (CBPC, IPCN, IPUM, MACN, MNHN), *L. angusticeps* (MHNS, MNHN, MZFC), *L. annulipes* (CBPC, CWOB, MHNS, MNHN, MZFC), *L. apicalis* (AMNH, BMNH, CMNC, MACN, MHNS, MLP, MZFC), *L. bimaculatus* (AMNH, BMNH, CMNC, CWOB, MACN, MHNS), *L. brevirostris* (MHNS), *L. brevisetis* (CBCP, DZUP, IPCN, MACN, MLP, MNHN), *L. bruchi* (CMNC, DZUP, FIML, IADIZA, MACN, MLP, MZFC), *L. charybdis* (MACN, MLP), *L. cinerarius* (BMNH, CMNC, CWOB, IADIZA, MHNS, MNHN, MZFC), *L. confusus* (DZUP, FIML, MACN, MLP, MNHN), *L. costirostris* complex (AMNH, BMNH, CBPC, CMNC, CWOB, DZUP, FIML, GJWC, MACN, MHNS, MLP, MNHN, MZFC, MZSP, USNM), *L. curvipes* (BMNH, CWOB, MHNS), *L. delaiguei* (BMNH, CADIC, CWOB, IPUM, MHNS, MZFC), *L. desertorum* (BMNH, CMNC, CWOB, MHNS, MZFC), *L. elegans* (GJWC, MACN, MLP, MNHN), *L. erinaceus* (MHNS), *L. fallax* (CWOB, MHNS, MZFC), *L. foveatus* (BMNH, CMNC, DZUP, FIML, GJWC, MACN, MZSP), *L. hoffmanni* (BMNH, CWOB, MHNS, MZFC), *L. howdenae* (CMNC, MLP, MZFC), *L. leviculus* (BMNH), *L. montanus* (MHNS, MZFC), *L. nodifer* (BMNH, CWOB, MACN, MHNS), *L. obrieni* (MHNS, MLP, MZFC), *L. paranensis* (DZUP, MNHN), *L. punicola* (CMNC, MHNS, MZFC), *L. pusillus* (CBPC, MLP, MNHN, MZFC), *L. robustior* (BMNH, CMNC, CWOB, MHNS, MLP, MZFC), *L. robustus* (CMNC, CWOB, MHNS), *L. scylla* (FIML, MLP), *L. trivialis* (MHNS), *L. tuberculifer* (CMNC, MHNS, MZFC), *L. uruguayensis* (BMNH, CMNC), *L. wagneri* (BMNH, MNHN) and *L. wittei* (MACN, MNHN).

***Methypora* Pascoe, 1865**

<http://species-id.net/wiki/Methypora>

Methypora Pascoe, 1865: 416.

Type species. *Methypora postica* Pascoe, 1865 (by indication, monotypy).

Diagnosis. Small (4.0–7.0 mm); vestiture of subcircular scales and setae; pronotum subcylindrical, lacking tubercles; scutellum visible; elytra not fused along interelytral suture, lacking series of declivital tubercles; female elytral apex produced; tibiae with spurs.

Relationships. *Methypora* is the sister genus to *Rupanius*.

Species included. *Methypora postica* Pascoe, 1865 and *M. tibialis* Lea, 1911.

Geographical distribution. Australia (Oberprieler 2010).

Material examined. *Methypora postica* (BMNH).

***Philippius* Germain, 1895**

<http://species-id.net/wiki/Philippius>

Fig. 35

Philippius Germain, 1895: 314.

Type species. *Listroderes superbus* Reed, 1872 (subsequent designation by Wibmer and O'Brien 1986).

Diagnosis. Large to very large (17.5–22.8 mm); vestiture of scales with finger-like processes and setae; mandible with 3-4 setae; pronotum wider than elytra; scutellum not visible; elytra subrectangular, fused along interelytral suture; tibiae lacking spurs; tarsomeres 3 subcylindrical.

Relationships. *Philippius* is the sister genus to *Lamiarhinus*.

Species included. *Philippius superbus* (Reed, 1872).

Geographical distribution. Andean region (Subantarctic subregion), in southern Argentina and southern Chile (Kuschel 1987; Morrone 1990).

Material examined. *Philippius superbus* (IADIZA, MACN, MHNS, MLP, USNM).

***Rupanius* Morrone, 1995**

<http://species-id.net/wiki/Rupanius>

Rupanius Morrone, 1995c: 604.

Type species. *Rupanius carinatus* Morrone, 1995c.

Diagnosis. Small (5.3–6.6 mm); vestiture of seta-like scales and setae; pronotum transverse; elytra subrectangular, with carina on apical declivity, disc slightly convex, lacking anteapical tubercle.

Relationships. *Rupanius* is the sister genus to *Methypora* (Australia), and both are placed in Listroderina. In a previous analysis (Morrone 1997a) *Rupanius* was placed in the *Macrostyphlus* generic group (= *Macrostyphlina*).

Species included. *Rupanius carinatus* Morrone, 1995.

Geographical distribution. South American Transition Zone (North Andean Paramo biogeographical province), in Colombia (Morrone 1995c).

Material examined. *Rupanius carinatus* (CMNC).

***Trachodema* Blanchard, 1849**

<http://species-id.net/wiki/Trachodema>

Trachodema Blanchard, 1849: pl. 24.

Type species. *Trachodema tuberculosa* Blanchard, 1849 (by indication, monotypy).

Diagnosis. Small to very small (2.5–5.3 mm); vestiture of scales with finger-like processes and setae; scape long (surpassing posterior margin of eye when resting in scrobe); pronotum transverse.

Relationships. *Trachodema* is the sister genus to *Lamiarhinus-Philippius*.

Species included. *Trachodema paolae* Alonso-Zarazaga, 2012 and *T. tuberculosa* Blanchard, 1849.

Host plants. *Trachodema tuberculosa*: *Atriplex semibaccata* R. Br. (Chenopodiaceae) (Morrone 1992c).

Geographical distribution. Andean region (Central Chilean subregion) (Morrone 1992c).

Material examined. *Trachodema paolae* (MHNS) and *T. tuberculosa* (CMNC, CWOB, DZUP, FIML, MHNS, MZFC, USNM).

Species inquirendae

***Listroderes bicallosus* (Boheman, 1859)**

Cryptorhynchus bicallosus Boheman, 1859: 139.

Listroderes bicallosus; Wibmer and O'Brien, 1986: 113.

Distribution. Ecuador and Peru (Wibmer and O'Brien 1986).

***Listroderes mus* Germain, 1895**

Listroderes mus Germain, 1895: 102.

Distribution. Chile (Wibmer and O'Brien 1986).

Biogeographical Account

The geographical distribution of the genera analysed indicates that Listroderini are basically a Gondwanan taxon, with *Listronotus* being the only genus distributed in North America. All the subtribes have Andean representatives (especially in the Subantarctic subregion), each showing a different pattern:

- 1 Macrostyphlina: exclusively Andean, in both the Andean region (Subantarctic and Central Chilean subregions) and the South American Transition Zone.
- 2 Falklandiina: distributed in the Andean region (Subantarctic subregion) and New Zealand.
- 3 Listroderina: distributed in the Andean region (Subantarctic, Patagonian and Central Chilean subregions), the South American and Mexican Transition Zones and the Neotropical and Australian Temperate regions.
- 4 Palaechthina: distributed in the Andean (Subantarctic and Central Chilean subregions), Neotropical and Nearctic regions, the South American Transition Zone, the Tristan da Cunha-Gough islands, New Zealand and the Australian Temperate region.

By replacing the genera for the areas where they are distributed, a taxon-area cladogram was obtained (Fig. 36). The paralogy-free subtrees that can be obtained from this taxon-area cladogram are mostly uninformative, and the informative ones cannot be combined into a general area cladogram. Geographical paralogy is particularly evident in the Subantarctic subregion, where representatives of the four subtribes are represented, suggesting that Listroderini are an ancient Gondwanan group. Several extinction events might have obscured the relationships among the areas.




Figure 36. Taxon-area cladogram of the tribe Listroderini, with the geographical distribution of the subtribes represented on maps. AUS, Australia; CEN, Central Chilean subregion; MTZ, Mexican Transition Zone; NEA, Nearctic region; NEO, Neotropical region; NZ, New Zealand; PAT, Patagonian subregion; SATZ, South American Transition Zone; SUB, Subantarctic subregion; TCF, Tristan da Cunha-Gough islands.

Acknowledgements

I thank Rolf Oberprieler for stimulating discussions on weevil systematics over the years and for donating specimens of *Anorthorhinus*, Samuel Brown for communicating some omissions and corrections to my 2011 checklist, Silvia Espinosa Matías for preparing

the Scanning Electron Microscope images at the Facultad de Ciencias, UNAM, and Sergio Roig-Juñent for the habitus illustrations. Rolf Oberprieler and one anonymous reviewer provided very useful comments that helped improve the manuscript. The image of *Gromilus veneris* is used with permission of the Bishop Museum, Honolulu.

References

- Alonso-Zarazaga MA (2012) *Trachodema paolae* nom. nov. (Coleoptera, Curculionidae, Cyclo-minae) and a correction to Morrone's list of Listroderini. *Graellsia* 68: 219. doi: 10.3989/graelessia.2012.v68.065
- Alonso-Zarazaga MA, Lyal CHC (1999) A world catalogue of families and genera of Curculo-noidae (Insecta: Coleoptera) (excluding Scolytidae and Platypodidae). Entomopraxis, Barcelona.
- Anderson RS (1992) Curculionoidea of southern Florida: An annotated checklist (Coleoptera: Curculionoidea [excluding Curculionidae; Scolytinae, Platypodinae]). *Insecta Mundi* 6: 193–248.
- Anderson RS (2002) Chapter 131. Curculionidae. In: Arnett RH Jr, Thomas MC, Skelley PE, Frank JH (Eds) American Beetles, Volume II: Polyphaga: Scarabaeoidea through Curculionoidea, CRC Press, Boca Raton, 722–815.
- Anderson RS, Morrone JJ (1996) A new genus of microptalmic Rhytidrhinini from Andean leaf litter (Coleoptera: Curculionidae). *Entomologica Scandinavica* 27: 259–278. doi: 10.1163/187631296X00098
- Angell CWJ (1893) Notes on *Macrops* and *Anthonomus*. *Journal of the New York Entomological Society* 1: 12–15.
- Ashworth AC, Kuschel G (2003) Fossil weevils (Coleoptera: Curculionidae) from latitude 85°S Antarctica. *Palaeogeography, Palaeoclimatology, Palaeoecology* 191: 191–202. doi: 10.1016/S0031-0182(02)00712-5
- Blackburn T (1890) Notes on Australian Coleoptera, with description of new species. Part VII. *Proceedings of the Linnean Society of New South Wales* 5: 303–366.
- Blackburn T (1893) Further notes on Australian Coleoptera, with descriptions of new genera and species. XIV. *Transactions of the Royal Society of New South Wales* 17: 294–315.
- Blanchard E (1849) Láminas. In: Gay C (Ed) Historia física y política de Chile, vol. 5, Zoología, author, Paris, and Museo Nacional de Historia Natural de Santiago, Santiago, 32 pl.
- Blanchard E (1851) Fauna chilena. Insectos. Coleópteros. In: Gay C (Ed) Historia física y política de Chile, vol. 5, Zoología, author, Paris, and Museo Nacional de Historia Natural de Santiago, Santiago, 285–563.
- Blanchard E (1853) Insectes. Voyage au Pole Sud et dans l'Océanie sur les corvettes l'Astrolabe et la Zélée; exécutée par ordre du Roi pendant les années 1837–1838–1839–1840 sous le commandement de M. J. Dumont-d'Urville, Capitaine de vaisseau. *Zoologie*, vol. 4, pt. 1. Baudry, Paris, 422 pp.
- Blatchley WS (1916) In: Blatchley WS, Leng CW, Rhynchophora or weevils of North Eastern America, The Nature Publishing Co., Indianapolis. doi: 10.5962/bhl.title.1557

- Blatchley WS (1920) Some new Rhynchophora from eastern North America with additions to and corrections of the “Rhynchophora of Northeastern North America”. *Journal of the New York Entomological Society* 28: 161–178.
- Blatchley WS (1922) Notes on the Rhynchophora of eastern North America, with characterizations of new genera and descriptions of new species. *Journal of the New York Entomological Society* 30: 95–106.
- Blatchley WS (1925) Notes on the Rhynchophora of eastern North America with descriptions of new species, III. *Journal of the New York Entomological Society* 33: 87–113.
- Blatchley WS (1928) Notes on the Rhynchophora of eastern North America with descriptions of new species, IV. *Journal of the New York Entomological Society* 36: 235–262.
- Blodgett SL, Denke PM, Ivie MA, O'Brien CW, Lenssen AW (1997) *Listronotus montanus* Dietz (Coleoptera: Curculionidae) damaging spring wheat in Montana. *Canadian Entomologist* 129: 377–378. doi: 10.4039/Ent129377-2
- Bohemian CH (1834) In: Schoenherr CJ, Genera et species curculionidum, cum synonymia hujus familiae, Vol. 2, pt. 1, Roret, Paris, 1–326.
- Bohemian CH (1842) In: Schönherr CJ, Genera et species curculionidum cum synonymia hujus familiae, Vol. 6, pt. 2, Roret, Paris and Fleischer, Leipzig, 1–495.
- Bohemian CH (1843) In: Schoenherr CJ, Genera et species curculionidum cum synonymia hujus familiae, Vol. 7, pt. 1, Roret, Paris and Fleischer, Leipzig, 1–479.
- Bohemian CH (1859) Coleoptera. Species novas descripsit. In: Kongliga Svenska Fregatten Eugenies resa omkring Jorden under befäl af C. A. Virgin aren 1851–1853. Vetenskapliga iakttagelser Pa H. Maj: t Konung Oscar den Förstes befallning utgifna af K. Svenska Vetenskaps-Akademien. Zoologi. III. Insekter. Norstedt & Söner, Stockholm, 113–217.
- Boivin GS, Cote M, Anciso JR (1990) Egg parasitoid of a carrot weevil, *Listronotus texanus* (Stockton), in the lower Rio Grande Valley, Texas. *Journal of the Rio Grande Valley Horticultural Society* 43: 91–92.
- Bosq JM (1935) Primera lista de los coleópteros de la República Argentina dañinos a la agricultura. *Boletín del Ministerio de Agricultura de la Nación* 36: 313–346.
- Bouchard P, Bousquet Y, Davies AE, Alonso-Zarazaga MA, Lawrence JF, Lyal CHC, Newton AF, Reid CAM, Schmitt M, Ślipiński SA, Smith ABT (2011) Family-group names in Coleoptera (Insecta). *Zookeys* 88: 1–972. doi: 10.3897/zookeys.88.807
- Brèthes J (1910) Coleópteros argentinos y bolivianos. *Anales de la Sociedad Científica Argentina* 69: 205–227.
- Brèthes J (1926) Descripción de un gorgojo que ataca la zanahoria: “*Aulametopius dauci*”, n. gen., n. sp. *Physis* 8: 414–416.
- Brinck P (1948) Coleoptera of Tristan da Cunha. *Results of the Norwegian Scientific Expedition to Tristan da Cunha, 1937–1938* 17: 1–121.
- Brookes AE (1951) The Coleoptera of the Auckland and Campbell Islands. *Cape Expedition Series, Bulletin* 5: 1–68.
- Broun T (1885) Abstract of paper on New Zealand Scydmaenidae. *New Zealand Journal of Science*, Dunedin 2: 384–387.
- Broun T (1893a) Manual of the New Zealand Coleoptera. Parts V, VI, VII, vol. 5. Wellington.

- Broun T (1893b) Descriptions of new Coleoptera from New Zealand. *The Annals and Magazine of Natural History* 6: 288–302. doi: 10.1080/00222939308677625
- Broun T (1903) Descriptions of new genera and species of New Zealand Coleoptera. *The Annals and Magazine of Natural History* 12: 69–86. doi: 10.1080/00222930308678830
- Broun T (1904) Descriptions of new genera and species of New Zealand Coleoptera. *The Annals and Magazine of Natural History* 14: 105–127. doi: 10.1080/03745480409442977
- Broun T (1905) Descriptions of a new genus and four species of New Zealand Coleoptera. *The Annals and Magazine of Natural History* 15: 543–547. doi: 10.1080/03745480509442849
- Broun T (1909) Descriptions of new genera and species of New Zealand Coleoptera. *The Annals and Magazine of Natural History* 4: 51–71. doi: 10.1080/00222930908692639
- Broun T (1910) Descriptions of new genera and species of Coleoptera. *New Zealand Institute Bulletin* 1: 1–78.
- Broun T (1913) Descriptions of new genera and species of New Zealand Coleoptera. *Transactions and Proceedings of the New Zealand Institute* 45: 97–164.
- Broun T (1914) Descriptions of new genera and species of Coleoptera (Part II). *New Zealand Institute Bulletin* 1: 79–142.
- Broun T (1915) Descriptions of new genera and species of Coleoptera (Part IV). *New Zealand Institute Bulletin* 1: 267–346.
- Broun T (1917) Descriptions of new genera and species of Coleoptera (Part V). *New Zealand Institute Bulletin* 1: 347–474.
- Broun T (1921) Descriptions of new genera and species of Coleoptera (Part VI). *New Zealand Institute Bulletin* 1: 475–590.
- Broun T (1923) Descriptions of new genera and species of Coleoptera. *New Zealand Institute Bulletin* 1: 1–708.
- Burke HR (1963) New species of Texas weevils, with notes on others (Coleoptera: Curculionidae). *The Southwestern Naturalist* 8: 162–172. doi: 10.2307/3669209
- Casey TL (1895) Coleopterological notices. VI. *Annals of the New York Academy of Sciences* 8: 438–838.
- Champion GC (1902) *Biologia Centrali-Americanana. Insecta. Coleoptera. Rhynchophora. Curculionidae. Curculioninae (part)*, vol. 4, pt. 4, 1–144.
- Champion GC (1918a) The Coleoptera of the Falkland Islands. *The Annals and Magazine of Natural History*, series 9, 1: 167–186.
- Champion GC (1918b) Notes on various South American Coleoptera collected by Charles Darwin during the voyage of the “Beagle”, with descriptions of new genera and species. *Entomological Monthly Magazine* 54: 43–55.
- Cockerell TDA (1906) Preoccupied generic names of Coleoptera. *Entomological News* 17: 240–244.
- Cordo HA, DeLoach CJ (1982) Weevils *Listronotus marginicollis* and *L. cinnamoneus* that feed on *Limnobium* and *Myriophyllum* in Argentina. *The Coleopterists Bulletin* 36: 302–308.
- Cordo HA, DeLoach CJ, Ferrer R (1982) The weevils *Lixellus*, *Tanysphiroideus*, and *Cyrtobagous* that feed on *Hydrocotyle* and *Salvinia* in Argentina. *The Coleopterists Bulletin* 36: 279–286.

- Cragnolini CI (1994) Observaciones sobre el ciclo biológico y comportamiento de *Listronotus dauci* (Brethes) (Coleoptera-Curculionidae) en cultivos de zanahoria en Córdoba (Argentina). *Agriscentia* 11: 83–86.
- Desbrochers des Loges J (1898) Description d'un Curculionide nouveau de France constituant une coupe generique nouvelle de la tribu des Erirhinides. *Le Frelon* 7: 52–54.
- Dietz WG (1889) On the species of *Macrops* Kirby, inhabiting North America. *Transactions of the American Entomological Society* 16: 28–54.
- Edelson JV (1985) Biology of a carrot weevil, *Listronotus texanus* (Coleoptera: Curculionidae): Range and seasonality of infestations. *Journal of Economic Entomology* 78: 895–897.
- Edwards MA, Hopwood AT (Eds) (1966) *Nomenclator Zoologicus*. The Zoological Society of London, London.
- Enderlein G (1907) Die Rüsselkäfer der Falklands-Inseln. 13. Beitrag zur Kenntnis der antarktischen Fauna. *Stettiner Entomologische Zeitung* 68: 36–69.
- Enderlein G (1912) Die Insekten des Antarkto-Archiplato-Gebietes (Feuerland, Falklands-Inseln, Süd-Georgien). 20. Beitrag zur Kenntnis der antarktischen Fauna. *Kungliga Svenska Vetenskapsakademiens Handlingar* 48: 1–170.
- Erichson WF (1842) Beitrag zur Insecten-Fauna von Vandiemensland, mit besonderer Berücksichtigung der geographischen Verbreitung der Insecten. *Archiv für Naturgeschichte* 8: 83–287.
- Erichson WF (1847) Conspectus Insectorum Coleopterorum, quae in Republica Peruana observata sunt. *Archiv für Naturgeschichte* 13: 67–185.
- Fairmaire L (1884) Note sur quelques Coléoptères de Magellan et de Santa-Cruz. *Annales de la Société Entomologique de France* 1883 (1884), ser. 6: 483–506.
- Fairmaire L (1885) Liste de coléoptères recueillis à la Terre de Feu par la mission de la Romanche et description des espèces nouvelles. *Annales de la Société Entomologique de France* 6: 33–62.
- Friedman ALL (2009) The vegetable weevil, *Listroderes costirostris* Schoenherr (Curculionidae: Cyclominae): A new invasive pest in Israel. *Phytoparasitica* 37: 331–332. doi: 10.1007/s12600-009-0039-2
- Germain P (1895–1896) Apuntes sobre los insectos de Chile. Estudio i descripción de los Listroderitos de Chile i tierras magallánicas de la colección del Museo Nacional i de la de Don Fernando Paulsen. *Anales de la Universidad de Chile* 90: 287–324, 467–505, 567–602, 91: 53–104 (1895); 93: 791–838, 94: 721–752 (1896).
- Germain P (1911) Informes de los jefes de Sección i otros empleados del Museo. 1-Informe del jefe de la Sección de Entomología. *Boletín del Museo Nacional de Chile* 3: 197–221.
- Goloboff P (1993) Estimating character weights during tree search. *Cladistics* 9: 83–91. doi: 10.1111/j.1096-0031.1993.tb00209.x
- Goloboff PA, Farris JS, Nixon KC (2008) TNT, a free program for phylogenetic analysis. *Cladistics* 24: 774–786. doi: 10.1111/j.1096-0031.2008.00217.x
- Guérin-Ménéville FE (1839) Description de quelques Coléoptères des côtes du Détroit de Magellan. *Revue de Zoologie* 2: 295–305.
- Gyllenhal L (1834) In: Schoenherr CJ, Genera et species curculionidum, cum synonymia hujus familiae, Vol. 2, pt. 1, Roret, Paris, 1–326.

- Hatch MH (1971) The beetles of the Pacific Northwest. University of Washington Publications in Biology 16: 1–662.
- Henderson LS (1940) A revision of the genus *Listronotus*. University of Kansas Science Bulletin 26: 215–337.
- Hustache A (1921) Deux Curculionides américains introduits en France. Bulletin de la Société Entomologique de France, 1921: 134.
- Hustache A (1926) Contribution à l'étude des Curculionides de la République Argentine (première note). Anales del Museo Nacional de Historia Natural "Bernardino Rivadavia" 34: 155–261.
- Hustache A (1938a) Curculionides nouveaux de l'Amérique méridionale, qui se trouvent dans le Deutsches Entomologisches Institut. (1ère note). Arbeiten über morphologische und taxonomische Entomologie aus Berlin-Dahlem 5: 174–184.
- Hustache A (1938b) Curculionides nouveaux de l'Amérique méridionale, qui se trouvent dans le Deutsches Entomologisches Institut. 2ième note. Arbeiten über morphologische und taxonomische Entomologie aus Berlin-Dahlem 5: 265–288.
- Hustache A (1939a) Curculionides nouveaux de l'Argentine et autres régions Sud-Américaines. Anales de la Sociedad Científica Argentina 128: 38–64, 99–124.
- Hustache A (1939b) Curculionides nouveaux de l'Amérique méridionale, qui se trouvent dans le Deutsches Entomologisches Institut. Suite de la 3ième note. Arbeiten über morphologische und taxonomische Entomologie aus Berlin-Dahlem 6: 50–68.
- Jekel H (1865) Recherches sur la classification naturelle des Curculionides. Annales de la Société Entomologique de France, 1864 [1865], ser. 4: 537–566.
- Kirby W (1837) Part the fourth and last. The insects. In: Richardson J, Fauna Boreali-Americanica; or the zoology of the northern parts of British America: containing descriptions of the objects of natural history collected on the late Northern Land Expeditions, under command of Captain Sir John Franklin, R.N., Fletcher, Norwich, 1–325.
- Kirby W (1885) Coleoptera. In: Rye EC (Ed) The Zoological Record for 1882; being the volume twentieth of the record of zoological literature, 16–123.
- Kirsch T (1877) In: Kiesenwetter H von, Kirsch T, Die Käferfauna der Auckland-Inseln, nach Herm. Krone's Sammlungen beschrieben. Deutsche Entomologische Zeitschrift 21: 153–174.
- Kirsch T (1889) Coleopteren gesammelt in den Jahren 1868–1877 auf einer Reise durch Süd Amerika von Alphons Stübel. Abhandlungen und Berichte des königlichen zoologischen und anthropologisch-ethnographischen Museum zu Dresden, 1888/89 (1889), 4: 1–58.
- [Klug F] (1829) Preis-Verzeichniss vorrätiger Insectendoubletten des Königl. Zoologischen Museums der Universität, Berlin.
- Kolbe HJ (1907) Coleopteren. Ergebnisse der Hamburger Magalhaensische Sammelreise 8: 1–125.
- Kuschel G (1949) Los "Curculionidae" del extremo norte de Chile (Coleoptera, Curcul. ap. 6). Acta Zoológica Lilloana 8: 5–54.
- Kuschel G (1950) Nuevas sinonimias, revalidaciones y combinaciones (9º aporte a Col. Curculionidae). Agricultura Técnica de Chile 10: 10–21.

- Kuschel G (1952) Cylydrorhininae aus dem Britischen Museum. (Col. Curculionidae, 8. Beitr.). The Annals and Magazine of Natural History, ser. 12, 5: 121–137. doi: 10.1080/00222935208654274
- Kuschel G (1955) Nuevas sinonimias y anotaciones sobre Curculionoidea (1) (Coleoptera). Revista Chilena de Entomología 4: 261–312.
- Kuschel G (1958) Nuevos Cylydrorhininae de la Patagonia (Col. Curculionoidea, Aporte 18). Investigaciones Zoológicas Chilenas 4: 231–252.
- Kuschel G (1962) The Curculionidae of Gough Island and the relationships of the weevil fauna of the Tristan da Cunha Group. Proceedings of the Linnean Society of London, 1960-61 (1962), 173: 69–78.
- Kuschel G (1964) Insects of Campbell Island. Coleoptera: Curculionidae of the Subantarctic islands of New Zealand. Pacific Insects Monographs 7: 416–493.
- Kuschel G (1971) Entomology of the Aucklands and other islands south of New Zealand: Coleoptera: Curculionidae. Pacific Insects Monographs 27: 225–259.
- Kuschel G (1972) The foreign Curculionoidea established in New Zealand (Insecta: Coleoptera). New Zealand Journal of Science 15: 273–289.
- Kuschel G (1986) In: Wibmer GJ, O'Brien CW, Annotated checklist of the weevils (Curculionidae sensu lato) of South America (Coleoptera: Curculionoidea), Memoirs of the American Entomological Institute 39: 1–563.
- Kuschel G (1987) The subfamily Molytinae (Coleoptera: Curculionidae): General notes and descriptions of new taxa from New Zealand and Chile. New Zealand Entomologist 9: 11–29. doi: 10.1080/00779962.1987.9722488
- Kuschel G (1990) Beetles in a suburban environment: A New Zealand case study. New Zealand Department of Scientific and Industrial Research, Auckland.
- Lanteri AA, Marvaldi AE, Suárez S (2002) Gorgojos de la Argentina y sus plantas huéspedes: Tomo I: Apionidae y Curculionidae. Publicación Especial de la Sociedad Entomológica Argentina, nro. 1, San Miguel de Tucumán.
- Lea AM (1899) Descriptions of Australian Curculionidae, with notes on previously described species. Transactions and Proceedings of the Royal Society of South Australia [23]: 137–197.
- Lea AM (1911) Notes on Australian Curculionidae in the Berlin Museum with descriptions of new species. Mitteilungen aus dem Zoologischen Museum in Berlin 5: 177–201.
- Lea AM (1928) New species of Australian Erirhinides (Curculionidae). Proceedings of the Linnean Society of New South Wales 53: 375–396.
- LeConte JL (1857) Report upon insects collected on the survey. In: Reports of explorations and surveys for a railroad route from the Mississippi River to the Pacific Ocean, Vol. 9, no. 1, 1–72.
- LeConte JL (1876) In: LeConte JL, Horn GH, The Rhynchophora of America, north of Mexico. Proceedings of the American Philosophical Society 15: 1–455.
- López G, Hermann M (Eds) (2004) El cultivo del ulluco en la sierra central del Perú. Centro Internacional de la Papa, Lima.
- Maes JM, O'Brien CW (1990) Lista anotada de los Curculionoidea (Coleoptera) de Nicaragua. Revista Nicaragüense de Entomología 12: 1–78.

- Marshall GAK (1914) Four new injurious weevils from Africa. *Bulletin of Entomological Research* 5: 235–239.
- Marshall GAK (1926) On new Neotropical Curculionidae (Col.). *The Annals and Magazine of Natural History* 18: 530–543. doi: 10.1080/00222932608633549
- Marshall GAK (1930) New Curculionidae, with notes on synonymy. *The Annals and Magazine of Natural History* 9: 551–577. doi: 10.1080/00222933008673252
- Marshall GAK (1935) New Curculionidae (Col.) from tropical Africa, with notes on synonymy etc. *The Annals and Magazine of Natural History*, series 10: 497–518.
- Marshall GAK (1937) New Curculionidae (Col.) from New Zealand. *Transactions of the Royal Society of New Zealand* 67: 316–340.
- Marshall GAK (1943) New Indian Curculionidae (Col.). *The Annals and Magazine of Natural History* 11: 105–119. doi: 10.1080/03745481.1943.9727999
- Marshall GAK (1953) Four new Curculionidae (Col.) from New Zealand. *Transactions of the Royal Society of New Zealand* 81: 67–70.
- Martel P, Svec HJ, Harris CR (1976) The life history of the carrot weevil, *Listronotus oregonensis* (Coleoptera: Curculionidae) under controlled conditions. *Canadian Entomologist* 108: 931–934. doi: 10.4039/Ent108931-9
- Marvaldi AE (1994) Estudio taxonómico del género *Neopachytychius* Hustache (Coleoptera: Curculionidae). *Revista de la Sociedad Entomológica Argentina* 53: 57–63.
- Marvaldi AE (1998) Larvae of South American Rhytirrhininae (Coleoptera: Curculionidae). *Coleopterists Bulletin* 52: 71–89.
- May BM (1970) Aquatic adaptation in the larva of *Desiantha ascita* (Coleoptera: Curculionidae). *New Zealand Entomologist* 4: 7–11. doi: 10.1080/00779962.1970.9723066
- May BM (1971) Entomology of the Aucklands and other islands south of New Zealand: Immature stages of Curculionoidea. *Pacific Insects Monographs* 27: 271–316.
- May BM (1977) Immature stages of Curculionidae: Larvae of the soil-dwelling weevils of New Zealand. *Journal of the Royal Society of New Zealand* 7: 189–228. doi: 10.1080/03036758.1977.10427160
- May BM (1993) Larvae of Curculionoidea (Insecta: Coleoptera): A systematic overview. *Fauna of New Zealand* 28: 1–221.
- May BM (1994) An introduction to the immature stages of Australian Curculionoidea. In: Zimmerman, E. C. *Australian weevils. Volume II. Brentidae, Eurhynchidae, Apionidae and a chapter on immature stages by Brenda May*. CSIRO, Melbourne.
- Morrone JJ (1990) *Philippius* Germain, a remarkable Listroderini from southern South America (Coleoptera: Curculionidae). *The Coleopterists Bulletin* 44: 429–436.
- Morrone JJ (1992a) Revisión sistemática y análisis cladístico del género *Antarctobius* Fairmaire (Coleoptera: Curculionidae). *Neotropica* 38: 3–20.
- Morrone JJ (1992b) Revisión sistemática, análisis cladístico y biogeografía histórica de los géneros *Falklandius* Enderlein y *Lanteriella* gen. nov. (Coleoptera: Curculionidae). *Acta Entomológica Chilena* 17: 157–174.
- Morrone JJ (1992c) Revision of *Trachodema* Blanchard with the description of an allied genus from central Chile (Insecta, Coleoptera, Curculionidae). *Zoologica Scripta* 21: 417–422. doi: 10.1111/j.1463-6409.1992.tb00341.x

- Morrone JJ (1993a) Revisión sistemática de un nuevo género de Rhytirrhinini (Coleoptera: Curculionidae), con un análisis biogeográfico del dominio Subantártico. Boletín de la Sociedad de Biología de Concepción 64: 121–145.
- Morrone JJ (1993b) Revisión sistemática del género *Hyperoides* Marshall (Coleoptera: Curculionidae). Neotropica 39: 17–26.
- Morrone JJ (1993c) Revisión de las especies de *Listroderes* Schoenherr del grupo *curvipes* (Coleoptera: Curculionidae). Revista Chilena de Entomología 20: 15–21.
- Morrone JJ (1993d) Systematic revision of the *costirostris* species group of the weevil genus *Listroderes* Schoenherr (Coleoptera: Curculionidae). Transactions of the American Entomological Society 119: 271–315.
- Morrone JJ (1993e) Revisión de las especies de *Listroderes* Schoenherr del grupo *nodifer* (Coleoptera: Curculionidae). Boletín del Museo Nacional de Historia Natural, Chile, 1992 [1993], 43: 117–130.
- Morrone JJ (1993f) Cladistic and biogeographic analyses of the weevil genus *Listroderes* Schoenherr (Coleoptera: Curculionidae). Cladistics 9: 397–411. doi: 10.1111/j.1096-0031.1993.tb00233.x
- Morrone JJ (1994a) Systematics of the Andean genus *Acrorius* Kirsch (Coleoptera: Curculionidae). The Coleopterists Bulletin 48: 101–114.
- Morrone JJ (1994b) Systematics of the Patagonian genus *Acrostomus* Kuschel (Coleoptera: Curculionidae). Annals of the Entomological Society of America 87: 403–411.
- Morrone JJ (1994c) Systematics, cladistics, and biogeography of the Andean weevil genera *Macrostyphlus*, *Adioristidius*, *Puranius*, and *Amathynetoides*, new genus (Coleoptera: Curculionidae). American Museum Novitates 3104: 1–63.
- Morrone JJ (1994d) Cladistic placement of the Subantarctic genus *Haversiella* (Coleoptera: Curculionidae). Journal of the New York Entomological Society 102: 299–302.
- Morrone JJ (1994e) Clarification of the taxonomic status of species formerly placed in *Listroderes* Schoenherr (Coleoptera: Curculionidae), with the description of a new genus. American Museum Novitates 3093: 1–11.
- Morrone JJ (1995a) Estudio taxonómico y biogeográfico del género subantártico *Falklandielus* Kuschel (Coleoptera: Curculionidae). Physis (Buenos Aires), Sección C, 1992 (1995), 50: 105–110.
- Morrone JJ (1995b) Revisión de las especies de *Listroderes* Schoenherr del grupo *robustus* (Coleoptera: Curculionidae). Physis (Buenos Aires), Sección C, 1992 (1995), 50: 73–80.
- Morrone JJ (1995c) A new genus of Rhytirrhinini from Colombia (Coleoptera: Curculionidae). Caldasia 17: 603–606.
- Morrone JJ (1997a) Cladistics of the New World genera of Listroderina (Coleoptera: Curculionidae: Rhytirrhinini). Cladistics 13: 247–266. doi: 10.1111/j.1096-0031.1997.tb00318.x
- Morrone JJ (1997b) Nomenclatural notes on the subfamily Cyclominae (Coleoptera: Curculionidae). Acta Entomológica Chilena 21: 101–102.
- Morrone JJ (2002a) Checklist of the species of Cyclominae (Coleoptera: Curculionidae) occurring in America south of the United States. Revista de la Sociedad Entomológica Argentina 61: 1–8.

- Morrone JJ (2002b) On the species of the *Listroderes costirostris* complex (Coleoptera: Curculionidae). *Neotropica* 48: 70–72.
- Morrone JJ (2006) Biogeographic areas and transition zones of Latin America and the Caribbean Islands based on panbiogeographic and cladistic analyses of the entomofauna. *Annual Review of Entomology* 51: 467–494. doi: 10.1146/annurev.ento.50.071803.130447
- Morrone JJ (2011) Annotated checklist of the tribe Listroderini (Coleoptera: Curculionidae: Cyclominae). *Zootaxa* 3119: 1–68.
- Morrone JJ, Anderson RS (1995) The *Falklandius* generic group: Cladistic analysis with description of new taxa (Coleoptera: Curculionidae: Rhytirrhinini). *American Museum Novitates* 3121: 1–14.
- Morrone JJ, Díaz NB, Loiácono MS (1992) Comparative morphology of mouthparts in the tribe Listroderini (Coleoptera: Curculionidae). *Elytron* 6: 47–59.
- Morrone JJ, Marvaldi AE (1998) *Listroderes abditus* or *Antarctobius abditus*? A simultaneous analysis of larval and adult characters (Coleoptera: Curculionidae). *European Journal of Entomology* 95: 229–236.
- Morrone JJ, Marvaldi AE, O'Brien CW (1995) *Lixellus* LeConte, a new synonym of *Listronotus* Jekel (Coleoptera: Curculionidae). *Entomological News* 106: 108–112.
- Morrone JJ, O'Brien CW (2000) The aquatic and semiaquatic weevils (Coleoptera: Curculionoidea: Curculionidae, Dryophthoridae and Eriphiniidae) of Argentina, with indication of their host plants. *Physis* (Buenos Aires) secc. C, 57: 25–37.
- Morrone JJ, Ocampo F (1995) *Acroriellus*, a new weevil genus of northern Andean Listroderina (Coleoptera: Curculionidae). *Zoologica Scripta* 24: 257–262. doi: 10.1111/j.1463-6409.1995.tb00403.x
- Oberprieler RG (1992) Preliminary report on the weevils (Curculionidae) collected on Inaccesible and Nightingale islands during September 1989 by the expedition of the Fitzpatrick Institute for African Ornithology, University of Cape Town, submitted by Mr. A. J. Gardiner, Zimbabwe. Manuscript distributed by the author, 1–2.
- Oberprieler RG (2010) A reclassification of the weevil subfamily Cyclominae (Coleoptera: Curculionidae). *Zootaxa* 2515: 1–35.
- Oberprieler RG (In press) Cyclominae Schoenherr, 1826. In: Beutel RG, Leschen RAB (Eds) *Handbook of Coleoptera*, Vol. 3, deGruyter Press, Berlin.
- O'Brien CW (1977) The semiaquatic weevil genus *Listronotus* in Mexico and Central America (Coleoptera: Curculionidae: Cylindrorhininae). *Transactions of the Entomological Society of America* 70: 804–814.
- O'Brien CW (1981) The larger (4.5+ mm) *Listronotus* of America, north of Mexico (Cylindrorhininae, Curculionidae, Coleoptera). *Transactions of the American Entomological Society* 107: 69–123.
- O'Brien CW, Wibmer GJ (1982) Annotated checklist of the weevils (Curculionidae sensu lato) of North America, Central America, and the West Indies (Coleoptera: Curculionoidea). *Memoirs of the American Entomological Institute* 34: 1–382.
- Ocampo F, Morrone JJ (1996) Two new species and cladistic analysis of *Acrorius* Kirsch (Coleoptera: Curculionidae: Rhytirrhinini). *Neotropica* 42: 23–28.

- Olliff AS (1891) Coleoptera (continued). In: Whymper, E., Supplementary appendix to travels amongst the Great Andes of the Equator. Murray, London, 58–81.
- Pascoe FP (1865) On some new genera of Curculionidae. Part I. Journal of Entomology, London 2: 413–432.
- Pascoe FP (1870) Descriptions of some genera and species of Australian Curculionidae. Transactions of the Entomological Society of London 2: 181–209.
- Pascoe FP (1873) Additions to the Australian Curculionidae. The Annals and Magazine of Natural History 11: 178–199. doi: 10.1080/00222937308696794
- Pascoe FP (1876) Descriptions of new genera and species of New Zealand Coleoptera. The Annals and Magazine of Natural History 18: 57–67. doi: 10.1080/00222937608682007
- Posadas P (2008) A preliminar overview of species composition and geographical distribution of Malvinian weevils (Insecta: Coleoptera: Curculionidae). Zootaxa 1704: 1–26.
- Posadas P (2012) Species composition and geographic distribution of Fuegian Curculionidae (Coleoptera: Curculionoidea). Zootaxa 3303: 1–36.
- Posadas P, Morrone JJ (2004) A new species of *Antarctobius* Fairmaire from Islas Malvinas (Coleoptera: Curculionidae: Cyclominae). Insect Systematics and Evolution 35: 353–359. doi: 10.1163/187631204788920194
- Reed EC (1872) (Apéndice) Parte zoológica. In: Vidal Gormaz F (Ed.) Exploración de la costa de Llanquihue, practicada por órden del Supremo Gobierno por el capitán de corbeta don Francisco Vidal Gormaz. Anales de la Universidad de Chile 41: 354–355.
- Rothwell NL (2003) Investigation into *Listronotus maculicollis* (Coleoptera: Curculionidae), a pest of highly maintained turfgrass. Electronic Doctoral Dissertations for UMass Amherst. Paper AAI3096313. <http://scholarworks.umass.edu/dissertations/AAI3096313>
- Say T (1824) Descriptions of Coleopterous insects collected in the late expedition to the Rocky Mountains, performed by order of Mr. Calhoun, Secretary of War, under the command of Mayor Long. Journal of the Academy of Natural Sciences of Philadelphia, 1823 (1824), 3: 298–331.
- Say T (1831) Descriptions of new species of Curculionites of North America, with observations on some of the species already known. New Harmony, 1–30.
- Schenkling S, Marshall GAK (1929) Coleopterorum Catalogus. Pars 106: Curculionidae: Byrsopinae, Rhytirrhininae, Thecesterninae, Hipporhininae, Rhyparosominae. Junk, s'Gravenhage.
- Schenkling S, Marshall GAK (1931) Coleopterorum Catalogus. Pars 114, Curculionidae: Cy-lindrorrhininae, pp. 1–23. Junk, Berlin.
- Schönherr CJ (1823) Curculionides. Isis Oken, heft X, columns 1132–1146; heft V, columns 581–588.
- Schönherr CJ (1826) Curculionidum dispositio methodica cum generum characteribus, descriptionibus atque observationibus variis, seu prodromus ad synonymiae insectorum. Partem 4. Fleischer, Leipzig. doi: 10.5962/bhl.title.9327
- Schweiger H (1959) Über einige von der Skottsbergexpedition im Antarkto-Archiplata-Gebiet aufgesammelte Koleopteren. Arkiv för Zoologi 12: 1–43.
- Sharp D (1883) Some new species and genera of Coleoptera from New Zealand. Entomologist's Monthly Magazine 20: 66–68.

- Sharp D (1890) Insecta. In: *Zoological Record for 1889*, Vol. 26, London, 1–320.
- Sharp D (1892) Insecta. In: *Zoological Record for 1891*, Vol. 28, London, 1–311.
- Stierlin [G] (1903) Beschreibung einiger neuen europäischen Rüsselkäfer. *Mittheilungen der Schweizerischen Entomologischen Gessellschaft Schaffhausen* 11: 56–57.
- Stockton WD (1963) New species of *Hyperodes* Jekel and a key to the Nearctic species of the genus (Coleoptera: Curculionidae). *Bulletin of the California Academy of Sciences* 62: 140–149.
- Suffrian E (1871) Verzeichniss der von Dr. Gundlach auf der Insel Cuba gesammelten Rüsselkäfer (Fortsetzung, Siehe Jahreg. XXXVI. S. 150). *Archiv für Naturgeschichte* 37: 122–184.
- Torres AN, Casey H (2002) Sample scheme for carrot weevil (Coleoptera: Curculionidae) in parsley. *Environmental Entomology* 31: 1251–1258. doi: 10.1603/0046-225X-31.6.1251
- Thompson RT (1992) Observations on the morphology and classification of weevils (Coleoptera, Curculionoidea) with a key to major groups. *Journal of Natural History* 26: 835–891. doi: 10.1080/00222939200770511
- Van Dyke EC (1929) Two n. sp. *Listronotus*. *Pan-Pacific Entomologist* 5: 106–108.
- Voss E (1943) Einige neue Rüsselarten aus Argentinien und Paraguay (Coleoptera: Curculionidae.) (106. Beitrag zur Kenntnis der Curculioniden.). *Arbeiten über morphologische und taxonomische Entomologie aus Berlin-Dahlem* 10: 225–236.
- Voss E (1954) Curculionidae (Col.). *Beiträge zur Fauna Perus* 4: 193–376.
- Waterhouse CO (1884) Coleoptera collected during the Expedition of H.M.S. ‘Challenger’. *The Annals and Magazine of Natural History* 13: 276–283. doi: 10.1080/00222938409459235
- Waterhouse GR (1841) (Descriptions of numerous species of coleopterous insects from the southern parts of South America). *Proceedings of the Zoological Society of London* 9: 105–128.
- Wibmer GJ, O’Brien CW (1986) Annotated checklist of the weevils (Curculionidae *sensu lato*) of South America (Coleoptera: Curculionoidea). *Memoirs of the American Entomological Institute* 39: 1–563.
- Wild CH, McFadyen RE, Tomley AJ, Wilson BW (1992) The biology and host specificity of the stem-boring weevil *Listronotus setosipennis* [Col.: Curculionidae]: A potential bio-control agent for *Parthenium hysterophorus* [Asteraceae]. *Entomophaga* 37: 591–598. doi: 10.1007/BF02372329
- Zimmerman EC (1994) Australian weevils. Volume I. Anthribidae to Attelabidae. Centre of Scientific and Industrial Research, Melbourne.