

## Comorbidity and Myocardial Dysfunction Are the Main Explanations for the Higher 1-Year Mortality in Acute Myocardial Infarction With Left Bundle-Branch Block

Ulf Stenestrand, MD, PhD; Fariborz Tabrizi, MD; Johan Lindbäck, MSc; Anders Englund, MD, PhD; Mårten Rosenqvist, MD, PhD; Lars Wallentin, MD, PhD

**Background**—The purpose of this study was to assess the independent contribution of left bundle-branch block (LBBB) on cause-specific 1-year mortality in a large cohort with acute myocardial infarction (MI).

**Methods and Results**—We studied a prospective cohort of 88 026 cases of MI from the Register of Information and Knowledge about Swedish Heart Intensive care Admissions in 72 hospitals in 1995 to 2001. Long-term mortality was calculated by Cox regression analysis, adjusted for multiple covariates that affect mortality by calculation of a propensity score. LBBB was present in 9% (8041 of 88 026) of the MI admissions. Patients with LBBB were older and had a higher prevalence of comorbid conditions than patients with no LBBB. The unadjusted relative risk of death within 1 year was 2.16 (95% CI, 2.08 to 2.24;  $P < 0.001$ ) for LBBB (42%, 3350 of 8041) compared with those with no LBBB (22%, 17 044 of 79 011). After adjustment for a propensity score that takes into account differences in risk factors and acute intervention, LBBB was associated with a relative risk of death of 1.19 (95% CI, 1.14 to 1.24;  $P < 0.001$ ). In a subgroup of 11 812 patients for whom left ventricular ejection fraction was available and could be added to the analysis, the contributing relative risk of LBBB for death was only 1.08 (95% CI, 0.93 to 1.25;  $P = 0.33$ ). The most common cause of death in both groups was ischemic heart disease.

**Conclusions**—MI patients with LBBB have more comorbid conditions and an increased unadjusted 1-year mortality. When adjusted for age, baseline characteristics, concomitant diseases, and left ventricular ejection fraction, LBBB does not appear to be an important independent predictor of 1-year mortality in MI. (*Circulation*. 2004;110:1896-1902.)

**Key Words:** bundle-branch block ■ mortality ■ myocardial infarction ■ patients ■ prognosis

Left bundle-branch block (LBBB) on ECG at presentation has been reported to occur in 1% to 15% of patients with myocardial infarction (MI).<sup>1-6</sup> The significance of LBBB in the setting of MI has been studied in both the thrombolytic<sup>6-10</sup> and prethrombolytic<sup>1-3,11,12</sup> eras, demonstrating an association of LBBB with increased short- and long-term mortality. Previous studies have reported a wide range of 1-year mortality in patients with MI and LBBB, varying between 19% and 61%.<sup>3,4,6-13</sup> However, although some studies suggest that LBBB is an independent predictor of short- and long-term mortality in patients with MI,<sup>6,7,9</sup> others have found no effect.<sup>8</sup> Some of these studies had small numbers of patients,<sup>1-3,5,6,8-10</sup> and others were done in the prethrombolytic era.<sup>1-5,11-13</sup> The survival benefit by reperfusion treatment in patients with MI and LBBB has been demonstrated previously.<sup>14,15</sup> The recently published Multi-center Automatic Defibrillator Implantation Trial (MADIT) II study reported the largest survival benefit with implantable

defibrillator therapy in patients with wide QRS complex (48% LBBB) in a population with previous MI and left ventricular dysfunction.<sup>16</sup> Whether this larger benefit is related to the higher risk of death or to a specific effect because of conduction abnormality per se in LBBB population has not been elucidated.

The reasons for the poor prognosis in the LBBB patients with MI and the potential for specific effects of antiarrhythmic, reperfusion, or other treatments have not been well studied. The purpose of this study was to assess the independent contribution of LBBB compared with ST-segment elevation and non-ST-segment elevation on 1-year cause-specific mortality in a large cohort with MI.

### Methods

The Register of Information and Knowledge about Swedish Heart Intensive care Admissions (RIKS-HIA) registers all patients admitted to the coronary care units of all participating hospitals. Information, including data on 100 variables, is reported on case record

Received March 23, 2004; de novo received June 1, 2004; revision received June 15, 2004; accepted June 21, 2004.

From the Department of Cardiology, University Hospital of Linköping, Linköping (U.S.); Department of Cardiology, South Hospital, Stockholm (F.T., M.R.); Uppsala Clinical Research Center, Uppsala University, Uppsala (J.L., L.W.); and Department of Cardiology, Örebro University Hospital, Örebro (A.E.), Sweden.

Correspondence to Ulf Stenestrand, MD, PhD, Department of Cardiology, University Hospital, SE 581 85 Linköping, Sweden. E-mail stenestrand@riks-hia.se

© 2004 American Heart Association, Inc.

*Circulation* is available at <http://www.circulationaha.org>

DOI: 10.1161/01.CIR.0000143136.51424.38

forms and has been described elsewhere.<sup>17</sup> Briefly, the register includes information on age, sex, smoking status, hypertension, diabetes mellitus, hyperlipidemia, previous angina pectoris, previous MI, previous coronary revascularization, previous medications (“previous” means events occurring or medication started before the current admission), symptoms, ECG at entry, biochemical markers, echocardiography, reperfusion treatment, pharmacological treatment, interventional procedures, major complications, and outcomes during the hospital stay; risk assessment with stress test, coronary angiography, and revascularization procedures; and medications at discharge. (The full protocol is available at <http://www.riks-hia.se>.) Source data were continuously verified by comparison of the registry information with the patient records of the hospital by an external monitor.

One-year mortality data were obtained by merging the RIKS-HIA database with the National Cause of Death Register, which includes the vital status of all Swedish citizens in 1995 through 2002. Previous history of stroke, dementia, renal failure, congestive heart failure, chronic pulmonary disease, and cancer were obtained by merging with the National Patient Register, which includes diagnoses on all patients hospitalized in Sweden from 1987 and forward. All patients for whom data were entered into RIKS-HIA were informed of their participation in the registry (patients could request to be excluded) and the long-term follow-up. The registry and the merging with registries were approved by the National Board of Health and Welfare and the Swedish Data Inspection Board.

We included all patients admitted with ST-segment elevation, non-ST-segment elevation, or LBBB on the ECG at entry and with acute MI as the final diagnosis. LBBB was diagnosed according to standard definitions.<sup>18</sup> Patients with agonal or pacemaker-induced QRS complexes were excluded. The criteria for the diagnosis of acute MI were standardized and identical for all participating hospitals through the use of the World Health Organization criteria.<sup>19</sup> The biochemical criterion was at least 1 measurement indicating twice the upper limit of normal of an appropriate biochemical marker (usually creatine kinase [CK]-MB protein concentration or troponin-T). In the analyses, biochemical markers were divided into 4 groups: (1) CK-MB  $\leq 50$   $\mu\text{g/L}$  or troponin-T  $< 1.0$   $\mu\text{g/L}$ , (2) CK-MB 51 to 150  $\mu\text{g/L}$  or troponin-T 1.0 to 3.0  $\mu\text{g/L}$ , (3) CK-MB  $> 150$   $\mu\text{g/L}$  or troponin-T  $> 3.0$   $\mu\text{g/L}$ , and (4) missing values. ECG was classified according to 3 variables: rhythm ([1] sinus, [2] atrial fibrillation or flutter, [3] pacemaker rhythm, [4] other rhythm); QRS complex ([1] normal, [2] LBBB/pacemaker complex, [3] new Q wave, [4] old Q wave, [5] other pathological QRS complex); and STT segment ([1] normal, [2] LBBB/pacemaker complex, [3] ST-segment elevation, [4] ST-segment depression, [5] pathological T wave, and [6] other pathological STT segment). Starting in a hierarchical order from alternative 1 in each of the 3 variables, the first applicable alternative that was correct for that part of the ECG was the coding for it. Left ventricular ejection fraction (LVEF) was determined by echocardiography before discharge and coded into RIKS-HIA in 6 groups: (1) not performed, (2) performed (no information about LVEF), (3) normal (LVEF  $> 50\%$ ), (4) light dysfunction (LVEF 40% to 50%), (5) moderate dysfunction (LVEF 30% to 39%), and (6) severe dysfunction (LVEF  $< 30\%$ ). Before 2001, LVEF was not part of the mandatory protocol, and echocardiography was registered only as performed or not performed. Hence, analyses including LVEF were available in only a minority of patients.

### Statistical Analysis

Different patient strata were compared by  $\chi^2$  tests for categorical variables and by the *t* test for continuous variables. A propensity score was calculated for each patient to estimate the probability of being an LBBB patient (compared with a patient with no LBBB) given the background characteristics and treatments received. Adjusting for the propensity score in the analysis was aimed at giving a better estimate of the relative risk (RR) of death between groups. Multiple logistic regression analysis was performed to calculate the first propensity score (A), including 25 covariates: age, sex, previous MI, history of diabetes mellitus, history of stroke, congestive heart failure, renal failure, chronic pulmonary disease, dementia, cancer

within 3 years, history of hypertension, history of coronary artery revascularization, medications before study entry (including ACE inhibitors or angiotensin II receptor blockers, anticoagulants,  $\beta$ -blockers, aspirin, or platelet inhibitors, calcium channel blockers, digitalis, diuretics, lipid-lowering drugs, and long-acting nitrates), circulatory arrest at arrival, presence of atrial fibrillation, signs of congestive heart failure, and reperfusion therapy. Logistic regression analyses were used to establish the relation between LBBB and in-hospital complications. Cox proportional-hazards regression analyses including the propensity scores were used to identify whether LBBB by itself significantly influenced mortality.

To evaluate the influence of in-hospital therapy, coronary interventions, and discharge medication, a Cox proportional-hazards regression analysis was also performed concerning the same outcome in survivors at discharge and including a second propensity score. The second propensity score (B) of the likelihood of LBBB on ECG included the previous 25 covariates and an additional 13 covariates—heparin therapy, intravenous nitroglycerin, intravenous  $\beta$ -blockers, coronary revascularization within 14 days, discharge medication (including ACE inhibitor, oral anticoagulants, aspirin or platelet inhibitors,  $\beta$ -blockers, calcium channel blockers, digitalis, diuretics, lipid-lowering drugs, and long-acting nitrates)—for a total of 38 covariates. In some Cox proportional-hazards regression analyses, the ordinal variable, biochemical marker peak level group, and the scale variable, Killip class, were included. In a subgroup for which LVEF was available, the previously mentioned 4 groups of LVEF were added as a scale variable to the Cox proportional-hazards regression analysis of 1-year mortality, also including the 2 variables of propensity score (A) and presence of LBBB on ECG. Statistical analyses were performed with the statistical program R<sup>20</sup> and SPSS version 11.5 software (SPSS Inc).

## Results

### Data Validity

When 1972 computer forms from 38 hospitals containing 161 280 variables were reviewed specially by an external monitor, there was 97% agreement between the registered information and the source data in the patient records among the 25 covariates included in logistic regression analysis A and 95% agreement for the 38 variables in analysis B. With regard to the ECG variable, in 0.6% of the cases, LBBB had been miscoded as another type of QRS complex, and in 2.2%, a QRS complex that really was not LBBB had been incorrectly coded as LBBB.

### Patient Material and Baseline Characteristics

Nineteen hospitals participated in the registry in 1995, which gradually increased to 72 of all 80 Swedish hospitals (90%) in 2001. During this period, a total of 88 026 admissions for MI were recorded. Of these, 974 were excluded because of pacemaker rhythm with pacemaker QRS complexes, and 45 265 with non-ST-segment elevation MI (NSTEMI), 33 746 with ST-segment elevation MI (STEMI), and 8041 with LBBB MI were included. Patients with STEMI or NSTEMI were gathered into 1 group of no LBBB ( $n=79\ 011$ ). The prevalence of LBBB in the MI population treated in intensive cardiac care units was 9% (8041 of 88 026). This prevalence increased with age, and the mean age was statistically significantly higher among MI patients with LBBB (mean  $\pm$  SD,  $77 \pm 9$  years) compared with the no-LBBB group ( $71 \pm 12$  years;  $P < 0.001$ ). Because age was a very important risk factor and associated with many comorbid conditions, we have chosen to present data stratified into 3 age groups;  $< 65$ , 65 to 74, and  $\geq 75$  years. Table

TABLE 1. Characteristics of Acute MI Patients at Arrival

	LBBB on Admission ECG					
	Age $\leq$ 64 y (n=23 453)		Age 65–74 y (n=23 612)		Age $\geq$ 75 y (n=39 987)	
	No	Yes	No	Yes	No	Yes
MI, n	22 712	741	21 684	1928	34 615	5372
Male, %	78	79	68	69	54‡	56
Concomitant risk factors/ history of disease, %						
Current smoker	42‡	32	22*	18	9*	8
Stroke	4‡	10	10‡	16	16‡	18
Renal failure	1*	2	1‡	3	2‡	3
Chronic pulmonary disease	3‡	6	6‡	10	7‡	9
Dementia	0.1	0.1	0.3*	0.5	0.7†	0.4
Cancer within 3 years	2	2	4	4	5	5
Congestive heart failure	4‡	23	11‡	34	22‡	43
Previous MI	21‡	41	31‡	52	40‡	55
Diabetes mellitus	18‡	31	24‡	34	23‡	29
Hypertension	28‡	36	36‡	43	38	38
Previous PCI	6‡	12	5‡	7	3	3
Previous CABG	5‡	15	7‡	17	4‡	6
Treatment on admission, %						
ACE inhibitor	14‡	34	19‡	39	21‡	35
Antiarrhythmics	1‡	3	1‡	3	2‡	3
Acetylsalicylic acid	28‡	44	41‡	51	51‡	59
Warfarin	3‡	10	5‡	12	6‡	8
Calcium channel inhibitor	12‡	19	18*	21	21	21
$\beta$ -Blocker	30‡	43	39‡	48	42‡	46
Digitalis	2‡	11	6‡	15	13‡	21
Diuretics oral	12‡	32	25‡	51	47‡	64
Lipid lowering	16‡	29	16‡	25	8	8
Nitroglycerin	11‡	26	21‡	34	34‡	46
Index events, %						
Circulatory arrest at entry	1‡	3	1‡	3	1†	2

n=87 052.

\* $P<0.05$ , † $P<0.01$ , ‡ $P<0.001$  within age group by  $\chi^2$ .

1 shows baseline characteristics of patients with no LBBB or LBBB stratified by age. Even after age stratification, the differences in comorbid conditions, medications on admission, and previous coronary interventions between the LBBB and no-LBBB groups were significant. Killip class on admission was significantly ( $P<0.001$ ) poorer in patients with LBBB compared with no-LBBB patients, revealing the following distribution: Killip class I, 44% versus 66%; class II, 26% versus 19%; class III, 15% versus 5%; and class IV, 15% versus 10%. When age stratification was performed, the differences in Killip class remained between the 2 ECG groups (data not shown).

### In-Hospital Intervention and Long-Term Medical Treatment

Patients with LBBB were less likely to receive acute reperfusion therapy and to undergo cardiac catheterization and

percutaneous coronary intervention (PCI) compared with no-LBBB patients (Table 2). A higher proportion of patients with LBBB received medication indicative of congestive heart failure (ACE inhibitors, diuretics, and digitalis) at discharge. In contrast, more no-LBBB patients were likely to receive  $\beta$ -blockade and antithrombotic agents compared with the LBBB population. However, in multiple logistic regression analyses, each adjusting for 25 covariates, the differences in long-term medical treatment at discharge were statistically significant only for ACE inhibitors ( $P<0.001$ ) and digitalis ( $P=0.007$ ). All other discharge medications had no statistical association with ECG appearance in these 2 MI groups.

### In-Hospital Complications and Mortality

The 1-year mortality rates in relation to biochemical marker peak level (divided into 4 groups: [1] CK-MB  $\leq$ 50

**TABLE 2. In-Hospital Intervention and Long-Term Medical Treatment in Acute MI Patients**

	LBBB on Admission ECG					
	Age ≤64 y (n=23 453)		Age 65–74 y (n=23 612)		Age ≥75 y (n=39 987)	
	No	Yes	No	Yes	No	Yes
IV β-blocker, %	40†	35	33‡	27	27‡	22
IV nitroglycerin, %	42	42	41	41	38	38
Reperfusion therapy, %	46‡	32	35‡	21	25‡	12
PCI, %	22‡	12	13‡	7	5‡	2
CABG, %	4.4	5.4	5.4‡	3.3	2.3‡	1.5
Echocardiography, %	60†	54	53‡	47	38‡	32
Medical treatment at discharge, %						
ACE inhibitor	35‡	57	46‡	63	44‡	54
Antiarrhythmics	1‡	5	2‡	4	2‡	4
Anticoagulant/ASA	94†	92	92‡	87	86‡	83
Calcium channel inhibitor	11‡	16	15*	17	17†	15
β-Blocker	87‡	81	81‡	76	71‡	64
Digitalis	3‡	12	8‡	18	15‡	24
Diuretics oral	19‡	46	39‡	65	60‡	75
Lipid lowering	56	54	44	42	20‡	16
Nitroglycerin	28‡	42	40‡	46	48‡	53

IV indicates intravenous; ASA, acetylsalicylic acid. n=87 052.

\**P*<0.05, †*P*<0.01, ‡*P*<0.001 within age groups by  $\chi^2$ .

μg/L or troponin-T <1.0 μg/L, [2] CK-MB 51 to 150 μg/L or troponin-T 1.0 to 3.0 μg/L, [3] CK-MB >150 μg/L or troponin-T >3.0 μg/L, and [4] missing values) were 38%, 42%, 45%, and 49% for the LBBB patients and 21%, 20%, 22%, and 26%, respectively, for the no-LBBB patients. The distributions of patients into the 4 biochemical marker groups were 44%, 23%, 20%, and 14% for LBBB compared with 37%, 23%, 25%, and 14% for no-LBBB patients. Patients with LBBB were at higher risk of developing new atrial fibrillation and clinical signs of heart failure in all age categories (Table 3). After adjustment for the 25 background covariates in a multiple logistic regression analysis, the difference remained statistically significant with regard to development of heart


failure (*P*<0.001). Unadjusted 30-day and 1-year mortality rates were 21% (n=1682) and 42% (n=3350) for the LBBB group compared with 12% (n=9591) and 22% (n=17 044) for the no-LBBB patients. One-year mortality was ≈4 times as high for LBBB compared with no-LBBB patients among those <65 years of age. There was no statistically significant difference in cause of death, which was ischemic heart disease during the first year in no-LBBB patients in 74% (n=12 670) compared with 76% of LBBB patients (n=2532) (*P*=0.13). The unadjusted RR of death within 1 year was 2.16 (95% CI, 2.08 to 2.24; *P*<0.001) for MI patients with LBBB compared with the no-LBBB group (Figure 1A). After adjustment for propensity score A (baseline characteristics and acute events), the

**TABLE 3. In-Hospital and Long-Term Complications in Acute MI Patients**

	LBBB on Admission ECG					
	Age ≤64 y (n=23 453)		Age 65–74 y (n=23 612)		Age ≥75 y (n=39 987)	
	No	Yes	No	Yes	No	Yes
Hypotension, %	7	13*	9	14*	12	15*
New atrial fibrillation, %	6	14*	15	25*	26	71*
Clinical signs of heart failure, %	22	47*	38	59*	54	71*
Reinfarction, %	3	4	4	4	5	4
Mortality 30 d, %	3	12*	9	16*	20	24*
Mortality 1 y, %	6	22*	16	30*	35	48*

n=87 052.

\**P*<0.001 within groups by  $\chi^2$ .



**Figure 1.** Estimated survival after admission. A, Unadjusted RR of death between MI patients presenting with either LBBB or no LBBB is estimated to be 2.2 ( $n=87\ 052$ ). B, Estimated survival adjusted for differences in background characteristics and treatments by propensity score, biochemical marker, and Killip class resulted in RR of 1.19 for LBBB ( $n=87\ 052$ ). C, In subgroup in which adjustments could be made for propensity score and LVEF, RR was 1.08 for LBBB ( $n=11\ 812$ ).


large difference in relative mortality risk was reduced to 1.28 (95% CI, 1.23 to 1.33;  $P<0.001$ ) for LBBB. Further adjustment, including propensity score A, Killip class, and biochemical marker group, resulted in an additional reduction in RR to 1.19 (95% CI, 1.14 to 1.24;  $P<0.001$ ) for 1-year mortality for LBBB (Figure 1B). When later interventions also were included (propensity score B), there was still an increased risk of death associated with LBBB on ECG for MI patients (RR, 1.21; 95% CI, 1.15 to 1.27;  $P<0.001$ ).

### LVEF and Mortality

Echocardiography was performed significantly less frequently in patients with LBBB (Table 2). In the 11 812 MI patients in whom it was performed, LVEF was worse in patients with LBBB. Among the LBBB patients, there were a significantly smaller proportion with normal LVEF (15% [ $n=131$ ] compared with 39% [ $n=4202$ ];  $P<0.001$ ), less with slight dysfunction (30% [ $n=282$ ] compared with 34% [ $n=3675$ ];  $P=0.012$ ), a larger proportion with moderate dysfunction (25% [ $n=234$ ] compared with 19% [ $n=2053$ ]), and 4 times as high a proportion with severe dysfunction (LVEF  $<30\%$ ) among LBBB (32% [ $n=299$ ]) compared with (8% [ $n=936$ ];  $P<0.001$ ) among no-LBBB patients. With decreased LVEF, the survival decreased for both ECG groups at the same time as the LBBB influence on mortality decreased. Patients with LVEF  $<30\%$  revealed 1-year mortalities for the 3 age groups of 19%, 33%, and 47% for the no-LBBB group and 24%, 29%, and 50% for the LBBB group. When Cox proportional-hazards regression analysis that included the variables propensity score (A), Killip class, biochemical marker group, presence of LBBB on ECG, and LVEF group was performed in the 11 812 cases with LVEF available, LBBB did not have any significant contribution to the RR of 1-year mortality (RR, 1.08; 95% CI, 0.93 to 1.25;  $P=0.33$ ; Figure 1C). When patients with normal to moderately decreased LVEF  $>30\%$  were analyzed, unadjusted 1-year mortality was 23% (146 of 647) for LBBB and 10% (988 of 9930) for no-LBBB patients. For patients with severely decreased LVEF  $<30\%$ , the corresponding figures were 41% (123 of 299) for LBBB and 38% (353 of 936) for no-LBBB patients. After adjustment for propensity score A, Killip class, biochemical marker, and LVEF, there was a nonsignificant increase in RR of 1.20 (95% CI, 0.99 to 1.45;  $P=0.066$ ) associated with LBBB among those with normal to moderately decreased LVEF (Figure 2A), whereas among those with severely decreased LVEF, there was no statistically significantly higher risk for 1-mortality associated with LBBB (RR, 0.93; 95% CI, 0.74 to 1.17;  $P=0.54$ ; Figure 2B).

### Discussion

This study is the first to include a large enough population to allow an extensive multivariable statistical analysis to evaluate the risk contribution by LBBB per se, not only the associated risk. The conclusion that the extent of comorbidity substantially reduces the independent prognostic impact of LBBB in acute MI is new information of considerable importance for the management of these patients. Furthermore, in light of the ongoing animated debate about wide



**Figure 2.** Estimated survival after admission adjusted for differences in background characteristics and treatment by propensity score methods related to LVEF. A, Patients with LVEF >30% had LBBB RR of 1.20 (95% CI, 0.99 to 1.45;  $P=0.066$ ) for 1-year mortality ( $n=10\,577$ ). B, Patients with LVEF <30% had LBBB RR of 0.93 (95% CI, 0.74 to 1.17;  $P=0.54$ ) for 1-year mortality ( $n=1235$ ).

QRS complex and implantable cardioverter-defibrillator therapy, we think this study has an important message. The most important finding in the present study is that after adjustment for differences in age, comorbidities, LVEF, and acute treatment, there is no significant difference in 1-year mortality in an MI population with LBBB compared with no LBBB. This finding is in contrast to that of previous reports, which, however, did not use the same extensive adjustment for covariates for mortality.<sup>6,7,9</sup> The absence of any specific effect of the LBBB conduction abnormality is also supported by the comparable causes of death in LBBB and no-LBBB patients.

The results also indicate that those with LBBB MI tend to have lower peak levels of biochemical markers than those with no-LBBB infarctions, despite the fact that the latter group consists of 57% NSTEMIs. Peak level of biochemical marker had no major influence on 1-year mortality in our study. This finding might be due to the lack of identification of the peak levels in a substantial number of patients because of early deaths, lack of serial measurements, influence of different kinetics of the markers in reperfused and nonreperfused patients, and the use of different markers in different hospitals.

**Reliability of Data**

The method used by current RIKS-HIA was first used in 1991 and has become a reliable source of information on consecutive patients admitted to participating units. ECG had a high validity with 97% correct coding. The patient cohort in this study included unselected consecutive patients with MI from a large number of different hospital types. There were no exclusions resulting from the presence or absence of specific risk factors, comorbidities, anticipated adverse effects, participation in clinical trials, or contraindications to certain medications. The representativeness of the cohort was also strengthened by the inclusion of all patients with MI from the general population at the centers with different levels of care from 90% of the hospitals within an entire country.

In the present study, the prevalence of LBBB in a population of intensive cardiac care unit admissions for MI was 9%, consistent with the LBBB prevalence (7%) in the study reported by Newby et al.<sup>6</sup> In the GUSTO-I trial,<sup>9</sup> the prevalence of LBBB was as low as 0.5%, probably representing selection bias based on the entry criteria. Because the register covers background factors such as age, risk factors, and previous diseases and medications, adjusting for differences in the composition of different population is possible.

**Mortality and Left Ventricular Dysfunction**

In the present study as in previous studies, LBBB patients were older, which could explain the higher mortality among MI patients with LBBB. Compared with the no-LBBB group, patients with LBBB on admission for MI had a higher prevalence of preexisting comorbid conditions such as previous MI, congestive heart failure, diabetes mellitus, hypertension, renal failure, chronic pulmonary disease, and history of stroke. Moreover, the LBBB population had more frequently undergone coronary intervention, further supporting the occurrence of more severe coronary artery disease. The increased use of ACE inhibitors, diuretics, and digitalis among LBBB patients on arrival emphasized the more common occurrence of poor left ventricular function before the new MI in LBBB compared with no-LBBB patients. These differences in clinical background are consistent with previous reports<sup>7,13,21</sup> and seem to explain the difference in mortality in our study. When differences in baseline characteristics like age and comorbidity were compensated for, the RR of 1-year mortality for LBBB patients was substantially reduced. The importance of the higher age and more frequent comorbidities is also supported by the absence of differences in most long-term treatments after adjustment for covariates

in the multivariable analysis. During in-hospital stay, LBBB patients were less likely than the no-LBBB patients to have PCI or CABG, therapies shown to improve long-term survival.<sup>22</sup> At discharge, the LBBB population compared with no-LBBB group was given less secondary prevention ( $\beta$ -blockade and antithrombotic agents), consistent with previous studies,<sup>7,10</sup> but was more likely to receive medication indicative of heart failure. Complications associated with worse outcomes such as atrial fibrillation, hypotension, and congestive heart failure were more common in the LBBB group.

Increased long-term mortality associated with LBBB in patients with MI has been reported in previous studies.<sup>8,10</sup> In the present study, unadjusted 1-year mortality was more than twice as high for patients with LBBB compared with no-LBBB patients. In our study, however, when 1-year mortality was adjusted step by step for baseline characteristics, comorbid conditions, therapy received, and LVEF by Cox regression analysis, LBBB seemed not to be an important contributor to 1-year mortality in MI patients. Consequently, our results indicate that baseline characteristics and concomitant diseases in the settings of MI are the main reasons for the higher 1-year mortality in patients with LBBB. The absence of any specific effect of the LBBB conduction abnormality is also supported by the comparable causes of death in LBBB MI and no-LBBB patients.

### Study Limitations

A limitation of this study was that LVEF was not evaluated by a core laboratory but was an on-site judgment during routine care by the physician performing the echocardiography. Another limitation was that LVEF was not available in most patients. Still, to the best of our knowledge, the subgroup of MI patients with LBBB and measured LVEF is the largest reported so far. In multiple regression analysis, 5% of the study population was excluded because of missing variables in the registry. However, when a sensitivity analysis was performed, there was no change in the main result. This study did not record any transient LBBB, nor could it differentiate between newly developed or old LBBB, so the conclusions should be applied primarily to persistent LBBB in patients with acute MI.

### Clinical Implications

LBBB does not appear to be an important independent predictor of 1-year mortality in MI but mainly reflects higher age, comorbid conditions, and left ventricular dysfunction. Therefore, there seems to be no specific reasons and risks for death related to the conduction abnormality in the LBBB MI population. The recently published MADIT II study reported the largest survival benefit with implantable defibrillator therapy in patients with LBBB in a population with previous MI and left ventricular dysfunction.<sup>16</sup> In our study, LBBB MI patients had a medical therapy on admission and discharge indicative of a relatively high prevalence of congestive heart failure; consequently, this population was fairly similar to the MADIT II patients with LBBB. Still, the adjusted mortality rate and the causes of death were similar in the LBBB MI and no-LBBB populations. Thus, the anticipated risk of death rather than a conduction abnormality should probably be used as a guideline for the selection of implantable defibrillator

therapy and other effective treatments such as reperfusion therapy and early revascularization in post-MI patients.

### References

1. Bauer GE, Julian DG, Valentine PA. Bundle-branch block in acute myocardial infarction. *Br Heart J*. 1965;27:724–730.
2. Dubois C, Pierard LA, Smeets JP, et al. Short- and long-term prognostic importance of complete bundle-branch block complicating acute myocardial infarction. *Clin Cardiol*. 1988;11:292–296.
3. Godman MJ, Lassers BW, Julian DG. Complete bundle-branch block complicating acute myocardial infarction. *N Engl J Med*. 1970;282:237–240.
4. Hindman MC, Wagner GS, JaRo M, et al. The clinical significance of bundle branch block complicating acute myocardial infarction, 2: indications for temporary and permanent pacemaker insertion. *Circulation*. 1978;58:689–699.
5. Norris RM, Croxson MS. Bundle branch block in acute myocardial infarction. *Am Heart J*. 1970;79:728–733.
6. Newby KH, Pisano E, Krucoff MW, et al. Incidence and clinical relevance of the occurrence of bundle-branch block in patients treated with thrombolytic therapy. *Circulation*. 1996;94:2424–2428.
7. Go AS, Barron HV, Rundle AC, et al. Bundle-branch block and in-hospital mortality in acute myocardial infarction: National Registry of Myocardial Infarction 2 Investigators. *Ann Intern Med*. 1998;129:690–697.
8. Hod H, Goldbourt U, Behar S. Bundle branch block in acute Q wave inferior wall myocardial infarction: a high risk subgroup of inferior myocardial infarction patients: the SPRINT Study Group: Secondary Prevention Reinfarction Israeli Nifedipine Trial. *Eur Heart J*. 1995;16:471–477.
9. Sgarbossa EB, Pinski SL, Topol EJ, et al. Acute myocardial infarction and complete bundle branch block at hospital admission: clinical characteristics and outcome in the thrombolytic era: GUSTO-I Investigators: Global Utilization of Streptokinase and t-PA [Tissue-Type Plasminogen Activator] for Occluded Coronary Arteries. *J Am Coll Cardiol*. 1998;31:105–110.
10. Brilakis ES, Wright RS, Kopecky SL, et al. Bundle branch block as a predictor of long-term survival after acute myocardial infarction. *Am J Cardiol*. 2001;88:205–209.
11. Jones ME, Terry G, Kenmure AC. Frequency and significance of conduction defects in acute myocardial infarction. *Am Heart J*. 1977;94:163–167.
12. Gould L, Ramana CV, Gomprecht RF. Left bundle-branch block: prognosis in acute myocardial infarction. *JAMA*. 1973;225:625–627.
13. Otterstad JE, Gundersen S, Anderssen N. Left anterior hemiblock in acute myocardial infarction: incidence and clinical significance in relation to the presence of bundle branch block and to the absence of intraventricular conduction defects. *Acta Med Scand*. 1978;203:529–534.
14. Ellis CJ, French JK, White HD. Thrombolytic eligibility. *Aust N Z J Med*. 1998;28:518–524.
15. Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients: Fibrinolytic Therapy Trialists' (FTT) Collaborative Group. *Lancet*. 1994;343:311–322.
16. Moss AJ, Zareba W, Hall WJ, et al. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med*. 2002;346:877–883.
17. Stenestrand U, Wallentin L. Early statin treatment following acute myocardial infarction and 1-year survival. *JAMA*. 2001;285:430–436.
18. Criteria Committee of the New York Heart Association. *Nomenclature and Criteria for Diagnosis of Diseases of the Heart and Great Vessels*. Boston, Mass: Little Brown and Company; 1973.
19. Tunstall-Pedoe H, Kuulasmaa K, Amouyel P, et al. Myocardial infarction and coronary deaths in the World Health Organization MONICA Project: registration procedures, event rates, and case-fatality rates in 38 populations from 21 countries in four continents. *Circulation*. 1994;90:583–612.
20. R Development Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Vienna. Available at: <http://www.R-project.org>. 2003. Accessed April 2, 2004.
21. Freedman RA, Alderman EL, Sheffield LT, et al. Bundle branch block in patients with chronic coronary artery disease: angiographic correlates and prognostic significance. *J Am Coll Cardiol*. 1987;10:73–80.
22. Stenestrand U, Wallentin L. Early revascularisation and 1-year survival in 14-day survivors of acute myocardial infarction: a prospective cohort study. *Lancet*. 2002;359:1805–1811.

**Comorbidity and Myocardial Dysfunction Are the Main Explanations for the Higher 1-Year Mortality in Acute Myocardial Infarction With Left Bundle-Branch Block**  
Ulf Stenestrand, Fariborz Tabrizi, Johan Lindbäck, Anders Englund, Mårten Rosenqvist and  
Lars Wallentin

*Circulation*. 2004;110:1896-1902; originally published online September 27, 2004;  
doi: 10.1161/01.CIR.0000143136.51424.38

*Circulation* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231  
Copyright © 2004 American Heart Association, Inc. All rights reserved.  
Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the  
World Wide Web at:

<http://circ.ahajournals.org/content/110/14/1896>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation* is online at:  
<http://circ.ahajournals.org/subscriptions/>