



<http://dx.doi.org/10.11646/zootaxa.3641.3.9>

<http://zoobank.org/urn:lsid:zoobank.org:pub:CE8B3DC6-C285-4A42-9425-AADE2F9A2A5E>

DNA-sequence data require revision of the parrot genus *Aratinga* (Aves: Psittacidae)

J. V. REMSEN, JR.¹, ERIN E. SCHIRTZINGER², ANNA FERRARONI³, LUÍS FÁBIO SILVEIRA³
& TIMOTHY F. WRIGHT²

¹Museum of Natural Science & Department of Biological Sciences, Louisiana State University, Baton Rouge, Louisiana 70803, USA.
E-mail: najames@LSU.edu

²Department of Biology, New Mexico State University, Las Cruces, New Mexico 88003, USA

³Seção de Aves, Museu de Zoologia da Universidade de São Paulo, Av. Nazaré 481, Ipiranga, São Paulo, CEP 04263-000, Brazil

The genus *Aratinga* von Spix, 1824, as treated since Peters (1937), consists of 20 to 21 species (Kremer 1989, Collar 1997, Juniper & Parr 1998, Silveira *et al.* 2005, Dickinson 2003, Forshaw 2010) of medium-sized, pointed-tailed, mostly green parakeets that range throughout the Neotropical region. All species currently included in the genus *Aratinga* had already been recognized by Salvadori (1891) and placed in the genus *Conurus*. Ridgway (1916) placed the species in four genera: *Aratinga*, *Eupsittula*, *Nandayus*, and *Thectocercus*; Ridgway provided a rationale for his treatment using morphological and plumage characters, and he included a dichotomous key. Cory (1918) followed Ridgway's (1916) classification. Miranda-Ribeiro (1920) placed the species in four genera: *Conurus*, "*Nendayus*" (= *Nandayus*), *Gymnopsittacus*, and "*Eupsittacula*" (= *Eupsittula*). Peters (1937) placed all members of these genera but *Nandayus nenday* into a single genus, *Aratinga* (Table 1), but provided no rationale for his classification. Nonetheless, his treatment has been followed in all subsequent classifications, including Meyer de Schauensee (1970), Sibley & Ahlquist (1990), Collar (1997), Dickinson (2003), and Forshaw (2010), although Marien & Koopman (1955) suggested retention of three subgenera.

Ribas & Miyaki's (2004) results called into question the monophyly of Peters' broadly defined *Aratinga* when they found that *A. leucophthalma* was only distantly related to the other *Aratinga* sampled and that *Nandayus nenday* was embedded within the sampled *Aratinga* species. Silveira *et al.* (2005) provided additional details delimiting at least three groups within broadly defined *Aratinga* (see also Whitney 1996) and proposed that it was not a monophyletic genus. These three groups also show consistent difference in vocalizations among the groups and similarities among species within each group (B. M. Whitney, pers. comm.). Subsequent studies with much broader taxon sampling (Kirchman *et al.* 2012, Schirtzinger *et al.* 2012) confirmed that the genus consisted of three separate lineages, corresponding in part to the 3-genus classification of Ridgway (1916) and Pinto (1938). Further, Ribas & Miyaki (2004) and Tavares *et al.* (2006) found that the monotypic genus *Nandayus* was embedded in one of the *Aratinga* lineages. Subsequently, Urantowka *et al.* (2012) also found that *Aratinga* species were distributed in the three clades found by previous authors, but also found that *A. acuticaudata* was even more distantly related to the other *Aratinga* and was actually the sister species to *Diopsittaca nobilis*.

These new data require that *Aratinga* sensu Peters (1937) be partitioned into four genera. The type species (by subsequent designation) for *Aratinga* is *solstitialis* Linnaeus, 1758. Two species have traditionally been considered closely related to *A. solstitialis*: *A. jandaya* and *A. auricapillus*. In fact, Meyer de Schauensee (1966), Sick (1993), and others considered them conspecific, and Sibley & Ahlquist (1990) treated them as members of a superspecies. Ribas & Miyaki (2004) included *solstitialis* in their taxon sampling, but Silveira *et al.* (2005) noted that the specimens used were actually of a taxon that they described as a new species, *Aratinga pinto*. Subsequently, Nemésio and Rasmussen (2009) determined that *A. pinto* had been previously described as *Psittacus luteus* Boddaert, 1783, which is, in turn, synonym of *Psittacus maculatus* (= *Aratinga maculata*) Statius Muller, 1776, a name that had been relegated to the synonymy *A. solstitialis* and had been overlooked by subsequent authors, including Peters (1937). The plumage characters associated with *maculata* had been dismissed as age variation in *A. solstitialis*, aviary artifacts, or hybrids by everyone except Pinto (1966), until Silveira *et al.* (2005) showed that they represented discrete characters that defined a geographical representative of the *A. solstitialis* group. Therefore, we think it is reasonable to include *A. maculata* in *Aratinga* sensu stricto. Tavares *et al.* (2006) included *solstitialis* in their sampling, but did not provide information on their vouchers and

did not cite Silveira *et al.* (2005); thus whether their specimen was true *solstitialis* or *maculata* is uncertain. Kirchman *et al.* (2012) included true *solstitialis* (fide G. R. Graves corroboration of identification of the voucher specimen, USNM 614252) in their sampling, however, and its phylogenetic position there was consistent with that in Ribas & Miyaki (2004), who found that the species *jandaya* and *auricapillus* were closely related to *maculata* and that these three formed a clade to which *Nandayus nenday* and *A. weddellii* were successive sisters. Therefore, *jandaya* and *auricapillus* may be safely retained in *Aratinga*. *Nandayus nenday* has been placed in a monotypic genus largely because of its black crown and elongated upper mandible, but structurally and in other aspects of its plumage, it seems not only to be a typical *Aratinga*, where placed by Short (1975), but also a member of this species group (Silveira *et al.* 2005). One additional species almost certainly also belongs in this group: *Aratinga weddellii* grouped with these species in Ribas & Miyaki (2004) and shares with all species in this group the distinctive and diagnostic coloration pattern of the rectrices and primaries, and black bill, as outlined by Silveira *et al.* (2005); it is also allopatric to the other members of the group. *Aratinga* as thus narrowly constituted consists of six allopatric or parapatric taxa restricted to lowland tropical and subtropical South America. An unresolved question is whether the extinct *Conuropsis carolinensis* should be included within this group. Kirchman *et al.* (2012) found it to be sister to the clade composed of *A. solstitialis*, *A. auricapillus* and *N. nenday* based on molecular characters. They did not, however, sample *A. jandaya* or *A. weddellii*, so it remains uncertain whether *C. carolinensis* belongs within the newly circumscribed genus *Aratinga* or is sister to it.

TABLE 1. History of generic assignment of currently (Dickinson 2003) recognized species in broadly defined *Aratinga*. Species names and sequence as in Dickinson (2003). Blank cells represent taxa not treated in that reference.

SPECIES	Salvadori (1891)	Miranda-Ribeiro (1920)	Ridgway (1916)	Cory (1918)	Peters (1937)	Pinto (1938)	Dickinson (2003)
<i>acuticaudata</i>	<i>Conurus</i>	<i>Conurus</i>	<i>Thectocercus</i>	<i>Thectocercus</i>	<i>Aratinga</i>	<i>Psittacara</i>	<i>Aratinga</i>
<i>holochlora</i>	<i>Conurus</i>		<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>strenua</i>			<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>wagleri</i>	<i>Conurus</i>		<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>mitrata</i>	<i>Conurus</i>			<i>Aratinga</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>erythrogegens</i>				<i>Aratinga</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>finschi</i>	<i>Conurus</i>		<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>leucophthalma</i>	<i>Conurus</i>	<i>Conurus</i>		<i>Aratinga</i>	<i>Aratinga</i>	<i>Psittacara</i>	<i>Aratinga</i>
<i>euops</i>	<i>Conurus</i>		<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>chloroptera</i>	<i>Conurus</i>		<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>solstitialis</i>	<i>Conurus</i>	" <i>Nendayus</i> "	<i>Eupsittula</i>	<i>Eupsittula</i>	<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>
<i>jandaya</i>	<i>Conurus</i>	" <i>Nendayus</i> "	<i>Eupsittula</i>	<i>Eupsittula</i>	<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>
<i>auricapillus</i>	<i>Conurus</i>	" <i>Nendayus</i> "	<i>Eupsittula</i>	<i>Eupsittula</i>	<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>
<i>weddellii</i>	<i>Conurus</i>	<i>Gymnopsittacus</i>		<i>Eupsittula</i>	<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>
<i>nana</i>	<i>Conurus</i>		<i>Eupsittula</i>	<i>Eupsittula</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>canicularis</i>	<i>Conurus</i>		<i>Eupsittula</i>	<i>Eupsittula</i>	<i>Aratinga</i>		<i>Aratinga</i>
<i>aurea</i>	<i>Conurus</i>	" <i>Eupsittacula</i> "	<i>Eupsittula</i>	<i>Eupsittula</i>	<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>
<i>pertinax</i>	<i>Conurus</i>	<i>Gymnopsittacus</i>	<i>Eupsittula</i>	<i>Eupsittula</i>	<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>
<i>cactorum</i>	<i>Conurus</i>	<i>Gymnopsittacus</i>		<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>	<i>Aratinga</i>
<i>nenday</i>	<i>Conurus</i>	" <i>Nendayus</i> "	<i>Nandayus</i>	<i>Nandayus</i>	<i>Nandayus</i>	" <i>Nendayus</i> "	<i>Nandayus</i>

Kirchman *et al.* (2012) found that seven species included in *Aratinga* by Peters (1937) formed a monophyletic group that corresponded to Silveira *et al.*'s (2005) group II. This group is not closely related to true *Aratinga* but rather is the sister to a group that consists of *Leptosittaca branickii*, *Diopsittaca nobilis*, and *Guarouba guarouba*. The species are *leucophthalmus*, *wagleri*, *mitrata*, *holochlora*, *finschi*, *chloroptera*, and *euops*. The oldest genus name available for this group is *Psittacara* Vigors, 1825. Silveira *et al.* (2005) included two other species in this group that were not sampled in the DNA-based studies: *Aratinga strenua* and *A. erythrogegens*. We agree that they can be safely if provisionally also placed in *Psittacara*. *Aratinga strenua* is often treated as a subspecies of *A. holochlora* (e.g., Juniper & Parr 1998), and *A. erythrogegens* is traditionally considered closely related to *A. mitrata* and *A. wagleri* because of similarities in plumage

and size. *Psittacara* also differs from *Aratinga* sensu stricto and *Eupsittula* (see below) in that most component species range high into montane areas, and at least three taxa are exclusively montane (*Psittacara holochlorus brewsteri*, *P. wagleri*, *P. mitratus*).

Silveira *et al.* (2005) also included *Aratinga acuticaudata* in this group, although it had not been included in the DNA-based studies. However, it is the only member of the latter group with red in the rectrices and blue in the crown. Ridgway (1916) placed it in a monotypic genus, *Thectocercus*, because it differed from all *Aratinga* in having: (a) a very wide bill, wider than it was deep; (b) a bill tip that was “attenuated, acute, and ridged”; (c) a cere lacking feathers posterior to nares; and (d) inner webs of rectrices extensively red. However, *acuticaudata* does share the following features with members of the *Psittacara* group, as outlined by Silveira *et al.* (2005): relatively large size compared to other members of broadly defined *Aratinga*; mostly green plumage; large, whitish eyering; pale maxilla; and remiges ventrally yellowish green. Nonetheless, Ferraroni and Silveira (unpubl. data) have found additional morphological characters that reinforce Ridgway’s assessment. Finally, Urantowka *et al.* (2012) found that *acuticaudata* was not a member of the *Psittacara* group, and that it was embedded in the *Leptosittaca-Guarouba-Diopsittaca* group. Thus, resurrecting *Thectocercus* for *acuticaudata* is required to maintain monotypic *Psittacara*.

Arndt (2006) described a new species, *Aratinga hockingi*, from the Andes of Peru, and a new subspecies, *A. mitrata tucumana*, from Argentina. He also elevated the Andean taxon *A. mitrata alticola* to species rank. *Aratinga hockingi* and *A. alticola* were noted as differing from *A. mitrata* primarily in the extent of red on the head and thighs, and as differing from each other only in extent of red on forecrown; as noted by Arndt (2006), these characters also show individual and age variation within each taxon. Agnolin (2009) considered *tucumana* to be a synonym of nominate *A. mitrata* because he considered *tucumana* to represent individual variation within Argentine populations of *A. m. mitrata*. Other authors have discussed the issue of plumage variation in broadly defined *Aratinga*, including Blake & Hocking’s (1947) on individual and seasonal variation in *Aratinga acuticaudata*, and Marien & Koopman (1955) on individual variation in *Aratinga* in general. Based on this and the seeming unlikelihood that three or four species of *Psittacara* would be sympatric in some Andean valleys, Remsen *et al.* (2012) did not recognize *hockingi* as a valid species or *alticola* as a separate species pending additional corroborating field and lab studies (Ferraroni *et al.*, in prep.). If these taxa are subsequently shown to be valid species, they would also be placed in *Psittacara*.

Kirchman *et al.* (2012) also found that another three species (*pertinax*, *nana*, and *aurea*) included in *Aratinga* by Peters (1937) formed a monophyletic group that was not closely related to true *Aratinga* but rather was the likely sister group, among the genera sampled, to the genus *Rhynchopsitta*. Tavares *et al.* (2006) also found that *A. aurea* was unlikely to be closely related to *A. solstitialis*. Together, these results suggest that *Aratinga*, even without the species assigned above to *Psittacara*, is not monophyletic. Two other species have been linked to this group by molecular studies: Ribas & Miyaki (2004) found that *Aratinga cactorum* was sister to *A. aurea* among the taxa they sampled, and Schirtzinger *et al.* (2012) found that *A. canicularis* was sister to *A. pertinax*. Sick (1993) considered *A. cactorum* to form a superspecies with *A. pertinax*. The oldest genus name available for the group is *Eupsittula* Bonaparte, 1853, for which the type species is *canicularis*.

We recommend that the five species above, the same five identified by Silveira *et al.* (2005) as forming a natural group, be included in a resurrected *Eupsittula*. All five are relatively small species that share grayish or olive breast plumage, and rectrices that are mostly green dorsally and olivaceous ventrally. They also share a similar pattern of coloration in the upper wing coverts: like *Aratinga* (and unlike *Psittacara*) they have blue greater wing coverts, but the blue is paler (reduced to a faint tinge in *cactorum*) and does not extend to the outer primary coverts. All five species are found primarily in semi-humid to dry regions in deciduous forest or scrub and even desert, or if in humid areas, in edge or riverine habitat. They are allopatric except for minor, perhaps recent (B. M. Whitney, pers. comm.) overlap between *E. aurea* and *E. cactorum* in eastern Brazil.

Taxonomic recommendation. We recommend the following reclassification of the species formerly placed in *Aratinga* by Peters (1937):

Aratinga (gender feminine; type species = *solstitialis*)

Aratinga weddellii (Deville, 1851)

Aratinga nenday (Vieillot, 1823)

Aratinga solstitialis (Linnaeus, 1758)

Aratinga maculata (Statius Muller, 1776)

Aratinga jandaya (Gmelin, 1788)

Aratinga auricapillus (Kuhl, 1820)

Eupsittula (gender feminine; type species = *canicularis*)

Eupsittula nana (Vigors, 1830)
Eupsittula canicularis (Linnaeus, 1758)
Eupsittula aurea (Gmelin, 1788)
Eupsittula pertinax (Linnaeus, 1758)
Eupsittula cactorum (Kuhl, 1820)

Thectocercus (gender masculine, thus requiring changes to the endings of adjectival species names; type species = *acuticaudatus*)

Thectocercus acuticaudatus (Vieillot, 1818)

Psittacara (gender masculine, thus requiring changes to the endings of adjectival species names; type species = *leucophthalmus*)

Psittacara holochlorus (Sclater, 1859)
Psittacara strenuus (Ridgway, 1915)
Psittacara wagleri (G.R. Gray, 1845)
Psittacara mitratus (von Tschudi, 1844)
Psittacara erythrogegens Lesson, 1844
Psittacara finschi (Salvin, 1871)
Psittacara leucophthalmus (Statius Muller, 1776)
Psittacara euops (Wagler, 1832)
Psittacara chloropterus Souancé, 1856

Note that species limits in the *Psittacara holochlorus/strenuus* group are controversial, in that some classifications treat *strenuus* as a subspecies of *holochlorus* (Juniper & Parr 1998, Forshaw 2010), and others treat *P. h. rubritorquis* and *P. h. brevipes* each as separate species (Collar 1997). We also provide the following simple key to the four genera based on coloration of the greater wing coverts and tail and including some of the characters noted by Ridgway (1916):

A1. Greater wing coverts and tail green: ***Psittacara***.

A2. Greater wing coverts green and tail green with red in the inner webs: ***Thectocercus***.

A3. Greater wing covert with blue or violet, and tail green or olive-green tipped violet: B

B1. Greater primary coverts violet blue, tail olive-green tipped violet: ***Aratinga***.

B2. Greater primary coverts mostly green with some blue (bluish green in *cactorum*) on inner primary coverts, tail green: ***Eupsittula***.

Acknowledgements

We thank Steven M. Gregory and Edward C. Dickinson for help with nomenclature, Normand David for determining gender of the genera, G. R. Graves for checking the identification of a voucher specimen, and Jessica Eberhard, Terry Chesser, Jeremy Kirchner, Dan Lane, Bret Whitney, and an anonymous reviewer for comments on the paper.

Literature cited

- Agnolin, F.L. (2009) Sobre el complejo *Aratinga mitrata* (Psittaciformes: Psittacidae) en el noroeste argentino. Comentarios sistemáticos. *Notúlas Faunísticas*, 31, 1–5.
- Arndt, T. (2006) A revision of the *Aratinga mitrata* complex, with the description of one new species, two new subspecies, and species-level status of *Aratinga alticola*. *J. Ornithology*, 147, 73–86.
- Blake, E.R., & Traylor, M.A., Jr. (1947) The subspecies of *Aratinga acuticaudata*. *Fieldiana (Zoology)*, 31, 163–169.
- Collar, N. (1997) Family Psittacidae (parrots). In: "*Handbook of the Birds of the World*. Vol. 4. Sandgrouse to Cuckoos" (J. del Hoyo *et al.*, eds.). Lynx Edicions, Barcelona, pp. 280–477.
- Cory, C.B. (1918) *Catalogue of Birds of the Americas*. Field Museum Natural History Publications, Zool. Series, vol. 13, part 2, no. 1.
- Dickinson, E.C. (Ed.) (2003) *The Howard and Moore Complete Checklist of the Birds of the World*. 3rd Edition.

- Christopher Helm, London, 1040 pp.
- Forshaw, J.M. (2010) *Parrots of the World*. A & C Black Publishers Ltd, London, 584 pp.
- Juniper, T. & Parr, M. (1998) *Parrots: a Guide to Parrots of the World*. Yale University Press, New Haven and London, 584 pp.
- Kirchman, J.J., Schirtzinger, E.E. & Wright, T.F. (2012) Phylogenetic relationships of the extinct Carolina Parakeet (*Conuropsis carolinensis*) inferred from DNA sequence data. *Auk*, 129, 197–204.
- Kremer, H. (1989) *Aratinga* s. Ornith. Noordbergum, 159 pp.
- Marien, D., & Koopman, K. (1955) The relationships of the West Indian species of *Aratinga* (Aves, Psittacidae). *American Museum Novitates*, 1712, 1–20.
- Meyer de Schauensee, R. (1966) *The Species of Birds of South America and their Distribution*. Livingston Publishing Co., Narberth, Pennsylvania. 577 pp.
- Miranda-Ribeiro, A. (1920) Revisão dos psittacideos brasileiros. *Revista do Museu Paulista*, 12, 1–82.
- Nemésio, A., & Rasmussen, C. (2009) The rediscovery of Buffon's "Guarouba" or "Perriche jaune"; two senior synonyms of *Aratinga pinto* Silveira, Lima & Höfling, 2005 (Aves: Psittaciformes). *Zootaxa*, 2013, 1–16.
- Pinto, O.M.O. (1938) Catálogo das aves do Brasil. Parte 1. *Revista do Museu Paulista*, 22, 1–566.
- Pinto, O.M.O. (1966) Estudo crítico e catálogo remissivo das aves do Território Federal de Roraima. *Cadernos de Amazonia*, 8, 1–176.
- Remsen, J.V., Jr., Cadena, C.D., Jaramillo, A., Nores, M., Pacheco, J.F., Pérez-Emán, J., Robbins, M.B., Stiles, F.G., Stotz, D.F., & Zimmer, K.J. (2013) A classification of the bird species of South America. American Ornithologists' Union. Available from <http://www.museum.lsu.edu/~Remsen/SACCBaseline.html> (Accessed 7 February 2013).
- Ribas, C.C., & Miyaki, C.Y. (2004) Molecular systematics in *Aratinga* parakeets: species limits and historical biogeography in the *solstitialis* group, and the systematic position of *Nandayus nenday*. *Molecular Phylogenetics and Evolution*, 30, 663–675.
- Ridgway, R. (1916) *The Birds of North and Middle America*. Part 7. Bulletin U.S. National Museum, no. 50.
- Salvadori, T. (1891) Catalogue of the Psittaci, or Parrots, in the collection of the British Museum. Longmans & Co., London, 658 pp.
- Schirtzinger, E.E., Tavares, E.S., Gonzales, L.A., Eberhard, J.R., Miyaki, C.Y., Sanchez, J.J., Hernandez, A., Mueller, H., Graves, G.R., Fleischer, R.C., & Wright, T.F. (2012) Multiple independent origins of mitochondrial control region duplications in the order Psittaciformes. *Molecular Phylogenetics and Evolution*, 64, 342–356.
- Short, L.L. (1975) A zoogeographic analysis of the South American Chaco avifauna. *Bulletin American Museum Natural History*, 154, 1–352.
- Sibley, C.G., and Ahlquist, J.E. (1990) *Phylogeny and Classification of Birds*. Yale University Press, New Haven, Connecticut. 1080 pp.
- Silveira, L.F., Lima, F.C. T., & Höfling, E. (2005) A new species of *Aratinga* parakeet (Psittaciformes: Psittacidae) from Brazil, with taxonomic remarks on the *Aratinga solstitialis* complex. *Auk*, 122, 292–305.
- Tavares, E.S., Baker, A.J., Pereira, S.L., & Miyaki, C.Y. (2006) Phylogenetic relationships and historical biogeography of Neotropical parrots (Psittaciformes: Psittacidae: Arini) inferred from mitochondrial and nuclear DNA Sequences. *Systematic Biology*, 55, 454–470.
- Urantowka, A. D., Grabowski, K. A. & Strzala, T. (2012) Complete mitochondrial genome of Blue-crowned Parakeet (*Aratinga acuticaudata*) -- phylogenetic position of the species among parrots group called conures. *Mitochondrial DNA, Early Online*, 1–3.
- Whitney, B.M. (1996) Flight behaviour and other field characteristics of the genera of Neotropical parrots. *Cotinga*, 5, 32–42.
- Wright, T.F., Schirtzinger, E.R., Matsumoto, T., Eberhard, J.R., Graves, G.R., Sanchez, J.J., Capelli, S., Muller, Scharpegge, H.J., Chambers, G.K., & R. C. Fleischer, R.C. (2008) A multilocus molecular phylogeny of the parrots (Psittaciformes): support for a Gondwanan origin during the Cretaceous. *Molecular Biology and Evolution*, 25, 2141–2156.