

Molecular phylogeny of *Acremonium* and its taxonomic implications

Anthony E. Glenn¹

*Department of Plant Pathology, University of Georgia,
Athens, Georgia 30602*

Charles W. Bacon

*USDA-ARS, Toxicology & Mycotoxin Research Unit,
Russell Research Center, Athens, Georgia 30613*

Robert Price

*Department of Botany, University of Georgia, Athens,
Georgia 30602*

Richard T. Hanlin

*Department of Plant Pathology, University of Georgia,
Athens, Georgia 30602*

Abstract: *Acremonium* is generally considered to be a highly polyphyletic form genus containing distantly related fungi. Sectional divisions within *Acremonium* distinguish the clavicipitaceous grass endophytes of sect. *Albolanosa* from the generally saprobic species of sections *Acremonium*, *Chaetomioides*, *Gliomastix*, and *Nectrioides*. In an effort to assess the possible number of lineages currently placed within *Acremonium* and to determine which groups of sexual ascomycetes are phylogenetically affiliated with *Acremonium* species, maximum parsimony and neighbor-joining analyses were performed using partial sequences of the nuclear small subunit ribosomal DNA (18S rDNA). *Acremonium* was shown to be a polyphyletic taxon with affiliations to at least three ascomycetous orders: 1) most of the examined species from the sections *Acremonium*, *Gliomastix*, and *Nectrioides* showed a relationship to the Hypocreaceae even though many of these species have never been associated with any teleomorph; 2) the grass endophytes of sect. *Albolanosa* and other taxa from the Clavicipitaceae formed a monophyletic group derived from within the Hypocreales; 3) the thermophilic *A. alabamense* of sect. *Chaetomioides* was derived from within the Sordariales. *Acremonium alternatum*, the type species of the genus, was one of the species showing affiliation to the Hypocreaceae. In order to eliminate some of the heterogeneity within *Acremonium* while also emphasizing the unique biological, morphological, and ecological characteristics of the grass endo-

phytes, we are proposing that the anamorphs of *Epichloë* and closely related asexual grass endophytes be reclassified into the new form genus *Neotyphodium*. Phylogenetic and taxonomic considerations are also presented for other taxa.

Key Words: Clavicipitaceae, endophytes, *Epichloë*, Hypocreales, *Neotyphodium*, rDNA

INTRODUCTION

Members of the family Clavicipitaceae (Hypocreales; Ascomycotina) are well known pathogens of a diverse assemblage of hosts including grasses, sedges, other ascomycetes, and insects. These fungi are found throughout the tropical and temperate regions of the world (Diehl, 1950; White, 1994b). Rogerson (1970) listed 31 genera in the family, but eight of these are considered to be synonyms. The family is divided into three subfamilies with the insect and fungal pathogens variously placed in the Oomycetoideae and Cordycipitoideae (Diehl, 1950). The plant pathogens are by far the most widely studied and economically important members of the family and are classified as the subfamily Clavicipitoideae (Diehl, 1950). Diehl (1950) further divides the taxa of the Clavicipitoideae into the three tribes Clavicipiteae, Balansieae, and Ustilaginoideae. The fungus-host interactions of these taxa can be broadly categorized as being epibiotic or endophytic, systemic or localized, and heavily pathogenic, moderately pathogenic, or mutualistic.

Most of the grass pathogens are in the tribe Balansieae (Diehl, 1950) and are either localized to epibiotic reproductive stromata on leaves or inflorescences or form systemic endophytic infections intercellularly within their hosts in addition to forming external reproductive stromata. The Balansieae include the teleomorphic taxa *Atkinsonella*, *Balansia*, *Balansiopsis*, *Epichloë*, and *Myriogenospora* (Diehl, 1950; Luttrell and Bacon, 1977; Rykard et al., 1984). *Myriogenospora* is completely superficial without any penetration of the host epidermis (Luttrell and Bacon, 1977; White and Glenn, 1994), but *Atkinsonella* has some localized intercellular hyphae associated with the stroma (Leuchtmann and Clay, 1988; Morgan-Jones and White, 1989). *Balansia* contains a few epibiotic species (Leuchtmann and Clay, 1988; Clay and Frenz, 1993), but most species have a well de-

veloped endophytic habit (White et al., 1995). All species of the *Epichloë typhina* (Pers. : Fr.) Tul. group, including the apparent asexual derivatives of *Epichloë*, are intercellular grass endophytes (White, 1994b). Asexual grass endophytes were largely overlooked as being related to *Epichloë* until Bacon et al. (1977) found intercellular hyphae growing within tall fescue. They referred to the fescue endophyte as a biotype of *E. typhina* based on concurrent *in vitro* examinations and the earlier report by Sampson (1933). The obscurity of these asexual endophytes was a result of the host grasses never showing any symptoms or signs of a fungal infection. For the anamorph of the fungus, Bacon et al. (1977) applied the name *Sphacelia typhina* Sacc. which was the binomial designated by Saccardo (1881) for the anamorph of *E. typhina*. However, *S. typhina* was later considered a *nomen dubium* by Morgan-Jones and Gams (1982) because of insufficient type material. *Typhodium* (Link, 1826) is another genus of uncertain application to the anamorph of *E. typhina* and is now considered a synonym of *Epichloë* (Hawksworth et al., 1983). However, as a "convention of convenience," Diehl (1950) applied *Typhodium* as a form genus and used the term "typhodial" when referring to the anamorph of *Epichloë*.

Based on *in vitro* similarities, Morgan-Jones and Gams (1982) placed the anamorphs of both the fescue endophyte and *E. typhina* in the form genus *Acremonium*. The authors erected the new sect. *Albolanosa* for these endophytic fungi with the *in vitro* characteristics of slow-growing white to yellow colonies, solitary phialides narrower at the base than the subtending hyphae, hyaline and smooth-walled conidia, and teleomorphs in the Clavicipitaceae. They stated that the exclusively solitary phialides of the endophytes preclude them from classification in *Verticillium* sect. *Prostrata* which was restricted to those taxa, such as *Cordyceps* spp., that possess verticillate as well as solitary phialides (see also Gams, 1971). Morgan-Jones and Gams (1982) reported slight *in vitro* morphological differences between the anamorph of *E. typhina* and those of *Holcus mollis* L., *Dactylis glomerata* L., and *Sphenopholis obtusata* (Michx.) Scribn., which they treated as *A. typhinum* Morgan-Jones & W. Gams, and the tall fescue endophyte, which they described as the separate species *A. coenophialum* Morgan-Jones & W. Gams.

Acremonium is a cosmopolitan, morphologically simple genus (see Gams, 1971). *Acremonium alternatum* Link : Fr. is the lectotype and produces conidia in chains or heads. The genus *Cephalosporium* has often been distinguished from *Acremonium* because of the accumulation of conidia in slimy heads. However, Gams (1971) concluded heads versus chains was

untenable as a generic character and merged the two under the older *Acremonium*. The form genus *Gliomastix* and monophialidic species of *Paecilomyces* were also merged into *Acremonium* (Gams, 1971), thereby expanding its definition to comprise generally slow-growing species with hyaline or pigmented conidia that are one-celled or exceptionally two-celled, in chains or slimy heads, and produced from orthophialides or basitonously branched conidiophores.

Acremonium is perceived to be a heterogeneous taxon because several morphologically distinct teleomorphic genera have *Acremonium*-like anamorphs. Most of the known teleomorphs of *Acremonium* are species of *Nectria* (Gams, 1971; Samuels, 1973, 1976a,b; Lowen, 1995), but teleomorphic genera such as *Hypocrea*, *Hypomyces*, *Thielavia*, *Pronectria*, *Nectriopsis*, *Epichloë*, *Emericellopsis*, *Mycocarachis*, and *Nigrosabulum* also have anamorphs classified in *Acremonium* (Malloch and Cain, 1970; Gams, 1971; Morgan-Jones and Gams, 1982; Samuels, 1988; Lowen, 1995). Gams (1971, 1975) divided *Acremonium* into three sections: *Acremonium*, *Nectrioidea*, and *Gliomastix*. The majority of the species in these sections are saprobic in a wide variety of habitats, but some are plant parasites. Some species of sect. *Nectrioidea* have teleomorphs in the genus *Nectria*, but most of the species in the three sections do not have any known association to teleomorphs. Along with the establishment of sect. *Albolanosa*, Morgan-Jones and Gams (1982) also erected the new sect. *Chaetomioides* for anamorphs of ascomycetes within the family Chaetomiaceae that have, among other characteristics, short-aculeate to lageniform phialides. *Acremonium alabamense* Morgan-Jones, the type species of sect. *Chaetomioides*, is a thermophilic fungus originally isolated from fallen pine needles (Morgan-Jones, 1974), and it is the anamorph of *Thielavia terrestris* (Apinis) Malloch & Cain (Morgan-Jones and Gams, 1982). Other species of this section are anamorphs of several *Chaetomium* species and cannot be distinguished without their teleomorphs (Morgan-Jones and Gams, 1982). Most recently, sect. *Lichenoida* was erected for lichenicolous species (Lowen, 1995), and some of these species have hypocreaceous teleomorphs while others are not associated with any teleomorph.

Because of the possible heterogeneity of *Acremonium*, the placement of the grass endophytes in sect. *Albolanosa* was done with no claim of phylogenetic relationship to the nonendophytic species in the other sections (Morgan-Jones and Gams, 1982; White and Morgan-Jones, 1987). This classification scheme was based strictly on *in vitro* morphological similarities. Some disagreement with this classification of the endophytes was initially expressed, and it continues

TABLE I. *Acremonium* taxa included in the analyses and the various sections in which they are classified^a

Sect. <i>Acremonium</i> ^{b,c}	Sect. <i>Nectrioidae</i> ^b	Sect. <i>Gliomastix</i> ^b	Sect. <i>Albolanosa</i> ^d	Sect. <i>Chaetomioides</i> ^d
<i>A. alternatum</i>	<i>A. rutilum</i>	<i>A. murorum</i>	<i>A. coenophialum</i>	<i>A. alabamense</i> (= <i>Thielavia terrestris</i>)
<i>A. kiliense</i>	<i>A. chrysogenum</i>		<i>A. uncinatum</i> <i>Epichloë festucae</i> (= <i>A. typhinum</i>)	
<i>A. strictum</i> <i>Emericellopsis terricola</i> (= <i>Acremonium</i> sp.) <i>Emericellopsis minima</i> (= <i>Acremonium</i> sp.)	<i>A. furcatum</i> <i>Nectria vilior</i> ^e (= <i>A. berkeleyanum</i>)		<i>Epichloë amarillians</i> (= <i>A. t.</i> var. <i>typhinum</i>)	

^a Teleomorphs are indicated if they are known, and precedence in the listing is given to the teleomorphs except for the case of *A. alabamense* which was obtained from ATCC (#26796) as *A. alabamense* and not as *T. terrestris*.

^b Gams (1971).

^c Gams (1975).

^d Morgan-Jones and Gams (1982).

^e See Samuels et al. (1991) and Gams and Van Zaayen (1982).

to be debated (Latch et al., 1984; Rykard et al., 1984; Morgan-Jones et al., 1992; Gams, 1995). The simple morphology of the anamorphs may have been derived multiple times in the evolution of fungi. However, the connection of so many *Acremonium* species to *Nectria* and other genera of the Hypocreaceae suggests that while there may be some heterogeneity created by sect. *Albolanosa* and sect. *Chaetomioides*, *Acremonium* may be more homogeneous than postulated.

The application of strict monophyly has been suggested for fungal systematics and taxonomy so as to reflect the phylogenetic history of a group of organisms (Vilgalys et al., 1993). If a monophyletic classification scheme is to be the ultimate goal for any group of fungi, then the current delimitation of *Acremonium* is open to question. Molecular phylogenetic data are particularly helpful in resolving relationships of morphologically simple organisms such as those in *Acremonium*. Therefore, using sequences of the nuclear encoded small subunit ribosomal DNA (18S rDNA), this project was undertaken to 1) assess the minimum number of separate lineages currently placed within the form genus *Acremonium*, 2) determine which groups of sexual ascomycetes are closely affiliated with *Acremonium* species in the current classification, 3) determine if *Acremonium* sect. *Albolanosa* is appropriate for classification of the grass endophytes, and if not, 4) propose a new taxon for this group.

MATERIALS AND METHODS

Fungal isolates.—A total of fifteen taxa having anamorphs currently classified in *Acremonium* were selected from sections *Acremonium*, *Albolanosa*, *Chae-*

tomioides, *Gliomastix*, and *Nectrioidae* for inclusion in the rDNA comparisons (TABLES I, II). Isolates from sect. *Lichenoidea* were not sampled because of the recent establishment of this newest section. All isolates of *Acremonium* were either type or authenticated cultures, and additional taxa were either personally collected or obtained from other researchers (TABLE II). The fungi were maintained on cornmeal-malt agar (CMM) (Rykard et al., 1982) at room temperature.

Nucleic acid extraction.—For extraction of total genomic DNA from mycelium, all isolates were grown in M102 liquid medium (Rykard et al., 1982) on a rotary shaker (200 rpm) at room temperature until adequate growth occurred (usually 1–2 wk). The mycelium was collected by centrifugation, and the pelleted tissue was washed once with sterile distilled water to remove excess medium. The tissue was then ground in liquid nitrogen and stored at –80 C until ready for nucleic acid extraction. Extractions were made from 0.2–0.5 g (wet weight) of ground mycelium.

The nucleic acid extraction procedure was a modification of Lee and Taylor (1990) in which the chloroform:phenol step was repeated at least once so as to remove as much cellular debris as possible. The extracted nucleic acid samples were diluted to provide solutions with a DNA concentration range of 0.1 to 1.0 ng μL^{-1} .

Polymerase chain reaction and sequencing.—Fifty μL of each diluted sample was used as template for polymerase chain reactions (PCR) (Mullis and Fallona, 1987; Saiki et al., 1988). For each amplification, a total reaction volume of 100 μL was made containing diluted template, 10 mM Tris-HCl (pH 8.3), 50 mM

TABLE II. Taxa included in the analyses, source of isolates, and accession numbers for 18S rDNA sequences

Taxon	Isolate source ^a	GenBank accession #
<i>Acremonium alabamense</i>	ATCC 26796 ^b	U43969
<i>Acremonium alternatum</i>	CBS 223.70 & 406.66	U43970
<i>Acremonium chrysogenum</i>	ATCC 14615 ^b	U43971
<i>Acremonium coenophialum</i>	ATCC 52274 ^b ; <i>Festuca arundinacea</i>	U45942
<i>Acremonium furcatum</i>	CBS 122.42 ^b & 508.65	U43972
<i>Acremonium kiliense</i>	ATCC 34716 ^b	U43973
<i>Acremonium murorum</i>	CBS 214.69	U43966
<i>Acremonium rutilum</i>	CBS 225.70 & 394.66	U43967
<i>Acremonium strictum</i>	ATCC 34717 ^b	U43968
<i>Acremonium uncinatum</i>	J. F. White; <i>Festuca pratensis</i>	U45943
<i>Ascospaera apis</i>	UCB 78-018	M83264
<i>Atkinsonella hypoxylon</i>	A. E. Glenn; <i>Danthonia spicata</i>	U44034
<i>Balansia aristidae</i>	J. F. White; <i>Aristida</i> sp.	U44035
<i>Balansia henningsiana</i>	A. E. Glenn; <i>Andropogon</i> sp.	U44036
<i>Balansia obtecta</i>	J. F. White; <i>Cenchrus echinatus</i>	U44037
<i>Balansia sclerotica</i>	ATCC 16582	U32399
<i>Balansia strangulans</i>	J. F. White; <i>Panicum</i> sp.	U44038
<i>Candida tropicalis</i>	MUCL 30002	M55527
<i>Ceratocystis fimbriata</i>	T. C. Harrington C89	U32418
<i>Chaetomium globosum</i>	A. E. Glenn	U44039
<i>Claviceps purpurea</i>	R. T. Hanlin; <i>Dactylis glomerata</i>	U44040
<i>Cordyceps capitata</i>	A. E. Glenn; <i>Elaphomyces</i> sp.	U44041
<i>Cryphonectria parasitica</i>	R. T. Hanlin 5806	Denise Silva, unpubl.
<i>Daldinia concentrica</i>	ATCC 36659	U32402
<i>Diaporthe phaseolorum</i>	F. A. Uecker 458	L36985
<i>Echinodothis tuberiformis</i>	J. F. White; <i>Arundinaria tecta</i>	U44042
<i>Emericellopsis minima</i>	ATCC 16216	U44043
<i>Emericellopsis terricola</i>	CBS 120.40 ^b	U44112
<i>Epichloë amarillians</i>	J. F. White; <i>Agrostis hiemalis</i> (Ah3)	U35034
<i>Epichloë festucae</i>	J. F. White; <i>Festuca rubra rubra</i>	U44113
<i>Eurotium rubrum</i>	UCB 88-016	U00970
<i>Hypocrea pallida</i>	G. J. Samuels 89-83	U32408
<i>Hypomyces polyporinus</i>	ATCC 46844	U32410
<i>Microascus trigonosporus</i>	RSA 1942	L36987
<i>Monascus purpureus</i>	ATCC 16365	M83260
<i>Myriogenospora atramentosa</i>	A. E. Glenn; <i>Andropogon</i> sp.	U44114
<i>Myriogenospora atramentosa</i>	A. E. Glenn <i>Erianthus brevibarbis</i>	U44115
<i>Nectria cinnabarina</i>	G. J. Samuels 89-107	U32412
<i>Nectria vilior</i>	ATCC 16217	U44116
<i>Neocosmospora vasinfecta</i>	ATCC 28867	U44117
<i>Saccharomyces cerevisiae</i>	—	M27607
<i>Taphrina deformans</i>	ATCC 34556	U00971
<i>Xylaria hypoxylon</i>	ATCC 42768	U20378

^a ATCC = American Type Culture Collection, USA; CBS = Centraalbureau voor Schimmelcultures, Netherlands; MUCL = Mycothèque de l'Université Catholique, Louvain-la-Neuve, Belgium; RSA = Rancho Santo Ana, USA; UCB = University of California, Berkeley Microgarden, USA.

^b Type culture.

KCl, 2.5 mM MgCl₂, 0.5 μM of each primer, 200 μM of each dNTP, and 2.5 units of AmpliTaq[®] DNA polymerase (Roche Molecular Systems, Inc., Branchburg, NJ). Each reaction mixture was topped off with a thin layer of mineral oil and amplified using a Perkin-Elmer DNA Thermal Cycler 480 (Norwalk, CT).

A segment of the small subunit ribosomal RNA

gene (18S rDNA) was amplified using primers NS1 and NS6 (White et al., 1990). Thermal cycling parameters for amplification consisted of one initial cycle with denaturation at 95 C for 5 min, annealing at 54 C for 30 s, and extension at 72 C for 45 s. This cycle was followed by 38 cycles with denaturation at 95 C for 30 s, annealing at 54 C for 30 s, and exten-

sion at 72 C for 45 s. (plus 4 sec addition to extension segment per cycle). A final cycle was performed with an extension segment of 72 C for 10 min.

Amplified products were separated from unincorporated nucleotides and primers using either minicolumns (WizardTM PCR Preps, Promega Corp., Madison, WI) or microconcentrators (MicroconTM 100, Amicon, Inc., Beverly, MA) following each manufacturer's protocol. Purified samples were sequenced by the Molecular Genetics Facility of the University of Georgia (Athens, GA) using an Applied Biosystems automated sequencer (model 373A, version 1.2.1). Primers NS1, NS2, and NS3 (White et al., 1990) were used for sequencing the 18S rDNA. Primers NS1 and NS2 provide complementary sequences, but the complementary sequence to NS3 was not determined. The rDNA sequences were easily aligned by direct examination.

Data analysis.—Maximum parsimony analysis of the aligned sequences was conducted using PAUP v. 3.1.1 (Swofford, 1993) on a Macintosh Performa 6115CD. Alignment gaps were treated as missing data (GAPMODE=MISSING). However, one gap was included as an additional character in the data matrix (absence and presence of gap were coded as 0 and 1, respectively). In total, 43 isolates were included in the analysis (TABLE II). As a result of the large data set, only heuristic searches were performed with the following options in effect: tree-bisection-reconnection (TBR) swapping algorithm, collapsing zero length branches, and saving all minimal length trees (MULPARS). Ten replications with random addition of taxa were performed for each heuristic search in order to find any additional islands of minimum length trees (Maddison, 1991). To measure the relative support and stability of the resulting clades, bootstrap values (Efron, 1982; Felsenstein, 1985) and decay indices (Bremer, 1988; Donoghue et al., 1992) were calculated using PAUP v. 3.1.1. The computational intensity of these two values is directly dependent on the number of taxa and characters. Executing these computations on all of the taxa included was beyond the capability of the computer. Therefore, selected taxa were excluded, and the bootstrap values and decay indices were calculated on the smaller representative selection. Bootstrapping was performed with 250 replications. Decay indices up to 4 steps longer than the most parsimonious trees were determined. *Saccharomyces cerevisiae* E. C. Hansen, *Taphrina deformans* (Berk.) Tul., and *Candida tropicalis* (Castellani) Berkhout were used as outgroup taxa based on the results of previous analyses at a broader taxonomic scale by Berbee and Taylor

(1992), Bruns et al. (1992), and Spatafora and Blackwell (1993).

Phenetic analyses were also performed on the aligned 18S rDNA sequences using MEGA v. 1.01 (Kumar et al., 1993). Since MEGA isn't capable of evaluating binary code, the alignment gap was not included as an additional character in the data set. For neighbor-joining analyses (Saitou and Nei, 1987), the gamma (Kimura 2-parameter) distance method was used with complete deletion of all sites containing gaps or missing information. Bootstrapping of the neighbor-joining tree was performed with 500 replications. Again, *S. cerevisiae*, *T. deformans*, and *C. tropicalis* were used as outgroup taxa.

RESULTS

Segments of the *ssrRNA* gene (18S rDNA) encompassing 937 bp were analyzed. Alignment of sequences was easily accomplished by direct examination due to the gene's conserved nature. An alignment gap of one base pair, common to the outgroup taxa and the Clavicipitaceae, was added to the data set as an additional character. Of the 43 taxa included in this study (TABLE II), sequences from 16 of these were obtained from either GenBank or other researchers. The remaining 27 were sequenced for this study. Our sequencing efforts focused primarily on the form genus *Acremonium* and the ascomycete family Clavicipitaceae. Aligned sequences are available from the authors upon request.

A maximum parsimony analysis with all taxa included yielded eight equally parsimonious trees of 656 steps. For each of these trees, the consistency index (CI) was 0.585 with autapomorphies included, the retention index (RI) was 0.712, and the rescaled consistency index (RC) was 0.417. The strict consensus of the eight trees is presented in FIG. 1. In order to determine bootstrap and decay values for the broad range of taxa included in this study, a smaller subset of selected taxa was chosen to represent the various clades, and the support values were computed on the smaller collection of taxa. The maximum parsimony strict consensus cladogram of this subset is presented in FIG. 2 along with the bootstrap and decay indices. This same subset of taxa was also analyzed using the neighbor-joining method, and the results are presented in FIG. 3.

Based on these results, *Acremonium* is a polyphyletic taxon having species associated with at least three or four currently recognized ascomycete orders (FIG. 1). First, *A. furcatum* F. & V. Moreau : W. Gams is potentially associated with *Microascus* and *Ceratocystis* of the order Microascales. The bootstrap values for this connection are 67% using maximum parsimony.




FIG. 1. Strict consensus of eight cladograms (each of 656 steps, CI = 0.585, RI = 0.712, RC = 0.417) resulting from maximum parsimony analysis of 18S rDNA sequences of all taxa included in the study. Orders of the Euascomycetes are indicated. Black dots indicate species having *Acremonium*-like anamorphs.

mony (FIG. 2) and 83% with the neighbor joining analysis (FIG. 3). Second, *A. alabamense* is the known anamorph of *T. terrestris* of the Sordariales (see TABLE I), and our results definitively show this anamorph is strongly associated with *Chaetomium* from the Sordariales. Both maximum parsimony and neighbor-joining analyses resulted in a 100% bootstrap value for the *A. alabamense*-*Chaetomium* clade (FIGS. 2 and 3, respectively). Third, *Epichloë* and related asexual endophytes are well known members of the order Clavicipitales (FIG. 1), and their anamorphs are currently classified in *Acremonium* sect. *Albolanosa* (see TABLE I). Fourth, a large number of *Acremonium* species show affiliation to the Hypocreales (FIG. 1). Six conidial fungi classified in *Acremonium*, two *Emericellopsis* species which have *Acremonium* anamorphs, and *Nectria vilior* Starbäck, which also has an *Acremonium* anamorph (see TABLE I), are all placed within the moderately well supported clade comprising the Hypocreales. Based on maximum parsimony analysis, the Hypocreales had a 76% bootstrap value and a decay index of +2 (FIG. 2). MEGA's neighbor-joining analysis gave a bootstrap value of 87% for the comparable clade (FIG. 3).

Among the *Acremonium* species having affiliation to the Hypocreales is the type species of the genus, *A. alternatum*. It was weakly associated with other taxa such as *A. kiliense* Grütz, *A. strictum* W. Gams, *Emericellopsis minima* Stolk, and *E. terricola* J. F. H. Beyma. The parsimony clade of these five taxa had a bootstrap of only 59% and a decay index of +1 (FIG. 2). The phylogenetic placement of the cleistothecial *Emericellopsis* in the Hypocreales is supported here by the 18S rDNA sequences of both *E. terricola* and *E. minima*.

The monophyletic Clavicipitales appears to have been derived from within the Hypocreales (FIGS. 1-3). Monophyly of the Clavicipitaceae, the only family in the order Clavicipitales, was supported by neighbor-joining analysis with an 89% bootstrap value (FIG. 3), and support from maximum parsimony analysis was only slightly weaker with a bootstrap of 85% and a decay index of +3 (FIG. 2).

As indicated in FIGS. 2 and 3, *Epichloë* and its asexual derivatives in *Acremonium* sect. *Albolanosa* formed a well supported, monophyletic group with 84% and 96% bootstrap values, respectively. *Atkinsonella*, *Balansia*, and *Myriogenospora* formed a sister group to




FIG. 2. Strict consensus of two cladograms (each of 527 steps, CI = 0.639, RI = 0.727, RC = 0.465) resulting from maximum parsimony analysis of 18S rDNA sequences of a selected subset of taxa. Consensus cladogram is depicted with proportional branch lengths. Scale bar indicates the number of nucleotide changes per unit of measure. Bootstrap values greater than 50% are indicated above internodes. Decay indices are indicated below internodes. Black dots indicate species having *Acremonium*-like anamorphs. E = grass endophyte; e = grass epibiont.




FIG. 3. Neighbor-joining phenogram resulting from analysis of 18S rDNA sequences of a selected subset of taxa. Bootstrap confidence measures greater than 50% are indicated at internodes. Symbols are the same as in FIG. 2.

the *Epichloë* clade (FIGS. 1), but little support was available for resolving these relationships.

In general, there was a high degree of congruence between the resulting phylogenies of the cladistic (FIG. 2) and phenetic (FIG. 3) analyses. Relationships within the Hypocreales differed only in the placement of *N. vilior* and *A. chrysogenum* (Thirum. & Sukapure) W. Gams. Placement of the Xylariales also differed between the two analyses. Bootstrap values for comparable clades in the two resulting phylogenies were generally equivalent. However, three clades did show distinctly different values: the monophyletic clade comprising the Sordariales and Diaporthales, the Hypocreales clade, and the clade containing *A. furcatum* and the Microascales. These differences are probably inherent in the individual algorithms of the synapomorphy-based cladistical analysis and the similarity-based phenetic analysis and how they each affect the areas of the trees where the degree of support for the topology is limited.

DISCUSSION

New taxon and combinations.—Based on the 18S rDNA sequence analyses presented here, the present classification of the anamorphs of *Epichloë* and related mutualists in *Acremonium* is untenable. *Acremonium*, as typified by *A. alternatum*, appears to be restricted to the family Hypocreaceae. *Epichloë* and related genera of the Clavicipitaceae form a well circumscribed, monophyletic family that appears to be derived from within the Hypocreales. The monophyly of this family is supported by the unique morphology, ecology, and obligate parasitism of the Clavicipitaceae. The anamorphs of this family are unusual in that they are associated with stromatic tissue formed on a host (Diehl, 1950; Luttrell, 1980; Rykard et al., 1984; White and Morgan-Jones, 1987; Leuchtmann and Clay, 1988; Morgan-Jones and White, 1989). Therefore, we are proposing the erection of a new genus to accommodate anamorphs of clavicipitaceous fungi that form a palisade of simple phialidic conidiogenous cells over the surface of a stroma and in culture produce simple, strictly aculeate phialides often lacking a basal septum and usually arising as lateral branches from aerial hyphae. As thus defined, this genus should include the conidial fungi currently classified in *Acremonium* sect. *Albolanosa*.

Neotyphodium Glenn, Bacon & Hanlin gen. nov.

Coloniae albae vel flavidae, lente vel modice crescentes, hyphae aerae saepe abundantes, byssaceae, typice non fasciculatae; phialides solitariae, raro verticillatae, aculeatae, orthotropicae, exoriundae ex hyphis aeriis, basi plerumque

septo carente. Conidia oblonga, ellipsoidea vel cylindrica vel fusiformia vel lunata vel uncinata, hyalina, levia, amerospora. Fungi parasitae obligati graminum, aliqui facientes ectostroma hymenio phialidum obtectum. Genus continens anamorphoses Clavicipitacearum.

TYPUS: *Acremonium coenophialum* Morgan-Jones & W. Gams.

Colonies white to yellowish, very to moderately slow growth rate, aerial hyphae often abundant, cottony but usually not fasciculate; phialides solitary, rarely verticillate, aculeate, orthotropic, arising from aerial hyphae, bases frequently lacking a septum. Conidia oblong, ellipsoidal, cylindrical, fusiform, lunate, uncinata, hyaline, smooth-walled, amerosporous. Obligate parasites of grasses, some forming exposed ectostroma with palisade of phialides. Genus containing anamorphs of the Clavicipitaceae.

Etymology. Greek *neos* = new + *Typhodium*, a genus erected by Link (1826) that was of uncertain applicability to the anamorph of *Epichloë*.

Neotyphodium coenophialum (Morgan-Jones & W. Gams) Glenn, Bacon & Hanlin comb. nov.

Basionym. *Acremonium coenophialum* Morgan-Jones & W. Gams, *Mycotaxon* 15: 311. 1982.

No known teleomorph.

Neotyphodium typhinum (Morgan-Jones & W. Gams) Glenn, Bacon & Hanlin comb. nov.

Basionym. *Acremonium typhinum* Morgan-Jones & W. Gams, *Mycotaxon* 15: 311. 1982.

Teleomorphs are species of *Epichloë*.

Neotyphodium lolii (Latch, Christensen & Samuels) Glenn, Bacon & Hanlin comb. nov.

Basionym. *Acremonium lolii* Latch, Christensen & Samuels [as *Acremonium loliae*], *Mycotaxon* 20: 535. 1984.

No known teleomorph.

Neotyphodium chisosum (J. F. White & Morgan-Jones) Glenn, Bacon & Hanlin comb. nov.

Basionym. *Acremonium chisosum* J. F. White & Morgan-Jones, *Mycotaxon* 28: 179. 1987.

No known teleomorph.

Neotyphodium starrii (J. F. White & Morgan-Jones) Glenn, Bacon & Hanlin comb. nov.

Basionym. *Acremonium starrii* J. F. White & Morgan-Jones, *Mycotaxon* 30: 87. 1987.

No known teleomorph.

Neotyphodium huerfanum (J. F. White, G. T. Cole & Morgan-Jones) Glenn, Bacon & Hanlin comb. nov.

Basionym. *Acremonium huerfanum* J. F. White, G. T. Cole & Morgan-Jones, *Mycologia* 79: 148. 1987.

No known teleomorph.

Neotyphodium uncinatum (W. Gams, Petrini & D. Schmidt) Glenn, Bacon & Hanlin comb. nov.

Basionym. *Acremonium uncinatum* W. Gams, Petrini & D. Schmidt, *Mycotaxon* 37: 67. 1990.

No known teleomorph.

Neotyphodium chilense (Morgan-Jones, J. F. White & Piont.) Glenn, Bacon & Hanlin comb. nov.

Basionym. *Acremonium chilense* Morgan-Jones, J. F. White & Piont., *Mycotaxon* 39: 441. 1990.

No known teleomorph.

While not included in our analyses, *N. starrii* and *N. lolii* have been shown to be closely related to *N. coenophialum* based upon maximum parsimony analysis of the two internal transcribed spacer (ITS) regions of rDNA (Schardl et al., 1991). Unfortunately, *N. chilense*, *N. chisosum*, and *N. huerfanum* were also unavailable to us for examination, and these three have yet to be characterized by DNA-based phylogenetic analyses. We do, however, propose the transfer of these species into *Neotyphodium* based upon the examinations and personal communications of James F. White, Jr. and Walter Gams.

The conidial stromata of species of *Epichloë* are easily and logically addressed by the teleomorphic nomen even though anamorphic binomials and varieties have been erected for the various species (Morgan-Jones and Gams, 1982; White, 1992). However, the abundance of asexual endophytes that rarely or never produce teleomorphic stromata [clonal type-III endophytes of White (1988)] means that a dual system of nomenclature must be maintained whereby the type-III endophytes are classified in *Neotyphodium* and the sexually reproducing species are classified in *Epichloë*.

As defined, *Neotyphodium* is the most appropriate form genus for the typhodial stage of *Atkinsonella* (Rykard et al., 1984). However, the applicability of the nomen *Atkinsonella* to all stages of its development is empirical and conveys a holomorphic view of this fungus. Exclusively asexual species are not known. Therefore, since no binomial is currently defined for the typhodial stage of *Atkinsonella*, we do not see the necessity of defining one under *Neotyphodium*. The binomial *Ephelis borealis* Ellis & Everh. has been established for the sympodially proliferating synanamorph of *Atkinsonella* (Diehl, 1950).

The anamorph of *Echinodothis tuberiformis* (Berk. & Ravenel) G. F. Atk. presents a similar problem as found with *Atkinsonella*. The anamorph of this epibiont is similar to *Neotyphodium* in that a palisade of phialidic conidiogenous cells are produced over the surface of a host-associated stroma, but the conidia are different in being didymosporous (White, 1993b). White (1993b) suggested that *Echinodothis* and *Epichloë* were closely related and might deserve classification together in a separate tribe. While lacking statistical support, cladistical analysis suggested that *Echinodothis* may be intermediate between *Cordyceps* and *Claviceps* and is not closely related to *Epichloë* (FIG. 1). As with *Atkinsonella*, the holomorphic application of *E. tuberiformis* to its anamorph and teleomorph means there is no need to establish an anamorphic binomial.

The Clavicipitales-Hypocreales relationship.—The Clavicipitales is a morphologically well circumscribed order of mostly parasitic ascomycetes possessing morphologically distinctive characters such as deliquescent lateral paraphyses, cylindrical asci with a thickened apical cap, and filiform ascospores that are often septate and may disarticulate into part-spores (Rogerson, 1970). However, the ordinal affinity of the clavicipitaceous genera has been evaluated and interpreted differently by several researchers [see Rogerson (1970) for a thorough historical review of the taxonomic literature].

In recent years, the ordinal relationships among hypocreaceous and clavicipitaceous genera has again come into question as a result of phylogenetic evaluations using molecular data. Maximum parsimony analyses using the nuclear encoded 18S rDNA have indicated a monophyletic relationship exists between genera of the Clavicipitales and the Hypocreales, suggesting that all taxa may be classifiable under the single order Hypocreales with the monophyletic Clavicipitaceae being a separate family from the paraphyletic Hypocreaceae (Spatafora and Blackwell, 1993). Our results, which also utilized 18S rDNA, show the same general relationship (FIGS. 1–3). Additionally, gene phylogenies based on both the nuclear encoded large subunit ribosomal DNA (28S rDNA) and nuclear encoded orotidine-5'-monophosphate decarboxylase also support the derivation of the monophyletic Clavicipitaceae from within the Hypocreales (Rehner and Samuels, 1994b, 1995). All these gene phylogenies support the treatment by Kreisel (1969) with a single order incorporating the families Hypocreaceae (*sensu* Rogerson) and Clavicipitaceae.

The expanding Hypocreales.—Just as molecular systematics has aided in clarifying some of the taxonomic confusion concerning the ordinal relationships

