

Warfarin Anticoagulant Therapy: A Southern Italy Pharmacogenetics-Based Dosing Model

Cristina Mazzaccara^{1,2}, Valeria Conti³, Rosario Liguori^{1,2}, Vittorio Simeon⁴, Mario Toriello², Angelo Severini², Corrado Perricone⁵, Alfonso Meccariello⁶, Pasquale Meccariello⁶, Dino Franco Vitale⁷, Amelia Filippelli³, Lucia Sacchetti^{1,2*}

1 CEINGE– Biotecnologie Avanzate s.c.a.r.l., Napoli, Italy, **2** Dipartimento di Medicina Molecolare e Biotecnologie Mediche, Università di Napoli Federico II, Napoli, Italy, **3** Dipartimento di Medicina e Chirurgia, Università di Salerno, Salerno, Italy, **4** Laboratorio di Ricerca Pre-clinica e Traslazionale Oncologica, Centro di Riferimento Oncologico della Basilicata, Istituto di Ricovero e Cura a Carattere Scientifico, Rionero in Vulture (Pz), Italy, **5** A.O.R.N. Santobono Pausillipon, Napoli, Italy, **6** Dipartimento di Medicina Interna, Università di Napoli Federico II, Napoli, Italy, **7** Fondazione Salvatore Maugeri, IRCCS Istituto di Campoli Telese, Benevento, Italy

Abstract

Background and Aim: Warfarin is the most frequently prescribed anticoagulant worldwide. However, warfarin therapy is associated with a high risk of bleeding and thromboembolic events because of a large interindividual dose-response variability. We investigated the effect of genetic and non genetic factors on warfarin dosage in a South Italian population in the attempt to setup an algorithm easily applicable in the clinical practice.

Materials and Methods: A total of 266 patients from Southern Italy affected by cardiovascular diseases were enrolled and their clinical and anamnestic data recorded. All patients were genotyped for CYP2C9*2,*3, CYP4F2*3, VKORC1 -1639 G>A by the TaqMan assay and for variants VKORC1 1173 C>T and VKORC1 3730 G>A by denaturing high performance liquid chromatography and direct sequencing. The effect of genetic and not genetic factors on warfarin dose variability was tested by multiple linear regression analysis, and an algorithm based on our data was established and then validated by the Jackknife procedure.

Results: Warfarin dose variability was influenced, in decreasing order, by VKORC1-1639 G>A (29.7%), CYP2C9*3 (11.8%), age (8.5%), CYP2C9*2 (3.5%), gender (2.0%) and lastly CYP4F2*3 (1.7%); VKORC1 1173 C>T and VKORC1 3730 G>A exerted a slight effect (<1% each). Taken together, these factors accounted for 58.4% of the warfarin dose variability in our population. Data obtained with our algorithm significantly correlated with those predicted by the two online algorithms: Warfarin dosing and Pharmgkb ($p < 0.001$; $R^2 = 0.805$ and $p < 0.001$; $R^2 = 0.773$, respectively).

Conclusions: Our algorithm, which is based on six polymorphisms, age and gender, is user-friendly and its application in clinical practice could improve the personalized management of patients undergoing warfarin therapy.

Citation: Mazzaccara C, Conti V, Liguori R, Simeon V, Toriello M, et al. (2013) Warfarin Anticoagulant Therapy: A Southern Italy Pharmacogenetics-Based Dosing Model. PLoS ONE 8(8): e71505. doi:10.1371/journal.pone.0071505

Editor: Giuseppe Novelli, Tor Vergata University of Rome, Italy

Received: April 14, 2013; **Accepted:** June 30, 2013; **Published:** August 26, 2013

Copyright: © 2013 Mazzaccara et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The present work was supported by grants from CEINGE Regione Campania (DGRC 1901/2009); POR Campania FSE 2007–2013, project CREME. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: sacchett@unina.it

Introduction

Warfarin sodium [1] is the most frequently prescribed anticoagulant for the primary and secondary prevention of thromboembolic disorders worldwide [2–4]. Despite the advent of new oral antithrombotic agents such as dabigatran, rivaroxaban, apixaban, which have proven to be cost-effective compared with warfarin in some clinical conditions [5,6], warfarin remains the mainstay of treatment for patients with mechanical heart valves and patients noncompliant to new therapies because in these populations their efficacy have not been explored [7].

Warfarin inhibits the Vitamin K Epoxide Reductase Complex 1 (VKORC1) thus reducing the activities of vitamin K-dependent clotting factors II, VII, IX and X and coagulation. S-warfarin, the most active of the two (R- and S-) isomers in the administered

drug, is mainly metabolized by the cytochrome P450 2C9 isoenzyme (CYP2C9) [8].

Notwithstanding its wide use, warfarin has a narrow therapeutic range and a large interindividual variability in the dose needed (1–20 mg/day) to obtain an adequate anticoagulation effect [4]. The latter is generally measured by the prothrombin international normalised ratio (INR) and its range is 2.0–3.0 or higher in at-high risk patients [9]. Inappropriate INR levels may result in significant bleeding or stroke (INR levels greater or lower than the target range, respectively), particularly during the first weeks of therapy (induction phase) [9–14]. To date, most clinicians prescribe 3–10 mg/day for the first 2–5 days, then switch to a maintenance dose established based on frequent INR monitoring [2,11,14]. Warfarin-induced adverse effects account for over 10% of all adverse drug reactions leading to hospital admissions [15].

The large interindividual variation in warfarin dose requirement is attributable to clinical, demographic, environmental factors (age, gender, body mass index, daily vitamin K intake, concomitant diseases, interaction between drugs, and smoking), and to genetic factors, which account for 40–60% of the variability [16–18]. Among genetic factors, single nucleotide polymorphisms (SNPs) in the CYP2C9 (Gene Bank Accession Number AY702706; chr.10q24) and in VKORC1 (Gene Bank Accession Number AY587020; chr.16p11.2) genes were first described as major contributors to dose-response variability. Subjects bearing polymorphisms in one or both of these genes require lower or higher warfarin doses than subjects bearing the wild-type genes to obtain an adequate anticoagulant effect [1,8,16,19–22]. More recently, patients bearing a SNP (rs2108622) in the CYP4F2 gene (Gene Bank Accession Number AF22194; chr.19p13.12), which is the vitamin K₁ oxidase involved in vitamin K₁ metabolism, were found to require a warfarin dose slightly higher than normal [23–25] or similar to normal [26,27]. Moreover, a meta-analysis revealed a statistically significant association between rs2108622 and the interindividual warfarin dose variation [28,29]. However, it was annotated (www.pharmgkb.org) as a Level 1B clinical association, namely “a variant-drug combination where the preponderance of evidence shows an association. The association must be replicated in more than one cohort with significant p-values, and, preferably with a strong effect size”.

In 2007 and in 2010, the US Food and Drug Administration, Center for Drug Evaluation and Research, suggested that CYP2C9 and VKORC1 -1639 G>A gene polymorphisms be typed before starting warfarin therapy [30], and issued specific guidelines in this sense [31]. This prompted several clinical trials to evaluate the use of pharmacogenetic tests before starting warfarin therapy. It also prompted the development of warfarin-dosing algorithms that include genetic and non-genetic factors [32]. Notably, predictive algorithms must be based on data representative of the target population, and they should be validated. To date, few studies have evaluated the global effect of genetic and non-genetic factors on warfarin dosage in Italian subjects [1,33–36].

The aim of this study was to estimate, in a Southern Italy population of subjects affected by cardiovascular disorders undergoing warfarin therapy, the effect of the CYP2C9 (*2 and *3), CYP4F2*3, VKORC1 (-1639 G>A, 1173 C>T and 3730 G>A SNPs combined with clinical status, demographic and environmental factors on warfarin dosing.

Results

The clinical, anamnestic and demographic features of our warfarin-treated patients are shown in **Table 1**.

Our population did not differ in terms of gender (55.2% male). Similarly, there were no differences between men and women in terms of age, body mass index and the other parameters evaluated (data not shown).

Cardiac valve replacement and atrial fibrillation were the most frequent cardiovascular indications (43.9% and 38.1%, respectively). Most patients (43.2%) did not assume any drug in addition to warfarin.

Allelic and genotype frequencies of the CYP2C9*2, CYP2C9*3, CYP4F2*3, VKORC1-1639 G>A, VKORC1 1173 C>T and VKORC1 3730 G>A polymorphisms obtained in our patients and those reported in other Caucasian groups are reported in **Table 2**. Genotype frequencies, at the level of all tested genes, were in Hardy-Weinberg equilibrium. The comparison between the weekly warfarin dose assumed in the subjects bearing the wild-

Table 1. Clinical, anamnestic and demographic features of the warfarin-treated patients.

| | |
|--|-------------|
| Age (years) | 67.35±11.05 |
| Gender male | 55.2% |
| BMI (kg/m ²) | 26.90±4.24 |
| Indications for warfarin therapy | |
| Cardiac valve replacement | 43.9% |
| Atrial fibrillation | 38.1% |
| Dilatative cardiomyopathy | 8.5% |
| Deep venous thrombosis | 6.5% |
| Pulmonary embolism | 3.0% |
| Smoking | 8.7% |
| Liver disease | 15.5% |
| Dyslipidaemia | 65.2% |
| Hypertension | 62.5% |
| Drug assumption | 56.8% |
| Only drugs that increase the warfarin effect | 33.0% |
| Only drugs that decrease the warfarin effect | 17.0% |
| Both types of drugs | 6.8% |
| No drugs | 43.2% |
| Warfarin dose assumed (mg/week) | 28.73±13.22 |

Continuous variables are expressed as means ± standard deviation (SD) and categorical variables as percentages.

doi:10.1371/journal.pone.0071505.t001

allele or the polymorphic CYP2C9 variants is shown in **Figure 1A**. Patients bearing the CYP2C9*1/*3, *2/*3 and *3/*3 genotypes required a significantly lower warfarin dose than patients with the wild-type allele (22.03 mg/week±8.80; 13.4 mg/week±10.10; 9.74 mg/week±3.25; respectively *vs* 32.11 mg/week±13.98; *p*<0.001).

The mean weekly warfarin dose was also significantly lower in VKORC1 -1639 G>A mutated homozygotes and in heterozygotes than in patients with the wild-type allele (18.81 mg/week±7.98, 29.15 mg/week±11.79, and 37.80 mg/week±13.37, respectively, *p*<0.001) (**Figure 1B**). We also evaluated the additive effect of the CYP2C9 and VKORC1 -1639 G>A polymorphic genotypes on warfarin dose requirement. The simultaneous presence of these polymorphisms further significantly reduced the warfarin dose requirement as shown in **Figure 2**.

Slightly higher warfarin dosages were required by VKORC1 3730 G>A heterozygotes and homozygotes than by patients carrying the wild-type allele (32.56 mg/week±13.48 and 33.39 mg/week±16.08 *vs* 24.38 mg/week±11.12; *p*<0.05, respectively) (**Figure S1**). No difference in warfarin dosages was observed, by ANOVA analysis, in subjects bearing the VKORC1 1173 C>T (*p* = 0.72) or the CYP4F2*3 (*p* = 0.36) polymorphisms.

Haplotype software showed the lack of linkage disequilibrium between VKORC1-1639 G>A and 1173 C>T (*D'*: 0.186). Using multiple linear regression analysis we assessed the effect of the genetic and non genetic factors (see Table 1) on warfarin dose, with the actual weekly warfarin dose as dependent variable. Using the Jackknife procedure, we then validated the algorithm developed on our data set. The percentage contributions of the various factors on warfarin dose were in decreasing order: 29.7% VKORC1-1639 G>A, 11.8% CYP2C9*3, 8.5% age, 3.5%, CYP2C9*2, 2.0% gender and 1.7% CYP4F2*3. The effects of VKORC1 1173 C>T and VKORC1 3730 G>A were marginal

Table 2. Allele and genotype frequencies of CYP2C9*2, CYP2C9*3, VKORC1 (-1639 G>A, 1173 C>T, 3730 G>A) and CYP4F2 1297G>A polymorphisms obtained in our population and in other Caucasian populations.

| Gene | Genotype frequencies | | | Allelic frequencies | | |
|---------------------------|----------------------|------|----------------|---------------------|----------------|-----------|
| | Our data | | Other studies* | Our data | Other studies* | |
| | N | % | %(min-max) | % | %(min-max) | |
| CYP2C9 | | | | | | |
| *1/*1 | 159 | 60.2 | 56.4–66.9 | *1 | 77.5 | 50.3–83.0 |
| *1/*2 | 58 | 22.0 | 16.4–23.8 | *2 | 15.7 | 11.9–32.0 |
| *1/*3 | 33 | 12.5 | 8.9–12.7 | *3 | 9.8 | 5.7–17.2 |
| *2/*2 | 6 | 2.3 | 1.7–2.3 | | | |
| *2/*3 | 5 | 1.9 | 1.1–3.6 | | | |
| *3/*3 | 3 | 1.1 | 0.3–9.1 | | | |
| CYP4F2 1297G>A | | | | | | |
| G/G | 121 | 45.8 | 39.2–46.0 | G | 69.1 | 65.8–70.3 |
| G/A | 123 | 46.6 | 42.0–48.2 | A | 30.9 | 34.2–29.7 |
| A/A | 20 | 7.6 | 9.4–12.6 | | | |
| VKORC1 -1639G>A | | | | | | |
| G/G | 67 | 25.4 | 32.2–37.3 | G | 50.2 | 58.2–59.4 |
| G/A | 131 | 49.6 | 46.9–55.1 | A | 49.8 | 40.6–41.8 |
| A/A | 66 | 25.0 | 7.6–20.8 | | | |
| VKORC1 1173C>T | | | | | | |
| C/C | 114 | 43.2 | 26.4–40.8 | C | 65.1 | 57.8–62.2 |
| C/T | 116 | 43.9 | 43.2–50.8 | T | 34.9 | 37.8–42.2 |
| T/T | 34 | 12.9 | 8.3–25.0 | | | |
| VKORC1 3730G>A | | | | | | |
| G/G | 132 | 50.0 | 38.2–48.0 | G | 71.4 | 62.6–66.4 |
| G/A | 113 | 42.8 | 39.5–52.7 | A | 28.6 | 37.4–33.6 |
| A/A | 19 | 7.2 | 4.0–15.0 | | | |

*Refs. [33–39].

doi:10.1371/journal.pone.0071505.t002

(<1% each). In our population, the above factors accounted for 58.4% of the variance in warfarin dosage (**Table 3**) and 57.2% after the exclusion of VKORC1 1173 C>T and VKORC1 3730 G>A, which were associated with the lowest and highest doses, respectively.

To explore how our algorithm worked versus the two online algorithms (www.warfarindosing.org and www.pharmgkb.org) (accessed September 2011), we compared by Pearson analysis each patient's predicted warfarin dosage by the Jackknife procedure with those predicted by the two online algorithms (**Figure 3A and 3B**). The data obtained with our algorithm significantly correlated with those predicted by two online algorithms: Warfarin dosing ($p < 0.001$; $R^2 = 0.805$) and Pharmgkb ($p < 0.001$; $R^2 = 0.773$).

Discussion

We investigated the effect of genetic and not genetic factors on the mean weekly warfarin dose variability in an adult South Italian population to setup a simple algorithm easily applicable in clinical practice.

The allele and genotype frequencies of the CYP2C9*2, CYP2C9*3, CYP4F2*3 and VKORC1 (-1639 G>A, 3730 G>A and 1173 C>T) genes were similar to those found in other Caucasian populations, except for a slightly higher

prevalence of the VKORC1 -1639 A/A genotype (25% vs 7.6–20.8%) [33–39].

The effect of the CYP2C9, CYP4F2 and VKORC1 genotypes on warfarin dose was similar to those previously reported [10,12,40]. In particular, the warfarin dose was 17.0% and 32.0% respectively lower in subjects bearing the CYP2C9*2 or CYP2C9*3 polymorphic alleles versus wild type, which is similar to the previously reported reductions of 18–20% and 34–38%, respectively [10,38,40–42] in Caucasians. Furthermore, there was a 25% dose reduction in subjects bearing the VKORC1-1639 A allele, which is also in agreement with previously reported percentages (25–30%) [12]. The mean weekly warfarin dose was also lower in our patients bearing both the CYP2C9 and VKORC1-1639 G>A polymorphic genotypes, namely between 34.8% and 84.0% lower than in wild-type patients, which compares well with previously reported reductions (34%–75% and 41%–79%) [2,30]. In the two previous studies that typed smaller than our Italian populations (148/147 vs 266 patients) [1,35] for VKORC1 1173 C>T but not for -1639 G>A SNP, a different degree of association between VKORC1 1173C>T and warfarin dosing was observed, 0.8% (this study) vs 20% and 13.8% [1,35]. These data support a large population-based variability in gene polymorphism-dependent warfarin dosing.

In our population, we did not find any linkage disequilibrium (D' : 0.186) between VKORC1-1639 G>A and 1173 C>T, in




Figure 1. Relationship between the weekly warfarin dose and CYP2C9 (Panel A) or VKORC1 -1639G>A (Panel B) genotypes. Each box indicates the values from 25^o to 75^o percentile (interquartile range), the horizontal lines represent the median value of weekly warfarin dose, the maximum length of whisker is 1.5 fold the interquartile range. * p<0.05, ** p<0.001 at ANOVA test. doi:10.1371/journal.pone.0071505.g001

contrast with those reported in other ethnic groups [43,44]. Population differences in minor allele frequencies observed at level of the tested polymorphisms VKORC1 -1639G>A and 1173 C>T could drive interethnic differences detected among Caucasian populations, also from different Italian regions [1,35], and

these genetic factors together to different cultural and lifestyle factors could in part explain the above discrepancies. Higher warfarin doses were required by both the heterozygous and mutated homozygous VKORC1 3730 G>A patients with respect to subjects carrying the wild-type allele, which suggests that this




Figure 2. Combined effect of CYP2C9 and VKORC1 -1639 G>A polymorphic genotypes on stable weekly warfarin dose (mg/week).

Each box indicates the values from 25^o to 75^o percentile (interquartile range), the white lines represent the median value of weekly warfarin dose, the maximum length of whisker is 1.5 fold the interquartile range. In detail below are shown the specific statistical significances for each comparison.

CYP2C9 *1/*1+VKORC1 -1639 G/G vs CYP2C9 *1/*1+VKORC1 -1639A/A, $p < 0.001$.

CYP2C9 *1/*1+VKORC1 -1639 G/G vs CYP2C9 *1/*1+VKORC1 -1639G/A, $p < 0.05$.

CYP2C9 *1/*1+VKORC1 -1639 G/A vs CYP2C9 *1/*1+VKORC1 -1639A/A, $p < 0.001$.

CYP2C9 *1/*2+VKORC1 -1639 G/G vs CYP2C9 *1/*2+VKORC1 -1639A/A, $p < 0.001$.

CYP2C9 *1/*2+VKORC1 -1639 G/G vs CYP2C9 *1/*2+VKORC1 -1639G/A, $p < 0.05$.

CYP2C9 *1/*2+VKORC1 -1639 G/A vs CYP2C9 *1/*2+VKORC1 -1639A/A, $p < 0.001$.

CYP2C9 *1/*3+VKORC1 -1639 G/G vs CYP2C9 *1/*3+VKORC1 -1639A/A, $p < 0.05$.

CYP2C9 *1/*3+VKORC1 -1639 G/A vs CYP2C9 *1/*3+VKORC1 -1639A/A, $p < 0.05$.

doi:10.1371/journal.pone.0071505.g002

polymorphism has less impact on warfarin dosage than VKORC1-1639 G>A, in agreement with the meta-analysis reported by Yang et al. [38].

In the regression model, the variant CYP4F2*3 polymorphism entered with an R^2 of 1.7%, and the difference in warfarin dose between CYP4F2 A/A *vs* CYP4F2 wild type was 0.6 mg/day. This observation is in line with a previous finding that CYP4F2*3 has only a small effect on warfarin dose variability [23]. However, the effect of CYP4F2*3 on warfarin dose-response variability is debatable; in fact, it ranges from 1%–7% [23,33,34] to not significant [26,27].

Among non genetic factors, regression analysis revealed that age ($p < 0.0001$; $R^2 = 8.5\%$) and gender ($p = 0.001$; $R^2 = 2.0\%$) contributed to the overall variability in warfarin dose, which is in agreement with a previous report [34]. Warfarin dosages predicted by our algorithm significantly correlated with those predicted by the Warfarindosing and Pharmgkb algorithms. The algorithms explain

similar to ours the percentages of the warfarin response (47%–58%) in other Caucasian populations [45–46], but they use more data.

In conclusion, by exploring the most relevant genetic variants and by applying a user-friendly algorithm, our study contributes to the field of warfarin pharmacogenetics in a Southern Italy population. One may envisage that a genotype-guided and clinical-guided (versus clinical-guided) warfarin dosing algorithm could improve patient care in terms of dosage particularly in the initial phase of therapy, resulting in a decreased time below the therapeutic range and consequently in a reduction of adverse drug reactions.

Materials and Methods

Subjects

Two hundred and sixty-six warfarin-treated patients from Southern Italy, 45% female, were enrolled at the Department of

Table 3. Factors affecting weekly warfarin dose requirements in regression model*.

| Variable | P value | Partial R ² | Coefficient B (95% CI) |
|----------------------------|---------|------------------------|-------------------------|
| VKORC1 -1639 G>A | | | |
| -1639 G/A | 0.439 | | -0.033 (-0.116, -0.050) |
| -1639 A/A | <0.0001 | | -0.278 (-0.335, -0.221) |
| CYP2C9*3 | | | |
| (*1/*3) | <0.0001 | 11.8 | -0.149 (-0.201, -0.097) |
| (*3/*3) | <0.0001 | | -0.475 (-0.672, -0.278) |
| AGE | | | |
| | <0.0001 | 8.5 | -0.050 (-0.007, -0.003) |
| CYP2C9*2 | | | |
| (*1/*2) | <0.0001 | 3.5 | -0.090 (-0.132, -0.047) |
| (*2/*2) | 0.009 | | -0.218 (-0.381, -0.054) |
| GENDER | | | |
| | 0.001 | 2.0 | 0.061 (0.024,0.098) |
| CYP4F2*3 | | | |
| (*1/*3) | 0.121 | 1.7 | 0.030 (0.008,0.069) |
| (*3/*3) | 0.015 | | 0.087 (0.017,0.158) |
| VKORC1 1173 C>T | | | |
| 1173 C/T | 0.037 | 0.8 | -0.085 (-0.165, -0.050) |
| 1173 T/T | 0.437 | | -0.022 (-0.079,0.034) |
| VKORC1 3730 G>A | | | |
| 3730 G/A | 0.022 | 0.4 | 0.054 (0.008,0.100) |
| 3730 A/A | 0.100 | | 0.060 (-0.012,0.132) |

*(Total R² for the model 58.4%).

doi:10.1371/journal.pone.0071505.t003

Internal Medicine, University of Naples Federico II, at the Foundation Salvatore Maugeri IRCCS Institute of Campoli Teleso, Benevento, at the Santobono Pausilipon Hospital, Naples, and at the Department of Experimental Medicine, Second University of Naples, Italy. The study was performed according to the second Helsinki Declaration, all subjects provided written informed consent to participate in the study which was approved by the

Ethics Committees of the above institutions. At enrolment all patients had been taking a stable dose of warfarin for at least 3 months, which is warfarin dose to achieve INR 2–3. Anamnestic, clinical and lifestyle information were recorded on a structured interview form. Hypertension, systolic blood pressure above 130 mmHg and diastolic blood pressure above 85 mmHg, and body mass index (body weight [kg] divided by squared height [m²]) were also recorded. Liver dysfunction (aspartate aminotransferase >35 U/L women, >40 U/L men; alanine aminotransferase >35 U/L women, >40 U/L men), and dyslipidemia (serum total cholesterol and/or triglycerides levels above 190 mg/dL or 150 mg/dL, respectively) were also measured.

Samples and Methods

Three fasted blood samples (one with EDTA for DNA extraction, one with sodium citrate and one without anticoagulant for haematological and biochemical investigation, respectively) were collected from each patient. DNA was extracted with the Nucleon BACC2 kit (Amersham Life Science, England). Coagulation and biochemical tests were performed by routinely methods using reagent and equipment from Siemens, (Germany) and from Roche Diagnostics (Germany), respectively.

We genotyped patients for the CYP2C9 (CYP2C9*2, rs1799853, exon 3, c.430 C>T, p.Arg144Cys; CYP2C9*3, rs1057910, exon 7, c.1075 A>C, p.Ile359Leu), CYP4F2*3 (rs2108622, c.1297G>A, p.V433M) and VKORC1 -1639 G>A (rs9923231) (also known as 3673 G>A) polymorphisms, together with positive and negative quality control samples, using the Real-Time TaqMan method [47,48] and commercial kits, namely Pre-developed TaqMan Assay Reagents Human Allelic Discrimination (CYP2C9*2 and *3) (probe code 4312568 and 4312569) and TaqMan Drug Metabolism Genotyping Assay (CYP4F2*3 and VKORC1) Applied Biosystems, CA, USA.

The PCR was set up in a 96-well plate with a 25 µL mix reaction, 10–20 ng of genomic DNA per assay. The amplification protocol was performed according to the manufacturer's indications. Variants VKORC1 1173 C>T, (rs9934438) (also known as 6484 C>T) and VKORC1 3730 G>A, (rs7294) (also known as 9041G>A) were detected by denaturing high performance liquid chromatography on Wave 2.0 Transgenomic instruments (Oma-




Figure 3. Correlation analysis (Pearson coefficient) between our predicted warfarin dosage by Jackknife and that predicted by www.warfarindosing.org (panel A, $p < 0.001$, $R^2 = 0.805$) and www.pharmgkb.org (panel B, $p < 0.001$, $R^2 = 0.773$). The solid line represent a good correlation between the two doses. Open circles represent individuals for whom no variants in CYP2C9 and VKORC1 -1639 G>A were detected. Gray circles represent individuals with only one polymorphic gene (either VKORC1 -1639 G>A or CYP2C9*2 or *3). Black circles represent individuals with at least two polymorphic genes.
doi:10.1371/journal.pone.0071505.g003

ha, NE, USA). Each suspicious chromatogram was then sequenced. The PCR primers and conditions are listed in **Appendix S1**.

Statistical Analysis

The Hardy-Weinberg equilibrium was verified for all investigated polymorphisms by the χ^2 test.

The Kolmogorov-Smirnov test was performed to evaluate the distribution of continuous variables.

Data were expressed as average \pm standard deviation (SD) (continuous variables) or in percentage (categorical variables). We evaluated differences of clinical and genetic variables among groups by the Student *t* test and analysis of variance (ANOVA), followed by post hoc test with Bonferroni correction. A $p < 0.05$ was considered statistically significant. Linkage analysis was performed by Haploview 4.0 software [49]. Multivariate linear regression was performed to identify the factors associated with the weekly warfarin dose expressed on a logarithmic scale. Global and partial R² were measured, these latter assessing the percentage of the dose variability explained by the full model and by each factor included in the model. In order to obtain an unbiased estimate of the prediction ability of our algorithm we validated it using the Jackknife procedure [50], i.e., the predicted dose of each patient was obtained using the linear coefficients developed using the remaining patients in the data set, thus avoiding the bias introduced by scoring a patient with coefficients optimized with data of the patient himself. Weekly warfarin dose predictions were also obtained by two dosing algorithms published by the Warfarin Dose Refinement Collaboration (www.warfarindosing.org, accessed September

2011) and by the International Warfarin Pharmacogenetics Consortium (www.pharmgkb.org, accessed September 2011). These predicted doses were then correlated with those obtained by our validated algorithm. Statistical analysis was performed with the STATA 11.2 software (StataCorp LP).

Supporting Information

Appendix S1 Specific PCR primers (for: VKORC1 1173 C>T and VKORC1 3730G>A) and conditions. (DOC)

Figure S1 Relationship between the weekly warfarin dose and VKORC1 3730 G>A genotypes. Each box indicates the values from 25° to 75° percentile (interquartile range), the black central line represents the median value of weekly warfarin dose, the maximum length of whisker is 1.5 fold the interquartile range. * $p < 0.05$. (PDF)

Acknowledgments

We thank Jean Ann Gilder (Scientific Communication srl) for revising and editing the manuscript.

Author Contributions

Conceived and designed the experiments: LS AF CM PM. Performed the experiments: CM VC MT VS AS RL AM LS. Analyzed the data: CM DFV VC AF CP PM LS. Contributed reagents/materials/analysis tools: LS AF. Wrote the paper: CM LS. Performed the critical revision of the article for important statistical content: DFV.

References

- Borgiani P, Ciccacci C, Forte V, Romano S, Federici G, et al. (2007) Allelic variants in the CYP2C9 and VKORC1 loci and interindividual variability in the anticoagulant dose effect of warfarin in Italians. *Pharmacogenomics* 8(11):1545–50
- Kim M, Huang SM, Meyer UA, Rahman A, Lesko IJ (2009) A regulatory science perspective on warfarin therapy: a pharmacogenetic opportunity. *J Clin Pharmacol* 49(2):138–46
- Takeuchi F, McGinnis R, Bourgeois S, Barnes C, Eriksson N, et al. (2009) A genome-wide association study confirms VKORC1, CYP2C9, and CYP4F2 as principal genetic determinants of warfarin dose. *PLoS Genet* 5(3):1–9
- Johnson JA, Gong L, Whirl-Carrillo M, Gage BF, Scott SA, et al. (2011) Clinical Pharmacogenetics Implementation Consortium Guidelines for CYP2C9 and VKORC1 genotypes and warfarin dosing. *Clin Pharmacol Ther* 90(4):625–9
- Fareed J, Thethi I, Hoppensteadt D (2012) Old Versus New Oral Anticoagulants: Focus on Pharmacology. *Ann Rev Pharmacol Toxicol* 52:79–99
- Miller CS, Grandi SM, Shimony A, Filion KB, Eisenberg MJ (2012) Meta-Analysis of Efficacy and Safety of New Oral Anticoagulants (Dabigatran, Rivaroxaban, Apixaban) versus Warfarin in Patients with Atrial Fibrillation. *Am J Cardiol* 110:453–460
- Ansell J (2010) Warfarin Versus New Agents: Interpreting the Data. *Hematology Am Soc Hematol Educ Program* 2012:221–8
- Kamali F, Wynne H (2010) Pharmacogenetics of warfarin. *Annu Rev Med* 61:63–75
- Keeling D, Baglin T, Tait C, Watson, Perry D, et al. (2009) Pharmacogenomics of warfarin: uncovering a piece of the warfarin mystery. *Am J Health Syst Pharm* 66(2):123–33
- Gulseth MP, Grice GR, Dager WE (2009) Pharmacogenomics of warfarin: uncovering a piece of the warfarin mystery. *Am J Health Syst Pharm* 66(2):123–33
- Bon Homme M, Reynolds KK, Valdes R Jr, Linder MW (2008) Dynamic pharmacogenetic models in anticoagulation therapy. *Clin Lab Med* 28(4):539–52
- Kurnik D, Loebstein R, Halkin H, Gak E, Almog S (2009) 10 years of oral anticoagulant pharmacogenomics: what difference will it make? A critical appraisal. *Pharmacogenomics* 10(12):1955–65
- Teh LK, Langmia IM, Fazleen Haslinda MH, Ngow HA, Roziyah MJ, et al. (2012) Clinical relevance of VKORC1 (G-1639A and C1173T) and CYP2C9*3 among patients on warfarin. *J Clin Pharm Ther* 37(2):232–6
- Hill CE, Duncan A (2008) Overview of pharmacogenetics in anticoagulation therapy. *Clin Lab Med* 28(4):513–24
- Pirmohamed M, James S, Meakin S, Green C, Scott AK, et al. (2004) Adverse drug reactions as cause of admission to hospital: prospective analysis of 18820 patients. *BMJ* 329(7456):15–9
- Mahajan P, Meyer KS, Wall GC, Price HJ (2011) Clinical applications of pharmacogenomics guided warfarin dosing. *Int J Clin Pharm* 33(1):10–9
- Burmester JK, Berg RL, Glurich I, Yale SH, Schmelzer JR, et al. (2011) Absence of Novel CYP4F2 and VKORC1 Coding Region DNA Variants in Patients Requiring High Warfarin Doses. *Clin Med Res* 9(3–4):119–24
- You JH (2011) Pharmacoeconomic evaluation of warfarin pharmacogenomics. *Expert Opin. Pharmacoter* 12(3):435–441
- Wu AH (2011) Drug metabolizing enzyme activities versus genetic variances for drug of clinical pharmacogenomic relevance. *Clin Proteomics* Jul 28,8(1):12
- Lane S, Al-Zubied S, Hatch E, Matthews I, Jorgensen AL, et al. (2012) The Population Pharmacokinetics of R and S-Warfarin: Effect of Genetic and Clinical Factors. *Br J Clin Pharmacol* 73(1):66–76
- Pavani A, Naushad SM, Rupasree Y, Kumar TR, Malempati AR, et al. (2012) Optimization of warfarin dose by population-specific pharmacogenomic algorithm. *Pharmacogenomics J* 12(4):306–11
- Li T, Chang CY, Jin DY, Lin PJ, Khvorova A, et al. (2004) Identification of the gene for vitamin K epoxide reductase. *Nature* 427(6974):541–4
- Caldwell MD, Awad T, Johnson JA, Gage BF, Falkowski M, et al. (2008) CYP4F2 genetic variant alters required warfarin dose. *Blood* 111(8):4106–12
- Cooper GM, Johnson JA, Langae TY, Feng H, Stanaway IB, et al. (2008) A genome-wide scan for common genetic variants with a large influence on warfarin maintenance dose. *Blood* 112(4):1022–7.
- McDonald MG, Rieder MJ, Nakano M, Hsia CK, Rettie AE (2009) CYP4F2 is a vitamin K1 oxidase: An explanation for altered warfarin dose in carriers of the V433M variant. *Mol Pharmacol* 75(6):1337–46
- Kringen MK, Haug KB, Grimholt RM, Stormo C, Narum S, et al. (2011) Genetic variation of VKORC1 and CYP4F2 genes related to warfarin maintenance dose in patients with myocardial infarction. *J Biomed Biotechnol* 2011:739751
- Gong IY, Tirona RG, Schwarz UI, Crown N, Dresser GK, et al. (2011) Prospective evaluation of a pharmacogenetics-guided warfarin loading and maintenance dose regimen for initiation of therapy. *Blood* 118(11):3163–71
- Liang R, Wang C, Zhao H, Huang J, Hu D, et al. (2012) Influence of CYP4F2 genotype on warfarin dose requirement—a systematic review and meta-analysis. *Thromb Res* Jul,130(1):38–44
- Danese E, Montagnana M, Johnson JA, Rettie AE, Zamboni CF et al. (2012) Impact of the CYP4F2 p.V433M polymorphism on coumarin dose requirement: systematic review and meta-analysis. *Clin Pharmacol Ther* 92(6):746–56

30. Finkelman BS, Gage BF, Johnson JA, Brensinger CM, Kimmel SE (2011) Genetic warfarin dosing: tables versus algorithms. *J Am Coll Cardiol* 57(5): 612–8
31. Available: http://www.accessdata.fda.gov/drugsatfda_docs/label/2010/009218s108lbl.pdf. Accessed 2013 Mar.
32. Suarez-Kurtz G (2011) Population diversity and the performance of warfarin dosing algorithms. *Br J Clin Pharmacol* 72(3):451–3
33. Borgiani P, Ciccacci C, Forte V, Sirianni E, Novelli L, et al. (2009) CYP4F2 genetic variant (rs2108622) significantly contributes to warfarin dosing variability in the Italian population. *Pharmacogenomics* 10(2):261–6
34. Zambon CF, Pengo V, Padrini R, Basso D, Schiavon S, et al. (2011) VKORC1, CYP2C9 and CYP4F2 genetic-based algorithm for warfarin dosing: an Italian retrospective study. *Pharmacogenomics* 12(1):15–25
35. D'Andrea G, D'Ambrosio RL, Di Perna P, Chetta M, Santacroce R, et al. (2005) A polymorphism in the VKORC1 gene is associated with an interindividual variability in the dose-anticoagulant effect of warfarin. *Blood* 105(2):645–9
36. Cini M, Legnani C, Cosmi B, Guazzaloca G, Valdrè L, et al. (2012) A new warfarin dosing algorithm including VKORC1 3730 G > A polymorphism: comparison with results obtained by other published algorithms. *Eur J Clin Pharmacol* 68(8):1167–74
37. Scott SA, Khasawneh R, Peter I, Kornreich R, Desnick RJ (2010) Combined CYP2C9, VKORC1 and CYP4F2 frequencies among racial and ethnic groups. *Pharmacogenomics* 11(6):781–91
38. Yang L, Ge W, Yu F, Zhu H (2010) Impact of VKORC1 gene polymorphism on interindividual and interethnic warfarin dosage requirement—a systematic review and meta analysis. *Thromb Res* 125(4):e159–66
39. Limdi NA, Arnett DK, Goldstein JA, Beasley TM, McGwin G, et al. (2008) Influence of CYP2C9 and VKORC1 on warfarin dose, anticoagulation attainment and maintenance among European American and African Americans. *Pharmacogenomics* 9(5):511–526
40. Gage BF, Lesko LJ (2008) Pharmacogenetics of warfarin: regulatory, scientific, and clinical issues. *J Thromb Thrombolysis* 25(1):45–51
41. Ross KA, Bigham AW, Edwards M, Gozdzik A, Suarez-Kurtz G, et al. (2010) Worldwide allele frequency distribution of four polymorphisms associated with warfarin dose requirements. *J Hum Genet* 55(9):582–9
42. Sanderson S, Emery J, Higgins J (2005) CYP2C9 gene variants, drug dose, and bleeding risk in warfarin-treated patients: a HuGENet systematic review and meta-analysis. *Genet Med* 7(2):97–104
43. Limdi NA, Wadelius M, Cavallari L, Eriksson N, Crawford DC, et al. (2010) Warfarin pharmacogenetics: a single VKORC1 polymorphism is predictive of dose across 3 racial groups. *Blood* 115(18):3827–34
44. Leitner JM, Mannhalter C, Jilma B (2008) Genetic variations and their influence on risk and treatment of venous thrombosis. *Pharmacogenomics* 9(4):423–37
45. Lenzi P, Wadelius M, Kimmel S, Anderson JL, Jorgensen AL, et al. (2010) Integration of genetic, clinical, and INR data to refine warfarin dosing. *Clin Pharmacol Ther* 87(5):572–8
46. International Warfarin Pharmacogenetics Consortium, Klein TE, Altman RB, Eriksson N, Gage BF, et al. (2009) Estimation of the warfarin dose with clinical and pharmacogenetic data. *N Engl J Med* 360(8):753–64
47. Livak KJ (1999) Allelic discrimination using fluorogenic probes and the 5' nuclease assay. *Genet Anal* 14(5–6):143–9
48. Kutuyavin IV, Afonina IA, Mills A, Gorn VV, Lukhtanov EA, et al. (2000) 3'-minor groove binder-DNA probes increase sequence specificity at PCR extension temperatures. *Nucleic Acids Res* 28(2):655–61
49. Barrett JC, Fry B, Maller J, Daly MJ (2005) Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21(2):263–5
50. Gould WW (1995) Jackknife estimation. *Stata Technical Bulletin* 24, sg34: 25–29.