

THE GRONE-MERRIS CONJECTURE

HUA BAI

ABSTRACT. In spectral graph theory, the Grone-Merris Conjecture asserts that the spectrum of the Laplacian matrix of a finite graph is majorized by the conjugate degree sequence of this graph. We give a complete proof for this conjecture.

The Laplacian of a simple graph G with n vertices is a positive semi-definite $n \times n$ matrix $L(G)$ that mimics the geometric Laplacian of a Riemannian manifold; see §1 for definitions, and [2, 14] for comprehensive bibliographies on the graph Laplacian. The spectrum sequence $\lambda(G)$ of $L(G)$ can be listed in non-increasing order as

$$\lambda_1(G) \geq \lambda_2(G) \geq \cdots \geq \lambda_{n-1}(G) \geq \lambda_n(G) = 0.$$

For two non-increasing real sequences \mathbf{x} and \mathbf{y} of length n , we say that \mathbf{x} is *majorized* by \mathbf{y} (denoted $\mathbf{x} \preceq \mathbf{y}$) if

$$\sum_{i=1}^k x_i \leq \sum_{i=1}^k y_i \text{ for all } k \leq n, \text{ and } \sum_{i=1}^n x_i = \sum_{i=1}^n y_i.$$

This notion was introduced because of the following fundamental theorem.

Theorem 1 (Schur-Horn Dominance Theorem [18, 11]). *There exists a Hermitian matrix H with diagonal entry sequence \mathbf{x} and spectrum sequence \mathbf{y} if and only if $\mathbf{x} \preceq \mathbf{y}$.*

In particular, if $\mathbf{d}(G) = (d_1, d_2, \dots, d_n)^T$ is the non-increasing *degree sequence* of G , which coincides with the diagonal entry sequence of the Laplacian matrix $L(G)$, the Schur-Horn Dominance Theorem implies that $\mathbf{d}(G) \preceq \lambda(G)$. Grone [7] improves this majorization result: if G has at least one edge, then $(d_1 + 1, d_2, \dots, d_{n-1}, d_n - 1)^T \preceq \lambda(G)$.

For a non-negative integral sequence \mathbf{d} , we define its *conjugate degree sequence* as the sequence $\mathbf{d}' = (d'_1, d'_2, \dots, d'_n)^T$ where

$$d'_k := \#\{i : d_i \geq k\}.$$

Another important majorization relation is the following.

Theorem 2 (Gale-Ryser [6, 17]). *There exists a $(0, 1)$ -matrix A with row and column sum vectors \mathbf{r} and \mathbf{c} if and only if $\mathbf{r} \preceq \mathbf{c}'$.*

Received by the editors November 12, 2009 and, in revised form, December 11, 2009, January 1, 2010 and May 28, 2010.

2010 *Mathematics Subject Classification*. Primary 15A42; Secondary 05C50.

Key words and phrases. Grone-Merris Conjecture, Laplacian matrix, majorization, split graph, Courant-Fischer-Weyl Min-Max Principle, simplicial complex.

The author was partially supported by NSF grant DMS-0604866.

©2011 American Mathematical Society
Reverts to public domain 28 years from publication

Applying this to the adjacency matrix of G immediately gives that $\mathbf{d}(G) \preceq \mathbf{d}'(G)$.

In 1994, Grone and Merris [8, 9] raised the natural question whether the Laplacian spectrum sequence and the conjugate degree sequence are majorization comparable.

Grone-Merris Conjecture. For any graph G , the Laplacian spectrum is majorized by the conjugate degree sequence

$$\lambda(G) \preceq \mathbf{d}'(G).$$

In this paper, we give a complete proof to the Grone-Merris Conjecture. As a consequence, we have the double majorization $\mathbf{d}(G) \preceq \lambda(G) \preceq \mathbf{d}'(G)$.

See [3] for a partial result in this direction, as well as [19, 12, 13, 1] for proofs in the special cases. Also see [3] for a generalization to simplicial complexes, which is still open.

1. THE LAPLACIAN MATRIX AND THE MAJORIZATION RELATION

Let $G = (V, E)$ be a simple finite graph with $n = |V|$ vertices. We write $i \sim j$ when the i -th vertex is adjacent to the j -th vertex, and we let d_i denote the degree of the i -th vertex.

The *Laplacian matrix* $L(G)$ of the graph G is the $n \times n$ matrix defined by

$$L(G)_{ij} = \begin{cases} d_i & \text{if } i = j; \\ -1 & \text{if } i \sim j; \\ 0 & \text{otherwise.} \end{cases}$$

We can also express the Laplacian as $L(G) = D - A$, where D is the diagonal matrix defined by the degree sequence, and A is the adjacency $(0, 1)$ -matrix of the graph.

It is well known that $L(G)$ is positive semi-definite, since it corresponds to the quadratic form

$$\mathbf{x}^T L(G) \mathbf{x} = \sum_{i \sim j} (x_i - x_j)^2 \text{ for } \mathbf{x} = (x_1, \dots, x_n)^T \in \mathbb{R}^n.$$

Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)^T$ be the non-increasing *spectrum sequence* of the Laplacian matrix $L(G)$. The smallest eigenvalue is $\lambda_n = 0$, with eigenvector $\mathbf{1}_n = (1, 1, \dots, 1)^T$.

Given two vectors $\mathbf{x} = (x_1, \dots, x_n)^T$ and $\mathbf{y} = (y_1, \dots, y_n)^T$ in \mathbb{R}^n , rearrange their components in non-increasing order as

$$x_{[1]} \geq x_{[2]} \geq \dots \geq x_{[n]}, \quad y_{[1]} \geq y_{[2]} \geq \dots \geq y_{[n]}.$$

We say that \mathbf{x} is *majorized* by \mathbf{y} , and write $\mathbf{x} \preceq \mathbf{y}$, if

$$\sum_{i=1}^k x_{[i]} \leq \sum_{i=1}^k y_{[i]} \text{ for all } 1 \leq k \leq n, \text{ and } \sum_{i=1}^n x_i = \sum_{i=1}^n y_i.$$

We will make use of the following majorization inequality.

Theorem 3 (Fan [4]). *If H_1 and H_2 are Hermitian matrices, then*

$$\lambda(H_1 + H_2) \preceq \lambda(H_1) + \lambda(H_2).$$

2. SPLIT GRAPHS

A graph is *split* (also called *semi-bipartite* in [12]) if its vertices can be partitioned into a clique V_1 and a co-clique V_2 . This is equivalent to saying that the subgraph induced by V_1 is complete, and that the subgraph induced by V_2 is an independent set. See [5, 20, 15, 10] for many characterizations and properties of split graphs.

Given a split graph $G = (V, E)$, let $N = |V_1|$ be the size of the clique, and $M = |V_2|$ be the size of the co-clique. Let $\delta(G)$ be the maximum degree of vertices in V_2 . Clearly $\delta(G) \leq N$, and the Laplacian matrix of the split graph G is of the form

$$L(G) = \begin{pmatrix} K_N + D_1 & -A \\ -A^T & D_2 \end{pmatrix},$$

where K_N is the Laplacian matrix of the complete graph on N vertices, where D_1 and D_2 are diagonal matrices with diagonal entries the vertex degrees of V_1, V_2 , respectively, and where A is the adjacency matrix for edges between V_1 and V_2 .

The Laplacian matrix is symmetric, and therefore Hermitian.

Theorem 4 (Courant-Fischer-Weyl [16]). *Let the $n \times n$ matrix H be Hermitian, with eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. Then*

$$\lambda_k = \max_{\dim(S)=k} \min_{0 \neq x \in S} \frac{\langle Hx, x \rangle}{\langle x, x \rangle} = \min_{\dim(S)=n-k+1} \max_{0 \neq x \in S} \frac{\langle Hx, x \rangle}{\langle x, x \rangle},$$

where the max (resp. min) is taken over all k -dimensional (resp. $(n - k + 1)$ -dimensional) subspaces of \mathbb{R}^n .

We first investigate the Laplacian spectrum of a split graph.

Proposition 5. *If G is a split graph of clique size N , then*

$$\lambda_{N-1}(G) \geq N \geq \delta(G) \geq \lambda_{N+1}(G).$$

Moreover, if $\lambda_N(G) \geq N$, then

$$\sum_{i=1}^N d'_i = N^2 + \text{Tr}(D_1).$$

Proof. To prove the inequalities involving $\lambda_{N-1}(G)$ and $\lambda_{N+1}(G)$ by the Courant-Fischer-Weyl Min-Max Principle, it suffices to find an $(N - 1)$ -dimensional (resp. M -dimensional) subspace for which the action of $L(G)$ has a desirable lower (resp. upper) bound. There are natural candidates.

Let $P \subset \mathbb{R}^{M+N}$ be the $(N - 1)$ -dimensional subspace consisting of all vectors of the form $\begin{pmatrix} u \\ \mathbf{0}_M \end{pmatrix}$ with $u \in \mathbb{R}^N$ and $u \perp \mathbf{1}_N$. Then for any unit vector $u \in \mathbb{R}^n$,

$$\left\langle L(G) \begin{pmatrix} u \\ \mathbf{0}_M \end{pmatrix}, \begin{pmatrix} u \\ \mathbf{0}_M \end{pmatrix} \right\rangle = \langle (K_N + D_1)u, u \rangle = N + \langle D_1u, u \rangle \geq N.$$

Similarly, consider the M -dimensional subspace $Q \subset \mathbb{R}^{M+N}$ consisting of all vectors of the form $\begin{pmatrix} \mathbf{0}_N \\ u \end{pmatrix}$ with $u \in \mathbb{R}^M$. Then for any unit vector u ,

$$\left\langle L(G) \begin{pmatrix} \mathbf{0}_N \\ u \end{pmatrix}, \begin{pmatrix} \mathbf{0}_N \\ u \end{pmatrix} \right\rangle = \langle D_2u, u \rangle \leq \delta(G).$$

This proves the part of our first statement that $\lambda_{N-1}(G) \geq N \geq \delta(G) \geq \lambda_{N+1}(G)$.

When $\lambda_N(G) \geq N$, we assert that the degree of any vertex in the clique V_1 is at least N . For this, suppose that our assertion is false, namely that there exists a vertex $v_0 \in V_1$ with degree less than N . Then this vertex v_0 is adjacent to none of the vertices of the co-clique V_2 . Consequently G can be regarded as a split graph with new clique $V_1 \setminus \{v_0\}$ and new co-clique $V_2 \cup \{v_0\}$. The size of the new clique is $\tilde{N} = N - 1$. Applying the first part of the proposition, we obtain that

$$\lambda_N(G) = \lambda_{\tilde{N}+1}(G) \leq \tilde{N} = N - 1,$$

which is a contradiction.

For a conjugating pair of non-negative integral sequences, the partial sum of one sequence can be computed in a different way as

$$\sum_{i=1}^N d'_i = \sum_{i=1}^N \sum_{j=1}^{M+N} \chi(d_j \geq i) = \sum_{j=1}^{M+N} \min(d_j, N),$$

where χ is the characteristic function. The second part of the proposition now follows from the observation that

$$\sum_{j=1}^{M+N} \min(d_j, N) = \sum_{j \in V_1} N + \sum_{j \in V_2} d_j = N^2 + \text{Tr}(D_2) = N^2 + \text{Tr}(D_1). \quad \square$$

The next lemma will play an essential role in our proof of the Grone-Merris Conjecture. Its proof is presented in the next section.

Lemma 6. *Assume that G is a split graph of clique size N . If either $\lambda_N(G) > N$ or $\lambda_N(G) = N > \delta(G)$, then the N -th inequality of the Grone-Merris Conjecture holds, namely*

$$\sum_{i=1}^N \lambda_i \leq \sum_{i=1}^N d'_i.$$

3. THE HOMOTOPY METHOD

This section is devoted to proving Lemma 6. We adopt a homotopy method, following an idea of Katz [12] in his proof of the Grone-Merris Conjecture for 1-regular semi-bipartite graph.

Let $\alpha \in [0, 1]$. Define an $(M + N) \times (M + N)$ matrix L_α as

$$L_\alpha = (1 - \alpha) \begin{pmatrix} K_N + M & -J_{N \times M} \\ -J_{M \times N} & N \end{pmatrix} + \alpha \begin{pmatrix} K_N + D_1 & -A \\ -A^T & D_2 \end{pmatrix},$$

where $J_{M \times N}$ denotes the $M \times N$ matrix whose entries are all equal to 1.

Note that $L_1 = L(G)$ is the matrix we are interested in, and that L_0 is the Laplacian of a complete split graph. The spectrum of L_0 is well understood:

Lemma 7. *The Laplacian spectrum of the complete split graph of clique size N and co-clique size M is*

$$\{ (M + N)^{(N)}, N^{(M-1)}, 0^{(1)} \},$$

where $P^{(Q)}$ denotes Q copies of the number P . The eigenspace corresponding to the eigenvalue N consists of all vectors of the form $\begin{pmatrix} \mathbf{0}^N \\ v \end{pmatrix}$, where v is M -dimensional and $v \perp \mathbf{1}_M$; the eigenspace corresponding to the eigenvalue $(M + N)$ is spanned by the orthogonal vectors

$$(\mathbf{0}_{i-1}, M + N - i, -\mathbf{1}_{M+N-i})^T, \quad 1 \leq i \leq N.$$

Lemma 8. *If $\lambda_N(G) > N$ or $\lambda_N(G) = N > \delta(G)$, then*

$$\lambda_{N+1}^{(\alpha)} \leq N < \lambda_N^{(\alpha)} \text{ for all } 0 \leq \alpha < 1.$$

Proof. We again make use of the Courant-Fischer-Weyl Min-Max Principle. Recall that the M -dimensional subspace $Q \subset \mathbb{R}^{M+N}$ consists of all vectors of the form $\begin{pmatrix} \mathbf{0}^N \\ u \end{pmatrix}$ with $u \in \mathbb{R}^M$. Then for any unit vector u ,

$$\begin{aligned} \left\langle L_\alpha \begin{pmatrix} \mathbf{0}^N \\ u \end{pmatrix}, \begin{pmatrix} \mathbf{0}^N \\ u \end{pmatrix} \right\rangle &= (1 - \alpha)\langle Nu, u \rangle + \alpha\langle D_1(u), u \rangle \\ &\leq (1 - \alpha)N + \alpha\delta(G) \leq N. \end{aligned}$$

Therefore, the $(N + 1)$ -th largest eigenvalue $\lambda_{N+1}^{(\alpha)}$ is at most N .

For the eigenvalue $\lambda_N^{(\alpha)}$, let \tilde{P} be the N -dimensional subspace which is spanned by the eigenvectors of L_1 corresponding to the N largest eigenvalues. Clearly $\tilde{P} \perp \mathbf{1}_{M+N}$. For any unit vector $v \in \tilde{P}$, we know from Lemma 7 that $\langle L_0(v), v \rangle \geq N$. Moreover,

$$\begin{aligned} \langle L_\alpha(v), v \rangle &= \alpha\langle L_1(v), v \rangle + (1 - \alpha)\langle L_0(v), v \rangle \\ &\geq \alpha\lambda_N(G) + (1 - \alpha)N \geq N. \end{aligned}$$

Therefore, the N -th largest eigenvalue $\lambda_N^{(\alpha)}$ is at least N .

We next proceed to show that the inequality on $\lambda_N^{(\alpha)}$ is strict when $0 \leq \alpha < 1$. We already know that $\lambda_N^{(0)} = M + N$. If $\lambda_N^{(\alpha)} = N$ for some $0 < \alpha < 1$, then the above arguments show that necessarily

$$\lambda_N(G) = N, \quad \langle L_1 v, v \rangle = N, \text{ and } L_0(v) = Nv.$$

The first condition $\lambda_N(G) = N$ implies that $\delta(G) < N$, from our assumption on $\lambda_N(G)$; the third condition $L_0(v) = Nv$ implies that v is a unit vector in $\text{Ker}(L_0 - N)$, thus in turn a unit vector of Q . Then

$$\langle L_1 v, v \rangle \leq \delta(G) < N,$$

which contradicts the second condition $\langle L_1 v, v \rangle = N$. □

We now consider all possible N -dimensional subspaces $\begin{pmatrix} I_N \\ V^{(\alpha)} \end{pmatrix} \subseteq (\mathbf{1}_{M+N})^\perp$, where $V^{(\alpha)}$ is an $M \times N$ matrix. Here the notation of the subspace means that the subspace is spanned by the column vectors of the matrix $\begin{pmatrix} I_N \\ V^{(\alpha)} \end{pmatrix}$.

Lemma 9. *If the subspace $\begin{pmatrix} I_N \\ V^{(\alpha)} \end{pmatrix} \subseteq (\mathbf{1}_{M+N})^\perp$ is an invariant subspace of L_α , then the matrix $V^{(\alpha)}$ solves the quadratic matrix equation*

$$V^{(\alpha)} [(1 - \alpha)M + \alpha(N + D_1)] = - (1 - \alpha)J_{M \times N} - \alpha A^T + \alpha [D_2 - V^{(\alpha)}(J_{N \times M} - A)] V^{(\alpha)}.$$

In terms of matrix entries, this means that

$$(1) \quad v_{ji}^{(\alpha)} = \frac{-(1 - \alpha) - \alpha \chi(i \sim j) + \alpha \left(f_j v_{ji} - \sum_{i'=1}^N \sum_{j' \neq i'} v_{ji'}^{(\alpha)} v_{j'i}^{(\alpha)} \right)}{(1 - \alpha)M + \alpha(N + d_i)},$$

where the non-negative integers d_i, f_j are the entries of the diagonal matrices

$$D_1 = \text{Diag}(d_1, d_2, \dots, d_N), \quad D_2 = \text{Diag}(f_1, f_2, \dots, f_M).$$

Proof. It is easy to see that the orthogonal complement in \mathbb{R}^{M+N} of the subspace $\begin{pmatrix} I_N \\ V^{(\alpha)} \end{pmatrix}$ is the subspace $\begin{pmatrix} -V^{(\alpha)T} \\ I_M \end{pmatrix}$. If the subspace $\begin{pmatrix} I_N \\ V^{(\alpha)} \end{pmatrix}$ is an invariant subspace of L_α , then so is its orthogonal complement, since L_α is a symmetric matrix.

The L_α -invariance property is equivalent to the existence of two square matrices X_α and Y_α such that

$$L_\alpha \begin{pmatrix} I_N & -V^{(\alpha)T} \\ V^{(\alpha)} & I_M \end{pmatrix} = \begin{pmatrix} I_N & -V^{(\alpha)T} \\ V^{(\alpha)} & I_M \end{pmatrix} \begin{pmatrix} X_\alpha & 0 \\ 0 & Y_\alpha \end{pmatrix}.$$

By comparison of the corresponding four block matrices, we immediately obtain that

$$X_\alpha = K_N + (1 - \alpha)M + \alpha D_1 - [(1 - \alpha)J_{N \times M} + \alpha A]V^{(\alpha)},$$

$$Y_\alpha = (1 - \alpha)N + \alpha D_2 + [(1 - \alpha)J_{M \times N} + \alpha A^T]V^{(\alpha)T},$$

together with a quadratic matrix equation for $V^{(\alpha)}$:

$$V^{(\alpha)} [K_N + (1 - \alpha)M + \alpha D_1] + (1 - \alpha)J_{M \times N} + \alpha A^T = \left\{ (1 - \alpha)N + \alpha D_2 + V^{(\alpha)} [(1 - \alpha)J_{N \times M} + \alpha A] \right\} V^{(\alpha)}.$$

Because $\begin{pmatrix} I_N \\ V^{(\alpha)} \end{pmatrix} \perp \mathbf{1}_{M+N}$, the entries of $V^{(\alpha)}$ satisfy that

$$\sum_{j=1}^M v_{ji}^{(\alpha)} = -1 \text{ for any } 1 \leq i \leq N.$$

This condition, in terms of matrices, is equivalent to $J_{N \times M}V^{(\alpha)} = -J_{N \times N}$. This implies that $V^{(\alpha)}K_N = [N + V^{(\alpha)}J_{N \times M}]V^{(\alpha)}$, with which the above quadratic matrix equation can be simplified to

$$V^{(\alpha)}[(1 - \alpha)M + \alpha(N + D_1)] = - (1 - \alpha)J_{M \times N} - \alpha A^T + \alpha [D_2 - V^{(\alpha)}(J_{N \times M} - A)] V^{(\alpha)}. \quad \square$$

The quadratic matrix equation is complicated, and is almost impossible to be solved explicitly. Fortunately, we do not have to do so.

From Lemma 8 and the assumption on $\lambda_N(G)$, we know that

$$\lambda_{N+1}^{(\alpha)} \leq \lambda_N^{(\alpha)} \quad \text{for all } \alpha \in [0, 1].$$

Thus the subspace spanned by the eigenvectors of L_α corresponding to the N largest eigenvalues is unique. Assume that this subspace is given by $\begin{pmatrix} I_N \\ V^{(\alpha)} \end{pmatrix}$, so that the matrix $V^{(\alpha)}$ is well defined.

Lemma 10. *The map $V^{(\alpha)} : [0, 1] \rightarrow \mathbb{R}^{M \times N}$ is a continuous function of α , for the usual metric of $\mathbb{R}^{M \times N}$.*

Proof. Assume that α_n is a sequence in $[0, 1]$ such that $\alpha_n \rightarrow \alpha$ as $n \rightarrow \infty$.

According to the algebraic multiplicity of eigenvalues of L_α , there exist positive integers $l = l(\alpha)$ and i_1, \dots, i_l ($i_0 = 0$ by convention) such that $i_1 + i_2 + \dots + i_l = N$ and

$$\lambda_{i_1 + \dots + i_{k-1} + 1}^{(\alpha)} = \dots = \lambda_{i_1 + \dots + i_{k-1} + i_k}^{(\alpha)} > \lambda_{1 + i_1 + \dots + i_{k-1} + i_k}^{(\alpha)}, \quad \forall 1 \leq k \leq l.$$

Let $\{u_i^\beta\}_{i=1}^{M+N}$ be an orthonormal basis consisting of the eigenvectors corresponding to the eigenvalues $\lambda_i^{(\beta)}$ for any $\beta \in [0, 1]$, and $\{Z_k^{\alpha_n}\}_{k=1}^l, \{W_k^\alpha\}_{k=1}^l$ denote two sequences of monotonic subspaces of \mathbb{R}^{M+N} given by

$$Z_k^{\alpha_n} = \text{span}\{u_i^{\alpha_n} : i \leq i_1 + \dots + i_k\}, \quad W_k^\alpha = \text{span}\{u_i^\alpha : i > i_1 + \dots + i_{k-1}\}.$$

By the Courant-Fischer-Weyl Min-Max Principle,

$$\min_{0 \neq u \in Z_k^{\alpha_n}} \frac{\langle L_{\alpha_n}(u), u \rangle}{\langle u, u \rangle} = \lambda_{i_1 + \dots + i_k}^{(\alpha_n)} \rightarrow \lambda_{i_1 + \dots + i_k}^{(\alpha)} \quad \text{as } n \rightarrow \infty$$

and

$$\max_{0 \neq v \in W_{k+1}^\alpha} \frac{\langle L_\alpha(v), v \rangle}{\langle v, v \rangle} = \lambda_{1 + i_1 + \dots + i_k}^{(\alpha)} \leq \lambda_{i_1 + \dots + i_k}^{(\alpha)}.$$

It follows that $Z_k^{\alpha_n} \cap W_{k+1}^\alpha = \{0\}$ and $Z_k^{\alpha_n} \oplus W_{k+1}^\alpha = \mathbb{R}^{M+N}$ when n is sufficiently large. Moreover, we obtain that $Z_l^{\alpha_n} = \bigoplus_{k=1}^l (Z_k^{\alpha_n} \cap W_k^\alpha)$ from

$$\dim(Z_k^{\alpha_n} \cap W_k^\alpha) = \dim(Z_k^{\alpha_n}) + \dim(W_k^\alpha) - (M + N) = i_k.$$

Consider a basis of the subspace $Z_k^{\alpha_n} \cap W_k^\alpha$ which consists of unit vectors of the form

$$u_{k,n,s} = \cos(\theta_{k,n,s})u_{i_1 + \dots + i_{k-1} + s}^\alpha + \sin(\theta_{k,n,s})w_{k,s}, \quad 1 \leq s \leq i_k,$$

for some unit vector $w_{k,s} \in W_{k+1}^\alpha$. Necessarily $\lim_{n \rightarrow \infty} \sin(\theta_{k,n,s}) = 0$, since

$$\langle L_{\alpha_n}(u_{k,n,s}), u_{k,n,s} \rangle \geq \lambda_{i_1 + \dots + i_k}^{(\alpha_n)} \quad \text{and}$$

$$\begin{aligned} \langle L_\alpha(u_{k,n,s}), u_{k,n,s} \rangle &= \cos^2(\theta_{k,n,s})\lambda_{i_1 + \dots + i_k}^\alpha + \sin^2(\theta_{k,n,s})\langle L_\alpha(w_{k,s}), w_{k,s} \rangle \\ &\leq \cos^2(\theta_{k,n,s})\lambda_{i_1 + \dots + i_k}^{(\alpha)} + \sin^2(\theta_{k,n,s})\lambda_{i_1 + \dots + i_{k+1}}^{(\alpha)}. \end{aligned}$$

Any vector $u \in Z_l^{\alpha_n}$ can now be expressed as

$$u = \sum_{k=1}^l \sum_{s=1}^{i_k} c_{k,s} \left[\cos(\theta_{k,n,s})u_{i_1 + \dots + i_{k-1} + s}^\alpha + \sin(\theta_{k,n,s})w_{k,s} \right].$$

Assume that the maximum of $|c_{k,s}|$ is achieved at $|c_{k_0,s_0}|$. Due to the orthogonality of $\{u_i^\alpha\}_i$, the absolute value of the coefficient of $u_{i_1+\dots+i_{k_0-1}+s_0}^\alpha$ is at most $\|u\|$. But when n is sufficiently large, it is at least

$$|c_{k_0,s_0}| \cdot \left(\left| \cos(\theta_{k_0,n,s_0}) \right| - \sum_{k=1}^l \sum_{s=1}^{i_k} |\sin(\theta_{k,n,s})| \right) \geq \frac{|c_{k_0,s_0}|}{2}.$$

Hence $|c_{k_0,s_0}| \leq 2\|u\|$. For any given vector $v \in W_{l+1}^\alpha$, we see that

$$|\langle u, v \rangle| = \left| \sum_{k=1}^l \sum_{s=1}^{i_k} \langle c_{k,s} \sin(\theta_{k,n,s}) w_{k,s}, v \rangle \right| \leq 2\|u\| \cdot \|v\| \cdot \sum_{k=1}^l \sum_{s=1}^{i_k} |\sin(\theta_{k,n,s})|,$$

which goes to zero as n goes to infinity.

The subspace $Z_l^{\alpha_n}$ is nothing else but $\begin{pmatrix} I_N \\ V^{(\alpha_n)} \end{pmatrix}$, while W_{l+1}^α is nothing else but $\begin{pmatrix} -V^{(\alpha)T} \\ I_M \end{pmatrix}$. The inner product of the i -th column vector of the first matrix and the j -th column vector of the second matrix is equal to

$$V_{ji}^{(\alpha_n)} - V_{ji}^{(\alpha)},$$

which must go to zero as n goes to infinity. This proves the continuity of $V^{(\alpha)}$ on α . □

Lemma 11. *Let Ω be the subset*

$$\{(x_{ji}) : \sum_{k=1}^M x_{ki} = -1, \forall 1 \leq i \leq N, \text{ and } x_{ji} \leq 0, \forall 1 \leq j \leq M, 1 \leq i \leq N\}$$

of $\mathbb{R}^{M \times N}$. Then $V^{(\alpha)} \in \Omega$ for all $\alpha \in [0, 1]$.

Proof. Consider the subset

$$\Gamma = \{\alpha \in [0, 1) : V^{(\alpha)} \in \Omega\}$$

of the half-open half-closed interval $[0, 1)$.

When $\alpha = 0$, $v_{ji}^{(0)} \equiv -\frac{1}{M}$ (see Lemma 7 or equation (1)). As a consequence, $V^{(0)} \in \Omega$, so that $0 \in \Gamma$ and Γ is not empty.

Suppose there is a sequence of points $\alpha_n \in \Gamma$ and $\lim_{n \rightarrow \infty} \alpha_n = \alpha$ with α still in $[0, 1)$. By Lemma 10, $\lim_{n \rightarrow \infty} V^{(\alpha_n)} = V^{(\alpha)}$. Because Ω is a compact set, so $V^{(\alpha)} \in \Omega$ and $\alpha \in \Gamma$. Therefore, Γ is a closed subset of $[0, 1)$.

Suppose $\alpha \in \Gamma$, namely $V^{(\alpha)} \in \Omega$ for some $\alpha \in [0, 1)$. Because the quantities $\chi(i \sim j)$, f_j and $v_{j'i'}^{(\alpha)} v_{j'i}^{(\alpha)}$ in equation (1) are all non-negative, we see that

$$v_{ji}^{(\alpha)} \leq \frac{-(1-\alpha)}{(1-\alpha)M + \alpha(N+d_i)} < 0 \text{ for all } 1 \leq j \leq M, 1 \leq i \leq N.$$

Therefore $V^{(\alpha)}$ is contained in the interior of Ω . Since $V^{(\alpha)}$ depends continuously on α , it follows that Γ is an open subset of $[0, 1)$.

The interval $[0, 1)$ is connected, and Γ is an open closed non-empty subset of it, therefore Γ is equal to $[0, 1)$.

By continuity at $\alpha = 1$, $V^{(1)}$ is also in Ω . This proves that $V^{(\alpha)} \in \Omega$ for all $\alpha \in [0, 1]$. □

During the proof of Lemma 9, we have already known that

$$L_\alpha \left(\begin{matrix} I_N \\ V^{(\alpha)} \end{matrix} \right) = \left(\begin{matrix} I_N \\ V^{(\alpha)} \end{matrix} \right) X_\alpha$$

where

$$X_\alpha = K_N + (1 - \alpha)M + \alpha D_1 - [(1 - \alpha)J_{N \times M} + \alpha A]V^{(\alpha)}.$$

So the sum of the N largest eigenvalues of L_1 is equal to the trace of

$$X_1 = K_N + D_1 - AV^{(1)}.$$

But $V^{(1)} \in \Omega$ by Lemma 11, therefore

$$\text{Tr}(AV^{(1)}) = \sum_{i=1}^N \sum_{j:j \sim i} v_{ji} \geq \sum_{i=1}^N \sum_{j=1}^M v_{ji} = -N.$$

Then

$$\sum_{i=1}^N \lambda_i = N(N - 1) + \text{Tr}(D_1) - \text{Tr}(AV^{(1)}) \leq N^2 + \text{Tr}(D_1).$$

By Proposition 5, this completes the proof of Lemma 6.

4. PROOF OF THE GRONE-MERRIS CONJECTURE

For consistence we restate the Grone-Merris Conjecture here.

Grone-Merris Conjecture. For any graph G , its Laplacian spectrum is majorized by its conjugate degree sequence, namely $\lambda(G) \preceq \mathbf{d}'(G)$.

The Grone-Merris Conjecture behaves nicely under complementation, in the sense of the proposition below.

The *complement graph* of a graph G is a graph \overline{G} on the same vertices such that two vertices of \overline{G} are adjacent if and only if they are not adjacent in G . The Laplacian matrices of the graph G and of its complementary graph \overline{G} are related by the property that

$$L(G) + L(\overline{G}) + J_n = nI_n.$$

All these matrices commute with each other, so that

$$\begin{aligned} \lambda(\overline{G}) &= (n - \lambda_{n-1}(G), \dots, n - \lambda_1(G), 0); \\ \mathbf{d}'(\overline{G}) &= (n - d'_{n-1}(G), \dots, n - d'_1(G), 0). \end{aligned}$$

From these we see that

Proposition 12. For any $1 \leq k < n$, the k -th inequality holds for the graph G if and only if the $(n - k - 1)$ -th inequality holds for the complement graph \overline{G} ,

$$\sum_{i=1}^k \lambda_i(G) \leq \sum_{i=1}^k d'_i(G) \iff \sum_{j=1}^{n-1-k} \lambda_j(\overline{G}) \leq \sum_{j=1}^{n-1-k} d'_j(\overline{G}), \quad \forall 1 \leq k < n.$$

We are now ready to prove the Grone-Merris Conjecture.

Assume that the Grone-Merris Conjecture is not true, and the graph $G = (V, E)$ is a counterexample. Namely, there exists an integer k with $1 < k < n = |V|$, such that

$$\sum_{i=1}^k \lambda_i > \sum_{i=1}^k d'_i.$$

Without loss of generality, we can assume that this integer k is minimum over all counterexamples. Then we have

$$\sum_{i=1}^{k-1} \lambda_i \leq \sum_{i=1}^{k-1} d'_i, \quad \text{and} \quad \lambda_k > d'_k.$$

Moreover, we can further assume that the number $|E|$ of edges is minimum over all counterexamples with the same k . Under this assumption, we claim that

Lemma 13. *For any two vertices i, j in the graph G , if $d_i \leq k$ and $d_j \leq k$, then they are not adjacent in G .*

Proof. We will prove this by contradiction by assuming that the lemma is false. Namely there exists a pair of vertices such that

$$d_i \leq k, \quad d_j \leq k, \quad i \sim j.$$

Let \tilde{G} be the graph obtained from G by deleting the edge ij . Due to the minimum property of $|E|$, we must have

$$\sum_{i=1}^k \lambda_i(\tilde{G}) \leq \sum_{i=1}^k d'_i(\tilde{G}).$$

Two Laplacian matrices are related via $L(G) = L(\tilde{G}) + H$, where $H_{n \times n}$ is a positive semi-definite matrix whose only non-zero entries are $H_{ii} = H_{jj} = 1$ and $H_{ij} = H_{ji} = -1$. Applying Fan's Theorem 3, we see that

$$\begin{aligned} \sum_{i=1}^k \lambda_i(G) &\leq \sum_{i=1}^k \lambda_i(\tilde{G}) + \sum_{i=1}^k \lambda_i(H) \leq \sum_{i=1}^k d'_i(\tilde{G}) + \text{Tr}(H) \\ &= \left[\sum_{i=1}^k d'_i(G) - 2 \right] + 2 = \sum_{i=1}^k d'_i(G). \end{aligned}$$

This contradicts our assumption that G was a counterexample, and therefore concludes the proof. \square

Next, we add new edges to G to get a new graph \hat{G} . Add to G a new edge ij for any pair of vertices i, j in G such that

$$d_i \geq k, \quad d_j \geq k, \quad \text{and} \quad i \not\sim j.$$

The new graph \hat{G} so obtained is a split graph.

The clique of \hat{G} consists of all vertices of G whose degree is at least k , so the size of the clique is equal to $d'_k(G)$. Let $N = d'_k(G)$ denote this size. The co-clique consists of all vertices of G whose degree is less than k , so the maximum degree of vertices in the co-clique is $\delta(\hat{G}) \leq k - 1$.

Note that

$$d'_1(\hat{G}) = d'_1(G), \dots, d'_k(\hat{G}) = d'_k(G)$$

while $\lambda_i(\hat{G}) \geq \lambda_i(G)$ for all $1 \leq i \leq n$, so these two inequalities are still valid for the new graph \hat{G} , namely

$$\sum_{i=1}^k \lambda_i(\hat{G}) > \sum_{i=1}^k d'_i(\hat{G}) \quad \text{and} \quad \lambda_k(\hat{G}) > d'_k(\hat{G}) = N.$$

Let us discuss the relationship between N and k .

If $N < k$, then $\lambda_k(\widehat{G}) \leq \lambda_{N+1}(\widehat{G}) \leq N$, which leads to a contradiction. The second inequality comes from Proposition 5.

If $N = k$, then \widehat{G} is a split graph of clique size N , with the property that

$$\sum_{i=1}^N \lambda_i(\widehat{G}) > \sum_{i=1}^N d'_i(\widehat{G}) \quad \text{and} \quad \lambda_N(\widehat{G}) > N.$$

This contradicts Lemma 6.

So $k < N$. Note that \widehat{G} is a split graph of clique size N . In this graph \widehat{G} , the maximum degree of vertices in the co-clique is at most $(k - 1)$, while the minimum degree of vertices in the clique is at least $(N - 1)$. This means that

$$d'_{N-1}(\widehat{G}) = \dots = d'_{k+1}(\widehat{G}) = d'_k(\widehat{G}) = N.$$

Combining this with $\lambda_{k+1}(\widehat{G}) \geq \dots \geq \lambda_{N-1}(\widehat{G}) \geq N$ from Proposition 5, we see immediately that the inequality

$$\sum_{i=1}^k \lambda_i(\widehat{G}) > \sum_{i=1}^k d'_i(\widehat{G}) \quad \text{can be extended to} \quad \sum_{i=1}^{N-1} \lambda_i(\widehat{G}) > \sum_{i=1}^{N-1} d'_i(\widehat{G}).$$

Then we proceed to compare $\lambda_N(\widehat{G})$ with the clique size N .

First consider the case where $\lambda_N(\widehat{G}) \geq N$. Because $N = d'_{N-1}(\widehat{G}) \geq d'_N(\widehat{G})$, the split graph \widehat{G} has clique size N , with the additional property that

$$\sum_{i=1}^N \lambda_i(\widehat{G}) > \sum_{i=1}^N d'_i(\widehat{G}) \quad \text{and} \quad \lambda_N(\widehat{G}) \geq N > \delta(\widehat{G}).$$

This again contradicts Lemma 6.

In the other case, where $\lambda_N(\widehat{G}) < N$, we switch attention to the complement graph of \widehat{G} . This complement graph is another split graph $\overline{\widehat{G}}$. Its clique size is M , and

$$\lambda_M(\overline{\widehat{G}}) = (N + M) - \lambda_N(\widehat{G}) > M.$$

According to Proposition 12,

$$\sum_{i=1}^{N-1} \lambda_i(\widehat{G}) > \sum_{i=1}^{N-1} d'_i(\widehat{G}) \quad \implies \quad \sum_{i=1}^M \lambda_i(\overline{\widehat{G}}) > \sum_{i=1}^M d'_i(\overline{\widehat{G}}).$$

Therefore, $\overline{\widehat{G}}$ is a split graph of clique size M , with the additional property that

$$\sum_{i=1}^M \lambda_i(\overline{\widehat{G}}) > \sum_{i=1}^M d'_i(\overline{\widehat{G}}) \quad \text{and} \quad \lambda_M(\overline{\widehat{G}}) > M.$$

This again contradicts Lemma 6.

All possible cases are eliminated, and the Grone-Merris Conjecture is proved.

ACKNOWLEDGEMENTS

This work was started while the author was visiting the University of Southern California, whose support and hospitality is gratefully acknowledged. The author also thanks Francis Bonahon for his support and encouragement throughout the years, Jun Ying and Jie Ying for critical Matlab computations, Russell Merris for useful suggestions, and Andries Brouwer, Tao Li and the referee for many valuable comments.

REFERENCES

- [1] Ravindra B. Bapat, Arbind K. Lal and Sukanta Pati, *Laplacian spectrum of weakly quasi-threshold graphs*, *Graphs Comb.* **24**, no. 4 (2008), 273–290. MR2438859 (2009g:05099)
- [2] Fan R. K. Chung, *Spectral graph theory*, volume **92** of CBMS Regional Conference Series in Mathematics, American Mathematical Society, 1997. MR1421568 (97k:58183)
- [3] Art M. Duval and Victor Reiner, *Shifted simplicial complexes are Laplacian integral*, *Trans. Amer. Math. Soc.* **354** (2002), 4313–4344. MR1926878 (2003j:15017)
- [4] Ky Fan, *On a theorem of Weyl concerning eigenvalues of linear transformations I*, *Proc. Nat. Acad. Sci. USA* **35** (1949), 652–655. MR0034519 (11:600e)
- [5] Stéphane Földes and Peter L. Hammer, ‘*Split graphs*’, *Proceedings of the Eighth Southeastern Conference on Combinatorics, Graph Theory and Computing, 1977, Congressus Numerantium, XIX, Winnipeg: Utilitas Math.*, 311–315. MR0505860 (58:21844)
- [6] David Gale, *A theorem on flows in networks*, *Pacific J. Math.* **7** (1957), 1073–1082. MR0091855 (19:1024a)
- [7] Robert Grone, *Eigenvalues and degree sequences of graphs*, *Lin. Multilin. Alg.* **39** (1995) 133–136. MR1374475 (97a:05148)
- [8] Robert Grone and Russell Merris, *Coalescence, majorization, edge valuations and the Laplacian spectra of graphs*, *Lin. Multilin. Alg.* **27**, No.2 (1990), 139–146. MR1054137 (91c:05129)
- [9] Robert Grone and Russell Merris, *The Laplacian spectrum of a graph II*, *SIAM J. Disc. Math.* **7** (1994), 221–229. MR1271994 (95d:05085)
- [10] Peter L. Hammer and Bruno Simeone, *The spittance of a graph*, *Combinatorica* **1** (1981), 275–284. MR637832 (84c:05050)
- [11] Roger Alfred Horn, *Doubly stochastic matrices and the diagonal of a rotation matrix*, *Amer. J. Math.* **76** (1954), 620–630. MR0063336 (16:105c)
- [12] Nets Hawk Katz, *The Grone Merris conjecture and a quadratic eigenvalue problem*, preprint, 2005; arXiv:math.CA/0512647.
- [13] Steve Kirkland, *Near threshold graphs*, *Electron. J. Combin.* **16**, no. 1(2009), Research Paper R42. MR2491644 (2010g:05221)
- [14] Russell Merris, *Laplacian matrices of graphs: a survey*, *Linear Algebra and its Applications* **197 & 198** (1994), 143–176. MR1275613 (95e:05084)
- [15] Russell Merris, *Split graphs*, *European J. Combin.* **24**, **4** (2003), 413–430. MR1975945 (2004b:05064)
- [16] Michael Reed and Barry Simon, *Methods of Modern Mathematical Physics IV: Analysis of Operators*, Academic Press, 1978. MR0493421 (58:12429c)
- [17] Herbert J. Ryser, *Combinatorial properties of matrices of zeros and ones*, *Pacific J. Math.* **7** (1957), 1073–1082. MR0087622 (19:379d)
- [18] Issac Schur, *Über eine Klasse von Mittelbildungen mit Anwendungen die Determinanten*, *Sitzungsber. Berlin. Math. Gesellschaft* **22** (1923), 9–20.
- [19] Tamon Stephen, *On the Grone-Merris conjecture*, *DMTCS proc. AE* (2005), 187–192.
- [20] Regina I. Tyshkevich and Arkady A. Chernyak, *Canonical partition of a graph defined by the degrees of its vertices*, (in Russian), *Isv. Akad. Nauk BSSR, Ser. Fiz.-Mat. Nauk* **5** 1979, 14–26. MR554162 (81f:05098)

DEPARTMENT OF MATHEMATICS, BOSTON COLLEGE, CHESTNUT HILL, MASSACHUSETTS 02467
E-mail address: baihu@bc.edu
E-mail address: huabai@alumni.usc.edu